

# Lamellipodium is a myosin-independent mechanosensor

Patrick W. Oakes<sup>a,b,c,d,e,1</sup>, Tamara C. Bidone<sup>c,d,f</sup>, Yvonne Beckham<sup>c,d,e</sup>, Austin V. Skeeters<sup>a</sup>, Guillermina R. Ramirez-San Juan<sup>c,d,e</sup>, Stephen P. Winter<sup>g</sup>, Gregory A. Voth<sup>c,d,f</sup>, and Margaret L. Gardel<sup>c,d,e,1</sup>

<sup>a</sup>Department of Physics and Astronomy, University of Rochester, Rochester, NY 14627; <sup>b</sup>Department of Biology, University of Rochester, Rochester, NY 14627; <sup>c</sup>Institute for Biophysical Dynamics, University of Chicago, Chicago, IL 60637; <sup>d</sup>James Franck Institute, University of Chicago, Chicago, IL 60637; <sup>e</sup>Department of Physics, University of Chicago, Chicago, IL 60637; <sup>f</sup>Department of Chemistry, University of Chicago, Chicago, IL 60637; and <sup>g</sup>Interdisciplinary Scientist Training Program, University of Chicago, Chicago, IL 60637

Edited by David A. Weitz, Harvard University, Cambridge, MA, and approved February 5, 2018 (received for review September 8, 2017)

The ability of adherent cells to sense changes in the mechanical properties of their extracellular environments is critical to numerous aspects of their physiology. It has been well documented that cell attachment and spreading are sensitive to substrate stiffness. Here, we demonstrate that this behavior is actually biphasic, with a transition that occurs around a Young's modulus of ~7 kPa. Furthermore, we demonstrate that, contrary to established assumptions, this property is independent of myosin II activity. Rather, we find that cell spreading on soft substrates is inhibited due to reduced myosin-II independent nascent adhesion formation within the lamellipodium. Cells on soft substrates display normal leading-edge protrusion activity, but these protrusions are not stabilized due to impaired adhesion assembly. Enhancing integrin-ECM affinity through addition of Mn<sup>2+</sup> recovers nascent adhesion assembly and cell spreading on soft substrates. Using a computational model to simulate nascent adhesion assembly, we find that biophysical properties of the integrin-ECM bond are optimized to stabilize interactions above a threshold matrix stiffness that is consistent with the experimental observations. Together, these results suggest that myosin II-independent forces in the lamellipodium are responsible for mechanosensation by regulating new adhesion assembly, which, in turn, directly controls cell spreading. This myosin II-independent mechanism of substrate stiffness sensing could potentially regulate a number of other stiffness-sensitive processes.

nascent adhesion | myosin-II | mechanosensing | integrin | catch-bond

The ability of cells to sense mechanical forces and convert them into biochemical responses regulates a plethora of physiological functions (1–3). In particular, cells respond to changes in the stiffness of the extracellular matrix (ECM) by altering a number of adhesion-dependent behaviors, including spreading (4–12), migration (4, 13, 14), proliferation (15), differentiation (16, 17), and metastasis (18, 19). Matrix mechanosensing is thought to be mediated by focal adhesions, hierarchical organelles comprising ~150 proteins that facilitate dynamic and force-sensitive interactions between the ECM and the actin cytoskeleton (20–22). How these dynamic organelles mediate environmental sensing in a variety of physiological contexts, however, is still largely unknown.

Previous efforts have focused primarily on myosin II-mediated mechanisms for substrate stiffness sensing (23–28). Stresses generated by myosin motors on the actin cytoskeleton are transmitted to the ECM via focal adhesions. These stresses, coupled with the matrix rigidity, impact the deformation and binding affinity of proteins within the focal adhesion (29–32). Changes in the composition and kinetics of proteins within focal adhesions are thought to variably regulate force transmission from the actin cytoskeleton and the matrix (33–35), leading many to describe focal adhesions as molecular clutches. Initial adhesion formation, however, occurs in the leading edge of the lamellipodium and is a myosin-independent process (36, 37). These structures, known as nascent adhesions, are instead subject to forces that primarily originate from polymerization of actin filaments. The contribution of nascent adhesions to mechanisms of substrate stiffness sensing has not been thoroughly explored.

One of the best-characterized metrics of environmental sensing by adherent cells is their ability to attach and spread on ligand-coated

substrates. The extent of cell spreading is controlled by the density and spatial organization of matrix ligands (38–40), as well as the rigidity of the substrate to which these ligands are attached (4–12). It has also been suggested that the stress-relaxing properties of the matrix can contribute to cell spreading (41, 42). In the limit of soft substrates with a Young's modulus <500 Pa, cell spreading is inhibited. As the substrate stiffness increases, the spread area increases and ultimately plateaus (8–12). While previous reports have differed on the exact range of relevant stiffness which regulates this behavior, likely due to variances in experimental approaches (43), cell spreading remains a robust metric to study substrate stiffness sensing.

Here, we study the mechanism regulating substrate stiffness-dependent cell spreading. We found that NIH 3T3 cell spreading is acutely impacted as the Young's modulus of the substrate increases from 5 to 8 kPa. On substrates with a stiffness <5 kPa, cells spread poorly. Average cell spread area increased on substrates stiffer than 5 kPa, plateauing on substrates stiffer than 8 kPa. Above this threshold, cell spread area remained constant. Surprisingly, we found this stiffness-dependent change in cell spreading was independent of myosin II motor activity. Instead, we found that spreading on soft substrates is impaired by reduced assembly of nascent, myosin-independent adhesions at the cell periphery. Enhancing integrin-ligand affinity through the addition of Mn<sup>2+</sup> was sufficient both to stabilize nascent adhesions and increase cell spread area on soft substrates. We then implemented a computational model to determine how changes in integrin-substrate catch-bond kinetics affected integrin binding on substrates of different stiffness. We found that the biophysical properties of integrin-matrix catch-bonds were optimized to sense changes in substrate stiffness at ~6 kPa, consistent with our experimental results. Together, these results illustrate that nascent adhesion formation in the lamellipodium functions as a myosin

## Significance

Cell physiology can be regulated by the mechanics of the extracellular environment. Here, we demonstrate that cell spreading is a mechanosensitive process regulated by weak forces generated at the cell periphery and independent of motor activity. We show that stiffness sensing depends on the kinetics of the initial adhesion bonds that are subjected to forces driven by protein polymerization. This work demonstrates how the binding kinetics of adhesion molecules are sensitively tuned to a range of forces that enable mechanosensation.

Author contributions: P.W.O. and M.L.G. designed research; P.W.O., Y.B., A.V.S., G.R.R.-S.J., and S.P.W. performed research; P.W.O., T.C.B., and G.A.V. contributed new reagents/analytic tools; P.W.O., T.C.B., and A.V.S. analyzed data; and P.W.O., T.C.B., and M.L.G. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Published under the PNAS license.

<sup>1</sup>To whom correspondence may be addressed. Email: poakes@rochester.edu or gardel@uchicago.edu.

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1715869115/-DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1715869115/-DCSupplemental).


Published online February 27, 2018.

## II-independent mechanosensor to control cell adhesion and spreading.

### Results

**Spread Area Is a Biphasic Response of Substrate Stiffness, Independent of Myosin Activity.** To investigate the mechanisms that drive substrate stiffness sensing, we chose to measure the spread area of adherent cells. We first plated NIH 3T3 fibroblasts on a series of polyacrylamide gels covalently coupled with fibronectin and with Young's moduli ranging from 0.6 to 150 kPa (Fig. 1A). Cells were also plated on glass absorbed with fibronectin as a control. Consistent with previous reports (4, 6, 8–11), we found that cells spread area was sensitive to substrate stiffness (Fig. 1A). In contrast, however, we found that this response could be broken down into two regimes: There was poor spreading on soft (less than ~5 kPa) substrates and high spreading on stiff (more than ~8 kPa) substrates, with a transition region between these values and no statistical difference in spread area between populations within each regime (Fig. 1A). Furthermore, the morphology of cells on soft and stiff substrates was noticeably different (5). Cells on soft substrates were more rounded with a disorganized actin cytoskeleton (Fig. 1B). In contrast, cells on stiff substrates exhibited more polarized shapes and tended to have prominent stress fibers (Fig. 1B).

Because myosin II activity has been widely implicated in mechanosensing (28), we next hypothesized that its inhibition would eliminate any change in spread area as a function of substrate stiffness. Surprisingly, cells incubated with 50  $\mu$ M blebbistatin, a myosin II ATPase inhibitor, continued to exhibit a biphasic response to substrate stiffness (Fig. 1C). Cells treated



**Fig. 1.** Spread area is a biphasic response of substrate stiffness, independent of myosin activity. (A) Boxplots of the spread area of NIH 3T3 fibroblasts plated on fibronectin-coated polyacrylamide gels of varying stiffness. Cells can be grouped into soft ( $\leq 4.5$  kPa) and stiff ( $\geq 8.4$  kPa) regimes. From left to right,  $n = 182, 674, 205, 155, 254, 400, 205, 487$ , and 170. (B) Representative images of control cells on soft and stiff substrates. (C) Boxplots of the spread area of cells treated with 50  $\mu$ M blebbistatin to inhibit myosin activity. While blebbistatin-treated cells spread more than control cells, they exhibited the same biphasic response as a function of substrate stiffness. From left to right,  $n = 169, 228, 329, 67, 159, 125, 119, 183$ , and 56. (D) Representative images of blebbistatin-treated cells on soft and stiff substrates. (E) Boxplots of the spread area of cells treated with 20  $\mu$ M Y-27632, which inhibits ROCK activity. Cells treated with Y-27632 still exhibited a difference in spread area on soft ( $n = 148$ ) and stiff ( $n = 203$ ) substrates. (F) Representative images of Y-27632-treated cells on soft and stiff substrates. Boxplots represent 25th, 50th, and 75th percentiles, while whiskers extend to the 10th and 90th percentiles.  $*P < 0.01$ .

with blebbistatin had an increased spread area compared with control cells across all stiffnesses, but exhibited the same soft and stiff regimes. Morphologically, myosin-inhibited cells on all substrates showed more protrusions, but on stiff substrates, the cells exhibited more spindle-like projections (Fig. 1D). Similar phenotypes were seen when cells were incubated with Rho-Kinase inhibitor (Y-27632; Fig. 1E and F) and when cells were plated on other ECM proteins (Fig. S1). Thus, the change in cell spread area that occurred between the soft and stiff regimes did not require myosin II activity.

### Substrate Stiffness Does Not Inhibit Lamellipodia Protrusion Dynamics.

To understand how substrate stiffness impacts cell spread area, we investigated the effects of substrate stiffness on protrusion dynamics. We tracked lamellipodia formation by taking time-lapse images of cells transiently transfected with a fluorescent membrane marker (GFP-stargazin) and treated with 20  $\mu$ M Y-27632 on representative soft (2.1 kPa) and stiff (48 kPa) substrates 30 min after plating (Fig. 2A and B and Movies S1 and S2). Cells on soft substrates exhibited repeated cycles of protrusion and retraction, as seen in the kymograph (Fig. 2A), reducing their ability to spread. Cells on stiff substrates, however, exhibited continuous and steady protrusions that resulted in leading-edge advance (Fig. 2B). Using cell contours derived from the fluorescence images, we identified protrusive regions and measured their morphology and characteristics (Fig. 2C). We found no statistically significant difference between soft and stiff substrates for measurements of the average protrusion area (Fig. 2D) or the average protrusion width (Fig. 2E). These data indicate that substrate stiffness affects the stability of leading-edge protrusions, but not the protrusion dynamics themselves. Arp2/3-mediated lamellipodium formation is still required for spreading, as cells on both soft and stiff substrates that were treated with CK-869, an Arp2/3 inhibitor, were indistinguishable from control cells on soft substrates (Fig. 2F and G). Together, these results illustrate that it is the stabilization, not the formation, of Arp2/3-dependent lamellipodial protrusions that is hindered on soft substrates.

**Soft Substrates Impair Nascent Adhesion Formation.** To explore the mechanism of substrate stiffness-dependent changes in stabilization of myosin II-independent protrusions, we examined the assembly of myosin II-independent, nascent adhesions that form at the base of the lamellipodium. Two hours after plating, cells were treated with 20  $\mu$ M Y-27632 for 30 min and then fixed and stained for actin, p34 (a subunit of Arp 2/3), and the focal adhesion protein paxillin (Fig. 3A and B). On both soft and stiff substrates, p34 localized to the cell periphery, indicative of the Arp2/3-dependent lamellipodium (Fig. 3A and B). On stiff substrates, paxillin formed small punctate nascent adhesions near the leading edge, which is characteristic of nascent adhesion formation on glass substrates (44). By contrast, on soft substrates, paxillin-rich nascent adhesions were seen at a lower density and formed further away from the leading edge. To quantify these differences in protein localization, we measured the average actin, p34, and paxillin intensity in  $\sim 0.5$ - $\mu\text{m}$  bands measured radially from the edge of the cell (Fig. 3C). We found that the peak of p34 intensity was localized right at the edge of the cell on all substrates. On stiff substrates, paxillin was located within  $\sim 0.5$   $\mu\text{m}$  of the p34 peak (Fig. 3D). On soft substrates, there was a significantly reduced accumulation of paxillin, and its peak was found  $\sim 5$   $\mu\text{m}$  behind the leading edge (Fig. 3D). These data suggest that cells have reduced nascent adhesion formation on soft substrates.

### Activation of Integrins via $\text{Mn}^{2+}$ Is Sufficient to Promote Spreading on Soft Substrates.


Given the reduced density of nascent adhesions on soft substrates, we sought to explore the extent to which changes in integrin–ligand affinity could stimulate their formation. The presence of 3  $\mu\text{M}$   $\text{Mn}^{2+}$  increased the lifetime of integrin–fibronectin bonds (45), but did not affect the contractility of the cell (Fig. S2). When cells were plated on soft substrates in the presence of 3  $\mu\text{M}$   $\text{Mn}^{2+}$ , they exhibited a greater






than twofold increase in spread area on soft substrates, similar to their spread area on stiff substrates either in the presence or absence of  $Mn^{2+}$  (Fig. 4A). To directly compare the effect of  $Mn^{2+}$  on adhesion assembly on soft substrates, we performed immunofluorescence of paxillin and actin. Addition of  $Mn^{2+}$  to cells on soft substrates stimulated the formation of paxillin-rich adhesions near the cell periphery and even the formation of lamellar actin bundles (Fig. 4B).

To determine how rapidly  $\text{Mn}^{2+}$  could induce changes in adhesion formation and cell spread area, we performed live cell imaging of EGFP–paxillin and mApple–actin in cells plated on a soft substrate during addition of  $\text{Mn}^{2+}$  to the medium. (Fig. 4 C and D and [Movie S3](#)). Before addition of  $\text{Mn}^{2+}$ , there was significant protrusive activity on soft substrates, but no change in area or cell shape. Upon addition of  $3\ \mu\text{M}\ \text{Mn}^{2+}$ , protrusions stabilized, new focal adhesions formed, and the cell increased in




Oakes et al.



Collectively, these results illustrate that rigidity sensing in the lamellipodium is determined by catch-bond kinetics of integrin–fibronectin bonds and that the fraction of bound integrins is sensitive to both the unloaded lifetime and the maximum lifetime of the catch-bond curve. Addition of  $Mn^{2+}$  resulted in longer integrin lifetimes on soft substrates, thereby increasing the average fraction of bound integrins. This change in integrin binding kinetics allows cells to spread on soft substrates.





**Fig. 5.** Computational model of integrin-based adhesion dynamics. (A and B) Schematics of the computational model from the top (A) and side (B) perspectives. Two quasi-2D surfaces are placed 20 nm apart. The bottom surface represents the substrate and consists of a random distribution of ideal springs with stiffness,  $k_{\text{sub}}$ , proportional to the substrate Young's modulus. The top surface mimics a representative unit of the ventral surface of a fibroblast, with integrins diffusing (green particles) with diffusion coefficient,  $D$ , and establishing interactions with the substrate springs (purple particles). Upon binding the substrate, a force is exerted on the integrin particle parallel to the substrate, building tension on the bond. This tension determines the integrin unbinding rate,  $k_{\text{off}}$ . (C and D) Role of unloaded lifetime on the fraction of bound integrin on different substrate stiffnesses. In C, graphs of lifetime vs. force with unloaded lifetimes equal to the maximum lifetime,  $\tau_0 = \tau_{\text{max}}$ , at tensions  $\leq 30$  pN and zero otherwise, used to calculate the average fraction of ligand-bound integrins as a function of the substrate's Young's modulus for the curves shown in D. The average fraction of bound integrins is insensitive to substrate stiffness. (E and F) Role of  $\tau_{\text{max}}$  on the fraction of bound integrins at different substrate stiffnesses. In E, graphs indicating the bond lifetime vs. tension with fixed  $\tau_0$  and increasing  $\tau_{\text{max}}$  used to calculate the fraction of bound integrin as a function of substrate stiffness in F. In this case, the average fraction of bound integrins is weakly sensitive to substrate stiffness. (G) Lifetime vs. tension relationship for WT and  $\text{Mn}^{2+}$ -treated integrins, which amounts to both a shift in  $\tau_0$  and in  $\tau_{\text{max}}$ . (H) The average fraction of bound integrins for the curves shown in G. The number of bound integrins on soft substrates in the presence of  $\text{Mn}^{2+}$  is identical to the number of bound integrins for WT integrins on stiff substrates.

## Discussion

The ability to sense the stiffness of their extracellular environment is critical to cells' ability to regulate growth, viability, migration, and differentiation (1–3). Here, we show that fibroblasts exhibit a biphasic response in spreading on matrices of variable stiffness (Fig. 1). For matrices with a Young's modulus less than  $\sim 5$  kPa, cells were poorly spread with minimal adhesion assembly and few organized actin structures. Above  $\sim 8$  kPa, fibroblasts achieved a maximal spread area with typical adhesion assembly and highly organized actin cytoskeletons. This transition stiffness was comparable to physiological tissue stiffness (16) and was of the same order of magnitude as reported values (8–12). While it has been suggested that cell spread area as a function of substrate stiffness follows a power-law behavior (9, 12), using a larger number of substrates, we find that it is better described as biphasic.

Due to its overwhelming role in cellular force generation, myosin II has been presumed to be the predominant mechanism of substrate stiffness sensing by adherent cells (23–28). Here, however, we demonstrate a myosin-independent stiffness sensing mechanism that controls spread area and arises from forces generated by actin polymerization within the lamellipodium. Integrins, which connect and transmit stress between the cytoskeleton and the ECM, behave as catch-bonds whose lifetime is determined as a function of the applied load (45, 53). As the load on the integrin increases, the lifetime of the bond also increases (45). On stiff substrates, this increase in lifetime is sufficient to promote clustering and adhesion formation (Fig. 4). Conversely, on soft substrates, the reduction in stiffness leads to shorter bond lifetimes which inhibit the required clustering for adhesion formation (Fig. 5). Both our experimental and simulation data suggest that integrin force-dependent binding kinetics are most sensitive to substrates with a stiffness between  $\sim 5$  and 8 kPa. Addition of  $\text{Mn}^{2+}$ , which alters the kinetics of integrin–ECM bonds by increasing the unloaded and peak force lifetimes (45, 50–52), both increased the number of bound integrins and decreased the average spacing between bound integrins. Together, these effects promoted adhesion formation and enabled cells on soft substrates to spread and take on the morphology characteristics of cells on stiff substrates (Fig. 4).

Conceptually, this framework is similar to the general motor-clutch model (54) that has been suggested as a mechanism for understanding mechanosensitivity (23, 24, 26, 27). Instead of forces being generated by myosin motors, the force applied across the integrin bonds is generated by actin polymerization in the lamellipodium. These polymerization forces modulate the integrin–ECM bond kinetics and offer a surprisingly simple and elegant mechanism to understand substrate stiffness sensing. Previous work has established that there is a minimum spacing required between integrins for adhesion formation (38). Binding of integrins to their ligands also limits their diffusion in the membrane (47) and drives clustering at the nanoscale (36). Once a nanoscale cluster of integrins has formed, the force required to rupture the adhesion (i.e., the adhesion strength) is more than an order of magnitude greater than typical tensions generated in the cytoskeleton (55). Thus, by increasing the density of bound integrins, adhesion stabilization is increased, and the cell is able to spread.

Together, these results suggest that the lamellipodium acts as a myosin-independent mechanosensor, applying force to bound integrins via actin polymerization-driven retrograde flow. On soft substrates, the increased pliability of the matrix leads to a reduced load on the integrin–ECM bond, resulting in a shorter lifetime. This shorter lifetime prevents integrin clustering and thereby inhibits adhesion stabilization, leading to a poor ability to spread. On stiff substrates, integrin–ECM bonds experience greater loads and thus increased lifetimes, which promote adhesion stabilization and enable cells to spread. While these results do not exclude the possibility that myosin-generated forces may be one mechanism to probe substrate stiffness, they suggest that stiffness sensing emerges passively from the properties of the

**integrin binding kinetics.** Given that simply shifting these kinetics can induce spreading on soft substrates, it will be interesting in the future to explore whether this approach is sufficient to recover other functions found to be impaired by soft substrates, as in development, differentiation, and disease.

## Materials and Methods

Polyacrylamide gels of different stiffnesses were fabricated on glass substrates by altering the ratio of acrylamide to bisacrylamide as reported (5, 56). Images were obtained on an inverted Nikon Ti-E microscope by using either a Yokogawa CSU-X1 confocal scanhead or a Lumen 200Pro metal halide

light source. Detailed information about image analysis and the computational model can be found in *SI Materials and Methods*.

**ACKNOWLEDGMENTS.** This work was supported through internal funds from the University of Rochester (P.W.O.); National Institute of General Medical Sciences Grant GM085087 (to M.L.G.); and the Department of Defense/Army Research Office through a Multidisciplinary University Research Initiative Grant W911NF1410403 (to M.L.G. and G.A.V.). This work was also partially supported by the University of Chicago Materials Research Science and Engineering Center, which is funded by National Science Foundation Award DMR-1420709. The National Science Foundation XSEDE resources at the Pittsburgh Supercomputing Center provided computational time.

- Paluch EK, et al. (2015) Mechanotransduction: Use the force(s). *BMC Biol* 13:47.
- Iskratsch T, Wolfenson H, Sheetz MP (2014) Appreciating force and shape—The rise of mechanotransduction in cell biology. *Nat Rev Mol Cell Biol* 15:825–833.
- Charras G, Sahai E (2014) Physical influences of the extracellular environment on cell migration. *Nat Rev Mol Cell Biol* 15:813–824.
- Lo C-M, Wang H-B, Dembo M, Wang Y-L (2000) Cell movement is guided by the rigidity of the substrate. *Biophys J* 79:144–152.
- Yeung T, et al. (2005) Effects of substrate stiffness on cell morphology, cytoskeletal structure, and adhesion. *Cell Motil Cytoskeleton* 60:24–34.
- Prager-Khoutorsky M, et al. (2011) Fibroblast polarization is a matrix-rigidity-dependent process controlled by focal adhesion mechanosensing. *Nat Cell Biol* 13:1457–1465.
- Oakes PW, et al. (2009) Neutrophil morphology and migration are affected by substrate elasticity. *Blood* 114:1387–1395.
- Califano JP, Reinhart-King CA (2010) Substrate stiffness and cell area predict cellular traction stresses in single cells and cells in contact. *Cell Mol Bioeng* 3:68–75.
- Han SJ, Bielawski KS, Ting LH, Rodriguez ML, Sniadecki NJ (2012) Decoupling substrate stiffness, spread area, and micropost density: A close spatial relationship between traction forces and focal adhesions. *Biophys J* 103:640–648.
- Ghibaudo M, et al. (2008) Traction forces and rigidity sensing regulate cell functions. *Soft Matter* 4:1836–1843.
- Tee S-Y, Fu J, Chen CS, Janmey PA (2011) Cell shape and substrate rigidity both regulate cell stiffness. *Biophys J* 100:L25–L27.
- Engler A, et al. (2004) Substrate compliance versus ligand density in cell on gel responses. *Biophys J* 86:617–628.
- Breckenridge MT, Desai RA, Yang MT, Fu J, Chen CS (2014) Substrates with engineered step changes in rigidity induce traction force polarity and durotaxis. *Cell Mol Bioeng* 7:26–34.
- Raab M, et al. (2012) Crawling from soft to stiff matrix polarizes the cytoskeleton and phosphoregulates myosin-II heavy chain. *J Cell Biol* 199:669–683.
- Wang HB, Dembo M, Wang YL (2000) Substrate flexibility regulates growth and apoptosis of normal but not transformed cells. *Am J Physiol Cell Physiol* 279: C1345–C1350.
- Engler AJ, Sen S, Sweeney HL, Discher DE (2006) Matrix elasticity directs stem cell lineage specification. *Cell* 126:677–689.
- Fu J, et al. (2010) Mechanical regulation of cell function with geometrically modulated elastomeric substrates. *Nat Methods* 7:733–736.
- Paszek MJ, et al. (2005) Tensional homeostasis and the malignant phenotype. *Cancer Cell* 8:241–254.
- Mekhdjian AH, et al. (2017) Integrin-mediated traction force enhances paxillin molecular associations and adhesion dynamics that increase the invasiveness of tumor cells into a three-dimensional extracellular matrix. *Mol Biol Cell* 28:1467–1488.
- Zaidel-Bar R, Itzkovitz S, Ma'ayan A, Iyengar R, Geiger B (2007) Functional atlas of the integrin adhesome. *Nat Cell Biol* 9:858–867.
- Schiller HB, Friedel CC, Boulegue C, Fässler R (2011) Quantitative proteomics of the integrin adhesome show a myosin II-dependent recruitment of LIM domain proteins. *EMBO Rep* 12:259–266.
- Kanchanawong P, et al. (2010) Nanoscale architecture of integrin-based cell adhesions. *Nature* 468:580–584.
- Chan CE, Odde DJ (2008) Traction dynamics of filopodia on compliant substrates. *Science* 322:1687–1691.
- Macdonald A, Horwitz AR, Lauffenburger DA (2008) Kinetic model for lamellipodial actin-integrin “clutch” dynamics. *Cell Adhes Migr* 2:95–105.
- Bangasser BL, Odde DJ (2013) Master equation-based analysis of a motor-clutch model for cell traction force. *Cell Mol Bioeng* 6:449–459.
- Elosegui-Artola A, et al. (2014) Rigidity sensing and adaptation through regulation of integrin types. *Nat Mater* 13:631–637.
- Elosegui-Artola A, et al. (2016) Mechanical regulation of a molecular clutch defines force transmission and transduction in response to matrix rigidity. *Nat Cell Biol* 18: 540–548.
- Plotnikov SV, Pasapera AM, Sabass B, Waterman CM (2012) Force fluctuations within focal adhesions mediate ECM-rigidity sensing to guide directed cell migration. *Cell* 151:1513–1527.
- Yan J, Yao M, Goult BT, Sheetz MP (2015) Talin dependent mechanosensitivity of cell focal adhesions. *Cell Mol Bioeng* 8:151–159.
- Thievesten I, et al. (2013) Vinculin-actin interaction couples actin retrograde flow to focal adhesions, but is dispensable for focal adhesion growth. *J Cell Biol* 202:163–177.
- Sawada Y, et al. (2006) Force sensing by mechanical extension of the Src family kinase substrate p130Cas. *Cell* 127:1015–1026.
- Oakes PW, Banerjee S, Marchetti MC, Gardel ML (2014) Geometry regulates traction stresses in adherent cells. *Biophys J* 107:825–833.
- Hu K, Ji L, Applegate KT, Danuser G, Waterman-Storer CM (2007) Differential transmission of actin motion within focal adhesions. *Science* 315:111–115.
- Aratyn-Schaus Y, Gardel ML (2010) Transient frictional slip between integrin and the ECM in focal adhesions under myosin II tension. *Curr Biol* 20:1145–1153.
- Gardel ML, et al. (2008) Traction stress in focal adhesions correlates biphasically with actin retrograde flow speed. *J Cell Biol* 183:999–1005.
- Bachir AI, et al. (2014) Integrin-associated complexes form hierarchically with variable stoichiometry in nascent adhesions. *Curr Biol* 24:1845–1853.
- Alexandrova AY, et al. (2008) Comparative dynamics of retrograde actin flow and focal adhesions: Formation of nascent adhesions triggers transition from fast to slow flow. *PLoS One* 3:e3234.
- Cavalcanti-Adam EA, et al. (2007) Cell spreading and focal adhesion dynamics are regulated by spacing of integrin ligands. *Biophys J* 92:2964–2974.
- Reinhart-King CA, Dembo M, Hammer DA (2005) The dynamics and mechanics of endothelial cell spreading. *Biophys J* 89:676–689.
- Dubin-Thaler BJ, Giannone G, Döbereiner H-G, Sheetz MP (2004) Nanometer analysis of cell spreading on matrix-coated surfaces reveals two distinct cell states and STEPs. *Biophys J* 86:1794–1806.
- Chaudhuri O, et al. (2015) Substrate stress relaxation regulates cell spreading. *Nat Commun* 6:6364.
- Bauer A, et al. (2017) Hydrogel substrate stress-relaxation regulates the spreading and proliferation of mouse myoblasts. *Acta Biomater* 62:82–90.
- Denisin AK, Pruitt BL (2016) Tuning the range of polyacrylamide gel stiffness for mechanobiology applications. *ACS Appl Mater Interfaces* 8:21893–21902.
- Choi CK, et al. (2008) Actin and alpha-actinin orchestrate the assembly and maturation of nascent adhesions in a myosin II motor-independent manner. *Nat Cell Biol* 10: 1039–1050.
- Kong F, Garcia AJ, Mould AP, Humphries MJ, Zhu C (2009) Demonstration of catch bonds between an integrin and its ligand. *J Cell Biol* 185:1275–1284.
- Ponti A, Machacek M, Gupton SL, Waterman-Storer CM, Danuser G (2004) Two distinct actin networks drive the protrusion of migrating cells. *Science* 305:1782–1786.
- Rossier O, et al. (2012) Integrins  $\beta 1$  and  $\beta 3$  exhibit distinct dynamic nanoscale organizations inside focal adhesions. *Nat Cell Biol* 14:1057–1067.
- Xu X-P, et al. (2016) Three-dimensional structures of full-length, membrane-embedded human  $\alpha 5 \beta 1$  integrin complexes. *Biophys J* 110:798–809.
- Burnette DT, et al. (2011) A role for actin arcs in the leading-edge advance of migrating cells. *Nat Cell Biol* 13:371–381.
- Gailit J, Ruoslahti E (1988) Regulation of the fibronectin receptor affinity by divalent cations. *J Biol Chem* 263:12927–12932.
- Smith JW, Piotrowicz RS, Mathis D (1994) A mechanism for divalent cation regulation of beta 3-integrins. *J Biol Chem* 269:960–967.
- Mould AP, Akiyama SK, Humphries MJ (1995) Regulation of integrin alpha 5 beta 1-fibronectin interactions by divalent cations. Evidence for distinct classes of binding sites for  $Mn^{2+}$ ,  $Mg^{2+}$ , and  $Ca^{2+}$ . *J Biol Chem* 270:26270–26277.
- Choquet D, Felsenfeld DP, Sheetz MP (1997) Extracellular matrix rigidity causes strengthening of integrin-cytoskeleton linkages. *Cell* 88:39–48.
- Suter DM, Errante LD, Belotserkovsky V, Forscher P (1998) The Ig superfamily cell adhesion molecule, apCAM, mediates growth cone steering by substrate-cytoskeletal coupling. *J Cell Biol* 141:227–240.
- Coyer SR, et al. (2012) Nanopatterning reveals an ECM area threshold for focal adhesion assembly and force transmission that is regulated by integrin activation and cytoskeleton tension. *J Cell Sci* 125:5110–5123.
- Aratyn-Schaus Y, Oakes PW, Stricker J, Winter SP, Gardel ML (2010) Preparation of compliant matrices for quantifying cellular contraction. *J Vis Exp* (46):2173.