OCD Patients Severity & Clinical Insights

Analysis of Severity Distribution, Comorbid Depression, and Clinical Trends

Ramprasad M

INTRODUCTION:

Obsessive-Compulsive Disorder (OCD) is a chronic psychiatric condition characterized by intrusive thoughts (obsessions) and repetitive behaviours (compulsions), significantly affecting quality of life.

OBJECTIVE:

To analyse the severity distribution among OCD patients
To evaluate demographic patterns (age, gender, marital status)
To assess the relationship between severity and duration
To examine the prevalence of comorbid depression

DATA SOURCE:

OCD patient dataset consisting of 1,500 individuals

METHODLOGY:

- Data cleaning and preprocessing performed using SQL
- Key variables transformed into structured format
- Visualization and analysis conducted using Power BI

KEY VARIABLES:

- Age, Gender, Ethnicity, Marital Status
- Duration of symptoms (months)
- Y-BOCS score (Obsessions + Compulsions)
- Depression diagnosis

DEMOGRAPHIC INSIGHTS :

Key Observations:

- Gender distribution is nearly equal (Male: ~49.8%, Female: ~50.2%).
- Majority of patients are single, followed by married and divorced.
- Average age of patients: 46.78 years

Interpretation:

- OCD affects both genders equally.
- Higher proportion of single individuals may indicate early onset or social impact of the disease

CLINICAL SEVERITY OVERVIEW:

Findings:

- Average severity score: **39.67**
- Majority of patients fall under **Extreme severity category**
- Moderate and severe categories follow, while mild cases are least

Interpretation:

- Indicates a high burden of severe OCD cases
- Suggests delayed diagnosis or insufficient early intervention

SEVERITY vs DURATION ANALYSIS

Observations:

- Wide variation in severity across all durations
- No clear linear relationship between duration and severity
- Patients with long duration do not always show higher severity

Interpretation:

- OCD progression is highly variable
- Severity may depend on treatment adherence, coping mechanisms, and individual factors

DEPRESSION & COMORBIDITY INSIGHTS

Findings:

- Higher prevalence of depression in patients with **extreme severity**
- Depression cases decrease as severity reduces

Interpretation:

- Strong association between OCD severity and depression
- Highlights the need for integrated psychiatric evaluation

KEY FINDINGS:

- OCD severity is predominantly moderate to extreme
- Equal distribution across genders
- Significant number of patients remain single, indicating social impact
- No direct correlation between duration and severity
- High prevalence of comorbid depression in severe cases

CLINICAL IMPLICATIONS & CONCLUSION:

Clinical Implications:

- Early diagnosis and intervention are essential
- Routine screening for depression in OCD patients is recommended
- Treatment should be individualized based on severity

Conclusion:

Data-driven analysis provides valuable insights into OCD patterns, enabling better clinical decision-making and improved patient outcomes.

LinkedIn : https://www.linkedin.com/in/ramprasad-m-155b53214

OCD PATIENTS SEVERITY & CLINICAL INSIGHTS DASHBOARD

Analysis of severity distribution, comorbid depression, and duration trends

Total Patients

1,500

Average Age

46.78

Average Severity Score

39.67

Ethnicity

All

Marital Status Distribution



Gender Distribution



Severity Distribution



Severity vs Duration (by Gender)



Depression Prevalence Across Severity Levels