


Splinting



Advice adapted from IADT – Avulsion of permanent teeth 2020

Type of trauma	Splint type	Splinting time	Follow up
Root fracture	Passive flexible	a. 4 weeks b. If near cervical -up to 4months	a. 4 weeks splint removal 6-8 weeks, 4, 6, 12 months b. Cervical 4 month to remove splint and then 6months
Subluxation	Passive flexible	2 weeks	2 weeks remove splint. 12 weeks, 6, 12 months
Extrusive luxation	Passive flexible	2 weeks	2 weeks remove splint. 4, 8, 12 weeks, 6, 12 months
Lateral luxation	Passive flexible	4 weeks	Post 2 weeks endodontic evaluation. 4 weeks splint removal. 8,12 weeks, 6,12 months
Intrusive luxation	Passive flexible	4 weeks	<i>Incomplete root formation</i> -allow for re-eruption without intervention for 4 weeks. No re-eruption-orthodontic repositioning. <i>Complete root formation</i> - <3mm- allow for re-eruption. If no eruption within 8 weeks, then surgical repositioning, and splint. 3-7mm- reposition surgically/orthodontically. >7mm surgical repositioning
Alveolar fracture	Rigid/ flexible	4 weeks	4 weeks splint removal. 6-8 weeks, 4,6,12months