

Direct route vs old pipeline: 26 questions, scored against the printed page


Copy_of_0dea66c2_p009_q007


printed

7. The correct representation of wavelength intensity relationship of an ideal black body radiation at two different temperatures T_1 and T_2 is

(a) 


(b) 


(c) 


(d) 


direct route (PDF text layer + judged pictures)

The correct representation of wavelength intensity relationship of an ideal black body radiation at two different temperatures T1 and T2 is

(A) 

(B) 

(C) 


(D) 


stem ✓ options ✓ pictures ✓ ok


old pipeline (OCR + DeepSeek + Haiku, OCR-box crops)

The correct representation of wavelength intensity relationship of an ideal black body radiation at two different temperatures T_1 and T_2 is

(A) [figure option]
(B) [figure option]
(C) [figure option]
(D) [figure option]

(a) 

(b) 


(c) 

stem ✓ options ✗ pictures ✗ option (d) missing; options are stem images with labels, no option_images

Copy_of_0dea66c2_p028_q014

printed

14. The figure shows the change in concentration of species A and B as a function of time. The equilibrium constant K_c for the reaction




$A(g) \rightleftharpoons 2B(g)$ is :

(a) $K_c > 1$ (b) $K < 1$
(c) $K = 1$ (d) data insufficient

direct route (PDF text layer + judged pictures)

The figure shows the change in concentration of species A and B as a function of time. The equilibrium constant K_c for the reaction $A(g) \rightleftharpoons 2B(g)$ is:

(A) $K_c > 1$
(B) $K < 1$
(C) $K = 1$
(D) data insufficient




stem ✓ options ✓ pictures ✓ ok

old pipeline (OCR + DeepSeek + Haiku, OCR-box crops)

The figure shows the change in concentration of species A and B as a function of time. The equilibrium constant K_c for the reaction $A(g) \rightleftharpoons 2B(g)$ is:

(A) $K_c > 1$
(B) $K < 1$
(C) $K = 1$
(D) data insufficient



stem ✓ options ✓ pictures ✓ ok

Copy_of_0dea66c2_p042_q012

printed

12. In the below mentioned compounds the decreasing order of reactivity towards electrophilic substitution is

(i)  (ii) 


(iii)  (iv) 

(a) (iv) > (i) > (ii) > (iii)
(b) (ii) > (iii) > (i) > (iv)
(c) (iii) > (i) > (iv) > (ii)
(d) (i) > (ii) > (iii) > (iv)

direct route (PDF text layer + judged pictures)

In the below mentioned compounds the decreasing order of reactivity towards electrophilic substitution is

(A) (iv) > (i) > (ii) > (iii)
(B) (ii) > (iii) > (i) > (iv)
(C) (iii) > (i) > (iv) > (ii)
(D) (i) > (ii) > (iii) > (iv)






stem ✓ options ✓ pictures ✓ ok

old pipeline (OCR + DeepSeek + Haiku, OCR-box crops)

In the below mentioned compounds the decreasing order of reactivity towards electrophilic substitution is

(A) (iv) > (i) > (ii) > (iii)
(B) (ii) > (iii) > (i) > (iv)
(C) (iii) > (i) > (iv) > (ii)
(D) (i) > (ii) > (iii) > (iv)

(i)  (ii) 

(iii)  (iv) 

(a) (iv) > (i) > (ii) > (iii)
(b) (ii) > (iii) > (i) > (iv)
(c) (iii) > (i) > (iv) > (ii)
(d) (i) > (ii) > (iii) > (iv)

stem ✓ options ✓ pictures ✗ fig includes option text; label debris

Copy_of_0dea66c2_p042_q013

printed

13. The correct order of stability of the following carbanions is

(i) $\overline{R}CH_2$ (ii) 

(iii)  (iv) 

(a) III > IV > I > II (b) I > II > III > IV

(c) I > III > II > IV (d) IV > III > II > I

direct route (PDF text layer + judged pictures)


The correct order of stability of the following carbanions is

(A) III > IV > I > II

(B) I > II > III > IV

(C) I > III > II > IV

(D) IV > III > II > I




stem ✓ options ✓ pictures ✓ ok

old pipeline (OCR + DeepSeek + Haiku, OCR-box crops)

The correct order of stability of the following carbanions is

(A) (i) $\overline{R}CH_2$

(B) (ii) 

(C) (iii) 

(a) III > IV > I > II

(c) I > III > II > IV

(D) (iv) 

(b) I > II > III > IV


(d) IV > III > II > I



stem ✓ options ✗ pictures ✗ structures mislabelled as option images with option-text debris



Copy_of_0dea66c2_p042_q014

printed

14. Which of the following is not a resonance form of the enolate ion formed in the following acid base equilibrium?





(a)  (b) 


(c)  (d) 


direct route (PDF text layer + judged pictures)


Which of the following is not a resonance form of the enolate ion formed in the following acid base equilibrium?

(A) 

(B) 

(C) 

(D) 



stem ✓ options ✓ pictures ✓ ok


old pipeline (OCR + DeepSeek + Haiku, OCR-box crops)

Which of the following is not a resonance form of the enolate ion formed in the following acid base equilibrium?

(A)

(a)


se equilibrium?



(B)

(C)

(c)


(D) 

stem ✓ options ✗ pictures ✗ option images clipped/garbled, label debris

Copy_of_0dea66c2_p042_q016


printed

16. The major product of the following reaction is –



direct route (PDF text layer + judged pictures)

The major product of the following reaction is –



stem ✓ options ✓ pictures ✓ ok

old pipeline (OCR + DeepSeek + Haiku, OCR-box crops)


The major product of the following reaction is - (reagents: PhSNa in dimethylformamide)

(A) [figure option]

(B) [figure option]

(C) [figure option]

(D) [figure option]



stem ✗ options ✗ pictures ✗ extra reagent text in stem; options as one stem image with debris

Copy_of_0dea66c2_p048_q023

printed

23. In which of the following reaction, Markovnikov's rule is violated?

(a) $CH_3-O-CH=CH_2 \xrightarrow[CCl_4]{HBr}$

(b) $CH_3-NH-CH=CH_2 \xrightarrow[CCl_4]{HBr}$

(c) $CH_3-S-CH=CH_2 \xrightarrow[CCl_4]{HBr}$

(d) $O_2N-CH=CH_2 \xrightarrow[CCl_4]{HBr}$

direct route (PDF text layer + judged pictures)

In which of the following reaction, Markovnikov's rule is violated? HBr

(A) $CH_3-O-CH=CH_2 \xrightarrow[CCl_4]{HBr}$

(B) $CH_3-NH-CH=CH_2 \xrightarrow[CCl_4]{HBr}$

(C) $CH_3-S-CH=CH_2 \xrightarrow[CCl_4]{HBr}$

(D) $O_2N-CH=CH_2 \xrightarrow[CCl_4]{HBr}$

stem ✗ options ✗ pictures ✓ stem has stray HBr; option text mojibake

old pipeline (OCR + DeepSeek + Haiku, OCR-box crops)


In which of the following reaction, Markovnikov's rule is violated?

(A) $CH_3-O-CH=CH_2 \xrightarrow[CCl_4]{HBr}$

(B) $CH_3-NH-CH=CH_2 \xrightarrow[CCl_4]{HBr}$

(C) $CH_3-S-CH=CH_2 \xrightarrow[CCl_4]{HBr}$

(D) $O_2N-CH=CH_2 \xrightarrow[CCl_4]{HBr}$



stem ✓ options ✓ pictures ✗ stem image is neighbour-question fragment

Copy_of_0dea66c2_p048_q025

printed

25. The compound X in the following reaction is

$CH_3C\equiv CCH_3 \xrightarrow[(ii)\ Br_2]{(i)\ H_2/Pt} X$

(a) *d*-2, 3-Dibromobutane

(b) *l*-2, 3-Dibromobutane

(c) *dl*-2, 3-Dibromobutane

(d) *meso*-2, 3 Dibromobutane.

direct route (PDF text layer + judged pictures)

The compound X in the following reaction is (i) H /Pt2 CH_3C º CCH_3¾¾¾¾¾® X (ii) Br2

(A) d-2, 3-Dibromobutane

(B) l-2, 3-Dibromobutane

(C) dl-2, 3-Dibromobutane

(D) meso-2, 3 Dibromobutane.

stem ✗ options ✓ pictures ✓ stem garbled formula text

old pipeline (OCR + DeepSeek + Haiku, OCR-box crops)

The compound X in the following reaction is

$CH_3C\equiv CCH_3 \xrightarrow[(ii)\ Br_2]{(i)\ H_2/Pt} X$


(A) *d*-2,3-Dibromobutane

(B) *l*-2,3-Dibromobutane

(C) *dl*-2,3-Dibromobutane


(D) meso-2,3 Dibromobutane


26.



$\xrightarrow[(ii)\ CH_3Br]{(i)\ NaNH_2,\ NH_3}$ (A) $\xrightarrow[\text{Linder catalyst}]{H_2}$ (B) ;

Product (B) is:






stem ✓ options ✓ pictures ✗ 3 stray neighbour figures

Copy_of_0dea66c2_p048_q027

printed

27. Among the following compounds, the decreasing order of reactivity towards electrophilic substitution is




(a) III > I > II > IV (b) IV > I > II > III
(c) I > II > III > IV (d) II > I > III > IV

direct route (PDF text layer + judged pictures)

Among the following compounds, the decreasing order of reactivity towards electrophilic substitution is


(A) III > I > II > IV
(B) IV > I > II > III
(C) I > II > III > IV
(D) II > I > III > IV




stem ✓ options ✓ pictures ✓ ok

old pipeline (OCR + DeepSeek + Haiku, OCR-box crops)

Among the following compounds, the decreasing order of reactivity towards electrophilic substitution is

(A) 

(a) III > I > II > IV

(B) 

(b) IV > I > II > III

(C) (c) I > II > III > IV

(D) (d) II > I > III > IV


stem ✓ options ✗ pictures ✗ structures as option images with option-text debris


Copy_of_0dea66c2_p048_q028

printed

28. Bromination of ethyl benzene in presence of light gives

(a) 


(b) 

(c) 

(d) Both (a) and (b)

direct route (PDF text layer + judged pictures)

Bromination of ethyl benzene in presence of light gives

(A) 


(B) [figure option]
(C) [figure option]
(D) Both (a) and (b)

stem ✓ options ✗ pictures ✗ options B,C have no images

old pipeline (OCR + DeepSeek + Haiku, OCR-box crops)

Bromination of ethyl benzene in presence of light gives

(A) [figure option]
(B) [figure option]
(C) [figure option]
(D) Both (a) and (b)





stem ✓ options ✗ pictures ✗ whole question as stem image; options placeholders; debris

Copy_of_0dea66c2_p081_q010


printed


10. Pick up the final product in the following reaction.



(a) 


(b) 


(c) 


(d) 


direct route (PDF text layer + judged pictures)


Pick up the final product in the following reaction.

(A) 

(B) 

(C) 


(D) 





stem ✓ options ✓ pictures ✓ ok


old pipeline (OCR + DeepSeek + Haiku, OCR-box crops)

Pick up the final product in the following reaction. (reagents: C2H5MgBr, then H2O)

(A) 

(B) 

(C) 


(D) 


stem ✗ options ✗ pictures ✗ extra parenthetical in stem; option images wrong/labels debris

Copy_of_0dea66c2_p081_q011

printed


11. The structure of the major product formed in the following reaction





(a) 


direct route (PDF text layer + judged pictures)


The structure of the major product formed in the following reaction CH Cl 2 NaCN is DMF I CH Cl 2

(A) 

(B) 

(C) 

(D) 





stem ✗ options ✗ pictures ✓ stem garbled; option text garbage


old pipeline (OCR + DeepSeek + Haiku, OCR-box crops)


The structure of the major product formed in the following reaction (reagents: NaCN, DMF) is


(A) [figure option]



(a) 

(b) 

(c) 

(d) 

stem ✗ options ✗ pictures ✗ extra reagents in stem; only option A; debris

Copy_of_0dea66c2_p081_q012

printed

12. Which of the following is fast de-brominated ?

(a) 


(b) 


(c) 


(d) 


direct route (PDF text layer + judged pictures)

Which of the following is fast de-brominated ?

(A) 

(B) 


(C) 


(D) 


stem ✓ options ✓ pictures ✓ ok


old pipeline (OCR + DeepSeek + Haiku, OCR-box crops)

Which of the following is fast de-brominated ?

(A) (a) 

(B) (b) 

(C) (c) 


(D) (d) 


stem ✓ options ✗ pictures ✗ label debris, clipped fragments


Copy_of_0dea66c2_p081_q013


printed

13. Which of the following is least reactive towards nucleophilic substitution with aq. KOH ?

(a) 


(b) 


(c) 


(d) 


direct route (PDF text layer + judged pictures)

Which of the following is least reactive towards nucleophilic substitution with aq. KOH?

(A) 

(B) 


(C) 

(D) 


stem ✓ options ✓ pictures ✓ ok


old pipeline (OCR + DeepSeek + Haiku, OCR-box crops)

Which of the following is least reactive towards nucleophilic substitution with aq. KOH?


(A) (a) 

(b)

(B) 

(C) (c) 

(d)

(D) 

stem ✓ options ✗ pictures ✗ label debris, neighbour text fragment

Copy_of_0dea66c2_p081_q014


printed

14. Which of the following gives white precipitate when treated with silver nitrate solution ?

(a) $CHCl_3$

(b) $CH_2=CHCH_2Cl$

(c) $C_6H_5CH_2Cl$

(d) 

WWW.JEEBOOKS.IN


direct route (PDF text layer + judged pictures)

Which of the following gives white precipitate when treated with silver nitrate solution ?

(A) $CHCl_3$

(B) $CH_2=CHCH_2Cl$

(C) $C_6H_5CH_2Cl$

(D) 

stem ✓ options ✓ pictures ✓ ok

old pipeline (OCR + DeepSeek + Haiku, OCR-box crops)


Which of the following gives white precipitate when treated with silver nitrate solution?

(A) $CHCl_3$

(B) $CH_2 = CHCH_2Cl$

(C) $C_6H_5CH_2Cl$

(D) [figure option]


(d) 

stem ✓ options ✗ pictures ✗ option D only as stem image with (d) label

Copy_of_0dea66c2_p043_q020

printed

20. Arrange the following (w, x, y, z) in decreasing order of their boiling points:



(w) (x) (y) (z)

(a) w > x > z > y (b) w > x > y > z

(c) w > z > y > x (d) w > z > x > y

direct route (PDF text layer + judged pictures)


Arrange the following (w, x, y, z) in decreasing order of their boiling points:

(A) w > x > z > y

(B) w > x > y > z

(C) w > z > y > x

(D) w > z > x > y





(w) (x) (y) (z)

stem ✓ options ✓ pictures ✓ ok

old pipeline (OCR + DeepSeek + Haiku, OCR-box crops)

Arrange the following (w, x, y, z) in decreasing order of their boiling points:

(A) 

(B) 

(w) (x)

(C) (a) w > x > z > y

(c) w > z > y > x

(y) (z)

(D) (b) w > x > y > z


(d) w > z > x > y

stem ✓ options ✗ pictures ✗ option_images mislabelled, debris

Copy_of_0dea66c2_p043_q021


printed


21. Which of the following compounds has most acidic hydrogen?

(a) 

direct route (PDF text layer + judged pictures)

Which of the following compounds has most acidic hydrogen?

(A) 

(B) 

(C) N CH N H


(D) O CH N H


stem ✓ options ✗ pictures ✗ options C,D garbage and no images; B/C/D images missing


old pipeline (OCR + DeepSeek + Haiku, OCR-box crops)


Which of the following compounds has most acidic hydrogen?


(A) [figure option]

(a) 

(b) 

(c) 

(d) 




(III)

stem ✓ options ✗ pictures ✗ only option A; neighbour fragments, label debris

Copy_of_0dea66c2_p043_q023

printed

23. X and Y



X Y


are

(a) enantiomers
(b) diastereomers
(c) constitutional isomers
(d) conformers

direct route (PDF text layer + judged pictures)

X and Y are

(A) enantiomers
(B) diastereomers
(C) constitutional isomers
(D) conformers




X Y

stem ✓ options ✓ pictures ✓ ok

old pipeline (OCR + DeepSeek + Haiku, OCR-box crops)

X and Y are

(A) enantiomers
(B) diastereomers
(C) constitutional isomers
(D) conformers




X Y

stem ✓ options ✓ pictures ✗ sliver of stem text at top of figure


Copy_of_0dea66c2_p092_q008


printed


8. The major organic product formed from the following reaction :




is

(a) 


(b) 


(c) 


(d) 


direct route (PDF text layer + judged pictures)


The major organic product formed from the following reaction: O (i) CH NH32 ¾¾¾¾¾¾¾®..... (ii) LiAlH, (iii) H O42 is O – $NHCH_3$

(A) 

(B) 

(C) 

(D) 


(A) 

stem ✗ options ✗ pictures ✓ stem and option text garbled


old pipeline (OCR + DeepSeek + Haiku, OCR-box crops)


The major organic product formed from the following reaction ((i) CH3NH2; (ii) LiAlH4; (iii) H2O) is


(A) [figure option]
(B) [figure option]
(C) [figure option]
(D) [figure option]





is


(a) 

(b) 

(c) 

(d) 

9. 

(a) 

(b) 

(c) 


(d) 

stem ✗ options ✗ pictures ✗ extra reagents in stem; options placeholders; neighbour Q9 in image

Copy_of_0dea66c2_p092_q010


printed

10. Predict the product:



direct route (PDF text layer + judged pictures)

Predict the product: NHCH + NaNO + HCl 3 2 ¾¾¾®Product CH_3 NNO_2




stem ✗ options ✗ pictures ✓ stem and option text garbled

old pipeline (OCR + DeepSeek + Haiku, OCR-box crops)

Predict the product:
$C_6H_5NHCH_3 + NaNO_2 + HCl \rightarrow$ Product

(A) [figure option]





stem ✓ options ✗ pictures ✗ only option A; options in stem images with debris


Copy_of_0dea66c2_p092_q011


printed


11. In a set of reactions *m*-bromobenzoic acid gave a product D. Identify the product D.



(a) 


(b) 


(c) 


(d) 


direct route (PDF text layer + judged pictures)


In a set of reactions m-bromobenzoic acid gave a product D. Identify the product D.

(A) 

(B) 

(C) 

(D) 



stem ✓ options ✓ pictures ✓ ok

old pipeline (OCR + DeepSeek + Haiku, OCR-box crops)


In a set of reactions m-bromobenzoic acid gave a product D. Identify the product D. (SOCl2 gives B; NH3 gives C; NaOH/Br2 gives D)


(A) [figure option]


(B) [figure option]


(C) [figure option]


(D) [figure option]





(a) 

(b) 

(c) 

(d) 






stem ✗ options ✗ pictures ✗ extra reagent parenthetical; options placeholders; neighbour fragment

Copy_of_0dea66c2_p044_q025

printed

25. The most acidic proton and the strongest, nucleophilic nitrogen in the following compound



respectively, are

(a) N^a – H; N^b (b) N^b – H; N^c

(c) N^a – H; N^c (d) N^c – H; N^a

direct route (PDF text layer + judged pictures)


The most acidic proton and the strongest, nucleophilic nitrogen in the following compound respectively, are

(A) N^a – H; N^b

(B) N^b – H; N^c

(C) N^a – H; N^c

(D) N^c – H; N^a



stem ✓ options ✗ pictures ✓ options text mojibake

old pipeline (OCR + DeepSeek + Haiku, OCR-box crops)


The most acidic proton and the strongest nucleophilic nitrogen in the following compound respectively, are

(A) N^a – H; N^b

(B) N^b – H; N^c

(C) N^a – H; N^e

(D) N^e – H; N^a




respectively, are

stem ✓ options ✗ pictures ✗ options C,D use N^e instead of N^c; figure has fragments and caption text

Copy_of_0dea66c2_p044_q026

printed

26. The chlorine atom of the following compound



that reacts most readily with $AgNO_3$ to give a precipitate is

(a) Cl^a (b) Cl^b
(c) Cl^c (d) Cl^d

direct route (PDF text layer + judged pictures)


The chlorine atom of the following compound that reacts most readily with AgNO3 to give a precipitate is

(A) Cl^a

(B) Cl^b

(C) Cl^c

(D) Cl^d



stem ✓ options ✓ pictures ✓ ok

old pipeline (OCR + DeepSeek + Haiku, OCR-box crops)


The chlorine atom of the following compound that reacts most readily with $AgNO_3$ to give a precipitate is

(A) Cl^a

(B) Cl^b

(C) Cl^c

(D) Cl^d



stem ✓ options ✓ pictures ✓ ok

Copy_of_0dea66c2_p044_q028

printed

28. IUPAC name of the following compound



is

(a) 1 -hydroxycyclohex-4-en-3-one
(b) 1 -hydroxycyclohex-3-en-5-one
(c) 3-hydroxycyclohex-5-en-1-one
(d) 5-hydroxycyclohex-2-en-1 one

direct route (PDF text layer + judged pictures)


IUPAC name of the following compound is

(A) 1 -hydroxycyclohex-4-en-3-one

(B) 1 -hydroxycyclohex-3-en-5-one

(C) 3-hydroxycyclohex-5-en-1-one

(D) 5-hydroxycyclohex-2-en-1 one



stem ✓ options ✓ pictures ✓ ok

old pipeline (OCR + DeepSeek + Haiku, OCR-box crops)


IUPAC name of the following compound is

(A) 1-hydroxycyclohex-4-en-3-one

(B) 1-hydroxycyclohex-3-en-5-one

(C) 3-hydroxycyclohex-5-en-1-one

(D) 5-hydroxycyclohex-2-en-1-one





stem ✓ options ✓ pictures ✓ ok

Copy_of_0dea66c2_p044_q029

printed

29. The absolute configurations of the following compounds






respectively, are


(a) *R* and *R* (b) *S* and *S*
(c) *R* and *S* (d) *S* and *R*

direct route (PDF text layer + judged pictures)

The absolute configurations of the following compounds HCH SH_2HOH H C_3CH OH_2H C_3CH SH_2 respectively, are

(A) R and R
(B) S and S
(C) R and S
(D) S and R a







stem ✗ options ✗ pictures ✓ stem garbled; option D has stray a

old pipeline (OCR + DeepSeek + Haiku, OCR-box crops)

The absolute configurations of the following compounds respectively, are


(A) R and R
(B) S and S
(C) R and S
(D) S and R





respectively, are

30.




,

stem ✓ options ✓ pictures ✗ figure has caption text and neighbour Q30

Copy_of_0dea66c2_p084_q011

printed


11. Which carbocation is more likely to be formed in




direct route (PDF text layer + judged pictures)

Which carbocation is more likely to be formed in OH + the dehydration of H +


(A)




(B)




(C)




(D)




(A)



(C)







stem ✗ options ✗ pictures ✓ stem and option text garbled

old pipeline (OCR + DeepSeek + Haiku, OCR-box crops)


Which carbocation is more likely to be formed in the dehydration of 1-(1-methylcyclopentyl)ethan-1-ol with H^+?




(a)




(b)



(c)



(d)



stem ✗ options ✗ pictures ✗ stem invents compound name; no options; debris