



## Review

Denitrification and N<sub>2</sub>O:N<sub>2</sub> production in temperate grasslands: Processes, measurements, modelling and mitigating negative impacts

Surinder Saggar<sup>a,\*</sup>, N. Jha<sup>b</sup>, J. Deslippe<sup>a</sup>, N.S. Bolan<sup>c,d</sup>, J. Luo<sup>e</sup>, D.L. Giltrap<sup>a</sup>, D.-G. Kim<sup>a,1</sup>, M. Zaman<sup>f</sup>, R.W. Tillman<sup>b</sup>

<sup>a</sup> Ecosystems & Global Change Team, Landcare Research, Private Bag 11052, Palmerston North 4442, New Zealand

<sup>b</sup> Soil & Earth Sciences Group, Institute of Natural Resources, Massey University, Private Bag 11222, Palmerston North 4442, New Zealand

<sup>c</sup> Centre for Environmental Risk Assessment and Remediation, University of South Australia, SA-5095, Australia

<sup>d</sup> Cooperative Research Centre for Contaminants Assessment and Remediation of the Environment (CRC CARE), University of South Australia, SA-5095, Australia

<sup>e</sup> Climate, Land & Environment Section, AgResearch, Ruakura Research Centre, Private Bag 3123, Hamilton 3240, New Zealand

<sup>f</sup> Ballance Agri-Nutrients Limited New Zealand, Private Bag 12503, Tauranga 3143, New Zealand

## HIGHLIGHTS

- We review denitrification, denitrifier populations and enzyme activities, nitrous oxide and dinitrogen production in soils.
- We describe molecular techniques to assess gene diversity and reveal enzymes that consume nitrous oxide.
- We consider a process-based approach to quantify the nitrous oxide:dinitrogen ratio and nitrous oxide emissions.
- We discuss the management strategies to enhance nitrous oxide reduction during denitrification.

## ARTICLE INFO

## Article history:

Received 11 September 2012

Received in revised form 11 November 2012

Accepted 12 November 2012

Available online 20 December 2012

## Keywords:

Denitrifier genes and enzymes

Regulation

Quantification

Modelling

Nitrous oxide reduction

Management practices

## ABSTRACT

In this review we explore the biotic transformations of nitrogenous compounds that occur during denitrification, and the factors that influence denitrifier populations and enzyme activities, and hence, affect the production of nitrous oxide (N<sub>2</sub>O) and dinitrogen (N<sub>2</sub>) in soils. Characteristics of the genes related to denitrification are also presented. Denitrification is discussed with particular emphasis on nitrogen (N) inputs and dynamics within grasslands, and their impacts on the key soil variables and processes regulating denitrification and related gaseous N<sub>2</sub>O and N<sub>2</sub> emissions. Factors affecting denitrification include soil N, carbon (C), pH, temperature, oxygen supply and water content. We understand that the N<sub>2</sub>O:N<sub>2</sub> production ratio responds to the changes in these factors. Increased soil N supply, decreased soil pH, C availability and water content generally increase N<sub>2</sub>O:N<sub>2</sub> ratio. The review also covers approaches to identify and quantify denitrification, including acetylene inhibition, <sup>15</sup>N tracer and direct N<sub>2</sub> quantification techniques. We also outline the importance of emerging molecular techniques to assess gene diversity and reveal enzymes that consume N<sub>2</sub>O during denitrification and the factors affecting their activities and consider a process-based approach that can be used to quantify the N<sub>2</sub>O:N<sub>2</sub> product ratio and N<sub>2</sub>O emissions with known levels of uncertainty in soils. Finally, we explore strategies to reduce the N<sub>2</sub>O:N<sub>2</sub> product ratio during denitrification to mitigate N<sub>2</sub>O emissions. Future research needs to focus on evaluating the N<sub>2</sub>O-reducing ability of the denitrifiers to accelerate the conversion of N<sub>2</sub>O to N<sub>2</sub> and the reduction of N<sub>2</sub>O:N<sub>2</sub> ratio during denitrification.

© 2012 Elsevier B.V. All rights reserved.

## Contents

1. Introduction	174
2. Overview of denitrification	175
2.1. Denitrification reactions	177
2.2. Denitrifiers	177
2.3. Denitrification enzymes	177

\* Corresponding author. Tel.: +64 6 353 4934; fax: +64 6 353 4801.

E-mail address: [Saggar@landcareresearch.co.nz](mailto:Saggar@landcareresearch.co.nz) (S. Saggar).

<sup>1</sup> Present address: Wondo Genet College of Forestry & Natural Resources, Hawassa University, PO Box 128, Shshemene, Ethiopia.

2.4.	Denitrification genes	178
3.	Factors affecting denitrification	179
3.1.	Soil and plant factors	180
3.1.1.	Soil mineral nitrogen	180
3.1.2.	Carbon availability	180
3.1.3.	Oxygen supply and water content	181
3.1.4.	Soil pH	181
3.2.	Environmental factors	182
3.2.1.	Soil temperature	182
3.2.2.	Rainfall	182
3.2.3.	Soil drying–rewetting and freezing–thawing	182
3.2.4.	Availability of trace metals	183
4.	Approaches to identify and quantify denitrification and N <sub>2</sub> O and N <sub>2</sub> emissions	183
4.1.	Acetylene inhibition (AI) technique	183
4.2.	<sup>15</sup> N tracer technique	184
4.3.	Direct N <sub>2</sub> quantification	184
4.4.	Molecular techniques	184
5.	Modelling the denitrification process and emissions of N <sub>2</sub> O and N <sub>2</sub>	185
5.1.	Modelling approaches	185
5.2.	Model validation	186
5.3.	Future requirements	187
6.	Management practices to control denitrification and N <sub>2</sub> O and N <sub>2</sub> emissions	187
6.1.	Soil C availability	187
6.2.	N availability	187
6.3.	Plant composition	188
6.4.	Oxygen supply	188
6.5.	Soil pH	188
6.6.	Microbial enzymes	188
7.	Conclusions and future research needs	188
	Acknowledgements	189
	References	189

## 1. Introduction

Grasslands cover about 25% of the earth's surface, occupy 117 million km<sup>2</sup> of vegetated land, and provide feed for over 1800 million live-stock units (Saggar et al., 2009). Nitrogen (N) dynamics in temperate grasslands, occupying ca. 9 million km<sup>2</sup>, have been extensively studied, particularly for managed grazed pasture systems (Haynes and Williams, 1993; Ledgard et al., 1999; Bolan et al., 2004; Saggar et al., 2009, 2011a). The direct effects of grazing grassland include consumption of above-ground biomass, removal of nutrients in animal products (milk, meat and wool), redistribution of nutrients through uneven deposition of excreta and gaseous and leaching/run-off losses of nutrients. Managed grasslands have a high demand for N to support plant growth, which is provided by symbiotic N fixation in association with legumes such as white clover (*Trifolium repens* L.) or/and by N fertiliser. The relative contributions of symbiotic N fixation and N fertiliser to the total N input to pastoral systems vary widely according to the intensity of the production system. Globally, about 14–15 Tg N [1 Tg = 10<sup>12</sup> g = 1 million metric tonne] is applied as fertiliser in pastoral farms annually (Saggar et al., 2009), which is 14–15% of the total 103 Tg global fertiliser N use in 2010 (IFA, 2011).

Intensively managed pasture systems reach N saturation when plants, microorganisms and soils can no longer assimilate or retain excess N, which is lost from the system through leaching of nitrate (NO<sub>3</sub><sup>-</sup>) and gaseous emissions (Saggar et al., 2011a). In grazed grassland soils, the N associated with animal excreta (both urine and faeces) is an additional important component and critical source contributing to and affecting the N cycle. In New Zealand grazing systems' daily excretal N ranges between 182 and 398 g N (mean 290 g N) for dairy cattle (Saggar et al., 2004a), and between 26 and 50 g N (mean 40 g N) for sheep (Saggar et al., 2007b). About 60–75% of the dietary N ingested is excreted in the urine consisting of over 70% urea, while the bulk of N in dung is in organic form. Nitrogen deposition ranges from 20 to

80 g m<sup>-2</sup> in dung patches and 50 to 200 g m<sup>-2</sup> in urine patches (Oenema et al., 1997; Bolan et al., 2004). The estimated amount of N excreted in animal manures on grazing grassland is as large as, or larger than, the current global annual N fertiliser consumption of 103 Tg N (IFA, 2011). For example, in New Zealand, animals void almost 5 times more N annually (1.5 Tg) than the N applied in fertilisers (0.31 Tg) (Saggar et al., 2005). Consequently, urine makes a much greater contribution to gaseous and leaching losses than N fertiliser. Although N is the major nutrient limiting pasture production, it is also a major contributor to environmental degradation where it occurs in excess of plant and microbial requirements in soil (Saggar et al., 2005). Accordingly, the challenge is to maximise the use of N for production in agricultural systems, while limiting the release of reactive N (biologically, chemically and radiatively active N) into the environment. Fig. 1 provides a schematic of N transformations in a leguminous pasture. A detailed description of the biotic and abiotic N transformations in grazed pastures is given by Bolan et al. (2004).

In the context of environmental pollution and global climate change, NO<sub>3</sub><sup>-</sup> leaching attracts attention because of its potential effects on human and animal health, and on water pollution (Parliamentary Commissioner for the Environment, 2004). Gaseous N, such as ammonia (NH<sub>3</sub>), nitrous oxide (N<sub>2</sub>O) and nitric oxide (NO), cause concern because of their radiative or chemical effects on the atmosphere (Bolan et al., 2004). Denitrification completes the N-cycle by returning N<sub>2</sub> to the atmosphere (Fig. 1). Nitric oxide and N<sub>2</sub>O are produced at intermediate steps during the denitrification process and may escape to the atmosphere before being reduced to N<sub>2</sub>. While, N<sub>2</sub> is an environmentally benign gas, NO is a recognised pollutant that contributes to stratospheric ozone destruction and radiative forcing in the troposphere (Ravishankara et al., 2009). Nitrous oxide is also formed in soils during the microbial processes of nitrification. At low oxygen (O<sub>2</sub>) concentrations, nitrifying bacteria may contribute to N<sub>2</sub>O emissions through nitrifier-denitrification. However, biological denitrification is




Fig. 1. Schematic representation of nitrogen transformations in legume-based pastures.


the primary process producing  $N_2O$  in temperate grassland soils (Saggar et al., 2009) and it accounts globally for about 60% of total  $N_2O$  emissions to the atmosphere (Mosier et al., 1998; Kroeze et al., 1999).

This review explores the biotic and abiotic transformations of nitrogenous compounds that occur during denitrification, and the factors that influence denitrifier populations and their enzyme activities, and hence affect the production of  $N_2O$  and  $N_2$ . We address denitrification with emphasis on temperate grasslands, and provide an overview of N inputs and dynamics and their impacts on the key soil processes regulating denitrification and related gaseous emissions. We also consider difficulties and limitations in measurement techniques, and describe a process-based approach that can be used to quantify  $N_2O$  emissions and the  $N_2O:N_2$  product ratio with known levels of uncertainty. Finally, we discuss options for mitigating  $N_2O$  losses, identifying current gaps and limitations in our knowledge, and conclude by presenting the main research needs to develop future measurement, verification and mitigation strategies for  $N_2O$  emissions from managed temperate grasslands.

Only a brief description of N dynamics is given here, and the reader is referred to appropriate reviews that provide in-depth coverage of sources of N inputs, N transformations, gaseous and leaching losses, the environmental impacts of N losses, and their mitigation (Haynes and Williams, 1993; Ledgard et al., 1999; Bolan et al., 2004; de Klein and Eckard, 2008; Ledgard and Luo, 2008; Saggar et al., 2009, 2011a; Luo et al., 2010).

## 2. Overview of denitrification

Denitrification is considered as a much more potent source of  $N_2O$  than nitrification in grassland soils (Saggar et al., 2009). For the purposes of this review, denitrification is a microbially mediated process of dissimilatory reduction of one or both of the ionic nitrogen (N) oxides ( $NO_3^-$ , and nitrite  $NO_2^-$ ) that may ultimately produce  $N_2$  through a series of gaseous N oxide products (Eq. (1)).



The general requirements for biological denitrification are:

- the presence of bacteria, fungi, other denitrifying eukaryote or archaea possessing metabolic activity
- suitable electron donors such as available organic C compounds
- anaerobic conditions or restricted supply of  $O_2$
- availability of N oxides:  $NO_3^-$ ,  $NO_2^-$ , NO, or  $N_2O$  as terminal electron acceptors.

The rate of denitrification, and the relative proportions of NO,  $N_2O$ , and  $N_2$  produced, depend on the outcomes of complex interactions between soil properties, soil micro-organisms, climatic factors, and management practices. The availability of mineral N (both  $NH_4^+$  and  $NO_3^-$ ) and labile C, together with processes that affect reaction rates, such as temperature, pH and redox potential of soil microsites, controls the rate of denitrification in soils. These factors influence both the number of denitrifiers and the activity of denitrification enzymes (Luo et al., 1996; Cavigelli and Robertson, 2000; Wallenstein et al., 2006; Zaman et al., 2007, 2008a, 2008b; Liu et al., 2010).

The rates and percentage of N-input that was denitrified for 36 temperate grassland soils are shown in Table 1. This review of 241 datasets suggests a median daily loss of  $0.017 \text{ kg N ha}^{-1}$  through denitrification (Table 2) for a total mean loss of approximately 2.5% of available N. Thus globally each year, temperate grasslands could lose 5.6 Tg of N through denitrification.

Despite a qualitative understanding of the multiple factors that regulate the rates of denitrification in pastoral soils, we have only limited ability to integrate this knowledge to construct and validate robust and predictive process-based models of denitrification (Boyer et al., 2006). We lack knowledge of reliable quantitative denitrification rates for grazed pasture soils and how these rates and the  $N_2O:N_2$  product ratios vary spatially and temporally. Issues that hinder this understanding include: difficulties in measuring denitrification rates (Groffman et al., 2006), the large spatial and temporal heterogeneity of denitrification products, and the large background atmospheric  $N_2$  concentration (Davidson and Seitzinger, 2006). An understanding of the biological processes, and the soil and environmental factors that have the potential to reduce the production of  $N_2O$  during denitrification, is vital for the development of novel and effective  $N_2O$  mitigation technologies.

**Table 1**  
Summary of denitrification studies in temperate grasslands.

Location	Texture	Treatments	Denitrification method	Denitrification (kg N ha <sup>-1</sup> d <sup>-1</sup> )	N loss by denitrification (% of total N input)	Reference
				mean (min.–max.)	mean (min.–max.)	
Australia	<sup>a</sup>	AN + U (200 kg N ha <sup>-1</sup> )	Soil core	0.040 (0.018–0.061)	0.73 (0.32–1.11)	Eckard et al. (2003)
Belgium	Loamy, sandy loam	NK (200 kg N ha <sup>-1</sup> )	Soil core	0.038 (0.024–0.052)	7.52 (3.7–11.31)	Vermoesen et al. (1996)
Belgium	Sandy loam	NK (200 kg N ha <sup>-1</sup> )	Chamber	0.329 (0.008–0.654)	3.06 (1.2–5.7)	Vermoesen et al. (1997)
Belgium	Sandy loam	AN (295 kg N ha <sup>-1</sup> )	Soil core	0.240	22.8	Van Cleemput et al. (1994)
Belgium	Sandy loam	AN (215 kg N ha <sup>-1</sup> )	Soil core	0.059	4.9	Kaiser et al. (1996)
China	Sandy loam	Grazing intensity	Soil core	0.0001 (0.0000076–0.00013)	<sup>a</sup>	Xu et al. (2008)
Denmark	Loamy sand	U, ammonium solution (529 kg N ha <sup>-1</sup> )	Soil core	0.0140	<sup>a</sup>	Carter (2007)
France	Sandy	Grazing intensity	Soil Slurry	1.4 (1.2–1.8)	<sup>a</sup>	Patra et al. (2005)
Germany	Sand	CAN (78 kg N ha <sup>-1</sup> )	Chamber	<sup>a</sup>	2.3	Mogge et al. (1999)
Germany	Loam	AN, PS, CS (450 kg N ha <sup>-1</sup> )	Chamber	0.003 (0.001–0.008)	0.32 (0.1–0.7)	Schwarz et al. (1994)
New Zealand	Silt loam	U (500 kg N ha <sup>-1</sup> )	N balance	1.414 (0.863–1.974)	27.0 (16.4–37.5)	Clough et al. (1996)
New Zealand	Silt loam	Natural rainfalls (26 mm)	Soil core	0.020 (0.003–0.247)	<sup>a</sup>	Luo et al. (2000)
New Zealand	Fine sandy loam	U (500 kg N ha <sup>-1</sup> )	Soil core	0.014 (0.005–0.031)	3.61 (2.6–4.8)	Ruz-Jerez et al. (1994)
New Zealand	Silt loam	KN (200 kg N ha <sup>-1</sup> ), Lime, Zeolite	Sieved soil	1.48 (0.80–2.33)	24.7 (20.0–32.0)	Zaman et al. (2008a)
New Zealand	Silt loam	U, KN (400 kg N ha <sup>-1</sup> ), Lime, Zeolite	Soil core	0.0145 (0.0009–0.03)	0.591 (0.39–0.84)	Zaman and Nguyen (2010)
Netherlands	Sand	CAN, AS, U, KN, CA, O (kg N ha <sup>-1</sup> )	Soil core	0.152 (0.006–0.382)	4.99 (0.2–22.8)	Velthof et al. (1996)
Netherlands	Sand, loam, peat	Irrigation (0, 10 mm), temperature (10, 20 °C), KN (40, 80 kg N ha <sup>-1</sup> )	Soil core	1.413 (0.03–3.6)	<sup>a</sup>	de Klein and Van Logtestijn (1996)
Netherlands	Heavy clay	CS, KN, deposition (33–264 kg N ha <sup>-1</sup> )	Soil core	0.276	20	van der Salm et al. (2007)
Switzerland	Sandy loam	AN (85 kg N ha <sup>-1</sup> )	Soil core	0.078 (0.065–0.091)	18.1 (14.6–21.6)	Rudaz et al. (1999)
USA	Sandy loam to loamy sand	*	Chamber	0.015 (0.07–0.024)	<sup>a</sup>	Hixson et al. (1990)
USA	<sup>a</sup>	U + AS (240 kg N ha <sup>-1</sup> )	Soil core	0.142 (0.011–0.280)	10.6 (10.4–10.8)	Horwath et al. (1998)
USA	Sandy loam, loam, silt loam	Poultry litter (103–254 kg N ha <sup>-1</sup> )	Soil core	0.016 (0.002–0.042)	1.85 (0.4–4.8)	Marshall et al. (1999)
USA	<sup>a</sup>	KN (300 kg N ha <sup>-1</sup> )	N balance	5.67 (4.49–6.84)	56.7 (44.9–68.4)	Rolston et al. (1976)
USA	Sandy loam to loamy sand	Dry/wet meadow	Chamber	0.026	<sup>a</sup>	Walker et al. (1992)
USA	<sup>a</sup>	PS, AN (112 kg N ha <sup>-1</sup> )	Soil core	0.0003 (0.000072–0.00048)	0.03 (0.007–0.052)	Sullivan et al. (2005)
USA	<sup>a</sup>	U (510 kg N ha <sup>-1</sup> )	Soil core	0.0008 (0.00025–0.0015)	0.002 (0.0058–0.002)	Frank and Groffman (1998)
UK	Clay, clay loam	CAN, U (100, 200, 300 kg N ha <sup>-1</sup> )	Soil core	0.071 (0.004–0.216)	11.4 (3.3–26.3)	Jordan (1989)
UK	Clay	AN (210 kg N ha <sup>-1</sup> )	Chamber	0.009	1.0	Colbourn et al. (1984)
UK	Sandy loam over clay loam	CS, KN (700 kg N ha <sup>-1</sup> )	Chamber	0.071 (0.018–0.174)	5.04 (1.0–9.1)	Egginton and Smith (1986)
UK	<sup>a</sup>	AN (200 kg N ha <sup>-1</sup> )	Soil core	0.168 (0.080–0.24)	44.7 (41.3–48.1)	Elliott et al. (1991)
UK	Silty clay loam	CS (45 kg N ha <sup>-1</sup> ), AN (60 kg N ha <sup>-1</sup> )	Soil core	0.117 (0.06–0.151)	1.41 (0.9–1.7)	Ellis et al. (1998)
UK	Loam, silty clay loam	CS + (DCD, acid, nitrapyrin) (200–296 kg N ha <sup>-1</sup> )	Soil core	0.065 (0.003–0.237)	4.12 (0.1–15.6)	Pain et al. (1990)
UK	Loam over clay	AN (250, 500 kg N ha <sup>-1</sup> )	Chamber	0.038 (0.004–0.080)	5.13 (4.4–5.8)	Ryden (1983)
UK	Sandy loam	CS (44–72 kg N ha <sup>-1</sup> ), PS (223–278 kg N ha <sup>-1</sup> )	Soil core	0.059 (0.002–0.101)	5.59 (0.6–14.0)	Misselbrook et al. (1996)
UK	Coarse sandy loam	NK (280 kg N ha <sup>-1</sup> )	Soil core	0.645 (0.02–0.37)	<sup>a</sup>	Jarvis et al. (2001)
UK	Sandy loam	Clover, grass	Soil core	0.141 (0.024–0.464)	<sup>a</sup>	Šimek et al. (2004)

NK: Not known fertiliser, AN: ammonium nitrate, CAN: calcium ammonium nitrate, PS: Pig slurry, CS: Cattle slurry, U: urea, AS: ammonium sulphate, KN: potassium nitrate.

<sup>a</sup> No data available.



**Table 2**

Summarised values for denitrification rate and percent N loss for temperate grassland soils.

Site	Number study	Min	Max	Median	Mean $\pm$ s.e.	95% confidence interval of mean
Denitrification rate (kg N ha <sup>-1</sup> d <sup>-1</sup> )	241	0.0001	6.84	0.0171	0.258 $\pm$ 0.050	0.098
Nitrogen loss (% of total N input)	236	0.002	68.4	2.45	7.97 $\pm$ 1.26	2.51

Mitigation approaches should focus on ways to reduce the production of N<sub>2</sub>O during denitrification and enhancing the reduction of N<sub>2</sub>O to N<sub>2</sub> thus lowering the N<sub>2</sub>O:N<sub>2</sub> product ratio.

### 2.1. Denitrification reactions

In grazed pastures, autotrophic nitrification of various ammonium (NH<sub>4</sub><sup>+</sup>) inputs, including those released from animal excreta, fertiliser, and soil organic matter, leads to continuous formation of NO<sub>3</sub><sup>-</sup>-N, which is either taken up by plants, lost from soils as leachate or denitrified. Microbially mediated denitrification is an energy-yielding process catalysed by a range of intracellular enzymes. During denitrification, N-oxides are reduced through the addition of two electrons per N-atom. The product of each reduction forms the substrate for the subsequent step in the process, as follows:



These reactions differ with regard to the amounts of energy they yield. The relative redox potential (E'o) of the reduction of NO<sub>3</sub><sup>-</sup> to NO<sub>2</sub><sup>-</sup> is +420 mV; of NO<sub>2</sub><sup>-</sup> to NO is +375 mV; of NO to N<sub>2</sub>O is +1175 mV; and of NO to N<sub>2</sub> gas is +1355 mV (Zannoni, 2004).

### 2.2. Denitrifiers

Organisms capable of all reaction steps 2–5, as described above, are facultative-anaerobes that utilise NO<sub>3</sub><sup>-</sup> (dissimilative reduction) when O<sub>2</sub> becomes limiting in the environment. Denitrifiers utilise N oxides as the terminal electron acceptor in the generation of ATP and are phylogenetically diverse, spanning three kingdoms: bacteria, archaea, and eukarya. While ubiquitous in soil (Payne, 1981; Firestone, 1982; Tiedje, 1988) denitrifiers are not numerically dominant, typically representing only 0.5–5% of the total bacterial community (Tiedje, 1988). More than 60 genera of bacteria and archaea, as well as many saprobic fungi, and protozoans, are capable of denitrifying N oxides. Most prokaryotic, and all eukaryotic denitrifiers are heterotrophs, and many can grow by fermenting organic compounds. Chemolithotrophic denitrifying bacteria and archaea, which couple the reduction of N-oxides to the oxidation of inorganic compounds, are also known (Van Cleemput and Samater, 1996; Zumft, 1997; Fernández et al., 2008). All these microorganisms can use NO<sub>2</sub><sup>-</sup> instead of NO<sub>3</sub><sup>-</sup> for denitrification.

Many denitrifiers lack one or more enzymes to catalyse reduction steps 2–5, and are thus often said to be 'incomplete'. In soil, complete denitrification of NO<sub>3</sub><sup>-</sup> to N<sub>2</sub> predominantly occurs through the synergistic activities of microbial consortia (Zumft, 1997; Wallenstein et al., 2006). The denitrification systems of most fungi (Kobayashi et al., 1996, and references therein) and approximately one third of sequenced bacterial denitrifiers (Philippot et al., 2011) lack N<sub>2</sub>O reductase, and consequently


evolve N<sub>2</sub>O as the final denitrification product. In fact the composition of microbial communities is an important factor in determining the N<sub>2</sub>O:N<sub>2</sub> product ratio emitted from temperate pasture soils. Indeed, increased N<sub>2</sub>O emissions that were reported resulted from the addition of bacteria lacking N<sub>2</sub>O reductase to soils (Philippot et al., 2011) and also from fungi dominated temperate grassland soil (Laughlin and Stevens, 2002) where the mole fraction of N<sub>2</sub>O was 0.7 of total N<sub>2</sub>O + N<sub>2</sub>-N.

### 2.3. Denitrification enzymes

Denitrification is catalysed by the action of four, independent intracellular enzymes: Nitrate reductase (Nar), Nitrite reductase (Nir), Nitric-oxide reductase (Nor) and Nitrous oxide reductase (Nos) (Eq. (1)). Denitrification genes are named for their enzyme products. To distinguish the enzyme from the gene, it is conventional to use an italic font so that, for example, the respiratory Nar is encoded by the *Nar* genes (Fig. 2). The denitrification proteome (Nar, Nir, Nor and Nos plus several other proteins) is synthesised in response to oxygen depletion, and is blocked by high oxygen concentrations (both transcriptional and post-transcriptional control) (Van Spanning et al., 2007). With the onset of anaerobic conditions and addition of C substrate, these enzymes are usually sequentially induced (Holtan-Hartwig et al., 2000; Tavares et al., 2006). Nar is activated within 2–3 h, Nir between 2 and 12 h, and Nor between 24 and 48 h, which explains the sequence of production of these enzymes with developing anaerobic conditions in soil (Holtan-Hartwig et al., 2000).

Under anaerobic growth conditions, a NO<sub>3</sub><sup>-</sup> sensor protein controls the expression of the membrane-bound respiratory Nar. The Nar enzymes are typically composed of three subunits. The large alpha subunit, which varies in mass from 112 to 145 kDa (kilodalton), carries the molybdenum cofactor, the active site of the reductase (Hille, 1996). The smaller, beta subunit of the enzyme binds four Fe–S clusters, which are important for maintaining enzyme structure and the retention of the cofactor. The catalytic site of Nar is oriented inwards, and its action releases NO<sub>2</sub><sup>-</sup> into the cell cytoplasm (Fig. 2). During denitrification, NO<sub>2</sub><sup>-</sup> must cross the cell membrane before reduction by Nir, which is localized in the periplasmic space (Zumft, 1997). Where Nar and Nap co-occur within an organism, it is likely that Nar catalyses the first step of anaerobic denitrification since this process is energetically more favourable (Richardson et al., 2001).

Denitrifying bacteria contain one of two forms of Nir: a cytochrome cd<sub>1</sub> form or a Cu-containing form. The cytochrome cd<sub>1</sub> form is a homodimer composed of two 60 kDa subunits (Nurizzo et al., 1998). Each of the subunits contains a heme C and a heme D<sub>1</sub> prosthetic group. Heme D<sub>1</sub> is the catalytic site of the enzyme, and both the substrate NO<sub>2</sub><sup>-</sup> and the product NO, bind via their N-atoms to the Fe<sup>2+</sup> in heme D<sub>1</sub> (Nurizzo et al., 1998). At the catalytic site, NO<sub>2</sub><sup>-</sup> is protonated before water is cleaved to yield NO. The Cu-containing form of Nar in bacteria is similar to the fungal Nar. The Cu-Nir in bacteria and fungi are trimeric enzymes. The three 40 kDa subunits are tightly associated around a three-fold axis with a central channel. At the intersection of the subunits are six copper atoms, two at each junction. Copper is believed to bind the oxygen atom in NO<sub>2</sub><sup>-</sup>, and subsequent hydrolysis of the Cu-ONO intermediate, yields NO. Both forms of bacterial Nir are periplasmic and, analogous with fungi, Cu-Nir is located in the intermembrane space of the mitochondria (Kobayashi et al., 1996; Zumft, 1997).



**Fig. 2.** Sequential reductive pathway of denitrification showing the location of enzymes relative to the cytoplasmic membrane. The enzymes are nitrate reductase Nar (encoding genes *narG*, *napA*), nitrite reductase Nir (encoding genes *nirK*, *nirS*), nitric oxide reductase Nor (encoding genes *norB*, *norZ*, *qnor*), and nitrous oxide reductase Nos (encoding gene *nosZ*). Only one gene is required to carry out each step. The genes frequently targeted by molecular approaches are shaded in grey. Adapted from Ye et al. (1994) and Wallenstein et al. (2006).

Nitric oxide is membrane-labile and highly toxic to most bacteria and fungi including those that denitrify. As a result, all denitrifiers synthesize Nor to rid themselves of the potentially lethal product of Nir (de Boer et al., 1996; Butland et al., 2001; Richardson et al., 2009). During steady-state denitrification, extracellular NO is in the low (1–30) nanomolar range, and NO toxicity occurs at concentrations exceeding 1 mM. The Nor enzyme is virtually identical among bacterial denitrifiers that use either the Cu-Nir or the cytochrome *cd*<sub>1</sub> form of Nir. Bacterial Nor are complexes of at least two subunits of 17 and 53 kDa. The large subunit NorB is a hydrophobic cytochrome b, located within the cell membrane. The small subunit, NorC, is a c-type cytochrome containing a large heme-binding domain that protrudes into the periplasm. Nor reduces NO to N<sub>2</sub>O and requires two electrons to catalyse the formation of the N–N bond. These electrons are transferred from a periplasmic electron donor to the heme group of NorC, and subsequently to the catalytic site of NorB, thought to be a binuclear heme B- and non-heme Fe<sup>2+</sup> site, which would allow for the close proximity of two NO molecules (Zumft, 1997). The reaction mechanism of P450<sub>nor</sub>, the fungal N<sub>2</sub>O reductase, differs from that in bacteria. In fungi, a 44 kDa monoprotochrome exists in the ferric-state. NO-binding induces a conformational change in the enzyme, allowing it to accept electrons directly from NADH, to fuel the production of N<sub>2</sub>O by reaction with a second NO molecule (Zumft, 1997). The activity of isolated P450<sub>nor</sub> is more than five times higher than the activity of bacterial NO reductases (reviewed in Morozkina and Kurakov, 2007).

The reduction of N<sub>2</sub>O to N<sub>2</sub> is a respiratory process in its own right, and several bacteria have been identified that can use N<sub>2</sub>O as a sole electron acceptor to fuel heterotrophic growth. Conversely, not all complete denitrifiers will reduce exogenous N<sub>2</sub>O. Nos is similar in all the heterotrophic, phototrophic and chemolithotrophic bacteria where it has been studied, but is often absent in fungi. Located in the periplasm, it is a dimer of identical 66 kDa subunits. Each subunit contains up to four Cu atoms, which have been proposed as the catalytic sites of the enzyme (Zumft, 1997). Persistence of Nos was reported to be lower under aerated conditions in pastoral soil microcosms, and once anaerobic conditions were induced, N<sub>2</sub>O reduction occurred only after 16 h (Dendooven and Anderson, 1994), suggesting that the activity of this enzyme may be a strong short-term control on the N<sub>2</sub>O:N<sub>2</sub> ratio.

#### 2.4. Denitrification genes

Due to the phylogenetic diversity of denitrifiers, studies that seek to identify these organisms in natural habitats often target genes that

encode denitrification enzymes. The target gene usually codes for a key structural component of the enzyme of interest. For example, the *narG* or *napA* genes, which encode the large, catalytic subunit of Nar and periplasmic Nar, respectively, have been targeted for quantification in soils (Bru et al., 2007). However, because these genes can also be present in dissimilatory reducers of NO<sub>3</sub><sup>−</sup> to NH<sub>3</sub>, *narG* and *napA* have not been widely used to characterise denitrifying bacterial communities (Philippot et al., 2002; Chèneby et al., 2003, 2004).

Molecular characterization of the organisms that drive the denitrification process often begins at the second step of denitrification, with the genes that encode Nir. *NirK* encodes the Cu-containing form of Nir, while *nirS* encodes the cytochrome *cd*<sub>1</sub> form, and both genes have been widely used to characterise bacterial denitrifiers in soil (Sharma et al., 2005; Kandeler et al., 2006; Wertz et al., 2006, 2007; Bremer et al., 2007; Yoshida et al., 2009; Liu et al., 2010; Smith, 2010).

While polymerase chain reaction (PCR) primers to target two types of *norB* genes, which code for the large subunit of nitric oxide reductase, are available (Braker and Tiedje, 2003), relatively few studies have examined the presence of these genes as a marker for denitrifying bacteria in soils. Like the *narG* and *napA* genes, the *qnorB*-type genes have been discovered in a variety of non-denitrifying organisms and may be involved in the detoxification of exogenous NO (Richardson, 2000). Consistent with its radically different product, the fungal P450<sub>nor</sub> gene shows marked contrast to the bacterial *nor* genes (Kobayashi et al., 1996; Zhang et al., 2001). As yet, studies that target fungal P450<sub>nor</sub> genes have been limited to cultured organisms (Zhang et al., 2001, 2002; Kaya et al., 2004), and their role in denitrification in pasture soils requires further study.

*NosZ* encodes N<sub>2</sub>O reductase, and is unique in being the only known gene whose product catalyses the final step of denitrification (Burger and Matiassek, 2009). Like the *nir* genes, it is widely used to describe denitrifier communities (Chèneby et al., 1998; Rich and Myrold, 2004; Henry et al., 2006; Philippot et al., 2009, 2011; Čuhel et al., 2010). Primers for the amplification of *nosZ* from soils have improved over time. In an early study Chèneby et al. (1998) found that *nosZ* was unamplified from 35% of the N<sub>2</sub>O-reducing bacteria isolated from an agricultural soil. Later, updated primers allowed for the detection of *nosZ* in all but one N<sub>2</sub>O-reducing organism in a riparian soil (Rich and Myrold, 2004). The addition of serial dilutions of a bacterium lacking *nosZ* to experimental microcosms recently proved the principle that the genetic composition of the denitrifying community can influence the nature of the denitrification end products (Philippot et al., 2011), a finding that is likely to stimulate research in this area.

Denitrification genes are typically held in clusters within genomic DNA. Adjacent to genes that encode the structural gene products are genes that encode associated proteins such as those involved in co-factor synthesis, electron donation and protein assembly processes (Zumft, 1997; Demanèche et al., 2009). An interesting feature of *Nir* and *Nos* gene clusters is the occurrence of individual putative binding sites for cyclic AMP receptor proteins (Crp) and fumarate/nitrate reductase (Fnr) regulatory proteins (Demanèche et al., 2009). These proteins are thought to suppress the transcription of *Nir* and *Nos* in the presence of oxygen and NO (Vollack et al., 1999) and suggest that the production of enzymes to catalyse each step in the denitrification process may be under independent environmental control in some organisms. Conversely, in other organisms, gene clusters coding for denitrification enzymes may be linked to one another, forming denitrification “gene islands”. For example, *Nir*, *Nor* and *Nos* genes are closely associated in *Brucella melitensis* and share a single *NnrA* binding site for Crp/Fnr regulators (Haine et al., 2006), suggesting that the production of these three enzymes is under a single regulatory control.

In bacteria, the organization of denitrification gene clusters is often variable (Demanèche et al., 2009), and denitrification genes may be held in multiple copies within a single organism (Cramm et al., 1997; Etchebehere and Tiedje, 2005; Jones et al., 2008; Demanèche et al., 2009). Where denitrification genes occur in multiple copies, one or more copies are often plasmid borne (Cramm et al., 1997; Schwintner et al., 1998). These observations, together with the wide distribution of denitrification genes among phylogenetically unrelated bacteria, suggest that horizontal gene transfer is an important mechanism for the acquisition of the ability to denitrify (Philippot, 2002; Braker and

Tiedje, 2003; Demanèche et al., 2009). Accordingly, affiliating denitrification gene sequences to bacterial taxa is difficult because of poor congruence between denitrification gene trees and the 16S rRNA gene phylogeny (Jones et al., 2008; Demanèche et al., 2009). In a recent review, Bakken et al. (2012) established that denitrification phenotypes differ strongly in their tendency to emit  $N_2O$  and suggested that characterisation of community structure and diversity of denitrifier genes have a limited value in revealing the potential  $N_2O$  production. These authors proposed to identify regulatory gene sequences that characterise various denitrification regulatory phenotype (DRP) across phyla.

### 3. Factors affecting denitrification

Environmental factors that can affect denitrification have been categorised as proximal and distal regulators (Tiedje, 1988; Groffman and Tiedje, 1989; Groffman, 1991; de Klein et al., 2001; Wallenstein et al., 2006) (Fig. 3). Proximal regulators of denitrification are those that immediately affect denitrifying communities leading to instantaneous changes in denitrification rates. These include factors such as  $NO_3^-$  concentrations, C availability,  $O_2$  concentration, and temperature (Beauchamp et al., 1980; Knowles, 1982; Weier et al., 1993; Thomas et al., 1994; de Klein et al., 2001, 2010; Mosier et al., 2002; Dalal et al., 2003; Bolan et al., 2004; Wallenstein et al., 2006; Drury et al., 2008; Saggar et al., 2009). Distal regulators control the composition and diversity of denitrifying communities over larger spatial and temporal scales for a longer term and on a larger scale than proximal regulators (Tiedje, 1988; Groffman and Tiedje, 1989; Groffman, 1991; de Klein et al., 2001; Wallenstein et al., 2006). These regulators include factors such as plant growth, management practices (i.e. cultivation and animal grazing), soil




Fig. 3. Factors affecting denitrification in soils (red shaded: Proximal factors; blue shaded: Distal factors). Adapted from Tiedje (1988); Groffman (1991); and de Klein et al. (2001).

texture, soil pH, and water availability. In general, complete denitrification is promoted by high soil water content, neutral to slightly basic soil pH, high soil temperature, low rates of  $O_2$  diffusion and the presence of labile C.

While considerable experimental work has led to the generalizations about the effects of distal and proximal factors on denitrification and on the  $N_2O:N_2$  ratio (Knowles, 1982; Weier et al., 1993; de Klein and Van Logtestijn, 1994; Thomas et al., 1994; Mosier et al., 2002; Dalal et al., 2003; Bolan et al., 2004; Wallenstein et al., 2006; Zaman et al., 2007; 2008a, 2008b; Saggar et al., 2009; Zaman and Nguyen, 2010), the interactions among these factors are likely to control denitrification and the  $N_2O:N_2$  ratio under field conditions, and these interactions remain poorly understood. We understand the general trends of change in the  $N_2O:N_2$  ratio with changes in some of the individual factors. But how these factors interact to affect denitrification and  $N_2O:N_2$  ratios under various edaphic and environmental conditions remains poorly understood.

### 3.1. Soil and plant factors

#### 3.1.1. Soil mineral nitrogen

The availability of N, especially in the form of  $NO_3^-$ , for denitrifying bacteria is the primary requirement for denitrification and therefore the  $NO_3^-$  concentration in the soil solution can be one of the principal factors limiting denitrification. Soil  $NO_3^-$  concentrations depend on mineralisation and nitrification rates, plant N uptake, microbial immobilisation and  $NO_3^-$  movement by leaching and diffusion (Tiedje, 1988; Zaman et al., 2007). Diffusion of  $NO_3^-$ -N to active microsites is the primary limiting factor for denitrification in grazed pasture, especially in dry soils (Luo et al., 1999b). This positive relationship between denitrification rate and  $NO_3^-$  concentration has been reported when the  $NO_3^-$ -N concentration is  $<1$  mmol (Ogilvie et al., 1997; Zhong et al., 2010). Luo et al. (1996) found that the increase in  $NO_3^-$ -N concentration beyond  $50 \mu g\ g^{-1}$  soil is detrimental for denitrification activity. This negative or 'toxic' effect of high  $NO_3^-$  concentration on denitrification has also been identified in a recent study by Senbayram et al. (2012) who found a significant decline in denitrification rate with increasing soil  $NO_3^-$  concentration from 2 to 20 mM.


The  $NO_3^-$  concentration is one of the key factors that influence the  $N_2O:N_2$  ratio of denitrification, with higher  $NO_3^-$  concentrations usually resulting in a higher  $N_2O:N_2$  ratio (Bremner and Blackmer, 1978; Mosier et al., 1983; Arah and Smith, 1990; Weier et al., 1993; Zaman et al., 2007). In diary grazed pastures,  $NO_3^-$  concentrations under urine patches are usually high (up to  $418\ mg\ N\ kg^{-1}$  soil) compared with non-urine areas ( $1$  to  $11\ mg\ N\ kg^{-1}$  soil) (Saggar et al., 2004a). A higher level of  $NO_3^-$  in soil is also known to result in incomplete denitrification and thus higher  $N_2O:N_2$  due to suppression of Nos activity, the enzyme responsible for the conversion of  $N_2O$  to  $N_2$  (Blackmer and Bremner, 1978; Sahrawat and Keeney, 1986; Cho et al., 1997; Scholefield et al., 1997; Stevens and Laughlin, 1998). Because denitrifiers obtain more energy by reducing  $NO_3^-$  than by reducing  $N_2O$ , high  $NO_3^-$  concentration under the urine patches may favour  $N_2O$  production. This is supported by the results of Weier et al. (1993), who measured both  $N_2O$  and  $N_2$  emissions from four soils treated with three different rates of applied  $NO_3^-$  and found that high  $NO_3^-$  concentrations inhibited the conversion of  $N_2O$  to  $N_2$  and resulted in a higher ratio of  $N_2O:N_2$ .

The relatively few simultaneous measurements of  $N_2O$  and  $N_2$  emissions currently limit our ability to quantify changes in the  $N_2O:N_2$  product ratio in response to changes in  $NO_3^-$  concentration. Cai et al. (2002) measured both  $N_2O$  and  $N_2$  emissions and found an increase in  $N_2O:N_2$  ratio with increasing  $NO_3^-$  concentration, but the actual emission of  $N_2$  did not change. As  $NO_3^-$  concentration decreases, a greater proportion of the N is emitted as  $N_2$  due to the stimulation of Nos, where  $N_2O$  acts as a major electron acceptor (Swerts et al., 1996; Dendooven et al., 1997). Senbayram et al. (2009) observed low  $N_2O$  emissions following a large input of labile C, without significant  $NO_3^-$

input in soils amended with organic matter and suggested that most of the  $N_2O$  could have been converted to  $N_2$  — although they did not measure  $N_2$  emissions. Difficulties in accurately measuring  $N_2$  production under an  $N_2$ -rich atmosphere is the bottle-neck in further understanding the influence of  $NO_3^-$  on  $N_2O:N_2$  ratio during denitrification research. Recent developments (Molstad et al., 2007) in simultaneous quantification procedure for  $N_2O$  and  $N_2$  measurements are likely to improve understanding and estimation of gaseous losses of N during denitrification.

#### 3.1.2. Carbon availability

Denitrifiers require a readily available C before reduction of added  $NO_3^-$  can occur (Burford and Bremner, 1975; Delwiche, 1981).



The heterotrophic denitrification as described in Eq. (6) shows that 5 mol of glucose (30 mol of C) is required for the denitrification of 24 mol of  $NO_3^-$ . The theoretical ratios of COD (chemical oxygen demand) during denitrification are 2.86 for  $NO_3^-$  and 1.71 for  $NO_2^-$  and Akunna et al. (1993) found higher experimental ratios 3.7–4.1 for  $NO_3^-$  and 2.0–2.8 for  $NO_2^-$  in acetic and lactic acid media. The difference between theoretical and experimental ratios is due to the assimilation of C for cell synthesis. Heterotrophic denitrification is, therefore, often limited by labile C in agricultural soils, and any process that influences the rate of C mineralisation in soils (e.g., temperature, drying–wetting, tillage, liming, incorporation of crop residues, organic or inorganic fertiliser application, root exudates) can have a major impact on the denitrification rate.

Permanent pasture soils develop surface layers rich in organic material with high potential for denitrification when fertilised or when urine and dung are deposited during grazing (Ryden, 1986). The amendment of soils with farm effluents, slurries and organic manures containing readily available C as an energy source for denitrifiers creates anaerobic conditions through enhanced respiration and thereby trigger denitrification under anoxic conditions (Burford and Bremner, 1975; Beauchamp et al., 1989; Bhandral et al., 2007a; Nishina et al., 2009; Cayuela et al., 2010). Application of farm effluents such as dairy and piggery effluents has been shown to enhance  $N_2O$  emissions by increasing soil C availability (Bhandral et al., 2007a), and increased  $N_2O$  emissions following the application of animal waste slurries are often attributed to the increased labile C availability (Barton et al., 1999; Chadwick et al., 2000). Likewise, increased rates of denitrification following application of urea fertilisers compared with ammonium-based fertilisers (e.g., DAP) have often been attributed to the direct supply of dissolved organic C (DOC) from the urea and the solubilisation of soil C resulting from an increase in soil pH caused by urea hydrolysis (Barton et al., 1999). Interestingly, applying urea fertiliser with a C source (wheat straw and green manure) reduced  $N_2O$  emissions substantially compared with applying urea fertiliser alone (Aulakh et al., 2001). This is possibly due to microbial immobilisation of some of the added N (Tiedje, 1988) or dissimilatory nitrate reduction to ammonium (DNRA) (Matheson et al., 2002).

High denitrification rates are found in soils where the plants have been cut or damaged, and this has been attributed to the leaking of available organic C into the soil from the roots remaining in the soil (Beck and Christensen, 1987; Robertson et al., 1987). Thus the increases in rhizosphere available C following a winter grazing event may enhance denitrification when the soil moisture levels are high.

The accessibility of available C to microorganisms is, therefore, an important factor in controlling denitrification, especially under field conditions. The rate of C supply is likely to be faster in the case of DOC when compared with other more stable C sources such as lignin and cellulose in soils (Myrold and Tiedje, 1985). However, the presence of low molecular weight DOC fractions rather than total DOC



concentrations is an important factor controlling  $\text{N}_2\text{O}$  release in upland and peat soils after disturbance (Beauchamp et al., 1980; Drury et al., 2008; Saari et al., 2009). Indeed, Vallejo et al. (2006) noticed a decrease in  $\text{N}_2\text{O}:\text{N}_2$  with increasing concentration of DOC in various sources of effluents. Organic C and DOC concentrations decrease with depth in most mineral soils. Thus, although leaching of  $\text{NO}_3^-$ -N into lower horizons is a common phenomenon, the availability of C is usually limiting denitrification activity in subsoils (Parkin and Meisinger, 1989; Luo et al., 1996; Brye et al., 2001; Barton and Schipper, 2001). However, the drainage conditions that favour DOC losses may enhance denitrification in subsoil.

It is generally considered that increasing C availability decreases the ratio of  $\text{N}_2\text{O}:\text{N}_2$  (Smith and Tiedje, 1979; Arah and Smith, 1990; Dendooven et al., 1998). However, the effect of available C on the absolute and relative amounts of  $\text{N}_2\text{O}$  and  $\text{N}_2$  produced and emitted from soils also varies with  $\text{NO}_3^-$  concentration and aeration status (Water-filled pore-space; WFPS). It has been suggested that in anaerobic zones of fertilised soils,  $\text{NO}_3^-$  concentration may control the denitrification product ratio ( $\text{N}_2\text{O}:\text{N}_2$ ), while labile C concentration controls the denitrification rate (Tiedje, 1988; Weier et al., 1993). The change in the  $\text{N}_2\text{O}:\text{N}_2$  product ratio from denitrification due to the changing labile C: $\text{NO}_3^-$  ratio could be explained by changes in enzyme status, and/or the diffusion rate of  $\text{NO}_3^-$  into denitrifying microsites (Swerts et al., 1996; Weymann et al., 2010). The  $\text{N}_2\text{O}:\text{N}_2$  ratio is generally lower in the rhizosphere, which provides more available C in the form of root exudates and root debris, has generally low  $\text{NO}_3^-$  concentrations due to plant N uptake, and is characterised by low partial pressures of  $\text{O}_2$  (due to  $\text{O}_2$  consumption by plant roots) (Casella et al., 1984; Henry et al., 2008). On the other hand, a high  $\text{NO}_3^-$  concentration immediately after fertiliser application or animal urine deposition may increase the  $\text{N}_2\text{O}:\text{N}_2$  ratio in the rhizosphere (Gift et al., 2010).

In addition to its effect on denitrification and the  $\text{N}_2\text{O}:\text{N}_2$  ratio, the nature and availability of soil C affect the denitrification enzyme activity (DEA) and microbial community composition. In comparison with glucose, addition of liquid manure to soil increases the abundance of *NosZ*. This in turn enhances denitrification (Paul and Beauchamp, 1989; Van Kessel and Reeves, 2000) due to the ability of the manure to provide both labile and complex forms of C for the soil organisms.

### 3.1.3. Oxygen supply and water content

As denitrification is an anaerobic reaction,  $\text{O}_2$  availability is one of the most important factors inhibiting this process in soil (Knowles, 1982; Lloyd, 1993). The soil water content, together with the rate of  $\text{O}_2$  consumption by plant roots and microorganisms (respiration), determines the  $\text{O}_2$  availability in the root zone (Tiedje, 1988). Field studies have shown an increase in denitrification rate under restricted soil aeration associated with high soil water content (e.g., de Klein and van Logtestijn, 1994; Ledgard et al., 1999; Luo et al., 2000). Studies (e.g., de Klein and Van Logtestijn, 1994; Hefting et al., 2003) have shown the existence of a soil water threshold above which denitrification rates increased sharply with increasing soil water content. Below this critical soil water threshold value, denitrification rates appeared to be unrelated to soil water content. The soil water threshold value differs according to soil type and is expressed as water-filled pore space (WFPS). de Klein and van Logtestijn (1996) suggested that the critical WFPS for denitrification in many soils is equivalent to field capacity or above. Although the presence of small quantities of  $\text{O}_2$  (0.02 atm) can cause large decreases in DEA (Firestone et al., 1979), denitrification and DEA have also been reported during an oxic phase following an anoxic phase (Robertson et al., 1989; Baumann et al., 1996; Morley et al., 2008). Among N-reductase enzymes, *Nos* was more severely inhibited by oxygen than *Nar*, *Nir* and *Nor* indicating denitrification may continue for some time during drainage following flooding even if the  $\text{O}_2$  supply is partly or fully restored (Morley et al., 2008). However, because of higher inhibition of *Nos* these anoxic/oxic phases may result in higher production of  $\text{N}_2\text{O}$  than  $\text{N}_2$ .

Oxygen supply is also affected by soil compaction. In grazed pastures compaction can be caused by animal treading especially with high stocking rates (Bilotta et al., 2007; Drewry et al., 2008). Soil compaction restricts  $\text{O}_2$  diffusion within the soil, thereby increasing the rate of denitrification (Rosswall et al., 1989; Luo et al., 1999a). McTaggart et al. (1997) and Hansen et al. (1993) observed that soil compaction increased  $\text{N}_2\text{O}$  emission by 100% from an imperfectly drained clay loam and by 36% from a sandy loam soil. Soil compaction also affects the nitrification process and thereby influences the supply of  $\text{NO}_3^-$  for denitrification. For example, Bhandral et al. (2007b) monitored  $\text{N}_2\text{O}$  emissions from various N sources in a pasture soil affected by compaction. While  $\text{N}_2\text{O}$  emissions from a  $\text{NO}_3^-$  source ( $\text{KNO}_3$ ) were 14 times more in the compacted than the uncompacted soils,  $\text{N}_2\text{O}$  emission from ammonium and urea sources was only 3–4 times higher in compacted than uncompacted soil. The different  $\text{N}_2\text{O}$  emission rates were attributed to different transformations of the added N substrates and to the consequently different soil  $\text{NO}_3^-$  supplies. These findings highlight an important interaction among N-source and compaction that acts to control in situ  $\text{N}_2\text{O}$  emission rates.

The  $\text{N}_2\text{O}:\text{N}_2$  ratio has often been found to decrease with increasing soil water content (Colbourn and Dowdell, 1984; Davidson, 1992; Rudaz et al., 1999), particularly when the soil water content exceeds 75% WFPS (Davidson, 1992; Weier et al., 1993). Likewise, the measured  $\text{N}_2\text{O}:\text{N}_2$  ratio was highest ( $\geq 1$ ) under dry conditions during summer and early autumn when denitrification was relatively inactive (Ruz-Jerez et al., 1994). Changes in  $\text{N}_2\text{O}:\text{N}_2$  ratio with increased WFPS will also depend on the ability of denitrifiers to produce  $\text{N}_2$  or  $\text{N}_2\text{O}$  as an end product (Sigman et al., 2001; Wood et al., 2001; Casciotti et al., 2002; Morley et al., 2008).

### 3.1.4. Soil pH

Soil pH is another key regulator of the microbiological processes that affect  $\text{N}_2\text{O}$  and  $\text{N}_2$  production. It is generally agreed that denitrification is slower in acid conditions (Bremner and Shaw, 1958; Bryan, 1981; Nägele and Conrad, 1990; Fageria and Baligar, 2008), but denitrification can still occur at pH values as low as 3.5 and can account for significant N losses in naturally acid soils (Parkin et al., 1985; Weier and Gilliam, 1986). Šimek et al. (2000) found no significant relationship between soil pH and denitrifying enzyme activity, although denitrification potential was significantly correlated with soil pH. This and other work led Šimek and Cooper (2002) to conclude that the expression “optimum pH for denitrification has little or no meaning” unless it is qualified by specifying the particular aspect of denitrification that is being considered.

It is also generally accepted that the level of soil acidity influences the  $\text{N}_2\text{O}:\text{N}_2$  product ratio of denitrification. At lower soil pH the  $\text{N}_2\text{O}:\text{N}_2$  ratio is increased (Koskinen and Keeney, 1982; Nägele and Conrad, 1990; Struwe and Kjeller, 1994; Liu et al., 2010), with  $\text{N}_2\text{O}$  frequently appearing as the dominant product in acid soil (Christensen, 1985; Parkin et al., 1985). Several laboratory experiments using pH modifying treatments, and field experiments where acidifying fertilisers were applied, have shown that at lower soil pH, denitrification produces more  $\text{N}_2\text{O}$  and the ratio  $\text{N}_2\text{O}$  to  $\text{N}_2$  is increased (Koskinen and Keeney, 1982; Struwe and Kjeller, 1994; Šimek and Cooper, 2002; Wolf and Brumme, 2003; Venterea, 2007; Zaman et al., 2007, 2008a; Zaman and Nguyen, 2010; Čuhel and Šimek, 2011). Rochester (2003) derived a negative exponential function between  $\text{N}_2\text{O}:\text{N}_2$  ratio and soil pH from a number of laboratory and field studies across a range of soil types. Approximately equivalent amounts of each gas were emitted from the soil at pH 6.0 but for the alkaline grey clays (pH 8.3–8.5) the  $\text{N}_2\text{O}:\text{N}_2$  ratio was 0.02.

The mechanism of pH control of the  $\text{N}_2\text{O}:\text{N}_2$  ratio of denitrification is not fully understood. The increased  $\text{N}_2\text{O}:\text{N}_2$  product ratio in acidic soils may reflect an enhanced post-transcriptional sensitivity of bacterial *Nos* to low pH, relative to the other denitrification enzymes (Liu et al., 2010). The lower activity of *Nos* in acidic soils (Knowles,

1982; Thomsen et al., 1994) may also reflect an increased dominance of fungi relative to bacteria in these systems, as fungi generally lack Nos. Acidic soils may also favour N<sub>2</sub>O production by autotrophic and heterotrophic nitrifiers (Martikainen and de Boer, 1993). It has been observed that in *Pseudomonas mandelii* expression of *nirS* and *cnorB* was unchanged over the pH range of 6–8; however, it reduced substantially with a decrease in pH to 5 (Saleh-Lakha et al., 2009a). Furthermore, the presence of increasing amounts of NO<sub>2</sub><sup>−</sup> at lower pH levels due to enhancement of abiotic transformations of NO<sub>3</sub><sup>−</sup> may contribute to the increased proportion of N<sub>2</sub>O produced (Koskinen and Keeney, 1982).

Increasing soil pH above 6.0 may offer a mechanism to mitigate N<sub>2</sub>O emissions by shifting the balance between N<sub>2</sub>O and N<sub>2</sub> (Stevens and Laughlin, 1998; Šimek and Cooper, 2002; Zaman et al., 2007, 2008a; Liu et al., 2010; Zaman and Nguyen, 2010). While there is a little doubt that the N<sub>2</sub>O:N<sub>2</sub> product ratio decreases with increasing pH, reduction may be partially offset by the tendency toward increased total denitrification rates with increasing pH.

### 3.2. Environmental factors

#### 3.2.1. Soil temperature

Denitrification can occur over a wide range of soil temperatures ranging from sub-zero to 75 °C (Knowles, 1982). The minimum temperature for N<sub>2</sub>O emissions during denitrification is generally associated with the occurrence of free water, while the maximum temperature is approximately that value limiting biological activity and NO<sub>3</sub><sup>−</sup> availability in the soil. Within normal physiological range for the cell, DEA follows Michaelis–Menten kinetics with a typical Q<sub>10</sub> value of approximately 2 in soils (Stanford et al., 1975) and with optimal measured denitrification rates converging at 30 °C. Dobbie and Smith (2001) observed that the temperature response of N<sub>2</sub>O emissions was greater over a lower range of temperature than over a higher range, with Q<sub>10</sub> values of 3.7 and 2.3 for 5–12 °C and 12–18 °C, respectively. Temperature is one of the main factors causing temporal fluctuations in denitrification (Ryden, 1983).

In addition to the direct effects of temperature on denitrifier activity, indirect effects include changes in gas solubility, and gas diffusion (Craswell, 1978), as well as temperature limitation of substrate supply through direct effects on N-mineralisation, C, NO<sub>3</sub><sup>−</sup> availability and nitrification. Several researchers have found that N-losses through denitrification during the winter period were limited by soil temperature (Ryden, 1986; Jarvis et al., 1991; de Klein and Van Logtestijn, 1996), although rapid losses of N through denitrification were also reported after the addition of mineral fertilisers and organic manure at relatively low soil temperatures (Egginton and Smith, 1986; Thompson et al., 1987; Schwarz et al., 1994).

In temperate countries, generally winter application of fertilisers to grasslands is limited/prohibited. However, denitrification loss could be higher in winter (soil temperature below 10 °C) (Ruz-Jerez et al., 1994; Luo et al., 2000; Saggar et al., 2004a). The high winter N<sub>2</sub>O fluxes and denitrification loss have been attributed to a combination of WFPS exceeding 0.60, a high soil NO<sub>3</sub><sup>−</sup> N content and low plant uptake of N.

Denitrifying bacteria acclimate and denitrifying communities adapt to the historic range of soil temperatures (Schimel and Guldge, 1998), resulting in different temperature optimums for denitrification in different regions (Powelson et al., 1988; Malhi et al., 1990). Dorland and Beauchamp (1991) reported that C amendment lowered the temperature threshold at which denitrification can occur in soil. Temperature changes affect the substrate availability and thus control the expression of specific enzymes involved in transformation, however, abundances and community structure of denitrifiers remain unaffected (Stres et al., 2008). However, Saleh-Lakha et al. (2009a) observed a negative effect of low temperature (below 10 °C) on denitrification activity

and the sensitivity of *nirS* and *cnorB* expression which increased positively with increasing the temperature from 10 to 30 °C.

Because the activation energy of N<sub>2</sub>O reduction is higher than the activation energy of N<sub>2</sub>O production (Holtan-Hartwig et al., 2002), it has been suggested that more N<sub>2</sub>O is produced at low temperatures and the N<sub>2</sub>O:N<sub>2</sub> product ratio is increased (Keeney et al., 1979; Avalakki et al., 1995). While some studies have reported a decrease in the N<sub>2</sub>O:N<sub>2</sub> ratio with increasing soil temperature (Bailey, 1976; McKenney et al., 1984; Maag and Vinther, 1996), no relationship between denitrification rate and soil temperature could be detected in others (Focht, 1974; Rudaz et al., 1999). The variable effects of temperature observed in these studies likely reflect the many interacting ways that temperature may affect denitrification rates; many of these have not been fully elucidated.

#### 3.2.2. Rainfall

Enhanced denitrification rates, denitrification activity and N<sub>2</sub>O emissions following rainfall commonly reported for agro-ecosystems have been generally attributed to the dual effects of bringing applied substrates into contact with soil micro-organisms and a reduction in soil O<sub>2</sub> levels (Ellis et al., 1998; Van Kessel et al., 1993; Luo et al., 2000). Peak N<sub>2</sub>O emissions or denitrification rates in temperate grassland soils generally occur following rainfall or irrigation events (Velthof et al., 1996; Luo et al., 2000; Saggar et al., 2004a). The emissions are much shorter lived in coarse-textured soils than in fine-textured soils (e.g., 12 h in a sandy loam soil and 48 h in a clay loam soil) (Sexston et al., 1985).

Local topography also plays an important role in influencing the effect of rainfall on denitrification and N<sub>2</sub>O emission rates. Corré et al. (1990) observed a better correlation between rainfall levels and emission rates at the foot of a slope (R<sup>2</sup> = 0.73) than from a flat area (R<sup>2</sup> = 0.57). The time taken for emissions to return to background levels was also longer (48 h) from the foot of the slope than from the flat area (24 h). This likely resulted from differences in the rate of soil drying between these two areas. Luo et al. (2000) also found differing denitrification rates on various topographic areas in a dairy grazed pasture, possibly due to differences in soil moisture contents.

#### 3.2.3. Soil drying–rewetting and freezing–thawing

Plant water uptake followed by rapid rewetting during precipitation generally creates drying–wetting cycles in the rooting zone. For up to several days following the wetting of dry soil, organic matter mineralisation is enhanced by i) growth and turnover of microorganisms, ii) rhizosphere release of osmotically active substances such as sugar, and iii) the disruption of bonds that physically protect soil organic matter (Adu and Oades, 1978; Lundquist et al., 1999; Fierer and Schimel, 2002; Muhr et al., 2008). For example, increased soil denitrification rates and N<sub>2</sub>O emissions following wetting of dry soil by rainfall or irrigation have been reported for various agricultural systems (Kessavalou et al., 1998; Kim et al., 2009; Nobre et al., 2001) including grazed pastures (Luo et al., 1998; Garcia-Montiel et al., 2003; Saggar et al., 2004a; Kim et al., 2010). Increased soil N<sub>2</sub>O emissions resulting from denitrification following thawing of frozen soils have also been observed in various agricultural systems including grasslands (Flessa et al., 1995; Kammann et al., 1998; van Bochove et al., 2000; Priemé and Christensen, 2001; Ludwig et al., 2004; Öquist et al., 2004; Mørkved et al., 2006; Holst et al., 2008; Wagner-Riddle et al., 2008; Wolf et al., 2010; Virkajärvi et al., 2010).

The mechanisms considered responsible for increased denitrification and N<sub>2</sub>O emission following rewetting and thawing are:

- increased availability and accessibility of the substrate (Kieft et al., 1987; Edwards and Cresser, 1992; Davidson et al., 1993; Schimel and Clein, 1996; van Bochove et al., 2000; Priemé and Christensen, 2001; Pesaro et al., 2003; Grogan et al., 2004; Saetre and Stark, 2005; Yergeau and Kowalchuk, 2008)

- creation of anaerobic conditions (Nyborg et al., 1997; Li et al., 2000; de Bruijn et al., 2009).
- release of trapped N<sub>2</sub>O and N<sub>2</sub> (Burton and Beauchamp, 1994; Koponen et al., 2006; de Bruijn et al., 2009; Goldberg et al., 2010; Virkajärvi et al., 2010).

### 3.2.4. Availability of trace metals

Denitrification enzymes require several metal ions as cofactors, including iron, copper and molybdenum (Ferguson, 1998). Thus soils that for reasons of parent material composition, or biological, or chemical immobilisation, lack these metals may have reduced rates of denitrification. For example, molybdate ion (Mo<sup>6+</sup>) acts as a component of Mo-cofactor for Nar and ferric ion (Fe<sup>3+</sup>) is essential for the cytochrome subunits of both Nar and Nir. The addition of Mo<sup>6+</sup> and Fe<sup>3+</sup> usually affects the growth and denitrification rates of microbial communities by accelerated NO<sub>3</sub><sup>-</sup> utilisation and NO<sub>2</sub><sup>-</sup> accumulation (Labbe et al., 2003; Zhou et al., 2007; Pintathong et al., 2009). For other copper-dependent enzymes like Nar or Nir, there is one more non-copper dependent variant available, but Nos is the only enzyme with no alternative non-copper type available to carry out reduction of N<sub>2</sub>O in the absence of Cu. Richardson et al. (2009) consider that the failure of the reduction of N<sub>2</sub>O to N<sub>2</sub> in bacterial denitrification process could be attributed to incomplete assembly of NosZ due to lack of sufficient Cu to provide full complement of Cu co-factors. However, the Cu limitation will directly impact only on release of NosZ. The limitation of other trace metals such as Fe or Mo that form co-factors in Nar and Nir would lead to inhibition of NO<sub>2</sub><sup>-</sup> and NO fluxes so that N<sub>2</sub>O may not be formed (Richardson et al., 2009).

## 4. Approaches to identify and quantify denitrification and N<sub>2</sub>O and N<sub>2</sub> emissions

Accurately measuring N<sub>2</sub> and N<sub>2</sub>O emissions, identifying denitrifiers, estimating the N<sub>2</sub>O:N<sub>2</sub> ratio, and predicting denitrification rates from grazed pastures are always problematic. This is because of:

- i) natural, and livestock induced spatial and temporal variability (uneven excretal-N deposition and animal treading) where denitrification occurs and also in the factors driving the process, and the complex way in which they interact (Dandie et al., 2008)

- ii) methodological limitations such as associated change in substrate concentration, disturbances to the physical environment, lack of sensitivity, and high costs
- iii) difficulty in accurately measuring N<sub>2</sub>, the final product of denitrification due to the high background concentration in the atmosphere.

The two most commonly used approaches for determining denitrification rate from measurements of N<sub>2</sub> and N<sub>2</sub>O production include a technique based on the acetylene (C<sub>2</sub>H<sub>2</sub>) inhibition (AI) of N<sub>2</sub>O reduction (Balderston et al., 1976; Yoshinari and Knowles, 1976; Tiedje, 1988) and an isotopic method using substrates enriched in <sup>15</sup>N that allows subsequent <sup>15</sup>N gases to be determined by isotope-ratio mass spectrometry (Mosier and Klemetsson, 1994). More recently, direct quantification of N<sub>2</sub> has been attempted using airtight systems containing He or Ar either with continuous flow of He + O<sub>2</sub> (Butterbach-Bahl et al., 2002) or a closed system with periodic head-space sampling and replacing the sampled volume with He. Only a brief description of these three methods is given here, and the reader is referred to the recent review of Groffman et al. (2006) on the methods available to measure and calculate denitrification in terrestrial systems. Here we also discuss the potential of using molecular techniques to improve our understanding of the denitrification processes. All the techniques, together with their advantages and disadvantages, have been summarised in Table 3.

### 4.1. Acetylene inhibition (AI) technique

The AI method has been widely used for more than 40 years and is still the most commonly used method to study denitrification rates and denitrification enzyme activities in both aquatic and terrestrial systems. The limitations include the use of C<sub>2</sub>H<sub>2</sub> as a microbial substrate (Yeomans and Beauchamp, 1982; Terry and Duxbury, 1995), insufficient blockage of the Nos and incomplete inhibition of reduction of N<sub>2</sub>O to N<sub>2</sub> (Yeomans and Beauchamp, 1978; Oremland et al., 1984; Simarmata et al., 1993; Yu et al., 2010; Qin et al., 2011), inhibition of NO<sub>3</sub><sup>-</sup> production through nitrification (Seitzinger et al., 1993), interactions in the presence of sulphide (Soerensen et al., 1978; Dalsgaard and Bak, 1992; Jones and Knowles, 1992), or inhibition of denitrification that is tightly coupled to nitrification (Rysgaard et al., 1993; Nielsen and Revsbech, 1994) and

**Table 3**  
Summary of denitrification measurement techniques, their advantages and disadvantages.

Denitrification measurement method	Advantage	Disadvantage
Acetylene inhibition (AI) technique Approaches:  • Calcium carbide (CaC <sub>2</sub> ) granules. • In situ chambers in field. • Static cores	<ul style="list-style-type: none"> <li>• Simple to conduct</li> <li>• Can run large number of samples at a time</li> <li>• Removes the spatial and temporal variability of denitrification rate</li> <li>• Useful in studying the effect of soil and environmental factors on denitrification and denitrification enzyme activity assay</li> </ul>	<ul style="list-style-type: none"> <li>• Can only be used in NO<sub>3</sub>-dominated systems</li> <li>• Inhibits nitrification; can underestimate denitrification</li> <li>• Slow diffusion of C<sub>2</sub>H<sub>2</sub> into soil or sediments limits blockage of Nos</li> <li>• Rapid decomposition of C<sub>2</sub>H<sub>2</sub> by microbes</li> <li>• Contamination of C<sub>2</sub>H<sub>2</sub> with other gases can affect denitrification</li> <li>• Scavenging of NO leading to underestimation of denitrification</li> </ul>
<sup>15</sup> N tracer technique	<ul style="list-style-type: none"> <li>• Considered better than AR technique</li> <li>• Gives reliable estimates of denitrification</li> </ul>	<ul style="list-style-type: none"> <li>• Laborious process</li> <li>• Requires costly instruments</li> <li>• Addition of <sup>15</sup>N to the N limiting condition results in overestimation of denitrification</li> </ul>
Direct N <sub>2</sub> quantification	<ul style="list-style-type: none"> <li>• No labelled N or inhibitor is added</li> <li>• Highly sensitive method can even detect low denitrification changes</li> <li>• Can be used to study temperature and moisture effect</li> </ul>	<ul style="list-style-type: none"> <li>• Can only be used in enclosed incubation experiments</li> <li>• Complex and difficult system</li> <li>• Not suitable for longer period.</li> <li>• Challenge to measure N<sub>2</sub> against high atmospheric N<sub>2</sub> concentration</li> </ul>
Molecular approaches  • Polymerase chain reaction • DNA microarray technique • Immunological techniques	<ul style="list-style-type: none"> <li>• Give reliable and realistic qualitative indication of enzymes and encoding genes involved in denitrification</li> <li>• Can be used both in aquatic and terrestrial</li> </ul>	<ul style="list-style-type: none"> <li>• Require very expensive equipment</li> <li>• Require technical expertise to extract, analyse and interpret enzymes and encoding genes</li> <li>• Useful to study microbial diversity</li> </ul>



the  $C_2H_2$ -catalysed decomposition of NO (Bollmann and Conrad, 1997) that may result in under- or over-estimation of the denitrification rate under some conditions (Table 3). Murray and Knowles (2003) reported that  $C_2H_2$  does not scavenge for NO during denitrification and hence does not lead to underestimation of soil denitrification potential. In a recent study, Qin et al. (2011) observed incomplete inhibition of the reduction of  $N_2O$  to  $N_2$  by  $C_2H_2$  but little changes in soil denitrification potential, and proposed the use of a correction factor to account for the incomplete inhibition.

The technique has also been applied in studies that have led to better understanding of the spatial and temporal variability of denitrification (Groffman et al., 1999). Recently, Seo and DeLaune (2010) used AI in combination with selective antibiotics to quantify the contributions from denitrifying bacteria and fungi in a swamp forest. They reported that fungi were the main contributor to total denitrification (34% compared with 1% for denitrifying bacteria) under aerobic to weakly reducing conditions (Eh + 250 and Eh + 400 mV), whereas bacteria were the main contributor (65% compared with 17%) under strongly reducing conditions (Eh < -100 mV).

#### 4.2. $^{15}N$ tracer technique

The  $^{15}N$  tracer technique is used for direct measurements of the  $^{15}N$  labelled gases ( $^{15}N_2O$  and  $^{15}N_2$ ) produced following the addition of either  $^{15}NO_3^-$  alone or with  $^{15}NH_4^+$  to the soils, sediments, ground, or surface water (Hauck and Melsted, 1956; Hauck et al., 1958; Nishio et al., 1983; Aulakh et al., 1991; Stevens and Laughlin, 2001). Although small additions of labelled  $NH_4^+$  and  $NO_3^-$  do not alter the coupled nitrification/denitrification processes in fertilised agricultural soils, the  $^{15}N$  method may cause over estimation of denitrification in N-limited agricultural systems. The analysis of  $^{15}N$  in soils and  $^{15}N_2O$  and  $^{15}N_2$  gases produced during denitrification by isotope-ratio mass spectrometry (IRMS) can be fully automated (Stevens et al., 1993). Thus the fluxes of  $N_2$  and  $N_2O$  can be measured simultaneously in laboratory and field experiments by using the  $^{15}N$  stable isotope labelling method. When compared with the AI method, the  $^{15}N$  labelling method is considered to be better for direct quantification of denitrification. However, its application is often limited because of the high cost of  $^{15}N$ -substrates,  $^{15}N$  analyses, the expensive IRMS required, and the laborious procedures involved in sample preparation. The limitations associated with the use of the  $^{15}N$  technique are listed in Table 3. These include the difficulties in homogenous mixing and effects of denitrification, nitrification and plant N uptake on the uniform distribution of the labelled N in denitrifying zone during the experiment. However, despite reports of significant under- (Seitzinger et al., 1993) or over-estimation (Groffman et al., 2006) of denitrification by the  $^{15}N$  technique, it is still favoured for generating reliable estimates of denitrification rates in a range of environments, compared with the AI method. Use of this technique in conjunction with the AI method can provide improved estimates of denitrification rates in soils.

#### 4.3. Direct $N_2$ quantification

In this method intact soil cores or sediments are kept in air-tight systems with the air inside replaced by an inert gas like He or Ar, and the background  $N_2$  concentration is decreased from 79%  $N_2$  to approximately 1%  $N_2$  (Stefanson and Greenland, 1970; Wickramasinghe et al., 1978; Seitzinger et al., 1980). The  $N_2$  produced during denitrification within the air-tight container is then analysed by gas chromatography. In general, two types of incubation systems are used for direct  $N_2$  measurements: i) continuous air flow with He +  $O_2$  but no  $N_2$ , with changes in  $N_2$  gas concentration at the outlet used for calculating  $N_2$  emission or  $N_2/Ar$  ratio (Butterbach-Bahl et al., 2002; Cardenas et al., 2003) and ii) a closed system with the changes in the composition of gases in the headspace measured by periodically sampling small amounts of gases and replacing the sampled volume with He (Liu et al.,

2010). In both systems it is necessary to remove all  $N_2$  present in the test soil aggregates or cores before the measurements commence. The continuous flow incubation system has been improved by the development of a technique to measure  $N_2$  and  $N_2O$  emissions simultaneously (Scholefield et al., 1997; Butterbach-Bahl et al., 2002; Cardenas et al., 2003; Groffman et al., 2006; Khalil and Richards, 2011). An improvement in this technique is the use of Membrane Inlet Mass Spectrometer (MIMS) capable of measuring a large number of dissolved trace gases without any interference of high atmospheric background of  $N_2$ . This method can measure  $N_2$  at a precision of <0.05% for the  $N_2/Ar$  where Ar is considered as a tracer to minimise variations in  $N_2$  production. This technique is only suitable for short-term in-situ qualitative and quantitative denitrification measurements. For determining  $N_2O:N_2$  ratios  $N_2O$  measurements have to be conducted separately. Thus further studies are needed to test the suitability of MIMS technique.

A major advantage of this method is that no addition of substrate (labelled or otherwise) or inhibitor is required. Thus, there is no interference with soil microbial processes, including those involved in denitrification, during the measurements. However, the incubation system used is complex. As it is not possible to measure very small quantities of  $N_2$  gas, this method is not able to detect small changes in denitrification products, and the method appears to be only useful for measuring denitrification in highly fertilised agricultural and grassland soils (Scholefield et al., 1997). However, in spite of its limitations this method is increasingly being used for quantification of  $N_2$  emissions in both aquatic and terrestrial ecosystems (Butterbach-Bahl et al., 2002) and further improvements and automations are made using robotized sampling and analysing system to overcome some of the limitations.

#### 4.4. Molecular techniques

Recently, molecular techniques based on the identification and quantifications of the genes encoding enzymes which catalyse key denitrification processes (*NirS*, *NirK*, *NosZ*) have been used (Braker et al., 2000; Philippot, 2002; Henry et al., 2004). Many commercial kits are available for nucleic acid (DNA or RNA) extraction from soils. These and published protocols (e.g. Griffiths et al., 2000) typically involve physical cell lysis and the inactivation of nucleases, nucleic acid extraction from the soil matrix and purification followed either by targeted polymerase chain reaction (PCR)-based analyses or direct characterisation.

Molecular characterization of the micro-organisms that drive the denitrification process often begins at the second step of denitrification, with the genes that encode NIR. *NirK* encodes the copper-containing form of NIR while *nirS* encodes the cytochrome  $cd_1$  form, and both genes have been widely used to characterise bacterial denitrifiers in soil (Sharma et al., 2005; Kandeler et al., 2006; Wertz et al., 2006, 2007; Bremer et al., 2007; Yoshida et al., 2009; Liu et al., 2010; Smith et al., 2010). The desire to characterise only those organisms that are actively denitrifying, and thus potentially contributing to  $N_2O$  emissions or consumption, has motivated the use of reverse transcription polymerase chain reaction (RT-PCR) in studies of denitrifiers. In addition to studies of pure cultures of denitrifiers (Saleh-Ikha et al., 2009a, 2009b) RT-PCR has now been successfully applied to the study of denitrifier activity in sediments (Nogales et al., 2002; Smith et al., 2007) and more recently also in soils (Dandie et al., 2011; Shannon et al., 2011). Once the gene of interest is amplified the 'community profile' or 'fingerprint' of denitrifying organisms is obtained using terminal restriction fragment length polymorphism (T-RFLP), denaturing gradient gel electrophoresis (DGGE) or its variant temperature gradient gel electrophoresis (TGGE), and single stranded conformation polymorphism (SSCP) (e.g. Braker et al., 2001; Rich and Myrold, 2004; Rösch and Bothe, 2005; Bremer et al., 2007). Quantitative real-time PCR (qRT-PCR) is an evolution of PCR that allows for the detection and quantification of DNA template copy number in "real-time" as the PCR proceeds. It has been used to quantify several denitrification genes in soils including *nirK*, *nirS*, *narG* and *nosZ* (Grüntzig et al., 2001; López-Gutiérrez et al.,



2004; Henry et al., 2004; Kandeler et al., 2006; Siciliano et al., 2009; Levy-Booth and Winder, 2010; Chen et al., 2011). Gene copy number is calculated based on the rate of product accumulation in the exponential phase of the PCR. A critical issue in qRT-PCR is the construction of adequate controls so that samples and standards are carefully matched.

All of the molecular approaches for the characterization of denitrifiers in soils described above share a common dependence on PCR, which has several inherent biases, such as:

- PCR primer design, based on the sequences of known denitrifiers, represents only a small portion of all potential denitrifiers in soils and may not represent the gene's abundance (Acinas et al., 2005; Cardon and Gage, 2006)
- amplification errors may result in the production of sequence artefacts
- many community profiling methods saturate very quickly with regard to the complexity of the community that can be resolved and underestimate the diversity of the community
- it is not possible to link the characterization of a single phylotype of a functional gene to a species.

For direct characterisation of denitrifiers in soils immunological techniques, based on the identification of specific translated proteins, are used (Metz et al., 2003). This is accomplished by direct hybridization of a labelled antibody that can be detected fluorometrically, chemically or radioactively. While the efficacy of immunological techniques is well established in complex environments like soil (Metz et al., 2003; Maron et al., 2004) concerns exist about the sensitivity of the methods to very low concentrations of the target protein (Philippot and Hallin, 2005). Similarly, functional gene arrays, which consist of probes for a variety of functional genes from a range of species have been applied to quantify denitrification gene expression in soils with a varying degree of success (Chèneby et al., 1998; Henry et al., 2006; Babic et al., 2008; Hallin et al., 2009). Currently, while metagenomics, the culture-independent analysis of the metagenome, and the collective genomes of all organisms present in an environmental sample (Handelsman et al., 1998; Rondon et al., 2000; Gillespie et al., 2005), allow for the very rapid accumulation of microbial gene and genome data, an understanding of the functional and ecological relevance of these is evolving more slowly (Suenaga, 2012).

The above review suggests the application of a common methodology and molecular techniques for the identification and quantification of genes and characterisation of denitrifiers is a pre-requisite for comparisons across land use and land management systems under variable soil conditions (mineralogy, SOC levels, pH).

## 5. Modelling the denitrification process and emissions of N<sub>2</sub>O and N<sub>2</sub>

As a major source of N<sub>2</sub>O production from agricultural soils, the denitrification process must be an important part of any process-based soil N<sub>2</sub>O model. As discussed in the previous section, separate measurements of N<sub>2</sub>O and N<sub>2</sub> emissions from denitrification can be carried out at the field scale. However, emissions of both N<sub>2</sub>O and N<sub>2</sub> during denitrification are too variable, both spatially and temporarily, to be practically measured simultaneously at the field scale. Moreover, denitrification does not occur in isolation, and the success of a process-based denitrification model also depends upon how well the model simulates the other processes affecting the soil environment.

### 5.1. Modelling approaches

Simple empirical models can be used to estimate emissions, but these are limited to the types of soils and management practices used in the model parameterisation. More detailed, process-based models can be used to create emission scenarios and to examine the potential impacts of novel mitigation strategies. A recent review by Heinen

(2006) identified three types of denitrification models: i) microbial growth models; ii) soil structural models; and iii) simplified process models that represent the denitrification rate in terms of easily measurable parameters such as soil moisture, temperature and NO<sub>3</sub><sup>-</sup> concentration.

The microbial growth models consider the dynamics of microorganisms involved in N cycling processes. Grant (1991) used the maximum O<sub>2</sub> requirements for the growth and maintenance of microorganisms to calculate the maximum potential respiration and denitrification rates. But his model did not consider the relative production of N<sub>2</sub>O and N<sub>2</sub>. In an earlier attempt, Leffelaar and Wessel (1988) developed a denitrification and respiration model considering three pools of denitrifiers that respectively used NO<sub>3</sub><sup>-</sup>, NO<sub>2</sub><sup>-</sup>, and N<sub>2</sub>O as electron acceptors. This scheme has been used by more detailed process-based models such as DNDC (Denitrification–Decomposition) (Li et al., 1992) and NLOSS (N Loss) (Riley and Matson, 2000). DNDC also includes temperature and pH rate modifying factors in the denitrifier growth and electron acceptor consumption equations.

Soil structural models consider gaseous diffusion into and out of soil aggregates with denitrification only occurring in anoxic parts of soil aggregates. Arah and Smith (1989) considered steady-state denitrification in a spherical soil aggregate only occurring within the anaerobic part of the aggregate. In these models denitrification is calculated from the estimates of NO<sub>3</sub><sup>-</sup> diffusion into the aggregate using Michaelis–Menten kinetics with respect to the local NO<sub>3</sub><sup>-</sup> concentration at radius  $r$  within the pore. The model was then scaled up to an assembly of aggregates by assuming log-normal distribution of aggregate radii and oxygen reduction potentials. Vinten et al. (1996) subsequently used an approximation of the anaerobic volume fraction ( $\Phi$ ) based on the mean aggregate radius, the mean oxygen reduction potential, the O<sub>2</sub> concentration in the inter-aggregate air-filled pore space, the water content, the water content of saturated aggregates, the diffusion coefficients of O<sub>2</sub> in air and water, and the volume fraction of soil occupied by aggregates.

A simpler “anaerobic balloon” (i.e. volumetric fraction of anaerobic microsites) concept is included in the DNDC (denitrification–decomposition) model to account for the fact that both nitrification and denitrification could occur simultaneously within the soil (Li et al., 2000), and is based on the relative partial pressures of O<sub>2</sub> in the liquid and the air. Substrates such as soil N and DOC are partitioned between the aerobic and anaerobic soil fractions, with nitrification occurring in the aerobic fraction and denitrification occurring within the anaerobic fraction.

The simplified process-based models represent denitrification as a base rate multiplied by factors accounting for the effect of soil nitrate ( $f_N$ ), water content ( $f_s$ ), temperature ( $f_T$ ) and pH ( $f_{pH}$ ) (Heinen, 2006). This type of equation assumes that the effects of all the rate modifying influences are independent of each other. If the denitrification rate is independent of the reactant concentration then the relationship is called a zero-order process, while a first-order process is one where the denitrification rate is directly proportional to the reactant concentration. A Michaelis–Menten or Monod type relationship (Eq. (6)) can also be used to simulate both reactant-limited and non-reactant limited cases:

$$f_N \propto \frac{N}{N + K} \quad (6)$$

where  $N$  is the concentration of NO<sub>3</sub><sup>-</sup> and  $K$  is a constant. Eq. (6) becomes approximately constant (zero order) for high values of  $N$  and approximately linear (first order) for low values of  $N$ . Table 4 lists the form of  $f_N$  used in a selection of denitrification models.

Denitrification processes are highly sensitive to the O<sub>2</sub> status of the soil, as the denitrifying bacteria are anaerobic. Many models use some form of soil moisture measurement (such as WFPS) as a proxy for O<sub>2</sub> availability. As described earlier, there is a threshold WFPS

**Table 4**  
Form of  $f_N$  used in some denitrification models.

Model Name	Form of $f_N$	Reference
ANIMO	Zero-order	Rijtema and Kroes (1991)
FASSET	Michaelis–Menten	Chatskikh et al. (2005)
InfoCrop	First-order	Aggarwal et al. (2006)
NEMIS	Michealis–Menten	Hénault and Germon (2000)
NOE	Michealis–Menten	Hénault et al. (2005)
WNMM	First-order	Li et al. (2007)

level beneath which denitrification does not occur, which varies depending on soil type (de Klein and van Logtestijn, 1996). Most denitrification models therefore use  $f_s = 0$  below a threshold WFPS, although the level of this threshold varies between models. For WFPS above the threshold,  $f_s$  is commonly (Heinen, 2006) expressed as:

$$f_s = \begin{cases} \left( \frac{WFPS - WFPS_{th}}{WFPS_{max} - WFPS_{th}} \right)^w, & WFPS_{th} \leq WFPS_{max} \\ 1, & WFPS > WFPS_{max} \end{cases} \quad (7)$$

where  $WFPS_{th}$  is the denitrification threshold WFPS,  $WFPS_{max}$  is the WFPS above which  $f_s = 1$ , and  $w$  is an empirical parameter.

At higher WFPS values a greater proportion of the denitrified N appears as  $N_2$  rather than as  $N_2O$ . In most models  $D_a$  represents the total  $N_2O + N_2$  produced by denitrification but some models, such as InfoCrop (Aggarwal et al., 2006), calculate only  $N_2O$ . Models primarily aimed at N-availability for plant growth or water contamination risk usually do not disaggregate the separate  $N_2O$  and  $N_2$  emission rates. However, for greenhouse gas calculations the proportion of denitrified N released as  $N_2O$  is critical. The ratio of  $N_2O:N_2$  can be calculated in numerous ways; some models assume a constant  $N_2O:N_2$  ratio, while others use functions based on various soil properties. Table 5 shows the factors considered in calculating the  $N_2O:N_2$  ratio in a number of models.

In general, microbial activity and denitrification will increase with increasing temperature, although at very high temperatures microbes will begin to die off, reducing the denitrification rate. Most models assume this will not occur at temperatures usually found in field conditions. Some models, such as NOE (Hénault et al., 2005) and NEMIS (Hénault and Germon, 2000), use stepwise functions for  $f_T$ , representing different microbial populations that dominate in different temperature regimes.

There are many different relationships used to account for the temperature effect, but one commonly used is a  $Q_{10}$  response of the form:

$$f_T = Q_{10}^{(T-T_b)/10} \quad (8)$$

where  $Q_{10}$  represents the factor by which the reaction is increased given an increase of 10 °C from some base temperature  $T_b$ .

As discussed earlier, denitrification rates are lower in acidic soils than neutral or slightly alkaline soils, and low pH also influences the proportions of  $N_2O$  and  $N_2$  produced. However, many models neglect the pH factor as soil pH does not change rapidly, except around events such as liming, and urine or urea application and deposition. The DNDC model (Li et al., 1992) uses three different pH reduction factors

**Table 5**  
Calculation of  $N_2O:N_2$  ratio in some denitrification models.

Model name	Variables used in $N_2O:N_2$ ratio function	Reference
DayCent	$NO_3^-/CO_2$ ratio, WFPS	Stehfest and Müller (2004)
FASSET	Temperature, WFPS, clay, depth	Chatskikh et al. (2005)
NGAS	Soil $NO_3^-$ , respiration and WFPS	Parton et al. (1996)
NOE	Empirical site-based parameter	Hénault et al. (2005)
WNMM	WFPS	Li et al. (2007)

for the different stages of denitrification ( $NO_3^- \rightarrow NO_2^-$ ;  $NO_2^- \rightarrow N_2O$ ;  $N_2O \rightarrow N_2$ ).

In simplified process models, the microbial activity or population status of the soil in the denitrification rate calculation is usually incorporated into the base denitrification rate. Some models use soil organic C as an indicator of the relative size of the microbial pool or explicitly track the size of microbial pools, and use soil respiration as an indicator of microbial activity. Other models use site-specific empirical parameters that implicitly incorporate microbial activity. The approaches used to account for microbial activity in different models are summarised in Table 6.

## 5.2. Model validation

Testing denitrification models is complicated by the fact that under field conditions there are many interacting processes that will affect denitrification rates. e.g., heat and water transport through soil and other N transformation processes. Therefore, the success of a process-based denitrification model depends not only on the quality of the denitrification component but also on its ability to simulate soil conditions accurately. For example, denitrification rates are highly dependent on soil moisture and improvements to the soil water-modelling component have been found to improve predictions of  $N_2O$  emission (Saggar et al., 2007a; Norman et al., 2008). Therefore, there would be little point in improving the denitrification process in a model if the poor performance of some other component was the limiting factor. In grazed grasslands the spatial variability of  $N_2O$  emissions (due to the non-uniform nature of excretal N inputs) is an additional challenge to model validation (Giltrap et al., 2011).

The denitrification model of Leffelaar and Wessel (1988) (based on the growth, maintenance and death rates of denitrifier populations) was tested against laboratory incubation experiments where the model produced reasonable simulations of the progression  $NO_3^- \rightarrow NO_2^- \rightarrow N_2O \rightarrow N_2$ . The availability of appropriate datasets to fully parameterise the model was noted as a key limitation. However, most models are tested under field conditions. This means that the denitrification component itself is not explicitly tested, but rather the combined effect of all the model processes is tested. Models are frequently validated against  $N_2O$  measurements. This is not necessarily a good test of the denitrification model as  $N_2O$  is a product of both nitrification and denitrification, while denitrification also produces  $N_2$ . So it is possible for a model to predict the  $N_2O$  emissions correctly while incorrectly simulating the overall denitrification rate. Frolking et al. (1998) found in a comparison of four different process-based models (DNDC, DayCent, CASA, and Expert-N) that even when the models produced similar  $N_2O$  fluxes they often produced very different estimates of gaseous losses as  $NO$ ,  $N_2$  and  $NH_3$  indicating differences in the simulated N-transformation processes. Therefore, in order to have confidence that a model is simulating denitrification well it is necessary to ensure that other processes such as  $NO_3^-$ -leaching,  $NH_3$  volatilisation, and plant uptake are also well simulated. Even though it is not usually feasible to measure all the possible forms of N (e.g.  $N_2O$ ,  $N_2$ ,  $NH_3$ , soil  $NO_3^-$  and  $NH_4^+$ )

**Table 6**  
Approach used to account for microbial activity for a number of denitrification models.

Model name	Indicator of microbial activity	Reference
DayCent	$CO_2$ concentration	Stehfest and Müller (2004)
DNDC	Microbial biomass	Li et al. (1992)
FASSET	Mineralisation rate	Chatskikh et al. (2005)
InfoCrop	Microbial biomass	Aggarwal et al. (2006)
NEMIS/NOE	Site-specific constant	Hénault and Germon (2000), Hénault et al. (2005)
NGAS	Respiration rate	Parton et al. (1996)
PaSim	Decomposition rate	Schmid et al. (2001)
WNMM	Soil organic carbon	Li et al. (2007)

in a given experiment, the more that are measured the more valuable the data set will be for model validation.

### 5.3. Future requirements

All process-based models contain empirical approximations at some level, either from limitations of understanding the underlying processes or from practical difficulties in measuring all the required parameters. The different approaches to denitrification modelling discussed above differ in the point at which empirical approximations begin. Simplified process models can often give good results for situations for which they have been well-parameterised (e.g., Hénault et al., 2005) and may be more suitable for some applications. However, these simplifications limit the model's ability to only providing insight into the simplified processes. For example, a model that uses a fixed  $N_2O:N_2$  ratio for denitrification products is not capable of determining situations where the  $N_2O:N_2$  ratio changes.

Improved understanding of the underlying processes can lead to improvements in the models. However, as  $N_2O$  production is the result of many interacting processes, improving the simulated denitrification process alone will not necessarily improve the model's performance if other processes are poorly simulated. Another limitation to improving models is the availability of appropriate parameter values.

Many models are validated at field scale simply by comparing measured  $N_2O$  emissions with the modelled results. However, this does not indicate which processes are well modelled and where the model is performing poorly. Further modelling work that examines more closely the performance of the different component processes leading to changes in  $N_2O:N_2$  ratio is needed to improve model predictability.

## 6. Management practices to control denitrification and $N_2O$ and $N_2$ emissions

From an agricultural perspective denitrification results in the loss of a valuable plant nutrient and should therefore be minimised. From an environmental perspective denitrification presents both a threat and an opportunity. Where denitrification uses  $NO_3^-$  that would otherwise be leached into rivers or lakes this may be an environmental benefit. If, however, denitrification results in emission of  $N_2O$ , this poses an environmental threat. Management of agricultural systems should therefore minimise the accumulation of  $NO_3^-$  in soil and thereby restrict both leaching and denitrification. But where denitrification is inevitable, or even desirable to protect water quality, management should ensure that as much of the denitrified  $NO_3^-$  as possible is emitted as  $N_2$  rather than  $N_2O$ .

A range of on-farm management options to control denitrification by either emitting less  $N_2O$  or by shifting the balance between harmful  $N_2O$  and the non-greenhouse gas  $N_2$  in grazed pastures are summarised in Table 7. These management options include increasing C inputs, minimising N inputs from animal excreta and chemical fertilisers,

reducing animal feed crude protein, decreasing soil  $NO_3^-$  concentrations, and improving soil aeration and soil pH.

### 6.1. Soil C availability

Soil C availability has been shown to regulate denitrification rate in grassland soils (Section 3). Shepherd et al. (2010) found that sucrose addition immobilised urine N and decreased N leaching by up to 66%. This mitigation option could also be used to potentially reduce denitrification.

When  $NO_3^-$  N has leached below the root zone of pastures, there may still be opportunities to remove it by denitrification. Two options are wetlands and denitrification walls where C, such as in sawdust, is added in a trench (e.g., Schipper and Vojvodic-Vukovic, 1998), both having the effect of intercepting the hydrological flow of drainage water to surface waterways. Both options have shown potential to remove  $NO_3^-$  by denitrification. However, they are highly dependent on the fit to the landscape and the ability to intercept a significant component of the drainage water flows. Increasing soil C through effluent irrigation and (in future) biochar application may also be a management option to lower the  $N_2O:N_2$  ratio. However, not all the biochar created is equal and contradictory findings on the effect of biochar in reducing  $N_2O$  emissions or affecting  $N_2O:N_2$  ratios from denitrification are limiting its use (Clough et al., 2010; Saggar et al., 2011b). Aspects of biochar on soil denitrification community, enzyme activity and  $N_2O:N_2$  ratios need to be investigated.

### 6.2. N availability

The simplest approach for controlling denitrification in managed pasture systems is to reduce the  $NO_3^-$  concentration in the soil. This both reduces  $N_2O$  emissions and shifts the balance between  $N_2O$  and  $N_2$ . As discussed in Section 2.1, the major sources of  $NO_3^-$  in grazed pasture system are animal excreta, effluent, and chemical fertilisers. Soil generally remains wet during late-autumn/winter and early spring in the temperate region due to excessive precipitation, and during this time soil compaction by animal treading, low pasture N uptake, and low N immobilisation as a result of low soil temperature are likely to accelerate denitrification (Saggar et al., 2004b; de Klein et al., 2006; Hyde et al., 2006; Bhandral et al., 2007b;). The cut and carry systems that avoid direct excretal N input of grazing animals tend to be more N efficient than the grazed pasture systems (Ball and Ryden, 1984; Grignani and Laidlaw, 2002) during this high risk period of N losses. Minimising N inputs to the soil during these critical months can therefore potentially reduce N losses by denitrification. This can be achieved by:

- minimising the time animals spend grazing in the field (Ledgard et al., 2006; Luo et al., 2006) during the critical period. This practice enables farmers to have a better control over animal excreta which are collected in feed pads or animal houses and can be later

**Table 7**

Summary of management practices to control denitrification in temperate grassland.

Control factor	Goal	Impact	Management practices
Carbon availability	Increase	Lower $N_2O:N_2$	Effluent irrigation and biochar application
Nitrogen availability	Decrease	Lower denitrification and $N_2O:N_2$	Reducing excreta deposition during late-autumn/winter and early spring (use of stand-off/feed pads or animal houses) Reducing the concentration and amount of N in urine (increase urine volume and low protein feed) Increasing hippuric acid concentration by manipulating animals diet Nitrification and urease inhibitors Avoiding fertilisation of urine or dung patches
Oxygen supply	Increase	Lower denitrification	Preventing soil compaction Improving drainage Avoiding excessive amount of irrigation
Soil pH	Increase	Lower $N_2O:N_2$	Liming
Microbial enzymes	Increase	Increased $N_2O$ reduction	Supply of Cu as a cofactor for release of <i>NosZ</i>



applied to pastures when the risk of N losses is minimal (van der Meer, 2008). In several farmlet studies in New Zealand,  $\text{N}_2\text{O}$  emissions were reported to be reduced by up to 60% when animals were on stand-off/feed pads or in animal houses for 3–4 months during late-autumn/winter compared with year-round grazing (de Klein et al., 2006; Ledgard et al., 2006; Luo et al., 2008).

- adding salts to animal feed (to increase urine volume and spread) and feeding animals with low protein feed such as maize silage when on stand-off/feed pads or in animal houses, can be effective management tools for reducing the concentration and amount of N in urine (Mulligan et al., 2004; Ledgard et al., 2007). However, increasing urine volume through salt addition can only decrease N concentration in urine but may not change the total N in urine. Reduction of pasture N content through the use of controlled N release fertiliser or decreasing fertiliser application rate to match pasture N requirements can also be an effective option to reduce N contribution in urine thereby reducing denitrification rate and potentially  $\text{N}_2\text{O}$ : $\text{N}_2$  product ratios from denitrification (Luo et al., 2010).
- increasing hippuric acid concentration in urine by manipulating animals' diet may also reduce denitrification and  $\text{N}_2\text{O}$  emission (Kool et al., 2006; Bertram et al., 2009). However, a New Zealand field study (Clough et al., 2009) did not observe any effect of the increased hippuric or benzoic acid concentrations in cow urine on  $\text{N}_2\text{O}$  emissions.
- applying nitrification and/or urease inhibitors to retain N in the  $\text{NH}_4^+$  form and/or reduce the hydrolysis of urea-N (Di et al., 2007; Saggar et al., 2008; Majumdar, 2009; Zaman et al., 2009; Parkin and Hatfield, 2010; Zaman and Blennerhassett, 2010). The inhibitors, by controlling hydrolysis and nitrification processes, can reduce the amount of  $\text{NO}_3^-$  available for denitrification and may potentially reduce the  $\text{N}_2\text{O}$ : $\text{N}_2$  product ratio. Sun et al. (2008) found that the use of NI, dicyandiamide (DCD), reduced denitrification N losses by 22% in a grazed pasture soil.
- including deep rooted pasture species may reduce the amount of  $\text{NO}_3^-$  in the subsurface soil horizons where there is a greater potential for denitrification. Dear et al. (2009) found that incorporating perennial pasture species in swards reduced  $\text{NO}_3^-$  leaching and prevented a decline in surface soil pH.

### 6.3. Plant composition

Plants with specific constituents, such as tannins or saponins, can increase N utilisation by animals and increase N excretion in the low-risk dung N forms relative to urine N (e.g., Miller et al., 2001). Feeding animals with grass silage containing elevated concentrations of water-soluble carbohydrates increased the N use efficiency for microbial growth in the rumen from 46 to 68% (Merry et al., 2003). Similarly, Miller et al. (2001) found that dairy cows on a “high sugar” variety of perennial ryegrass excreted 18% less N in total and 29% less urine N. Thus, manipulation of plant composition offers a potential to reduce N excretion in the urine, thereby reducing subsequent denitrification and  $\text{N}_2\text{O}$  losses. Plants also have the potential to alter soil N cycling via altering the quality of their residues or releasing compounds which have nitrification inhibition characteristics. Ledgard et al. (1998) measured a 10-fold difference in gross N immobilisation rate between non-N-fertilised grassland dominated by *Agrostis* and *Holcus* spp. compared with ryegrass with a regular N fertiliser history. Such differences in immobilisation potential may be important controllers of denitrification and  $\text{N}_2\text{O}$  losses from N sources such as animal urine, N fertilisers, and effluent/manure.

### 6.4. Oxygen supply

As discussed earlier, denitrification becomes a major source of  $\text{N}_2\text{O}$  and  $\text{N}_2$  production at low  $\text{O}_2$  partial pressure (<0.5 vol.%) and high WFPS (>60%) (Section 3.1.3). Improving soil drainage conditions,

avoiding soil compaction and pugging in wet soil conditions, and avoiding over-irrigation can all reduce the amount of denitrification N losses. In addition, biochar application to soils could reduce the amount of denitrification N losses by decreasing bulk density and improving aeration status of the soil (Lehmann and Joseph, 2009; Sohi et al., 2010). Biochar application to soils also increases soil organic matter contents which could subsequently reduce the product ratio of  $\text{N}_2\text{O}$  and  $\text{N}_2$  through denitrification.

### 6.5. Soil pH

Liming has also been mooted as a mitigation option for lowering soil  $\text{N}_2\text{O}$  emissions. For example, Clough et al. (2004) showed that while liming is likely to decrease  $\text{N}_2\text{O}$  emission from urine patches where soils are at field capacity, the increased  $\text{NO}_3^-$  concentration from lime-induced nitrification would be susceptible to denitrification if the soil moisture subsequently exceeds the field capacity. However, Zaman and Nguyen (2010) recently noticed that lime had no effect on  $\text{N}_2\text{O}$  emission in either urine- or  $\text{KNO}_3$ -treated soils, but increased  $\text{N}_2$  emissions and lowered  $\text{N}_2\text{O}$ : $\text{N}_2$  ratios in a short-term field experiment. In soils without an added C source, transcription of denitrification genes was below detectable levels but the addition of glutamate showed sharp peaks in the transcription of denitrifying genes with the transcription of *nosZ* exceeding that of *nirS* at lower pH. This led Liu et al. (2010) to conclude that the increased  $\text{N}_2\text{O}$ : $\text{N}_2$  product ratio at low pH reflects a post transcriptional phenomenon, whereby *Nos* assembly is more sensitive to low pH relative to other denitrification enzymes. As discussed, adding biochar to soil generally raises the pH of acid soils and may consequently reduce  $\text{N}_2\text{O}$  emissions. Technologies for applying biochar to perennial grasslands need to be developed to ensure that the biochar is fully mixed with the surface soil for maximising effects on  $\text{N}_2\text{O}$  reduction. Application of biochar during the cultivation of perennial grasslands for pasture renewal could be an approach.

### 6.6. Microbial enzymes

Denitrification engages several proteins that require metal ions as a cofactor (Ferguson, 1998). For example, addition of  $\text{Mo}^{6+}$  and  $\text{Fe}^{3+}$  usually affects the growth and denitrification rates of microbial communities by accelerated  $\text{NO}_3^-$  utilisation and  $\text{NO}_2^-$  accumulation (Labbe et al., 2003; Zhou et al., 2007; Pintathong et al., 2009). The accumulation of  $\text{NO}_2^-$  results from the higher increase in the membrane-bound *Nar* activity and consumption of  $\text{NO}_3^-$  compared with the smaller increase in periplasmic *Nir* activity and utilisation of  $\text{NO}_2^-$ . *Nos* is the only enzyme with no alternative non-copper type available to carry out reduction of  $\text{N}_2\text{O}$  in the absence of Cu. Application of supplements/fertilisers rich in Cu will be helpful to boost the activity of *NosZ* and thus converting more  $\text{N}_2\text{O}$  to  $\text{N}_2$  (Richardson et al., 2009).

## 7. Conclusions and future research needs

Denitrification is the primary source of  $\text{N}_2\text{O}$  emissions and a key process in temperate pastoral soils that could annually denitrify 5.6 Tg of reactive N. The need both for removal of excess  $\text{NO}_3^-$  from soil for protection of ground and surface waters, and for reductions in  $\text{N}_2\text{O}$  emissions produced during denitrification to mitigate global warming and ozone depletion, is now urgent. The last three decades have witnessed a resurgence of interest in denitrification in soils. Molecular techniques involving PCR-based amplification of DNA extracted directly from soil have advanced our understanding of the composition and abundance of the microbial community involved in the denitrification process. However, the DNA only indicates the presence of the denitrifier from which it was extracted, and not its quantitative functional contribution towards denitrification. Therefore, the extrapolation of DNA gene copy abundance to nitrification and



denitrification rates must be confirmed by next generation sequencing (NGS) technologies. As understanding of the denitrification process improves, this knowledge can be used to develop effective mitigation technologies.

It is clear from research on ecology of denitrification that while the ability to denitrify is widely distributed among a taxonomic and phylogenetically diverse group of microorganisms, not all can reduce  $\text{N}_2\text{O}$  to  $\text{N}_2$ . Therefore, the amount of  $\text{N}_2\text{O}$  produced can vary from almost 0 to over 90% of the total N-gases produced by denitrification. The research suggests that a significant number ( $\sim 1/3$ ) of denitrifying organisms do not have the capacity (*nosZ*) to reduce  $\text{N}_2\text{O}$ . This may provide an opportunity to manipulate soil denitrifying organisms to enhance *nosZ* activity. It could be done through altering the interaction between mineral N ( $\text{NH}_4^+$  and  $\text{NO}_3^-$ ) or Cu supply, organic C, soil aeration and pH to stimulate the activity of Nos, which will help result in emissions of more  $\text{N}_2$  than  $\text{N}_2\text{O}$ .

Simultaneous information on soil and environmental parameters and bacterial processes is needed to identify and assess the mechanisms regulating the production and reduction of  $\text{N}_2\text{O}$ . The multiplicity of soil and environmental conditions that could influence the ability of denitrifiers to complete denitrification and result in reducing  $\text{N}_2\text{O}$  emissions may explain why to date no clear picture has emerged as to the efficacy of soil management practices in reducing  $\text{N}_2\text{O}$  emissions. Understanding those mechanisms and the denitrifiers that have the potential to reduce the production of  $\text{N}_2\text{O}$  during denitrification is critical to the development of novel and effective  $\text{N}_2\text{O}$  mitigation strategies. Thus the key research questions are:

- What is the relationship between soil, microbial and environmental parameters and processes and  $\text{N}_2\text{O}/\text{N}_2$  ratio of denitrification?
- Is the number of high capacity denitrifiers (*nosZ*) in soils sufficient to reduce the  $\text{N}_2\text{O}/\text{N}_2$  during denitrification?
- How can the denitrifier community be manipulated and the *nosZ* enhanced?
- What is the effectiveness of soil manipulations for reducing  $\text{N}_2\text{O}$  production during denitrification?
- What are the optimum soil and environmental conditions required for maximum  $\text{N}_2\text{O}$  mitigation using the most effective soil amendment?

There are substantial knowledge gaps that require both laboratory and field research to determine the best soil type and optimal environmental conditions to encourage the activity of Nos during denitrification. Future research, therefore, needs to focus on evaluating the  $\text{N}_2\text{O}$ -reducing ability of the denitrifiers to accelerate the conversion of  $\text{N}_2\text{O}$  to  $\text{N}_2$  and the reduction of  $\text{N}_2\text{O}:\text{N}_2$  ratio during denitrification. More research is also required to measure  $\text{N}_2\text{O}$  and  $\text{N}_2$  emissions at field scale and quantitatively determine the emissions under different land-use and land management systems and to improve model approaches to quantify the  $\text{N}_2\text{O}:\text{N}_2$  ratio and  $\text{N}_2$  emissions from denitrification with greater certainty.

## Acknowledgements

This work was supported by funding from the New Zealand Agricultural Greenhouse Gas Research Centre (NZAGRC) and the New Zealand Ministry of Science and Innovation (MSI, now Ministry of Business, Innovation and Employment, MBIE). We are grateful to Drs Kevin Tate and Mehmet Senbayram for providing very helpful suggestions and thorough review of this article, and to Anne Austin for internal editing.

## References

Acinas SG, Sarma-Rupavtarm R, Klepac-Ceraj V, Polz MF. PCR-induced sequence artifacts and bias: insights from comparison of two 16S rRNA clone libraries constructed from the same sample. *Appl Environ Microbiol* 2005;71:8966–9.

Adu JK, Oades JM. Physical factors influencing decomposition of organic materials in soil aggregates. *Soil Biol Biochem* 1978;10:109–15.

Aggarwal PK, Banerjee B, Daryaei MG, Bhatia A, Bala A, Rani S, et al. InfoCrop: a dynamic simulation model for the assessment of crop yields, losses due to pests, and environmental impact of agro-ecosystems in tropical environments. II. Performance of the model. *Agric Syst* 2006;89:47–67.

Akunna JC, Bizeau C, Moletta R. Nitrate and nitrite reductions with anaerobic sludge using various carbon sources: glucose, glycerol, acetic acid, lactic acid and methanol. *Water Res* 1993;27:1303–12.

Arah JRM, Smith KA. Steady-state denitrification in aggregated soils: a mathematical model. *J Soil Sci* 1989;40:139–49.

Arah JRM, Smith KA. Factors influencing the fraction of the gaseous products of soil denitrification evolved to the atmosphere as nitrous oxide. In: Bouwman AF, editor. *Soils and the greenhouse effect: the present status and future trends concerning the effect of soils and their cover on the fluxes of greenhouse gas*. John Wiley & Sons; 1990. p. 475–80.

Aulakh MS, Doran JW, Mosier AR. Field evaluation of four methods for measuring denitrification. *Soil Sci Soc Am J* 1991;55:1332–8.

Aulakh MS, Khera TS, Doran JW, Bronson KF. Denitrification,  $\text{N}_2\text{O}$  and  $\text{CO}_2$  fluxes in rice-wheat cropping system as affected by crop residues, fertilizer N and legume green manure. *Biol Fertil Soils* 2001;34:375–89.

Avalakki UK, Strong WM, Saffigna PG. Measurement of gaseous emissions from denitrification of applied nitrogen-15. II Effects of temperature and added straw. *Aust J Soil Sci* 1995;33:89–99.

Babic KH, Schauss K, Ha iB, Sikora S, Redžepovic S, Radl V, et al. Influence of different *Sinorhizobium meliloti* inocula on abundance of genes involved in nitrogen transformations in the rhizosphere of alfalfa (*Medicago sativa* L.). *Environ Microbiol* 2008;10:2922–30.

Bailey LD. Effects of temperature and root on denitrification in a soil. *Can J Soil Sci* 1976;56:79–87.

Bakken LR, Berguast L, Liu B, Frostegå A. Regulation of denitrification at the cellular level: a clue to the understanding of  $\text{N}_2\text{O}$  emissions from soils. *Philos Trans R Soc Biol Sci* 2012;367:1226–34.

Balderston WL, Sherr B, Payne WJ. Blockage by acetylene of nitrous oxide reduction in *Pseudomonas perfectomarinus*. *Appl Environ Microbiol* 1976;31:504–8.

Ball P, Ryden J. Nitrogen relationships in intensively managed temperate grasslands. *Plant Soil* 1984;76:23–33.

Barton L, Schipper LA. Regulation of nitrous oxide emissions from soils irrigated with dairy farm effluent. *J Environ Qual* 2001;30:1881–7.

Barton L, McLay CDA, Schipper LA, Smith CT. Annual denitrification rates in agricultural and forest soils: a review. *Aust J Soil Res* 1999;37:1073–93.

Baumann B, Snozzi M, Zehnder AJB, Van Meer JRD. Dynamics of denitrification activity of *Paracoccus denitrificans* in continuous culture during aerobic-anaerobic changes. *J Bacteriol* 1996;178:4367–74.

Beauchamp EG, Gale C, Yeomans JC. Organic matter availability for denitrification in soils of different textures and drainage classes. *Commun Soil Sci Plant Anal* 1980;11:1221–33.

Beauchamp EG, Trevors JT, Paul JW. Carbon sources for bacterial denitrification. *Adv Soil Sci* 1989;10:113–42.

Beck H, Christensen S. The effect of grass maturing and root decay on  $\text{N}_2\text{O}$  production in soil. *Plant Soil* 1987;103:269–73.

Bertram JE, Clough TJ, Sherlock RR, Condron LM, O'Callaghan M, Wells NS, et al. Hippuric acid and benzoic acid inhibition of urine derived  $\text{N}_2\text{O}$  emissions from soil. *Glob Chang Biol* 2009;15:2067–77.

Bhandral R, Bolan NS, Saggar S, Hedley MJ. Nitrogen transformations and nitrous oxide emissions from various types of farm effluents. *Nutr Cycl Agroecosyst* 2007a;79:193–208.

Bhandral R, Saggar S, Bolan NS, Hedley MJ. Transformation of nitrous oxide emission from grassland soils as affected by compaction. *Soil Tillage Res* 2007b;94:482–92.

Bilotta GS, Brazier RE, Haygarth PM. The impacts of grazing animals on the quality of soils, vegetation, and surface waters in intensively managed grasslands. *Adv Agron* 2007;94:237–80.

Blackmer AM, Bremner JM. Inhibitory effect of nitrate on reduction of  $\text{N}_2\text{O}$  to  $\text{N}_2$  by soil microorganisms. *Soil Biol Biochem* 1978;10:187–91.

Bolan NS, Saggar S, Luo JF, Bhandral R, Singh J. Gaseous emissions of nitrogen from grazed pastures: processes, measurements and modelling, environmental implications, and mitigation. *Adv Agron* 2004;84:37–120.

Bollmann A, Conrad R. Acetylene blockage technique leads to underestimation of denitrification rates in oxic soils due to scavenging of intermediate nitric oxide. *Soil Biol Biochem* 1997;29:1067–77.

Boyer EW, Alexander RB, Parton WJ, Li C, Butterbach-Bahl K, Donner SD, et al. Modeling denitrification in terrestrial and aquatic ecosystems at regional scales. *Ecol Appl* 2006;16:2123–42.

Braker G, Tiedje JM. Nitric oxide reductase (*norB*) genes from pure cultures and environmental samples. *Appl Environ Microbiol* 2003;69:3476–83.

Braker G, Zhou J, Wu L, Devol AH, Tiedje JM. Nitrite reductase genes (*nirK* and *nirS*) as functional markers to investigate diversity of denitrifying bacteria in Pacific northwest marine sediment communities. *Appl Environ Microbiol* 2000;66:2096–104.

Braker G, Ayala-Del-Rio HL, Devol AH, Fesefeldt A, Tiedje JM. Community structure of denitrifiers, bacteria, and archaea along redox gradients in Pacific northwest marine sediments by terminal restriction fragment length polymorphism analysis of amplified nitrite reductase (*nirS*) and 16S rRNA Genes. *Appl Environ Microbiol* 2001;67:1893–901.

Bremer C, Braker G, Matthies D, Reuter A, Engels C, Conrad R. Impact of plant functional group, plant species, and sampling time on the composition of *nirK*-type denitrifier communities in soil. *Appl Environ Microbiol* 2007;73:6876–84.

Bremner JM, Blackmer AM. Nitrous oxide: emission from soils during nitrification of fertilizer nitrogen. *Science* 1978;199:295–6.

- Bremner JM, Shaw K. Denitrification in soil. II. Factors affecting denitrification. *J Agric Sci* 1958;51:39–52.
- Bru D, Sarr A, Philippot L. Relative abundances of proteobacterial membrane-bound and periplasmic nitrate reductases in selected environments. *Appl Environ Microbiol* 2007;73:5971–4.
- Bryan BA. Physiology and biochemistry of denitrification. In: Delwiche CC, editor. Denitrification, nitrification, and atmospheric nitrous oxide. New York: John Wiley & Sons; 1981. p. 67–84.
- Brye KR, Norman JM, Bundy LG, Gower ST. Nitrogen and carbon leaching in agroecosystems and their role in denitrification potential. *J Environ Qual* 2001;30:58–70.
- Burford JR, Bremner JM. Relationships between the denitrification capacities of soils and total water-soluble and readily decomposable soil organic matter. *Soil Biol Biochem* 1975;7:389–94.
- Burger M, Matiassek M. New approaches to distinguish between nitrification and denitrification as sources of nitrous oxide. In: Sheldou AI, Barnhart EP, editors. Nitrous oxide emissions research progress. New York: Nova Science Publishers; 2009. p. 167–78.
- Burton DL, Beauchamp EG. Profile nitrous oxide and carbon dioxide concentrations in a soil subject to freezing. *Soil Sci Soc Am J* 1994;58:115–22.
- Butland G, Spiro S, Watmough NJ, Richardson DJ. Two conserved glutamates in the bacterial nitric oxide reductase are essential for activity but not assembly of the enzyme. *J Bacteriol* 2001;183:189–99.
- Butterbach-Bahl K, Willibald G, Papen H. Soil core method for direct simultaneous determination of  $N_2$  and  $N_2O$  emissions from forest soils. *Plant Soil* 2002;240:105–16.
- Cai G, Chen D, White RE, Fan XH, Pacholski A, Zhu ZL, et al. Gaseous nitrogen losses from urea applied to maize on a calcareous fluvo-aquic soil in the North China Plain. *Aust J Soil Res* 2002;40:737–48.
- Cardenas LM, Hawkins MB, Chadwick D, Scholefield D. Biogenic gas emissions from soils measured using a new automated laboratory incubation system. *Soil Biol Biochem* 2003;35:867–70.
- Cardon ZG, Gage DJ. Resource exchange in the rhizosphere: molecular tools and the microbial perspective. *Annu Rev Ecol Syst* 2006;37:459–88.
- Carter MS. Contribution of nitrification and denitrification to  $N_2O$  emissions from urine patches. *Soil Biol Biochem* 2007;39:2091–102.
- Casciotti KL, Sigman DM, Hastings MG, Böhlke JK, Hilkert A. Measurement of the oxygen isotopic composition of nitrate in seawater and freshwater using the denitrifier method. *Anal Chem* 2002;74:4905–12.
- Casella S, Leporini C, Nuti MP. Nitrous oxide production by nitrogen-fixing fast growing rhizobia. *Microb Ecol* 1984;10:107–14.
- Cavigelli MA, Robertson GP. The functional significance of denitrifier community composition in a terrestrial ecosystem. *Ecology* 2000;81:1402–14.
- Cayuela ML, Oenema O, Kuikman PJ, Bakker RR, Van Groenigen JW. Bioenergy by-products as soil amendments? Implications for carbon sequestration and greenhouse gas emissions. *GCB Bioenergy* 2010;2:201–13.
- Chadwick DR, Pain BF, Brookman SKE. Nitrous oxide and methane emissions following application of animal manures to grassland. *J Environ Qual* 2000;29:277–87.
- Chatskikh D, Olesen J, Berntsen J, Regina K, Yamulki S. Simulation of effects of soils, climate and management on  $N_2O$  emission from grasslands. *Biogeochemistry* 2005;76:395–419.
- Chen Z, Liu J, Wu M, Xie X, Wu J, Wei W. Differentiated response of denitrifying communities to fertilization regime in paddy soil. *Microb Ecol* 2011;1:1–14.
- Chèneby D, Hartmann A, Hénault C, Topp E, Germon JC. Diversity of denitrifying microflora and ability to reduce  $N_2O$  in two soils. *Biol Fertil Soils* 1998;28:19–26.
- Chèneby D, Hallet S, Mondon M, Martin-Laurent F, Germon JC, Philippot L. Genetic characterization of the nitrate reducing community based on narG nucleotide sequence analysis. *Microb Ecol* 2003;46:113–21.
- Chèneby D, Perez S, Devroe C, Hallet S, Couton Y, Bizouard F, et al. Denitrifying bacteria in bulk and maize-rhizospheric soil: diversity and  $N_2O$ -reducing abilities. *Can J Microbiol* 2004;50:469–74.
- Cho CM, Burton DL, Chang C. Denitrification and fluxes of nitrogenous gases from soil under steady oxygen distribution. *Can J Soil Sci* 1997;77:261–9.
- Christensen S. Denitrification in an acid soil: effects of slurry and potassium nitrate on the evolution of nitrous oxide and on nitrate-reducing bacteria. *Soil Biol Biochem* 1985;17:757–64.
- Clough TJ, Sherlock RR, Cameron KC, Ledgard SF. Fate of urine nitrogen on mineral and peat soils in New Zealand. *Plant Soil* 1996;178:141–52.
- Clough TJ, Kelliher FM, Sherlock RR, Ford CD. Lime and soil moisture effects on nitrous oxide emissions from a urine patch. *Soil Sci Soc Am J* 2004;68:1600–9.
- Clough TJ, Ray JL, Buckthought LE, Calder J, Baird D, O'Callaghan M, et al. The mitigation potential of hippuric acid on  $N_2O$  emissions from urine patches: an in situ determination of its effect. *Soil Biol Biochem* 2009;41:2222–9.
- Clough TJ, Bertram JE, Ray J, Condron LM, O'Callaghan M, Sherlock RR, et al. Unweathered wood biochar impact on nitrous oxide emissions from a bovine-urine-amended pasture soil. *Soil Sci Soc Am J* 2010;74:852–60.
- Colbourn P, Dowdell RJ. Denitrification in field soils. *Plant Soil* 1984;76:213–26.
- Colbourn P, Harper IW, Iqbal MM. Denitrification losses from  $^{15}N$ -labelled calcium nitrate fertilizer in a clay soil in the field. *J Soil Sci* 1984;35:539–47.
- Corré WJ, Dijkman W, Kieskamp WM. Denitrification in the top soil of production grassland. *Mitt Dtsch Bodenkd Ges* 1990;60:183–8.
- Cramm R, Siddiqui RA, Friedrich B. Two iso-functional nitric oxide reductases in *Alcaligenes eutrophus* H16. *J Bacteriol* 1997;179:6769–77.
- Craswell ET. Some factors influencing denitrification and nitrogen immobilisation in a clay soil. *Soil Biol Biochem* 1978;10:241–5.
- Čuhel J, Šimek M. Proximal and distal control by pH of denitrification rate in a pasture soil. *Agric Ecosyst Environ* 2011;141:230–3.
- Čuhel J, Šimek M, Laughlin RJ, Bru D, Chèneby D, Watson CJ, et al. Insights into the effect of soil pH on  $N_2O$  and  $N_2$  emissions and denitrifier community size and activity. *Appl Environ Microbiol* 2010;76:1870–8.
- Dalal RC, Wang W, Robertson GP, Parton WJ. Nitrous oxide emission from Australian agricultural lands and mitigation options: a review. *Aust J Soil Res* 2003;41:165–95.
- Dalsgaard T, Bak F. Effect of acetylene on nitrous oxide reduction and sulfide oxidation in batch and gradient cultures of *Thiobacillus denitrificans*. *Appl Environ Microbiol* 1992;58:1601–8.
- Dandie CE, Burton DL, Zebarth BJ, Henderson SL, Trevors JT, Goyer C. Changes in bacterial denitrifier community abundance over time in an agricultural field and their relationship with denitrification activity. *Appl Environ Microbiol* 2008;74:5997–6005.
- Dandie CE, Wertz S, Leclair CL, Goyer C, Burton DL, Patten CL, et al. Abundance, diversity and functional gene expression of denitrifier communities in adjacent riparian and agricultural zones. *FEMS Microbiol Ecol* 2011;77:69–82.
- Davidson EA. Sources of nitric oxide and nitrous oxide following wetting of dry soil. *Soil Sci Soc Am J* 1992;56:95–102.
- Davidson EA, Seitzinger S. The enigma of progress in denitrification research. *Ecol Appl* 2006;16:2057–63.
- Davidson EA, Matson PA, Vitousek PM, Riley R, Dunkin K, Garcia-Mendez G, et al. Processes regulating soil emissions of NO and  $N_2O$  in a seasonally dry tropical forest. *Ecology* 1993;74:130–9.
- de Boer APN, Van Der Oost J, Reijnders WNM, Westerhoff HV, Stouthamer AH, Van Spanning RJM. Mutational analysis of the nor gene cluster which encodes nitric-oxide reductase from *Paracoccus denitrificans*. *Eur J Biochem* 1996;242:592–600.
- de Bruijn AMG, Butterbach-Bahl K, Blagodatsky S, Grote R. Model evaluation of different mechanisms driving freeze–thaw  $N_2O$  emissions. *Agric Ecosyst Environ* 2009;133:196–207.
- de Klein CAM, Eckard RJ. Targeted technologies for nitrous oxide abatement from animal agriculture. *Aust J Exp Agric* 2008;48:14–20.
- de Klein CAM, Van Logtestijn RSP. Denitrification in the top soil of managed grasslands in The Netherlands in relation to soil type and fertilizer level. *Plant Soil* 1994;163:33–44.
- de Klein CAM, Van Logtestijn RSP. Denitrification in grassland soils in The Netherlands in relation to irrigation, N-application rate, soil water content and soil temperature. *Soil Biol Biochem* 1996;28:231–7.
- de Klein CAM, Sherlock RR, Cameron KC, van der Weerden TJ. Nitrous oxide emissions from agricultural soils in New Zealand — a review of current knowledge and directions for future research. *J R Soc N Z* 2001;31:543–74.
- de Klein CAM, Smith LC, Monaghan RM. Restricted autumn grazing to reduce nitrous oxide emissions from dairy pastures in Southland, New Zealand. *Agric Ecosyst Environ* 2006;112:192–9.
- de Klein CAM, Eckard RJ, van der Weerden TJ. Nitrous oxide emissions from the nitrogen cycle in livestock agriculture: estimation and mitigation. In: Smith K, editor. Nitrous oxide and climate change. London, UK: Earthscan Ltd.; 2010. p. 107–42.
- Dear BS, Virgona JM, Sandral GA, Swan AD, Morris S. Changes in soil mineral nitrogen, nitrogen leached, and surface pH under annual and perennial pasture species. *Crop Pasture Sci* 2009;60:975–86.
- Delwiche CC. The nitrogen cycle and nitrous oxide. In: Delwiche CC, editor. Denitrification, nitrification and atmospheric nitrous oxide. New York (US): Wiley-Interscience; 1981. p. 1–15.
- Demanèche S, Philippot L, David MM, Navarro E, Vogel TM, Simonet P. Characterization of nitrification gene clusters of soil bacteria via a metagenomic approach. *Appl Environ Microbiol* 2009;75:534–7.
- Dendooven L, Anderson JM. Dynamics of reduction enzymes involved in the denitrification process in pasture soil. *Soil Biol Biochem* 1994;26:1501–6.
- Dendooven L, Splatt P, Pemberton E, Ellis S, Anderson JM. Controls over denitrification and its gaseous products in a permanent pasture soil. In: Jarvis SC, Pain BF, editors. Gaseous nitrogen emissions from grasslands. Wallingford: CAB International; 1997. p. 9–25.
- Dendooven L, Bonhomme E, Merckx R, Vlassak K. Injection of pig slurry and its effects on dynamics of nitrogen and carbon in a loamy soil under laboratory conditions. *Biol Fertil Soils* 1998;27:5–8.
- Di HJ, Cameron KC, Sherlock RR. Comparison of the effectiveness of a nitrification inhibitor, dicyandiamide, in reducing nitrous oxide emissions in four different soils under different climatic and management conditions. *Soil Use Manag* 2007;23:1–9.
- Dobbie KE, Smith KA. The effects of temperature, water-filled pore space and land use on  $N_2O$  emissions from an imperfectly drained gleysol. *Eur J Soil Sci* 2001;52:667–73.
- Dorland S, Beauchamp EG. Denitrification and ammonification at low soil temperatures. *Can J Soil Sci* 1991;71:293–303.
- Drewry JJ, Cameron KC, Buchan GD. Pasture yield and soil physical property responses to soil compaction from treading and grazing — a review. *Aust J Soil Res* 2008;46:237–56.
- Drury CF, Myrold DD, Beauchamp EG, Reynolds WD. Denitrification techniques for soils. In: Carter MR, Gregorich EG, editors. Soil sampling and methods of analysis. 2nd edition. Boca Raton, FL (USA): CRC Press Taylor & Francis; 2008. p. 471–93.
- Eckard RJ, Chen D, White RE, Chapman DF. Gaseous nitrogen loss from temperate perennial grass and clover dairy pastures in south-eastern Australia. *Aust J Agr Res* 2003;54:561–70.
- Edwards AC, Cresser MS. Freezing and its effect on chemical and biological properties of soil. *Adv Soil Sci* 1992;18:59–79.
- Egginton GM, Smith KA. Losses of nitrogen by denitrification from a grassland soil fertilised with cattle slurry and calcium nitrate. *J Soil Sci* 1986;37:69–80.

- Elliott PW, Knight D, Anderson JM. Variables controlling denitrification from earthworm casts and soil in permanent pastures. *Biol Fertil Soils* 1991;11:24–9.
- Ellis S, Yamulki S, Dixon E, Harrison R, Jarvis SC. Denitrification and N<sub>2</sub>O emissions from a UK pasture soil following the early spring application of cattle slurry and mineral fertiliser. *Plant Soil* 1998;202:15–25.
- Etchebehere C, Tiedje J. Presence of two different active nirS nitrite reductase genes in a denitrifying *Thauera* sp. from a high-nitrate-removal-rate reactor. *Appl Environ Microbiol* 2005;71:5642–5.
- Fageria NK, Baligar VC. Ameliorating soil acidity of tropical oxisols by liming for sustainable crop production. *Adv Agron* 2008;99:345–99.
- Ferguson SJ. Nitrogen cycle enzymology. *Curr Opin Chem Biol* 1998;2:182–93.
- Fernández N, Sierra-Alvarez R, Field JA, Amils R, Sanz JL. Microbial community dynamics in a chemolithotrophic denitrification reactor inoculated with methanogenic granular sludge. *Chemosphere* 2008;70:462–74.
- Fierer N, Schimel JP. Effects of drying–rewetting frequency on soil carbon and nitrogen transformations. *Soil Biol Biochem* 2002;34:777–87.
- Firestone MK. Biological denitrification. In: Stevenson FJ, editor. *Nitrogen in agricultural soils*. Madison, WI: American Society of Agronomy; 1982. p. 289–326.
- Firestone M, Smith M, Firestone R, Tiedje J. The influence of nitrate, nitrite, and oxygen on the composition of the gaseous products of denitrification in soil. *Soil Sci Soc Am J* 1979;43:1140–4.
- Flessa H, Dorsch P, Beese F. Seasonal-variation of N<sub>2</sub>O and CH<sub>4</sub> fluxes in differently managed arable soils in southern Germany. *J Geophys Res* 1995;100:23115–24.
- Focht DD. The effect of temperature, pH and aeration on the production of nitrous oxide and gaseous nitrogen — a zero order kinetic model. *Soil Sci* 1974;118:173–9.
- Frank DA, Groffman PM. Denitrification in a semi-arid grazing ecosystem. *Oecologia* 1998;117:564–9.
- Frolking SE, Mosier AR, Ojima DS, Li C, Parton WJ, Potter CS, et al. Comparison of N<sub>2</sub>O emissions from soils at three temperate agricultural sites: simulations of year-round measurements by four models. *Nutr Cycl Agroecosyst* 1998;52:77–105.
- Garcia-Montiel DC, Steudler PA, Piccolo M, Neill C, Melillo J, Cerri CC. Nitrogen oxide emissions following wetting of dry soils in forest and pastures in Rondônia, Brazil. *Biogeochemistry* 2003;64:319–36.
- Gift DM, Groffman PM, Kaushal GS, Mayer PM. Denitrification potential, root biomass, and organic matter in degraded and restored urban riparian zones. *Restor Ecol* 2010;18:113–20.
- Gillespie DE, Rondon MR, Williamson LL, Handelsman J. Metagenomic libraries from uncultured microorganisms. In: Osborn AM, Smith CJ, editors. *Molecular microbial ecology*. New York, NY: Taylor and Francis Group; 2005. p. 261–79.
- Giltrap DL, Saggar S, Singh J, Harvey M, McMillan A, Laubach J. Field-scale verification of nitrous oxide emission reduction with DCD in dairy-grazed pasture using measurements and modelling. *Soil Res* 2011;49:696–702.
- Goldberg S, Borken W, Gebauer G. N<sub>2</sub>O emission in a Norway spruce forest due to soil frost: concentration and isotope profiles shed a new light on an old story. *Biogeochemistry* 2010;97:21–30.
- Grant RF. A technique for estimating denitrification rates at different soil temperatures, water contents, and nitrate concentrations. *Soil Sci* 1991;152:41–52.
- Griffiths RI, Whiteley AS, O'Donnell AG, Bailey MJ. Rapid method for coextraction of DNA and RNA from natural environments for analysis of ribosomal DNA- and rRNA-based microbial community composition. *Appl Environ Microbiol* 2000;66:5488–91.
- Grignani C, Laidlaw AS. Nitrogen economy in grasslands and annual forage crops: control of environmental impact. In: Durand JL, Emile JC, Huyghe C, Lemaire G, editors. *Grassland Science in Europe*, vol. 7, multi-function grasslands — quality forages, animal products and landscapes. Proceedings of the 19th General Meeting of the European Grassland Federation. La Rochelle, France, 27–30 May 2002/Versailles, France: AEPF; 2002. p. 625–33.
- Groffman P. Ecology of nitrification and denitrification in soil evaluated at scales relevant to atmospheric chemistry. In: Rogers JE, Whitman WB, editors. *Microbial production and consumption of greenhouse gases: methane, nitrogen oxides and halomethanes*. Washington, DC (USA): American Society of Microbiology; 1991. p. 201–17.
- Groffman PM, Tiedje JM. Denitrification in north temperate forest soils: relationships between denitrifying and environmental factors at the landscape scale. *Soil Biol Biochem* 1989;21:621–6.
- Groffman PM, Holland E, Myrold DD, Robertson GP, Zou X. Denitrification. In: Robertson GP, Bledsoe CS, Coleman C, Sollins P, editors. *Standard soil methods for long term ecological research*. New York: Oxford University Press; 1999. p. 272–88.
- Groffman PM, Altabet MA, Böhlke JK, Butterbach-Bahl K, David MB, Firestone MK, et al. Methods for measuring denitrification: diverse approaches to a difficult problem. *Ecol Appl* 2006;16:2091–122.
- Grogan P, Michelsen A, Ambus P, Jonasson S. Freeze–thaw regime effects on carbon and nitrogen dynamics in sub-arctic heath tundra mesocosms. *Soil Biol Biochem* 2004;36:641–54.
- Grüntzig V, Nold SC, Zhou J, Tiedje JM. *Pseudomonas stutzeri* nitrite reductase gene abundance in environmental samples measured by real-time PCR. *Appl Environ Microbiol* 2001;67:760–8.
- Haine V, Dozet M, Dornand J, Letesson JJ, De Bolle X. NnrA is required for full virulence and regulates several *Brucella melitensis* denitrification genes. *J Bacteriol* 2006;188:1615–9.
- Hallin S, Jones CM, Schlöter M, Philippot L. Relationship between n-cycling communities and ecosystem functioning in a 50-year-old fertilization experiment. *ISME J* 2009;3:597–605.
- Handelsman J, Rondon MR, Brady SF, Clardy J, Goodman RM. Molecular biological access to the chemistry of unknown soil microbes: a new frontier for natural products. *Chem Biol* 1998;5:245–9.
- Hansen S, Maehlum JE, Bakken LR. N<sub>2</sub>O and CH<sub>4</sub> fluxes in soil influenced by fertilization and tractor traffic. *Soil Biol Biochem* 1993;25:621–30.
- Hauck RD, Melsted SW. Some aspects of the problem of evaluating denitrification in soils. *Soil Sci Soc Am J* 1956;20:361–4.
- Hauck RD, Melsted SW, Yankwich PE. Use of N-isotope distribution in nitrogen gas in the study of denitrification. *Soil Sci* 1958;86:287–91.
- Haynes RJ, Williams PJ. Nutrient cycling and soil fertility in the grazed pasture ecosystem. *Adv Agron* 1993;49:119–99.
- Hefting MM, Bobbink R, De Caluwe H. Nitrous oxide emission and denitrification in chronically nitrate-loaded riparian buffer zones. *J Environ Qual* 2003;32:1194–203.
- Heinen M. Simplified denitrification models: overview and properties. *Geoderma* 2006;133:444–63.
- Hénault C, Germon JC, NEMIS, a predictive model of denitrification on the field scale. *Eur J Soil Sci* 2000;51:257–70.
- Hénault C, Bizouard F, Laville P, Gabrielle B, Nicoulaud B, Germon JC, et al. Predicting in situ soil N<sub>2</sub>O emission using NOE algorithm and soil database. *Glob Chang Biol* 2005;11:115–27.
- Henry S, Baudoin E, López-Gutiérrez JC, Martin-Laurent F, Brauman A, Philippot L. Quantification of denitrifying bacteria in soils by nirK gene targeted real-time PCR. *J Microbiol Methods* 2004;59:327–35.
- Henry S, Bru D, Stres B, Hallet S, Philippot L. Quantitative detection of the nosZ gene, encoding nitrous oxide reductase, and comparison of the abundances of 16S rRNA, narG, nirK, and nosZ genes in soils. *Appl Environ Microbiol* 2006;72:5181–9.
- Henry S, Texier S, Hallet S, Bru D, Dambreville C, Chêneby D, et al. Disentangling the rhizosphere effect on nitrate reducers and denitrifiers: insight into the role of root exudates. *Environ Microbiol* 2008;10:3082–92.
- Hille R. The mononuclear molybdenum enzymes. *Chem Rev* 1996;96:2757–816.
- Hixson SE, Walker RF, Skau CM. Soil denitrification rates in four subalpine plant communities of the Sierra Nevada. *J Environ Qual* 1990;19:617–20.
- Holst J, Liu C, Yao Z, Brüggemann N, Zheng X, Giese M, et al. Fluxes of nitrous oxide, methane and carbon dioxide during freezing–thawing cycles in an Inner Mongolian steppe. *Plant Soil* 2008;308:105–17.
- Holtan-Hartwig L, Dörsch P, Bakken LR. Comparison of denitrifying communities in organic soils: kinetics of NO<sub>3</sub><sup>−</sup> and N<sub>2</sub>O reduction. *Soil Biol Biochem* 2000;32:833–43.
- Holtan-Hartwig L, Dörsch P, Bakken LR. Low temperature control of soil denitrifying communities: kinetics of N<sub>2</sub>O production and reduction. *Soil Biol Biochem* 2002;34:1797–806.
- Horwath WR, Elliott LF, Steiner JJ, Davis JH, Griffith SM. Denitrification in cultivated and noncultivated riparian areas of grass cropping systems. *J Environ Qual* 1998;27:225–31.
- Hyde BP, Hawkins MJ, Fanning AF, Noonan D, Ryan M, O'Toole P, et al. Nitrous oxide emissions from a fertilized and grazed grassland in the South East of Ireland. *Nutr Cycl Agroecosyst* 2006;75:187–200.
- IFA. Fertilizer — use statistics. International Fertilizer Association; 2011 [[www.fertilizer.org/ifa/statistics.asp](http://www.fertilizer.org/ifa/statistics.asp), May 2011].
- Jarvis SC, Barraclough D, Williams J, Rock AJ. Pattern of denitrification loss from grazed grassland: effects of N fertilizer input at different sites. *Plant Soil* 1991;131:77–88.
- Jarvis SC, Hatch DJ, Lovell RD. An improved soil core incubation method for the field measurement of denitrification and net mineralization using acetylene inhibition. *Nutr Cycl Agroecosyst* 2001;59:219–25.
- Jones AM, Knowles R. Sulfide alleviation of acetylene inhibition of nitrous oxide reduction by *Flexibacter canadensis*. *Can J Microbiol* 1992;38:143–8.
- Jones CM, Stres B, Rosenquist M, Hallin S. Phylogenetic analysis of nitrite, nitric oxide, and nitrous oxide respiratory enzymes reveal a complex evolutionary history of denitrification. *Mol Biol Evol* 2008;61:1110–5.
- Jordan C. The effect of fertiliser type and application rate on denitrification losses from cut grassland in Northern Ireland. *Nutr Cycl Agroecosyst* 1989;19:45–55.
- Kaiser EA, Eiland F, Germon JC, Gispert MA, Heinemeyer O, Henault C, et al. What predicts nitrous oxide emissions and denitrification N-loss from European soils? *Z Pflanzenernähr Bodenkd* 1996;159:541–7.
- Kammann C, Grunhage L, Müller C, Jacobi S, Jäger HJ. Seasonal variability and mitigation options for N<sub>2</sub>O emissions from differently managed grasslands. *Environ Pollut* 1998;102:179–86.
- Kandeler E, Deiglmayr K, Tschirko D, Bru D, Philippot L. Abundance of narG, nirS, nirK, and nosZ genes of denitrifying bacteria during primary successions of a glacier foreland. *Appl Environ Microbiol* 2006;72:5957–62.
- Kaya M, Matsumura K, Higashida K, Hata Y, Kawato A, Abe Y, et al. Cloning and enhanced expression of the cytochrome P450nor gene (nirA; CYP55A5) encoding nitric oxide reductase from *Aspergillus oryzae*. *Biosci Biotechnol Biochem* 2004;68:2040–9.
- Keeney DR, Fillery IR, Marx GP. Effect of temperature on the gaseous nitrogen products of denitrification in a silt loam soil 1. *Soil Sci Soc Am J* 1979;43:1124–8.
- Kessavalou A, Doran JW, Mosier AR, Drijber RA. Greenhouse gas fluxes following tillage and wetting in a wheat-fallow cropping system. *J Environ Qual* 1998;27:1105–16.
- Khalil MI, Richards KG. Denitrification enzyme activity and potential of subsoils under grazed grasslands assayed by membrane inlet mass spectrometer. *Soil Biol Biochem* 2011;43:1787–97.
- Kieft TL, Soroker E, Firestone MK. Microbial biomass response to a rapid increase in water potential when dry soil is wetted. *Soil Biol Biochem* 1987;19:119–26.
- Kim D-G, Isenhardt TM, Parkin TB, Schultz RC, Loynachan TE, Raich JW. Nitrous oxide emissions from riparian forest buffers, warm-season and cool-season grass filters, and crop fields. *Biogeochem Discuss* 2009;6:607–50.
- Kim D-G, Mishurov M, Kiely G. Effect of increased N use and dry periods on N<sub>2</sub>O emission from a fertilized grassland. *Nutr Cycl Agroecosyst* 2010;88:397–410.
- Knowles R. Denitrification. *Microbiol Rev* 1982;46:43–70.
- Kobayashi M, Matsuo Y, Takimoto A, Suzuki S, Maruo F, Shoun H. Denitrification, a novel type of respiratory metabolism in fungal mitochondrion. *J Biol Chem* 1996;271:16263–7.



- Kool DM, Hoffland E, Hummelink EWJ, van Groenigen JW. Increased hippuric acid content of urine can reduce soil N<sub>2</sub>O fluxes. *Soil Biol Biochem* 2006;38:1021–7.
- Koponen HT, Jaakkola T, Keinänen-Toivola MM, Kaipainen S, Tuomainen J, Servomaa K, et al. Microbial communities, biomass, and activities in soils as affected by freeze thaw cycles. *Soil Biol Biochem* 2006;38:1861–71.
- Koskinen WC, Keeney DR. Effect of pH on the rate of gaseous products of denitrification in a silt loam soil. *Soil Sci Soc Am J* 1982;46:1165–7.
- Kroeze C, Mosier A, Bouwman L. Closing the global N<sub>2</sub>O budget: a retrospective analysis 1500–1994. *Global Biogeochem Cycles* 1999;13:1–8.
- Labbe N, Parent S, Villemur R. Addition of trace metals increases denitrification rate in closed marine systems. *Water Res* 2003;37:914–20.
- Laughlin RJ, Stevens RJ. Evidence for fungal dominance of denitrification and codenitrification in a grassland soil. *Soil Sci Soc Am J* 2002;66:1540–8.
- Ledgard SF, Luo J. Nitrogen cycling in intensively grazed pastures and practices to reduce whole-farm nitrogen losses. In: *Organizing Committee of 2008 IGC/IRC Conference (Ed.), Multifunctional Grasslands in a Changing World, Volume 1. Guangdong People's Publishing House; 2008. pp. 292–297.*
- Ledgard SF, Jarvis SC, Hatch DJ. Short-term nitrogen fluxes in grassland soils under different long-term nitrogen management regimes. *Soil Biol Biochem* 1998;30:1233–41.
- Ledgard SF, Sprosen MS, Penno JW. Nitrogen inputs and losses from clover/grass pastures grazed by dairy cows: temporal variation and effects of nitrogen fertilization. *Plant Soil* 1999;229:177–87.
- Ledgard SF, Sprosen MS, Judge A, Lindsey S, Jensen R, Clark DA, et al. Nitrogen leaching as affected by dairy intensification and mitigation practices in the resource efficient dairying (RED) trial. In: Currie LD, Hanly JA, editors. *Implementing sustainable nutrient management strategies in agriculture. Occasional report Palmerston North, New Zealand: Fertilizer and Lime Research Centre, Massey University; 2006. p. 263–8.*
- Ledgard SF, Menneer J, Welten B, Kear M, Dexter M, Lindsey S. New nitrogen mitigation technologies for evaluation in the lake Taupo catchment. In: Currie LD, Yates LJ, editors. *Designing sustainable farmers: critical aspects of soil and water management. Occasional report Palmerston North, New Zealand: Fertilizer and Lime Research Centre, Massey University; 2007. p. 19–24.*
- Leffelaar PA, Wessel WW. Denitrification in a homogeneous, closed system: experiment and simulation. *Soil Sci* 1988;146:335–49.
- Lehmann J, Joseph S. *Biochar for environmental management: science and technology*. London: Earthscan; 2009. p. 448.
- Levy-Booth DJ, Winder RS. Quantification of nitrogen reductase and nitrite reductase genes in soil of thinned and clear-cut Douglas fir stands by using real-time PCR. *Appl Environ Microbiol* 2010;76:7116–25.
- Li C, Frolking S, Frolking TA. A model of nitrous oxide evolution from soil driven by rainfall events: 1. Model structure and sensitivity. *J Geophys Res* 1992;97:9759–76.
- Li C, Aber J, Stange F, Butterbach-Bahl K, Papen H. A process-oriented model of N<sub>2</sub>O and NO emissions from forest soils: 1. Model development. *J Geophys Res* 2000;105:4369–84.
- Li Y, White R, Chen D, Zhang J, Li B, Zhang Y, et al. A spatially referenced water and nitrogen management model (WNNM) for (irrigated) intensive cropping systems in the North China Plain. *Ecol Model* 2007;203:395–423.
- Liu B, Mørkved PT, Frostegård A, Bakken LR. Denitrification gene pools, transcription and kinetics of NO, N<sub>2</sub>O and N<sub>2</sub> production as affected by soil pH. *FEMS Microbiol Ecol* 2010;72:407–17.
- Lloyd D. Aerobic denitrification in soils and sediments: from fallacies to facts. *Trends Ecol Evol* 1993;8:352–6.
- López-Gutiérrez JC, Henry S, Hallet S, Martin-Laurent F, Catroux G, Philippot L. Quantification of a novel group of nitrate-reducing bacteria in the environment by real-time PCR. *J Microbiol Methods* 2004;57:399–407.
- Ludwig B, Wolf I, Teepe R. Contribution of nitrification and denitrification to the emission of N<sub>2</sub>O in a freeze-thaw event in an agricultural soil. *J Plant Nutr Soil Sci* 2004;167:678–84.
- Lundquist EJ, Scow KM, Jackson LE, Uesugi SL, Johnson CR. Rapid response of soil microbial communities from conventional, low input, and organic farming systems to a wet/dry cycle. *Soil Biol Biochem* 1999;31:1661–75.
- Luo J, White RE, Ball PR, Tillman RW. Measuring denitrification activity in soils under pasture: optimizing conditions for the short-term denitrification enzyme assay and effects of soil storage on denitrification activity. *Soil Biol Biochem* 1996;22:1137–52.
- Luo J, Tillman RW, White RE, Ball PR. Variation in denitrification activity with soil depth under pasture. *Soil Biol Biochem* 1998;30:897–903.
- Luo J, Tillman RW, Ball PR. Grazing effects on denitrification in a soil under pasture during two contrasting seasons. *Soil Biol Biochem* 1999a;31:903–12.
- Luo J, Tillman RW, Ball PR. Factors regulating denitrification in a soil under pasture. *Soil Biol Biochem* 1999b;31:913–27.
- Luo J, Tillman RW, Ball PR. Nitrogen loss through denitrification in a soil under pasture in New Zealand. *Soil Biol Biochem* 2000;32:497–509.
- Luo J, Dinnison A, Ross C, Ledgard S, Longhurst RD. Control of pollutants using stand-off pads containing different natural materials. *Proc N Z Grassl Assoc* 2006;68:315–20.
- Luo J, Ledgard SF, Lindsey SB. A test of a winter farm management option for mitigating nitrous oxide emissions from a dairy farm. *Soil Use Manag* 2008;24:121–30.
- Luo J, de Klein CAM, Ledgard SF, Saggar S. Management options to reduce nitrous oxide emissions from intensively grazed pastures: a review. *Agric Ecosyst Environ* 2010;136:282–91.
- Maag M, Vinther FP. Nitrous oxide emission by nitrification and denitrification in different soil types and at different soil moisture contents and temperatures. *Appl Soil Ecol* 1996;4:5–14.
- Majumdar D. Past, present and future nitrous oxide emissions from rice fields: a treatise. In: Sheldon AI, Barnhart EP, editors. *Nitrous oxide emissions research progress*. New York: Nova Science Publishers; 2009. p. 179–252.
- Malhi SS, McGill WB, Nyborg M. Nitrate losses in soils: effects of temperature, moisture and substrate concentration. *Soil Biol Biochem* 1990;22:733–7.
- Maron PA, Richaume A, Potier P, Lata JC, Lensi R. Immunological method for direct assessment of the functionality of a denitrifying strain of *Pseudomonas fluorescens* in soil. *J Microbiol Methods* 2004;58:13–21.
- Marshall SB, Cabrera ML, Braun LC, Wood CW, Mullen MD, Guertal EA. Denitrification from fescue pastures in the southeastern USA fertilized with broiler litter. *J Environ Qual* 1999;28:1978–83.
- Martikainen PJ, de Boer W. Nitrous oxide production and nitrification in acidic soil from a dutch coniferous forest. *Soil Biol Biochem* 1993;25:343–7.
- Matheson FE, Nguyen ML, Cooper AB, Burt TP, Bull DC. Fate of <sup>15</sup>N-nitrate in unplanted, planted and harvested riparian wetland soil microcosms. *Ecol Eng* 2002;19:249–64.
- McKenney DJ, Johnson GP, Findlay WI. Effect of temperature on consecutive denitrification reactions in Brookston clay and Fox sandy loam. *Appl Environ Microbiol* 1984;47:919–26.
- McTaggart IP, Parker J, Swan L, Clayton H, Smith KA. Nitrous oxide emissions from grassland and spring barley, following N fertiliser application with and without nitrification inhibitors. *Biol Fertil Soils* 1997;25:261–8.
- Merry RJ, Lee MRF, Davies DR, Moorby JM, Dewhurst RJ, Scollan ND. Nitrogen and energy use efficiency in the rumen of cattle fed high sugar grass and/or red clover silage. *Asp Appl Biol* 2003;70:87–92.
- Metz S, Beisker W, Hartmann A, Schlöter M. Detection methods for the expression of the dissimilatory copper-containing nitrite reductase gene (*Dnirk*) in environmental samples. *J Microbiol Methods* 2003;55:41–50.
- Miller LA, Moorby JM, Davies DR, Humphreys MO, Scollan ND, MacRae JC, et al. Increased concentration of water-soluble carbohydrate in perennial ryegrass (*Lolium perenne* L.): milk production from late-lactation dairy cows. *Grass Forage Sci* 2001;56:383–94.
- Misselbrook TH, Laws JA, Pain BF. Surface application and shallow injection of cattle slurry on grassland: nitrogen losses, herbage yields and nitrogen recoveries. *Grass Forage Sci* 1996;51:270–7.
- Mogge B, Kaiser EA, Munch JC. Nitrous oxide emissions and denitrification N-losses from agricultural soils in the Bornhöved Lake region: influence of organic fertilizers and land-use. *Soil Biol Biochem* 1999;31:1245–52.
- Molstad L, Dörsch P, Bakken LR. Robotized incubation system for monitoring gases (O<sub>2</sub>, NO, N<sub>2</sub>O, N<sub>2</sub>) in denitrifying cultures. *J Microbiol Methods* 2007;71:202–11.
- Mørkved PT, Dörsch P, Henriksen TM, Bakken LR. N<sub>2</sub>O emissions and product ratios of nitrification and denitrification as affected by freezing and thawing. *Soil Biol Biochem* 2006;38:3411–20.
- Morley N, Baggs EM, Dörsch P, Bakken L. Production of NO, N<sub>2</sub>O and N<sub>2</sub> by extracted soil bacteria, regulation by NO<sub>2</sub><sup>-</sup> and O<sub>2</sub> concentrations. *FEMS Microbiol Ecol* 2008;65:102–12.
- Morozkina E, Kurakov A. Dissimilatory nitrate reduction in fungi under conditions of hypoxia and anoxia: a review. *Appl Biochem Microbiol* 2007;43:544–9.
- Mosier AR, Klemmedtsen L. Measuring denitrification in the field. In: Weaver RW, editor. *Methods of soil analysis part 2. SSSA book series 5; Madison, WI 53711 USA: Soil Science Society of America Inc.; 1994. pp. 1047–65.*
- Mosier AR, Parton WJ, Hutchinson GL. Modeling nitrous-oxide evolution from cropped and native soils. *Ecol Bull* 1983;35:229–41.
- Mosier AR, Duxbury JM, Freney JR, Heinemeyer O, Minami K. Assessing and mitigating N<sub>2</sub>O emissions from agricultural soils. *Clim Chang* 1998;40:7–38.
- Mosier AR, Doran JW, Freney JR. Managing soil denitrification. *J Soil Water Conserv* 2002;57:505–13.
- Muhr J, Goldberg SD, Borken W, Gebauer G. Repeated drying–rewetting cycles and their effects on the emission of CO<sub>2</sub>, N<sub>2</sub>O, NO, and CH<sub>4</sub> in a forest soil. *J Plant Nutr Soil Sci* 2008;171:719–28.
- Mulligan FJ, Dillon P, Callan JJ, Rath M, O'Mara FP. Supplementary concentrate type affects nitrogen excretion of grazing dairy cows. *J Dairy Sci* 2004;87:3451–60.
- Murray RE, Knowles R. Production of NO and N<sub>2</sub>O in the presence and absence of C<sub>2</sub>H<sub>2</sub> by soil slurries and batch cultures of denitrifying bacteria. *Soil Biol Biochem* 2003;35:1115–22.
- Myrold DD, Tiedje JM. Diffusional constraints on denitrification in soil. *Soil Sci Soc Am J* 1985;49:651–7.
- Nägele W, Conrad R. Influence of soil pH on the nitrate-reducing microbial populations and their potential to reduce nitrate to NO and N<sub>2</sub>O. *FEMS Microbiol Ecol* 1990;74:49–57.
- Nielsen TH, Revsbech NP. Diffusion chamber for nitrogen-15 determination of coupled nitrification–denitrification around soil–manure interfaces. *Soil Sci Soc Am J* 1994;58:795–800.
- Nishina K, Takenaka C, Ishizuka S. Spatial variations in nitrous oxide and nitric oxide emission potential on a slope of Japanese cedar (*Cryptomeria japonica*) forest. *Soil Sci Plant Nutr* 2009;55:179–89.
- Nishio T, Koike I, Hattori A. Estimates of denitrification and nitrification in coastal and estuarine sediments. *Appl Environ Microbiol* 1983;45:444–50.
- Nobre AD, Keller M, Crill PM, Harris RC. Short-term nitrous oxide profile dynamics and emissions response to water, nitrogen and carbon additions in two tropical soils. *Biol Fertil Soils* 2001;34:363–73.
- Nogales B, Timmis KN, Nedwell DB, Osborn AM. Detection and diversity of expressed denitrification genes in estuarine sediments after reverse transcription-PCR amplification from mRNA. *Appl Environ Microbiol* 2002;68:5017–25.
- Norman J, Jansson PE, Farahbakhshazad N, Butterbach-Bahl K, Li C, Klemmedtsen L. Simulation of NO and N<sub>2</sub>O emissions from a spruce forest during a freeze/thaw event using an N-flux submodel from the PnET-N-DNDC model integrated to CoupModel. *Ecol Model* 2008;216:18–30.
- Nurizzo D, Cutruzzola F, Aresè M, Bourgeois D, Brunori M, Cambillau C, et al. Conformational changes occurring upon reduction and NO binding in nitrite reductase from *Pseudomonas aeruginosa*. *Biochemistry* 1998;37:13987–96.



- Nyborg M, Laidlaw JW, Solberg ED, Malhi SS. Denitrification and nitrous oxide emissions from a black chernozemic soil during spring thaw in Alberta. *Can J Soil Sci* 1997;77:153–60.
- Oenema O, Velthof GL, Yamulki S, Jarvis SC. Nitrous oxide emissions from grazed grassland. *Soil Use Manag* 1997;13:288–95.
- Ogilvie B, Nedwell DB, Harrison RM, Robinson A, Sage A. High nitrate, muddy estuaries as nitrogen sinks: the nitrogen budget of the River Colne estuary (United Kingdom). *Mar Ecol Prog Ser* 1997;150:217–28.
- Öquist MG, Nilsson M, Sörensson F, Kasimir-Klemetsson A, Persson T, Weslien P, et al. Nitrous oxide production in a forest soil at low temperatures – processes and environmental controls. *FEMS Microbiol Ecol* 2004;49:371–8.
- Oremland RS, Umberger C, Culbertson CW, Smith RL. Denitrification in San Francisco Bay intertidal sediments. *Appl Environ Microbiol* 1984;47:1106–12.
- Pain BF, Thompson RB, Rees YJ, Skinner JH. Reducing gaseous losses of nitrogen from cattle slurry applied to grassland by the use of additives. *J Sci Food Agric* 1990;50:141–53.
- Parkin TB, Hatfield JL. Influence of nitrpyrin on N<sub>2</sub>O losses from soil receiving fall-applied anhydrous ammonia. *Agric Ecosyst Environ* 2010;136:81–6.
- Parkin TB, Meisinger JJ. Denitrification below the crop rooting zone as influenced by surface tillage. *J Environ Qual* 1989;18:12–6.
- Parkin TB, Sextone AJ, Tiedje JM. Adaptation of denitrifying populations to low soil pH. *Appl Environ Microbiol* 1985;49:1053–6.
- Parliamentary Commissioner for the Environment. Growing for good – intensive farming, sustainability and New Zealand's environment. Parliamentary Commissioner for the Environment; 2004 [236 pp. <http://www.pce.govt.nz>].
- Parton WJ, Mosier AR, Ojima DS, Valentine DW, Schimel DS, Weier K, et al. Generalized model for N<sub>2</sub> and N<sub>2</sub>O production from nitrification and denitrification. *Global Biogeochem Cycles* 1996;10:401–12.
- Patra AK, Abbadié L, Clays-Josserand A, Degrange V, Grayston SJ, Loiseau P, et al. Effects of grazing on microbial functional groups involved in soil N dynamics. *Ecol Monogr* 2005;75:65–80.
- Paul JW, Beauchamp EG. Denitrification and fermentation in plant-residue-amended soil. *Biol Fertil Soils* 1989;7:303–9.
- Payne WJ. Denitrification. New York: Wiley; 1981.
- Pesaro M, Widmer F, Nicollier G, Zeyer J. Effects of freeze–thaw stress during soil storage on microbial communities and methidathion degradation. *Soil Biol Biochem* 2003;35:1049–61.
- Philippot L. Denitrifying genes in bacterial and Archeal genomes. *Biochim Biophys Acta* 2002;1446:1–23.
- Philippot L, Hallin S. Molecular analyses of soil denitrifying bacteria. In: Cooper JE, Rao JR, editors. Molecular approaches to soil rhizosphere and plant microorganism analysis. Belfast (UK): CABI; 2005. p. 146–65.
- Philippot L, Piutti S, Martin-Laurent F, Hallet S, Germon JC. Molecular analysis of the nitrate-reducing community from unplanted and maize-planted soils. *Appl Environ Microbiol* 2002;68:6121–8.
- Philippot L, Čuhel J, Saby NPA, Chêneby D, Chroňáková A, Bru D, et al. Mapping field-scale spatial patterns of size and activity of the denitrifier community. *Environ Microbiol* 2009;11:1518–26.
- Philippot L, Andert J, Jones CM, Bru D, Hallin S. Importance of denitrifiers lacking the gene encoding the nitrous oxide reductase for N<sub>2</sub>O emissions from soil. *Glob Chang Biol* 2011;17:1497–504.
- Pintathong P, Richardson DJ, Spiro S, Choorit W. Influence of metal ions and organic carbons on denitrification activity of the halotolerant bacterium, *Paracoccus pantotrophus* P16 a strain from shrimp pond. *Electron J Biotechnol* 2009;12(2). <http://dx.doi.org/10.2225/vol12-issue2-fulltext-12>. [online]. 15 April 2009. Available from Internet: 16 December 2010 <http://www.ejbiotechnology.cl/content/vol12/issue2/full/12/index.html>. ISSN 0717-3458].
- Powlson DS, Saffigna PG, Kragt-Cottar M. Denitrification at sub-optimal temperature in soils from different climatic zones. *Soil Biol Biochem* 1988;20:719–23.
- Priemé A, Christensen S. Natural perturbations, drying–wetting and freezing–thawing cycles, and the emission of nitrous oxide, carbon dioxide and methane from farmed organic soils. *Soil Biol Biochem* 2001;33:2083–91.
- Qin S, Hu C, Oenema O. Quantifying the underestimation of soil denitrification potential as determined by the acetylene inhibition method. *Soil Biol Biochem* 2011;47:14–7.
- Ravishankara AR, Daniel JS, Portmann RW. Nitrous oxide (N<sub>2</sub>O): the dominant ozone-depleting substance emitted in the 21st century. *Science* 2009;326:123–5.
- Rich JJ, Myrold DD. Community composition and activities of denitrifying bacteria from adjacent agricultural soil, riparian soil, and creek sediment in Oregon, USA. *Soil Biol Biochem* 2004;36:1431–41.
- Richardson DJ. Bacterial respiration: a flexible process for a changing environment. *Microbiology* 2000;146:551–71.
- Richardson DJ, Berks BC, Russell DA, Spiro S, Taylor CJ. Functional, biochemical and genetic diversity of prokaryotic nitrate reductases. *Cell Mol Life Sci* 2001;58:165–78.
- Richardson DJ, Felgate H, Watmough N, Thomson A, Baggs E. Mitigating release of the potent greenhouse gas N<sub>2</sub>O from the nitrogen cycle – could enzymatic regulation hold the key? *Trends Biotechnol* 2009;27:388–97.
- Rijtema PE, Kroes JG. Some results of nitrogen simulations with the model ANIMO. *Fertil Res* 1991;27:189–98.
- Riley WJ, Matson PA. NLOSS: a mechanistic model of denitrified N<sub>2</sub>O and N<sub>2</sub> evolution from soil. *Soil Sci* 2000;165:237–49.
- Robertson GP, Vitousek PM, Matson PA, Tiedje JM. Denitrification in a clearcut Loblolly pine (*Pinus taeda* L.) plantation in the southeastern US. *Plant Soil* 1987;97:119–29.
- Robertson LA, Cornelisse R, De Vos P, Hadjioetomo R, Kuenen JG. Aerobic denitrification in various heterotrophic nitrifiers. *Antonie Van Leeuwenhoek* 1989;56:289–300.
- Rochester IJ. Estimating nitrous oxide emissions from flood-irrigated alkaline grey clays. *Aust J Soil Res* 2003;41:197–206.
- Rolston DE, Fried M, Goldhamer DA. Denitrification measured directly from nitrogen and nitrous oxide gas fluxes. *Soil Sci Soc Am J* 1976;40:259–66.
- Rondon MR, August PR, Bettermann AD, Brady SF, Grossman TH, Liles MR, et al. Cloning the soil metagenome: a strategy for accessing the genetic and functional diversity of uncultured microorganisms. *Appl Environ Microbiol* 2000;66:2541–7.
- Rösch C, Bothe H. Improved assessment of denitrifying, N<sub>2</sub>-fixing, and total-community bacteria by terminal restriction fragment length polymorphism analysis using multiple restriction enzymes. *Appl Environ Microbiol* 2005;71:2026–35.
- Rosswall T, Bak Baldocchi D, Cicerone RJ, Conard R, Ehrlh DH, Firestone MK, et al. What regulates production and consumption of trace gases in ecosystems. In: Andreae MO, Schimel DS, editors. Exchange of trace gases between terrestrial ecosystems and the atmosphere. Chichester: Wiley; 1989. p. 73–98.
- Rudaz J, Walti E, Kuburz G, Lehmann P, Fuhrer J. Temporal variation in N<sub>2</sub>O and N<sub>2</sub> fluxes from permanent pasture in Switzerland in relation to management, soil water content and soil temperature. *Agric Ecosyst Environ* 1999;73:83–91.
- Ruz-Jerez BE, White RE, Ball PR. Long term measurement of denitrification in three contrasting pastures grazed by sheep. *Soil Biol Biochem* 1994;26:29–39.
- Ryden JC. Denitrification loss from a grassland soil in the field receiving different rates of nitrogen as ammonium nitrate. *J Soil Sci* 1983;34:355–65.
- Ryden JC. Gaseous losses from a grassland soil and the atmosphere. *Nature* 1986;292:235–7.
- Rysgaard S, Risgaard-Petersen N, Nielsen LP, Revsbech NP. Nitrification and denitrification in lake and estuarine sediments measured by the <sup>15</sup>N dilution technique and isotope pairing. *Appl Environ Microbiol* 1993;59:2093–8.
- Saari P, Saarnio S, Kukkonen JVK, Akkanen J, Heinonen J, Saari V, et al. DOC and N<sub>2</sub>O dynamics in upland and peatland forest soils after clear-cutting and soil preparation. *Biogeochemistry* 2009;94:217–31.
- Saetre P, Stark JM. Microbial dynamics and carbon and nitrogen cycling following re-wetting of soils beneath two semi-arid plant species. *Oecologia* 2005;142:247–60.
- Saggar S, Andrew RM, Tate KR, Rodda NJ, Hedley CB, Townsend JA. Modelling nitrous oxide emissions from dairy-grazed pastures. *Nutr Cycl Agroecosyst* 2004a;68:243–55.
- Saggar S, Bolan NS, Bhandral R, Hedley CB, Luo J. A review of emissions of methane, ammonia, and nitrous oxide from animal excreta deposition and farm effluent application in grazed pastures. *N Z J Agric Res* 2004b;47:513–44.
- Saggar S, Giltrap DL, Hedley CB, Lambie SM, Tate KR, Li C. Nitrous oxide emissions from grazed perennial pastures in New Zealand. In: van Amstel A, editor. Non-CO<sub>2</sub> greenhouse gases (NCGG-4). Rotterdam: Millipress; 2005. p. 113–21.
- Saggar S, Giltrap DL, Li C, Tate KR. Modelling nitrous oxide emissions from grazed grasslands in New Zealand. *Agric Ecosyst Environ* 2007a;119:205–16.
- Saggar S, Hedley CB, Giltrap DL, Lambie SM. Measured and modelled estimates of nitrous oxide emission and methane consumption from a sheep-grazed pasture. *Agric Ecosyst Environ* 2007b;122:357–65.
- Saggar S, Tate KR, Giltrap DL, Singh J. Soil-atmosphere exchange of nitrous oxide and methane in New Zealand terrestrial ecosystems and their mitigation options: a review. *Plant Soil* 2008;309:25–42.
- Saggar S, Luo J, Giltrap DL, Maddena M. Nitrous oxide emission from temperate grasslands: process, measurements, modelling and mitigation. In: Sheldon AI, Barnhart EP, editors. Nitrous oxide emissions research progress. New York: Nova Science Publishers; 2009. p. 1–66.
- Saggar S, Luo J, Kim D-G, Jha N. Intensification in pastoral farming: impacts on soil attributes and gaseous emissions. In: Singh BP, Cowie A, Chan Y, editors. Soil health and climate change (soil biology series). Berlin, Heidelberg, Germany: Springer-Verlag; 2011a. p. 207–36.
- Saggar S, Camps M, Hedley M. Is biochar a possible solution for the mitigation of pastoral GHG emissions? Release: the newsletter of the New Zealand Agriculture Greenhouse Gas Research Centre, August 2011; 2011b. p. 6–7.
- Sahrawat KL, Keeney DRC. Nitrous oxide emission from soils. *Adv Soil Sci* 1986;4:103–48.
- Saleh-Lakha S, Shannon KE, Henderson SL, Goyer C, Trevors JT, Zebarth BJ, et al. Effect of pH and temperature on denitrification gene expression and activity in *Pseudomonas mandelii*. *Appl Environ Microbiol* 2009a;75:3903–11.
- Saleh-Lakha S, Shannon KE, Henderson SL, Zebarth BJ, Burton DL, Goyer C, et al. Effect of nitrate and acetylene on nirS, cnorB, and nosZ expression and denitrification activity in *Pseudomonas mandelii*. *Appl Environ Microbiol* 2009b;75:5082–7.
- Schimel JP, Clein JS. Microbial response to freeze–thaw cycles in tundra and taiga soils. *Soil Biol Biochem* 1996;28:1061–6.
- Schimel JP, Gulledge J. Microbial community structure and global trace gases. *Glob Chang Biol* 1998;4:745–58.
- Schipper L, Vojvodic-Vukovic M. Nitrate removal from groundwater using a denitrification wall amended with sawdust: field trial. *J Environ Qual* 1998;27:664–8.
- Schmid M, Neftel A, Riedo M, Fuhrer J. Process-based modelling of nitrous oxide emissions from different nitrogen sources in mown grassland. *Nutr Cycl Agroecosyst* 2001;60:177–87.
- Scholefield D, Hawkins JMB, Jackson SM. Use of a flowing helium atmosphere incubation, technique to measure the effects of denitrification controls applied to intact cores of a clay soil. *Soil Biol. Biochem.* 1997;29:1337–44.
- Schwarz J, Kapp M, Benckiser G, Ottow JCG. Evaluation of denitrification losses by the acetylene inhibition technique in a permanent ryegrass field (*Lolium perenne* L.) fertilized with animal slurry or ammonium nitrate. *Biol Fertil Soils* 1994;18:327–33.
- Schwintner C, Sabaty M, Berna B, Cahors S, Richaud P. Plasmid content and localization of the genes encoding the denitrification enzymes in two strains of *Rhodobacter sphaeroides*. *FEMS Microbiol Lett* 1998;165:313–21.
- Seitzinger SP, Nixon SW, Pilson MEQ, Burke S. Denitrification and N<sub>2</sub>O production in nearshore marine sediments. *Beochim Cosmochim Acta* 1980;44:1853–60.

- Seitzinger SP, Nielsen LP, Caffrey J, Christensen PB. Denitrification measurements in aquatic sediments: a comparison of three methods. *Biogeochemistry* 1993;23:147–67.
- Senbayram M, Chen R, Mühling KH, Dittert K. Contribution of nitrification and denitrification to nitrous oxide emissions from soils after application of biogas waste and other fertilizers. *Rapid Commun Mass Spectrom* 2009;23:2489–98.
- Senbayram M, Chen R, Budai A, Bakken L, Dittert K.  $N_2O$  emission and the  $N_2O/(N_2O + N_2)$  product ratio of denitrification as controlled by available carbon substrates and nitrate concentrations. *Agric Ecosyst Environ* 2012;147:4–12.
- Seo DC, DeLaune RD. Fungal and bacterial mediated denitrification in wetlands: influence of sediment redox condition. *Water Res* 2010;44:2441–50.
- Sexton AJ, Parkin TP, Tiedje JM. Temporal response of soil denitrification to rainfall and irrigation. *Soil Sci Soc Am J* 1985;49:99–103.
- Shannon KEM, Saleh-Lakha S, Burton DL, Zebarth BJ, Goyer C, Trevors JT. Effect of nitrate and glucose addition on denitrification and nitric oxide reductase (*cnorB*) gene abundance and mRNA levels in *Pseudomonas mandelii* inoculated into anoxic soil. *Antonie Van Leeuwenhoek* 2011;100:183–95.
- Sharma S, Aneja MK, Mayer J, Munch JC, Schlöter M. Diversity of transcripts of nitrite reductase genes (*nirK* and *nirS*) in rhizospheres of grain legumes. *Appl Environ Microbiol* 2005;71:2001–7.
- Shepherd M, Menneer J, Ledgard S, Sarathchandra U. Application of carbon additives to reduce nitrogen leaching from cattle urine patches on pasture. *N Z J Agric Res* 2010;53:263–80.
- Siciliano SD, Ma WK, Ferguson S, Farrell RE. Nitrifier dominance of Arctic soil nitrous oxide emissions arises due to fungal competition with denitrifiers for nitrate. *Soil Biol Biochem* 2009;41:1104–10.
- Sigman DM, Casciotti KL, Andreani M, Barford C, Galanter M, Böhlke JK. A bacterial method for the nitrogen isotopic analysis of nitrate in seawater and freshwater. *Anal Chem* 2001;73:4145–53.
- Simarmata T, Benckiser G, Ottow JCG. Effect of an increasing carbon: nitrate-N ratio on the reliability of acetylene in blocking the  $N_2O$ -reductase activity of denitrifying bacteria in soil. *Biol Fertil Soils* 1993;15:107–12.
- Šimek M, Cooper JE. The influence of soil pH on denitrification: progress towards the understanding of this interaction over the last 50 years. *Eur J Soil Sci* 2002;53:345–54.
- Šimek M, Cooper JE, Pícek T, Santrúcková H. Denitrification in arable soils in relation to their physico-chemical properties and fertilization practice. *Soil Biol Biochem* 2000;32:101–10.
- Šimek M, Elhottová D, Klimes F, Hopkins DW. Emissions of  $N_2O$  and  $CO_2$ , denitrification measurements and soil properties in red clover and ryegrass stands. *Soil Biol Biochem* 2004;36:9–21.
- Smith K. Nitrous oxide and climate change. London, UK: Earthscan Ltd.; 2010 [232 pp.].
- Smith MS, Tiedje JM. Phases of denitrification following oxygen depletion in soil. *Soil Biol Biochem* 1979;11:261–7.
- Smith CJ, Nedwell DB, Dong LF, Osborn AM. Diversity and abundance of nitrate reductase genes (*narG* and *napA*), nitrite reductase genes (*nirS* and *nrfA*), and their transcripts in estuarine sediments. *Appl Environ Microbiol* 2007;73:3612–22.
- Smith J, Wagner-Riddle C, Dunfield K. Season and management related changes in the diversity of nitrifying and denitrifying bacteria over winter and spring. *Appl Soil Ecol* 2010;44:138–46.
- Soerensen J, Rasmussen LK, Koike I. Micromolar sulfide concentrations alleviate acetylene blockage of nitrous oxide reduction by denitrifying *Pseudomonas fluorescens*. *Can J Microbiol* 1978;33:1001–5.
- Sohi S, Krull E, Lopez-Capel E, Bol R. A review of biochar and its use and function in soil. *Adv Agron* 2010;105:47–82.
- Stanford G, Dzienia S, Vander Pol RA. Effect of temperature on denitrification rate in soils. *Soil Sci Soc Am J* 1975;39:867–70.
- Stefanson RC, Greenland DJ. Measurement of nitrogen and nitrous oxide evolution from soil–plant systems using sealed growth chambers. *Soil Sci* 1970;109:203–6.
- Stehfest E, Müller C. Simulation of  $N_2O$  emissions from a urine-affected pasture in New Zealand with the ecosystem model DayCent. *J Geophys Res D Atmos* 2004;109:1–7.
- Stevens RJ, Laughlin RJ. Measurement of nitrous oxide and di-nitrogen emissions from agricultural soils. *Nutr Cycl Agroecosyst* 1998;52:131–9.
- Stevens RJ, Laughlin RJ. Lowering the detection limit for dinitrogen using the enrichment of nitrous oxide. *Soil Biol Biochem* 2001;33:1287–9.
- Stevens RJ, Laughlin RJ, Atkins GJ, Prosser SJ. Automated determination of nitrogen-15-labelled dinitrogen and nitrous oxide by mass spectrometry. *Soil Sci Soc Am J* 1993;57:981–8.
- Stres B, Danevčič T, Pal L, Fuka MM, Resman L, Leskovec S, et al. Influence of temperature and soil water content on bacterial, archaeal and denitrifying microbial communities in drained fen grassland soil microcosms. *FEMS Microbiol Ecol* 2008;66:110–22.
- Struwe S, Kjeller A. Potential for  $N_2O$  production from beech (*Fagus sylvatica*) forest soils with varying pH. *Soil Biol Biochem* 1994;26:1003–9.
- Suenaga H. Targeted metagenomics: a high-resolution metagenomics approach for specific gene clusters in complex microbial communities. *Environ Microbiol* 2012;14:13–22.
- Sullivan DG, Wood CW, Owsley WF, Norfleet ML, Wood BH, Shaw JN, et al. Denitrification following land application of swine waste to bermudagrass pasture. *Commun Soil Sci Plant Anal* 2005;36:1277–88.
- Sun W, Luo J, Lindsey S, Ledgard S. Nitrogen losses through denitrification from a grazed pasture as affected by the use of a nitrification inhibitor. In: Multifunctional Grasslands in a Changing World, Volume 1. Organising committee of 2008 IGC/IRC Conference (Ed.). Guangdong People's Publishing House; 2008. pp. 385.
- Swerts M, Merckx R, Vlassak K. Denitrification,  $N_2$  fixation and fermentation during anaerobic incubation of soils amended with glucose and nitrate. *Biol Fertil Soils* 1996;23:229–35.
- Tavares P, Pereira AS, Moura JGG, Moura I. Metalloenzymes of the denitrification pathway. *J Inorg Biochem* 2006;100:2087–100.
- Terry RE, Duxbury JM. Acetylene decomposition in soils. *Soil Sci Soc Am J* 1995;49:90–4.
- Thomas KL, Lloyd D, Boddy L. Effects of oxygen, pH and nitrate concentration on denitrification by *Pseudomonas* species. *FEMS Microbiol Lett* 1994;118:181–6.
- Thompson RB, Ryden JC, Lockyer DR. Fate of nitrogen in cattle slurry following surface application or injection to grassland. *Soil Biol Biochem* 1987;33:1287–9.
- Thomsen JK, Geest T, Cox RP. Mass spectrometric studies of the effect of pH on the accumulation of intermediates in denitrification by *Paracoccus denitrificans*. *Appl Environ Microbiol* 1994;60:536–41.
- Tiedje JM. Ecology of denitrification and dissimilatory nitrate reduction to ammonium. In: Zehnder AJB, editor. *Biology of anaerobic microorganisms*. New York: Wiley & Sons; 1988. p. 179–244.
- Vallejo A, Skiba UM, García-Torres L, Arce A, López-Fernández S, Sánchez-Martín L. Nitrogen oxides emission from soils bearing a potato crop as influenced by fertilization with treated pig slurries and composts. *Soil Biol Biochem* 2006;38:2782–93.
- van Bochove E, Prevost D, Pelletier F. Effects of freeze–thaw and soil structure on nitrous oxide produced in a clay soil. *Soil Sci Soc Am J* 2000;64:1638–43.
- Van Cleemput O, Samater AH. Nitrite in soils: accumulation and role in the formation of gaseous N compounds. *Nutr Cycl Agroecosyst* 1996;45:81–9.
- Van Cleemput O, Vermoesen A, Degroot CJ, Vanryckeghem K. Nitrous oxide emission out of grassland. *Environ Monit Assess* 1994;31:145–52.
- van der Meer HG. Optimising manure management for GHG outcomes. *Aust J Exp Agric* 2008;48:38–45.
- van der Salm C, Dolfing J, Heinen M, Velthof GL. Estimation of nitrogen losses via denitrification from a heavy clay soil under grass. *Agric Ecosyst Environ* 2007;119:311–9.
- Van Kessel JS, Reeves JBIII. On-farm quick tests for estimating nitrogen in dairy manure. *J Dairy Sci* 2000;83:1837–44.
- Van Kessel C, Pennock DJ, Farrell RE. Seasonal variations in denitrification and nitrous oxide evolution at the landscape scale. *Soil Sci Soc Am J* 1993;57:988–95.
- Van Spanning RJM, Richardson D, Ferguson S. Introduction to the biochemistry and molecular biology of denitrification. In: Bothe H, Ferguson SJ, Newton WE, editors. *Biology of the nitrogen cycle*, Ch. 1. Amsterdam, The Netherlands: Elsevier; 2007. p. 382–95.
- Velthof GL, Jarvis SC, Stein A, Allen AG, Oenema O. Spatial variability of nitrous oxide fluxes in mown and grazed grasslands on a poorly drained clay soil. *Soil Biol Biochem* 1996;28:1215–25.
- Venterea RT. Nitrite-driven nitrous oxide production under aerobic soil conditions: kinetics and biochemical controls. *Glob Chang Biol* 2007;13:1798–809.
- Vermoesen A, deGroot CJ, Nollet L, Boeckx P, vanCleemput O. Effect of ammonium and nitrate application on the  $NO$  and  $N_2O$  emission out of different soils. *Plant Soil* 1996;181:153–62.
- Vermoesen A, Cleemput OV, Hofman G. Contribution of urine patches to the emission of nitrous oxide. In: Jarvis SC, Pain BF, editors. *Gaseous nitrogen emissions from grasslands*. Wallingford, Oxon: CAB International; 1997. p. 179–88.
- Vinten AJA, Castle K, Arah JRM. Field evaluation of models of denitrification linked to nitrate leaching for aggregated soil. *Eur J Soil Sci* 1996;47:305–17.
- Virkajärvi P, Maljanen M, Saarijärvi K, Haapala J, Martikainen PJ.  $N_2O$  emissions from boreal grass and grass – clover pasture soils. *Agric Ecosyst Environ* 2010;137:59–67.
- Vollack KU, Harting E, Körner H, Zumft WG. Multiple transcription factors of the FNR family in denitrifying *Pseudomonas stutzeri*: characterisation of four Fnr-like genes, regulatory responses and cognate metabolic processes. *Mol Microbiol* 1999;31:1681–94.
- Wagner-Riddle C, Hu Q, Van Bochove E, Jayasundara S. Linking nitrous oxide flux during spring thaw to nitrate denitrification in the soil profile. *Soil Sci Soc Am J* 2008;72:908–16.
- Walker RF, Hixson SE, Skau CM. Soil denitrification rates in a sub-alpine watershed of the Eastern Sierra-Nevada. *For Ecol Manage* 1992;50:217–31.
- Wallenstein MD, Myrold DD, Firestone M, Voytek M. Environmental controls on denitrifying communities and denitrification rates: insights from molecular methods. *Ecol Appl* 2006;16:2143–52.
- Weier KL, Gilliam JW. Effect of acidity on denitrification and nitrous oxide evolution from Atlantic coastal plain soils. *Soil Sci Soc Am J* 1986;50:1202–5.
- Weier WC, MacRae IC, Myers RJK. Denitrification in a clay soil under pasture and annual crop: losses from  $^{15}N$ -labelled nitrate in the subsoil in the field using  $C_2H_2$  inhibition. *Soil Biol Biochem* 1993;25:999–1004.
- Wertz S, Degrange V, Prosser JI, Poly F, Commeaux C, Freitag T, et al. Maintenance of soil functioning following erosion of microbial diversity. *Environ Microbiol* 2006;8:2162–9.
- Wertz S, Degrange V, Prosser JI, Poly F, Commeaux C, Guillaumaud N, et al. Decline of soil microbial diversity does not influence the resistance and resilience of key soil microbial functional groups following a model disturbance. *Environ Microbiol* 2007;9:2211–9.
- Weymann D, Geistlinger RW, von der Heide C, Flessa H. Kinetics of  $N_2O$  production and reduction in a nitrate-contaminated aquifer inferred from laboratory incubation experiments. *Biogeosci Discuss* 2010;7:503–43.
- Wickramasinghe KN, Talibudeen O, Witty JF. A gas flow-through system for studying denitrification in soils. *J Soil Sci* 1978;29:527–36.
- Wolf I, Brumme R. Dinitrogen and nitrous oxide formation in beech forest floor and mineral soils. *Soil Sci Soc Am J* 2003;67:1862–8.
- Wolf B, Zheng X, Bruggemann N, Chen W, Dannenmann M, Han X, et al. Grazing-induced reduction of natural nitrous oxide release from continental steppe. *Nature* 2010;464:881–4.
- Wood DW, Setubal JC, Kaul R, Monks DE, Kitajima JP, Okura VK, et al. The genome of the natural genetic engineer *Agrobacterium tumefaciens* C58. *Science* 2001;294:2317–23.

- Xu YQ, Wan SQ, Cheng WX, Li LH. Impacts of grazing intensity on denitrification and  $\text{N}_2\text{O}$  production in a semi-arid grassland ecosystem. *Biogeochemistry* 2008;88: 103–15.
- Ye RW, Averill BA, Tiedje JM. Denitrification: production and consumption of nitric oxide. *Appl Environ Microbiol* 1994;60:1053–8.
- Yeomans JC, Beauchamp EG. Limited inhibition of nitrous oxide reduction in soil in the presence of acetylene. *Soil Biol Biochem* 1978;10:517–9.
- Yeomans JC, Beauchamp EG. Acetylene as a possible substrate in the denitrification process. *Can J Soil Sci* 1982;62:139–44.
- Yergeau E, Kowalchuk G. Responses of Antarctic soil microbial communities and associated functions to temperature and freeze–thaw cycle frequency. *Environ Microbiol* 2008;10:2223–35.
- Yoshida M, Ishii S, Otsuka S, Senoo K. Temporal shifts in diversity and quantity of nirS and nirK in a rice paddy field soil. *Soil Biol Biochem* 2009;41:2044–51.
- Yoshinari T, Knowles R. Acetylene inhibition of nitrous oxide reduction by denitrifying bacteria. *Biochem Biophys Res Commun* 1976;69:705–10.
- Yu K, Seo DC, DeLaune RD. Incomplete acetylene inhibition of nitrous oxide reduction potential in denitrification assay as revealed by using  $^{15}\text{N}$ -nitrate tracer. *Commun Soil Sci Plant Anal* 2010;41:2201–10.
- Zaman M, Blennerhassett JD. Effects of the different rates of urease and nitrification inhibitors on gaseous emissions of ammonia and nitrous oxide, nitrate leaching and pasture production from urine patches in an intensive grazed pasture system. *Agric Ecosyst Environ* 2010;136:236–46.
- Zaman M, Nguyen ML. Effect of lime or zeolite on  $\text{N}_2\text{O}$  and  $\text{N}_2$  emissions from a pastoral soil treated with urine or nitrate-N fertilizer under field conditions. *Agric Ecosyst Environ* 2010;136:254–61.
- Zaman M, Nguyen ML, Matheson F, Blennerhassett JD, Quin BF. Can soil amendments (zeolite or lime) shift the balance between nitrous oxide and dinitrogen emissions from pasture and wetland soils receiving urine or urea-N? *Aust J Soil Res* 2007;45: 543–53.
- Zaman M, Nguyen ML, Saggar S.  $\text{N}_2\text{O}$  and  $\text{N}_2$  emissions from pasture and wetland soils with and without amendments of nitrate, lime and zeolite under laboratory condition. *Aust J Soil Res* 2008a;46:526–34.
- Zaman M, Nguyen ML, Gold AJ, Groffman PM, Kellogg DQ, Wilcock RJ. Nitrous oxide generation, denitrification, and nitrate removal in a seepage wetland intercepting surface and subsurface flows from a grazed dairy catchment. *Aust J Soil Res* 2008b;46:565–77.
- Zaman M, Saggar S, Blennerhassett JD, Singh J. Effect of urease and nitrification inhibitors on N transformation, gaseous emissions of ammonia and nitrous oxide, pasture yield and N uptake in grazed pasture system. *Soil Biol Biochem* 2009;41:1270–80.
- Zannoni D. Respiration in archaea and bacteria. *Diversity of prokaryotic respiratory system*, vol. 2. New York: Springer; 2004. p. 169–206.
- Zhang L, Takaya N, Kitazume T, Kondo T, Shoun H. Purification and cDNA cloning of nitric oxide reductase cytochrome P450nor (CYP55A4) from *Trichosporon cutaneum*. *Eur J Biochem* 2001;268:3198–204.
- Zhang L, Kudo T, Takaya N, Shoun H. Distribution, structure and function of fungal nitric oxide reductase P450nor-recent advances. *Int Congr Ser* 2002;1233: 197–202.
- Zhong J, Fan Chengxin C, Liu G, Zhang L, Shang J, Gu X. Seasonal variation of potential denitrification rates of surface sediment from Meiliang Bay, Taihu Lake, China. *J Environ Sci* 2010;22:961–7.
- Zhou Q, Takenaka S, Murakami S, Seesuriyachan P, Kuntiya A, Aoki K. Screening and characterization of bacteria that can utilize ammonium and nitrate ions simultaneously under controlled cultural conditions. *J Biosci Bioeng* 2007;103:185–91.
- Zumft WG. Cell biology and molecular basis of denitrification? *Microbiol Mol Biol Rev* 1997;61:533–616.