

B - 456

分值: 200 分

Problem Statement

There are three dice, each with six faces.

The j -th face of the i -th die has $A_{i,j}$ written on it.

For each die, each face comes up with probability $\frac{1}{6}$.

有三个骰子，每个骰子有六个面。第 i 个骰子的第 j 面上写着 $A_{i,j}$ 。对于每个骰子，每个面朝上的概率为 $\frac{1}{6}$ 。

When these dice are rolled simultaneously, find the probability that the values 4, 5, 6 each appear exactly once.

同时掷出这三个骰子时，求每个 4, 5, 6 恰好出现一次的概率。

Constraints

- $A_{i,j}$ is an integer between 1 and 6, inclusive.
 - $A_{i,j}$ 是介于 1 和 6 之间（含两端）的整数。
-

Input

The input is given from Standard Input in the following format:

输入按照以下格式从标准输入给出：

```
A1,1 A1,2 A1,3 A1,4 A1,5 A1,6  
A2,1 A2,2 A2,3 A2,4 A2,5 A2,6  
A3,1 A3,2 A3,3 A3,4 A3,5 A3,6
```

Output

Output the answer. The answer is considered correct if the absolute or relative error from the true answer is at most 10^{-6} .

输出答案。如果你的输出与正确答案的绝对误差或相对误差不超过 10^{-6} ，则视为正确。

Sample Input 1

```
1 2 3 4 5 6
1 2 3 4 5 6
1 2 3 4 5 6
```

Sample Output 1

```
0.0277777778
```

The desired probability is $\frac{1}{36}$. An output such as 0.027778 is also accepted.

所求概率为 $\frac{1}{36}$ 。类似 0.027778 的输出也会被接受。

Sample Input 2

```
4 5 6 4 5 6
4 4 5 5 6 6
6 5 4 4 5 6
```

Sample Output 2

```
0.2222222222
```

The desired probability is $\frac{2}{9}$.

所求概率为 $\frac{2}{9}$ 。