

C - Not Adjacent

分值: 300 分

Problem Statement

You are given a string S consisting of a, b, c.

给定一个字符串 S ，其字符组成为 a、b、c。

Find the number of non-empty **substrings** of S in which no two adjacent characters are the same, modulo 998244353.

求 S 的所有非空 **子串** 中，满足相邻两个字符互不相同的子串数量，结果对 998244353 取模。

Two substrings are considered distinct if they are taken from different positions, even if they are identical as strings.

只要两个子串的位置不同，即使内容完全相同，也视为不同的子串。

What is a substring?

什么是子串?

A **substring** of S is a string obtained by deleting zero or more characters from the beginning and zero or more characters from the end of S . For example, ab is a substring of abc, but ac is not a substring of abc.

S 的 **子串** 是指从 S 的开头删除若干个（可以是0个）字符，再从末尾删除若干个（可以是0个）字符后得到的字符串。例如，ab 是 abc 的子串，但 ac 不是 abc 的子串。

Constraints

- S is a string of length between 1 and 3×10^5 , inclusive, consisting of a, b, c.
- S 是长度在 1 到 3×10^5 之间（包含两端）的字符串，由 a、b、c 组成。

Input

The input is given from Standard Input in the following format:

输入按照以下格式从标准输入给出：

S

Output

Output the answer.

输出答案。

Sample Input 1

abbc

Sample Output 1

6

The substrings in which no two adjacent characters are the same are the following six:

满足相邻字符互不相同的子串共有以下6个：

- a (from the 1st to the 1st character of S)
- b (from the 2nd to the 2nd character of S)
- b (from the 3rd to the 3rd character of S)
- c (from the 4th to the 4th character of S)
- ab (from the 1st to the 2nd character of S)
- bc (from the 3rd to the 4th character of S)
- a (取自 S 的第 1 个字符到第 1 个字符)
- b (取自 S 的第 2 个字符到第 2 个字符)
- b (取自 S 的第 3 个字符到第 3 个字符)

- Note that, as with the 2nd and 3rd entries, two substrings are considered distinct if they are taken from different positions, even if they are identical as strings.

Sample Input 2

Sample Output 2

760