



NEW

MITHRA

MEDICAL LABORATORY

ACCURACY • CARE • TRUST

Chettiyankinar Road, Kuttippala,
Opp. Govt. Ayurveda Dispensary
Ph : +91 81578 87311, 95266 77311
e-mail : mithralab12020@gmail.com

ashik .G.**Age / Sex : Male****Ref. No. : 51367**

Referred by :

Sample Collected At : 21-08-2026 02:31 PMIP/OP/SRF No :

Institution : MITHRA MEDICAL

Report On : 21-08-2026 02:32 PM

Phone No : 6282378262

LABORATORY KUTTIPPALA

Test Description	Value Observed	Reference Range
------------------	----------------	-----------------

DEPARTMENT OF CLINICAL BIOCHEMISTRY

HbA1C- Glycated Hb
[Nephelometry Method]

10.5%**Mean Blood Glucose : 255mg/dl**

Normal : < 5.7
Pre Diabetes : 5.7- 6.5
Diabetes : > 6.5

Good Control : < 6.5
Adequate Control : 6.5 - 7.5
Inadequate Control : 7.5 - 8.5
Poor Control : > 8.5

Notes:

Glycosylated hemoglobin values are used to assess long-term glucose control in diabetes, especially in insulin-dependent diabetics whose glucose levels are labile, and in whom blood and urine glucose measurements exhibit significant daily variation. GHb measurements reflect the level of control present over the preceding 100-120 days. In such patients, whose fasting glucose concentrations are fairly consistent from day to day, there is a correlation between glycosylated hemoglobin and single fasting glucose levels. Continued high levels of blood glucose are reflected in high GHb concentrations. Glycosylated hemoglobin predicts the progression of retinopathy.

Chronic blood loss, hemolytic anemia, or other setting for decrease in RBC life span, results in a decrease in the glycosylated hemoglobin level. Pregnancy may lower glycosylated hemoglobin.

**** End Of Report ****

SAHEED MOHAMED P
Technologist

ABDUNNASER MAYYERI
Lab Incharge