EMAKAL 19

Lorem ipsum is placeholder text commonly used in the graphic, print, and publishing industries for previewing layouts and visual mockups.

Call Now | Download Brochure

Specification

Name: Emakal 19

Composition: Emamectin Benzoate 1.9% EC

Category: Agrochemicals

Sub Category: Insecticides

Mode of Action: Stomach action

Packing: 50ml | 100ml | 250ml | 500ml | 1 liter | 5 liter

Features

FEATURES:
It is an insecticide with stomach action and should be ingested by the larvae to be most effective. Affected larvae become paralyzed and stop feeding shortly after exposure to Emakal 19 and subsequently die after 2-4 days.

Direction of Use:
Apply the insecticide when the incidence of larvae is first observed and repeat applications as necessary. Apply the insecticide in sufficient quantity of water to ensure thorough coverage of the foliage.

Symptoms of Poisoning:
Early symptoms of poisoning may be a combination of dilation of pupils, muscular in-coordination, ataxia, and muscle tremors.

Usage

Crop(s)	Common Name of Pest	AI (g)	Formulation (ml)	Dilution in Water (L)	Waiting Period between last spray to harvest (days)	Re-entry after each Application (In Hours)
Chickpea	Pod borer	7.13	375	500	14	-
Chilli	Fruit borer & Thrips	7.13	375	500	14	-
Cotton	Bollworms	7.13	375	500	14	-
Paddy	Leaf folder & hispa	7.13	375	500	14	-
Soybean	Green semilooper, Pod borer, Girdle beetle and Tobacco caterpillar	7.13	375	500	14	-
Tomato	Fruit borer	7.13	375	500	14	-

More products from **Insecticides**

EMAKAL 19 | Weedicide | Weedicide | Weedicide

Weedicide | Weedicide | Weedicide | Weedicide

Weedicide | Weedicide | Weedicide | Weedicide

2007-2024 All Rights Reserved with Kalyani Industries Limited.