

1. From Fourier transform we know, the test function is a convolution of 2 equal box function

$$f(t) = g(t) * g(t)$$

Also we know the FT of a box function is a Sinc function and using the Fourier transform we get?

$$\begin{aligned} F[f(t)] &= F[g(t) * g(t)] \\ &= \text{Sinc}(u) \cdot \text{Sinc}(u) \\ &= \text{Sinc}^2(u) \end{aligned}$$

2. a) $\delta(t) * \delta(t - t_0)$

Using Fourier transform we get:

$$= \int_{-\infty}^{\infty} \delta(\tau) \delta(t - t_0 - \tau) d\tau$$

We know that the FT of impulse $\delta(t)$ is 1 if $t=0$.
therefore, $t - t_0 - \tau = 0$ giving us $\tau = t - t_0$

Substituting the value of τ in the 1st equation we get:

$$= \int_{-\infty}^{\infty} \delta(t - t_0) \delta(t - t_0 - t - t_0) d\tau$$

$$= \delta(t - t_0) \int_{-\infty}^{\infty} \delta(0) d\tau$$

Giving us $\delta(t - t_0)$

2. b) $\delta(t-t_0)$ and $\delta(t+t_0)$

Using Fourier we get,

$$\delta(t-t_0) * \delta(t+t_0) = \int_{-\infty}^{\infty} \delta(\tau-t_0) \delta(t+t_0-\tau) d\tau$$

The Fourier transform $\delta(t)$ of impulse is 1 if $t=0$.

\therefore , $t+t_0-\tau=0$ giving us $\tau=t+t_0$

Substituting the value of τ in the 1st equation:

$$= \int_{-\infty}^{\infty} \delta(t+t_0-t_0) \delta(t+t_0-t-t_0) d\tau$$

$$= \int_{-\infty}^{\infty} \delta(t) \delta(0) d\tau$$

$$= \delta(t) \int_{-\infty}^{\infty} \delta(0) d\tau$$

$$= \delta(t)$$

6.1) π_{Mod}

2) π_{vir}

3) π_m

7.1) π_{us}

2) $R =$

$S = T$

$T =$

R_{ex}

3) $R =$

R_{ex}

4) R

S

R

6.1) $\pi_{Model, price} (\sigma_{Name='Aston Martin', (Car \bowtie Manufacturer)})$

2) $\pi_{VIN} (\sigma_{Name='James Bond' (Car \bowtie Customer)})$

3) $\pi_{model} (\sigma_{Price < 100000 \text{ AND } year > 2016 \text{ AND } Name='Ford' (Car \bowtie Manufacturer)})$

7.1) $\pi_{Username} (\pi_{Seller, Username (Auction)} \bowtie \pi_{Buyer, Username (Sold)})$

2) $R = Auction \bowtie Sold$

$S = \pi_{Seller, Username} (\sigma_{Auction.AuctionID = Sold.AuctionID} (R))$

$T = \pi_{Seller, Username (Auction)} - S$

Result = $\pi_{Username} (T)$

3) $R = Auction \bowtie (\pi_{AuctionID (Sold)})$

Result = $\pi_{Auction.AuctionID} (\sigma_{Auction.EndTime < CurrentTime \text{ AND}}$

$Auction.AuctionID \text{ NOT IN } (\pi_{Sold.AuctionID (Sold)}) (R))$

4) $R = Auction \bowtie Item$

$S = R - ((R \bowtie \rho_{MaxPrice} (\pi_{Price (Auction)})))$

$- (\pi_{ItemID, SellerUsername, Price} (R))$

Result = $\pi_{ItemName, Auction, SellerUsername} (S)$

5) Butterworth filter can be defined as,

$$H(u, v) = \frac{1}{1 + \left[\frac{D_0}{D(u, v)} \right]^{2n}}$$

Using this we can construct the homomorphic filter as:

$$H(u, v) = (y_H - y_L) \left[\frac{1}{1 + \left[\frac{D_0}{D(u, v)} \right]^{2n}} \right] + y_L$$

~~Bonus~~

a) $\sum_{x=0}^{M-1} \sum_{y=0}^{N-1} B(x, y)$

b) $\frac{1}{MN}$

c) $\frac{1}{2MN}$

d) $\frac{1}{AB}$

BONUS 1

$$a) \sum_{x=0}^{M-1} \sum_{y=0}^{N-1} \delta(x-y) e^{-j2\pi \left(\frac{xu}{M} + \frac{yv}{N} \right)}$$

$$e^0 = 1$$

$$\delta(0,0) = 1$$

$$x=y=0$$

$$b) \frac{1}{MN} \sum_{x=0}^{M-1} \sum_{y=0}^{N-1} MN \delta(u,v) e^{j2\pi \left(\frac{xu}{M} + \frac{yv}{N} \right)}$$

$$\text{We know } u=v=0$$

$$= 1$$

$$c) \sum_{x=0}^{a-1} \sum_{y=0}^{b-1} \delta(x-x_0, y-y_0) e^{j2\pi \left(\frac{xu}{a} + \frac{yv}{b} \right)}$$

$$x-x_0 = 0 ; x=x_0$$

$$y-y_0 = 0 ; y=y_0$$

$$\Rightarrow e^{-j2\pi \left(\frac{x_0 u}{a} + \frac{y_0 v}{b} \right)} \Rightarrow e^{-j2\pi \left(\frac{x_0 u}{M} + \frac{y_0 v}{N} \right)}$$

$$d) \frac{1}{AB} \sum_{x=0}^{A-1} \sum_{y=0}^{B-1} AB \delta(u-u_0, v-v_0) e^{j2\pi \left(\frac{xu}{A} + \frac{yv}{B} \right)}$$

$$u-u_0 = 0 ; u=u_0$$

$$v-v_0 = 0 ; v=v_0$$

$$= e^{j2\pi \left(\frac{xu_0}{A} + \frac{yv_0}{B} \right)}$$

$$\Rightarrow e^{j2\pi \left(\frac{xu_0}{M} + \frac{yv_0}{N} \right)}$$

haric filter a:

$$\left. \begin{matrix} + y \\ 2\pi \end{matrix} \right\} \frac{1}{M}$$

$$\begin{aligned}
 e) \quad & \frac{1}{AB} \sum_{x=0}^{A-1} \sum_{y=0}^{B-1} \frac{AB}{2} \left[\delta(u+u_0, v+v_0) + \delta(u-u_0, v-v_0) \right] \\
 & \quad e^{j2\pi \left(\frac{xu}{A} + \frac{yv}{B} \right)} \\
 = & \frac{1}{2} \sum_{x=0}^{A-1} \sum_{y=0}^{B-1} \left[\delta(u+u_0, v+v_0) e^{j2\pi \left(\frac{xu}{A} + \frac{yv}{B} \right)} + \delta(u-u_0, v-v_0) e^{j2\pi \left(\frac{xu}{A} + \frac{yv}{B} \right)} \right] \\
 & \quad \left. \begin{array}{l} u+u_0=0 \Rightarrow u=-u_0 \\ v+v_0=0 \Rightarrow v=-v_0 \\ u-u_0=0 \Rightarrow u=u_0 \\ v-v_0=0 \Rightarrow v=v_0 \end{array} \right\}
 \end{aligned}$$

$$\begin{aligned}
 \Rightarrow & \frac{1}{2} \left[e^{j2\pi \left(-\frac{xu_0}{A} + \frac{yv_0}{B} \right)} + e^{j2\pi \left(\frac{xu_0}{A} + \frac{yv_0}{B} \right)} \right] \\
 = & \frac{1}{2} \left[e^{j2\pi \left(\frac{xu_0}{A} + \frac{yv_0}{B} \right)} + e^{-j2\pi \left(\frac{xu_0}{A} + \frac{yv_0}{B} \right)} \right]
 \end{aligned}$$

$$\begin{aligned}
 & = \cos \left(\frac{2\pi x_0}{A} + \frac{2\pi y_0}{B} \right) \\
 & = \cos \left(\frac{2\pi x_0}{M} + \frac{2\pi y_0}{N} \right)
 \end{aligned}$$

2. BON
Dura
are
DFT,
a var
in the
Now,
and
the

As,
mi
wh

BONUS

2. During the process of obtaining the right side image we are taking the conjugate and performing the inverse DFT (J^{-1}) on it. The complex conjugate basically transforms a variable in its negative form, for example, a to $-a$ in the inverse transform.

Now, if $f(x, y)$ is the original image with $M \times N$ size and $F(u, v)$ is the DFT then in general the image of the right side can be given as:

$$J^{-1}[F^*(u, v)] = \sum_{x=0}^{M-1} \sum_{y=0}^{N-1} F(u, v) e^{-j2\pi \left(\frac{ux}{M} + \frac{vy}{N} \right)}$$

$$J^{-1}[F^*(u, v)] = \sum_{x=0}^{M-1} \sum_{y=0}^{N-1} F(u, v) e^{j2\pi \left(\frac{u(-x)}{M} + \frac{v(-y)}{N} \right)}$$

$$J^{-1}[F^*(u, v)] = f(-x, -y)$$

As, $f(x, y)$ is the original image then $f(-x, -y)$ is basically mirroring the original image with respect to the origin which is resulting the image on the right side.