

Welcome**Introduction****Review****Model and Cost Function**

- ✓ **Video:** Model Representation
8 min
- ✓ **Reading:** Model Representation
3 min
- ✓ **Video:** Cost Function
8 min
- ✓ **Reading:** Cost Function
3 min
- ▶ **Video:** Cost Function - Intuition I
11 min
- 📖 **Reading:** Cost Function - Intuition I
4 min
- ▶ **Video:** Cost Function - Intuition II
8 min
- 📖 **Reading:** Cost Function - Intuition II
3 min

Parameter Learning

- ▶ **Video:** Gradient Descent
11 min
- 📖 **Reading:** Gradient Descent
3 min
- ▶ **Video:** Gradient Descent Intuition
11 min
- 📖 **Reading:** Gradient Descent Intuition
3 min




Gradient Descent

So we have our hypothesis function and we have a way of measuring how well it fits into the data. Now we need to estimate the parameters in the hypothesis function. That's where gradient descent comes in.

Imagine that we graph our hypothesis function based on its fields θ_0 and θ_1 (actually we are graphing the cost function as a function of the parameter estimates). We are not graphing x and y itself, but the parameter range of our hypothesis function and the cost resulting from selecting a particular set of parameters.

We put θ_0 on the x axis and θ_1 on the y axis, with the cost function on the vertical z axis. The points on our graph will be the result of the cost function using our hypothesis with those specific theta parameters. The graph below depicts such a setup.



We will know that we have succeeded when our cost function is at the very bottom of the pits in our graph, i.e. when its value is the minimum. The red arrows show the minimum points in the graph.

The way we do this is by taking the derivative (the tangential line to a function) of our cost function. The slope of the tangent is the derivative at that point and it will give us a direction to move towards. We make steps down the cost function in the direction with the steepest descent. The size of each step is determined by the parameter α , which is called the learning rate.