

Mathematical Logic and Computability

Lecture 4: Sequent Calculus

Yunpyo An

Ulsan National Institute of Science and Technology

Mathematical Logic and Computability

Aug 28, 2023

Outline

Sequent Rules

Structural Rules and Connective
Rules

Derive Connective
Rules

Quantifier and
Equality Rules

Further Derivable
Rules

Consistency

Sequent Rules

Structural Rules and Connective Rules

Derive Connective Rules

Quantifier and Equality Rules

Further Derivable Rules

Consistency

Introduction to Sequent Calculus

In this lecture, we will learn about *Sequent Calculus* to develop a proof system for first-order logic.

To show that a proposition follows from the axioms, we need to supply a proof like follows:

$$\Phi \rightarrow \varphi_1 \rightarrow \dots \rightarrow \varphi_n \rightarrow \varphi$$

where the set of formulas Φ is the set of axioms and φ is the proposition to be proved.

How can we show that φ follows from Φ ? Also, how can we concrete the proof? For convenience, we fix a symbol set as S .

Our observation in proof, we can describe a proof by listing the corresponding assumptions and the respective claim. If we call a nonempty list (sequence) of formulas a *sequent*, then we can use sequents to describe "stage of a proof".

The stages with assumptions are notated as $\varphi_1, \dots, \varphi_n$ and our claim φ is rendered by the sequent $\varphi_1 \dots \varphi_n \varphi$.

The sequence $\varphi_1 \dots \varphi_n$ is called the *antecedent* and φ is called the *succedent* of the sequent.

Sequent Rules

Remind the Sequent Calculi notation in the first lecture.

Let's consider the proof by contradiction as follows:

$$\frac{\varphi_1 \dots \varphi_n \quad \neg\varphi \quad \psi \quad \varphi_1 \dots \varphi_n \quad \neg\varphi \quad \neg\psi}{\varphi_1 \dots \varphi_n \quad \varphi}$$

We can say that "Our assumptions $\varphi_1 \dots \varphi_n$ and the negation of claim $\neg\varphi$ lead to a contradiction $\psi \wedge \neg\psi$."

For convenience, let's use the letters Γ, Δ, \dots to denote finite sets of formulas. (of course, possibly empty) If we define $\Gamma = \varphi_1 \dots \varphi_n$, we can rewrite as follows:

$$\frac{\Gamma \quad \neg\varphi \quad \psi \quad \Gamma \quad \neg\varphi \quad \neg\psi}{\Gamma \quad \varphi}$$

If in the calculus C , there is a derivation of the sequent $\Gamma\varphi$, then we write $\vdash \Gamma\varphi$ and say that $\Gamma\varphi$ is *derivable*.

Definition

A formula φ is *formally provable* or *derivable* from a set Φ of formulas ($\Phi \vdash \varphi$), if and only if there are finitely many formulas $\varphi_1, \dots, \varphi_n$ in Φ such that $\vdash \varphi_1 \dots \varphi_n \varphi$

Antecedent and Assumption Rules

Antecedent Rule (Ant)

if every member of Γ is also a member of Γ' ($\Gamma \subseteq \Gamma'$)

$$\frac{\Gamma \quad \varphi}{\Gamma' \quad \varphi} \quad (1)$$

Assumption Rule (Assm)

if φ is a member of Γ

$$\frac{}{\Gamma \quad \varphi} \quad (2)$$

Question, show that correctness of the antecedent rule and the assumption rule.

Proof by Case and Contradiction Rules

Proof by Case Rule (PC)

$$\frac{\begin{array}{l} \Gamma \quad \psi \quad \varphi \\ \Gamma \quad \neg\psi \quad \varphi \end{array}}{\Gamma \quad \varphi} \quad (3)$$

Proof by Contradiction (Ctr)

$$\frac{\begin{array}{l} \Gamma \quad \neg\varphi \quad \psi \\ \Gamma \quad \neg\varphi \quad \neg\psi \end{array}}{\Gamma \quad \varphi} \quad (4)$$

Rule for Disjunction

\vee -Rule for the Antecedent ($\vee A$)

$$\frac{\begin{array}{ccc} \Gamma & \varphi & \xi \\ \Gamma & \psi & \xi \end{array}}{\Gamma \quad (\varphi \vee \psi) \quad \xi} \quad (5)$$

\vee -Rule for the Succedent ($\vee S$)

$$\frac{\Gamma \quad \varphi}{\Gamma \quad (\varphi \vee \psi)} \quad \frac{\Gamma \quad \varphi}{\Gamma \quad (\psi \vee \varphi)} \quad (6)$$

Chain Rule

Second Contradiction Rule (Ctr')

$$\frac{\begin{array}{l} \Gamma \quad \psi \\ \Gamma \quad \neg\psi \end{array}}{\Gamma \quad \varphi} \quad (7)$$

Chain Rule (Ch)

$$\frac{\begin{array}{l} \Gamma \quad \varphi \quad \psi \\ \Gamma \quad \varphi \quad \psi \end{array}}{\Gamma \quad \psi} \quad (8)$$

Equation 7 looks very weird. Question: How can we prove these?

Outline

Sequent Rules

Structural Rules
and Connective
Rules

Derive Connective
Rules

Quantifier and
Equality Rules

Further Derivable
Rules

Consistency

Contraposition Rules

Contraposition Rules (Cp)

$$\frac{\Gamma \quad \varphi \quad \psi}{\Gamma \quad \neg\psi \quad \neg\varphi} \qquad \frac{\Gamma \quad \neg\varphi \quad \neg\psi}{\Gamma \quad \psi \quad \varphi} \qquad (9)$$

$$\frac{\Gamma \quad \neg\varphi \quad \psi}{\Gamma \quad \neg\psi \quad \varphi} \qquad \frac{\Gamma \quad \varphi \quad \neg\psi}{\Gamma \quad \psi \quad \neg\varphi} \qquad (10)$$

Modus Ponens

Proposition

$$\frac{\begin{array}{l} \Gamma \quad (\varphi \vee \psi) \\ \Gamma \quad \neg\varphi \end{array}}{\Gamma \quad \psi} \quad (11)$$

Modus Ponens (MP)

$$\frac{\begin{array}{l} \Gamma \quad (\varphi \rightarrow \psi) \\ \Gamma \quad \varphi \end{array}}{\Gamma \quad \psi} \quad (12)$$

Rule for Universal Quantifier

Rule for \exists -Introduction in the Succedent (\exists S)

$$\frac{\Gamma \quad \varphi \frac{t}{x}}{\Gamma \quad \exists x \varphi} \quad (13)$$

Rule for \exists -Introduction in the Antecedent (\exists A)

$$\frac{\Gamma \quad \varphi \frac{y}{x} \quad \psi}{\Gamma \quad \exists x \varphi \quad \psi} \quad (14)$$

Rule for Equality



Figure: Equivalence of Equality

Reflexivity Rule for Equality (\equiv)

$$\overline{t \equiv t}$$

(15)

Rule for Equality



Figure: Equivalence of Equality

Substitution Rule for Equality (Sub)

$$\frac{\Gamma \quad \varphi \frac{t}{x}}{\Gamma \quad t \equiv t' \quad \varphi \frac{t'}{x}}$$

(16)

Further Derivable Rules

$$\frac{\Gamma \quad \varphi}{\Gamma \quad \exists x \varphi} \quad \frac{\Gamma \quad \varphi \quad \psi}{\Gamma \quad \exists x \varphi \quad \psi} \quad \text{if } x \text{ is not free in } \Gamma \psi \quad (17)$$

$$\frac{\Gamma \quad \varphi}{\Gamma \quad x \equiv t \quad \varphi \frac{t}{x}} \quad (18)$$

$$\frac{\Gamma \quad t_1 \equiv t_2}{\Gamma \quad t_2 \equiv t_1} \quad \frac{\Gamma \quad t_1 \equiv t_2 \quad \Gamma \quad t_2 \equiv t_3}{\Gamma \quad t_1 \equiv t_3} \quad (19)$$

$$\frac{\begin{array}{c} \Gamma \quad R t_1 \dots t_n \\ \Gamma \quad t_1 \equiv t'_1 \\ \vdots \quad \vdots \\ \Gamma \quad t_n \equiv t'_n \end{array}}{\Gamma \quad R t'_1 \dots t'_n} \quad \frac{\begin{array}{c} \Gamma \quad t_1 \equiv t'_1 \\ \vdots \quad \vdots \\ \Gamma \quad t_n \equiv t'_n \end{array}}{\Gamma \quad f t_1 \dots t_n \quad f t'_1 \dots t'_n} \quad (20)$$

Lemma

For all Φ and φ , if and only if there is a finite subset Φ_0 of Φ such that $\Phi_0 \vdash \varphi$.

Theorem

For all Φ and φ , if $\Phi \vdash \varphi$, then $\Phi \models \varphi$.

Inverse of the theorem (completeness) is shown in the next lecture.

Exercise

Prove formal proof of the existence of left inverse from the group axioms as following

$$\varphi_0 : \forall x \forall y \forall z (x \circ y) \circ z \equiv x \circ (y \circ z)$$

$$\varphi_1 : \forall x x \circ e \equiv x$$

$$\varphi_2 : \forall x \exists y x \circ y \equiv e$$

Definition

$\text{Con}\Phi$: A set Φ is *consistent* if and only if there is no formula φ such that $\Phi \vdash \varphi$ and $\Phi \vdash \neg\varphi$.

$\text{Inc}\Phi$: A set Φ is *inconsistent* if and only if it is not consistent.

Consistency Lemmas

Lemma

For a set of formulas Φ the following are equivalent:

- ▶ Φ is inconsistent.
- ▶ For all φ , $\Phi \vdash \varphi$.

Lemma

For a set of formulas Φ the following are equivalent:

- ▶ Φ is consistent.
- ▶ There is a formula φ which is not derivable from Φ .

Consistency Lemmas

In next lecture, we will prove the completeness theorem, which states that for all Φ and φ , $\Phi \vdash \varphi$ if and only if $\Phi \models \varphi$.

Lemma

For all Φ , $\text{Con}\Phi$ if and only if $\text{Con}\Phi_0$ for all finite subsets Φ_0 of Φ .

Lemma

Every satisfiable set of formula is consistent

Lemma

For all Φ and φ the following holds:

- ▶ $\Phi \vdash \varphi$ if and only if $\text{Inc}\Phi \cup \{\neg\varphi\}$.
- ▶ $\Phi \vdash \neg\varphi$ if and only if $\text{Inc}\Phi \cup \{\varphi\}$.
- ▶ If $\text{Con}\Phi$, then $\text{Con}\Phi \cup \{\varphi\}$ or $\text{Con}\Phi \cup \{\neg\varphi\}$.

[Outline](#)[Sequent Rules](#)[Structural Rules
and Connective
Rules](#)[Derive Connective
Rules](#)[Quantifier and
Equality Rules](#)[Further Derivable
Rules](#)[Consistency](#)

Consistency Lemmas

Lemma

For $n \in \mathbb{N}$ let S_n be symbol sets such that

$$S_0 \subseteq S_1 \subseteq S_2 \subseteq \dots$$

and let Φ_n be set of S_n -formulas such that $\text{Con}_{S_n} \Phi_n$ and

$$\Phi_0 \subseteq \Phi_1 \subseteq \Phi_2 \subseteq \dots$$

let $S = \bigcup_{n \in \mathbb{N}} S_n$ and $\Phi = \bigcup_{n \in \mathbb{N}} \Phi_n$. Then $\text{Con}_S \Phi$.