

# Environmental drivers of *Plasmodium vivax* malaria in Manaus: a climate- and land-use-driven SEIRS-SEI model

Raphael Felberg Levy<sup>a</sup>, Flavio Codeço Coelho<sup>a</sup>

<sup>a</sup>School of Applied Mathematics, EMAP FGV, Praia de Botafogo 190, Rio de Janeiro, 22250-145, RJ, Brazil

## ARTICLE INFO

### Keywords:

Malaria transmission  
*Plasmodium vivax*  
*Anopheles darlingi*  
SEIRS-SEI model  
Mathematical epidemiology  
Brazilian Amazon  
Environmental drivers  
Climate-dependent transmission  
Model calibration

## ABSTRACT

This paper aims to analyze the behavior of malaria transmission in the Amazon region based on climatic and environmental changes, such as temperature, precipitation, humidity, deforestation and forest fires, through proposed modifications to the SIR and SEI models, in order to contribute to the study of applications of external effects on the evolution of the disease. The Trajetorias Project, developed by the Synthesis Center on Biodiversity and Ecosystem Services (SinBiose/CNPq) was used as an initial reference for the study.

This work employs a modified SEIRS/SEI methodology, based on the SIR/SEI model developed by Parham and Michael (2010) which takes into account rainfall and temperature, with further modifications.

## 1. Introduction

Malaria remains one of the most significant vector-borne diseases worldwide, caused by protozoa of the genus *Plasmodium* and transmitted by female mosquitoes of the genus *Anopheles*. Although present in multiple regions, the burden of malaria in the Americas is highly concentrated in the Amazon basin, which spans nine countries and accounts for a substantial proportion of reported cases [1]. The persistence of malaria in this region is closely linked to ecological, climatic, and socioeconomic factors that influence both vector dynamics and human exposure.

Environmental changes play a central role in shaping malaria transmission. In the Amazon, processes such as deforestation, agricultural expansion, and infrastructure development alter local ecosystems, affecting vector habitats and increasing human–vector contact. These changes often promote the proliferation of mosquito breeding sites and concentrate human populations in high-risk areas, particularly among non-immune migrants [4, 5]. In addition to land-use change, climatic variables such as temperature and rainfall, as well as disturbances such as forest fires and mining activities, further influence vector abundance, pathogen development, and transmission intensity.

Despite the recognized importance of environmental drivers, integrating these factors into mathematical models of malaria transmission remains challenging. In particular, distinguishing between their effects on mosquito biology and their impact on human–vector interaction is essential for developing models that are both biologically consistent and identifiable from available data.

In this work, we develop and analyze a modified SIR–SEI

model for malaria transmission in the Amazon, incorporating environmental variables such as temperature, rainfall, deforestation, and fire activity. Climatic factors are modeled as drivers of mosquito life-history parameters, while land-use variables are introduced as modifiers of transmission intensity. Using epidemiological and environmental data from the Trajetorias Project [6], we investigate how these factors influence disease dynamics and assess their relative contributions through analytical and numerical methods.

## 2. Environmental Drivers of Malaria Transmission in the Amazon

Environmental changes, particularly deforestation and forest fires, have been widely associated with the resurgence and spatial redistribution of malaria transmission in the Amazon region [7]. The removal of forest canopy alters local microclimatic conditions, increasing sunlight exposure and ambient temperature, which in turn favor the development of mosquito larvae and pupae. In particular, the primary malaria vector in the Amazon, *Anopheles darlingi*, is known to thrive in partially deforested environments, where sunlit water bodies promote larval growth and increase the availability of food sources such as algae [8, 9].

Beyond ecological effects, deforestation also influences human exposure to malaria. Activities such as logging, agriculture, and infrastructure development lead to increased human presence in recently altered environments, often in close proximity to mosquito breeding sites. This results in higher human–vector contact rates and facilitates transmission, especially in areas where both vector density and environmental suitability are elevated. Additionally, land-use changes such as agriculture may contribute to river sedimentation and the formation of artificial breeding sites, further enhancing mosquito proliferation.

Forest fragmentation has been shown to accelerate mosquito

 raphael.levy147@gmail.com (R.F. Levy);

flavio.codeco.coelho@fgv.br (F.C. Coelho)

ORCID(s):

development cycles by increasing temperature and light availability, reducing the time from larval to adult stages and increasing vectorial capacity and survivorship [9, 10]. Similarly, forest fires, often used as a land-clearing mechanism, remove vegetation and modify habitat structure, contributing to long-term ecological changes that can influence mosquito populations and disease transmission dynamics [11, 12]. These processes may also increase the likelihood of subsequent fires by promoting understory drying and fuel accumulation [13].

In this work, deforestation and fire activity are not embedded as mechanistic drivers within the epidemiological model itself. Instead, they are treated as environmental variables that describe the context in which malaria transmission occurs, and are used together with temperature, precipitation and humidity as predictors in the regression of the weekly transmission probabilities  $b_1$  and  $b_2$  (see Section 6.4). In that methodological framework, the freed per-bite transmission probabilities  $b_{1,w}$  and  $b_{2,w}$ , recovered from the observed case dynamics, are regressed against a set of environmental covariates that includes the land-cover and fire variables described below. This enables deforestation and fire activity to modulate transmission indirectly through the empirical relation between environmental conditions and the effective transmission coefficients, rather than by directly altering the mechanistic mosquito demographic or physiological rates.

Deforestation was obtained from the DETER system, which provides daily records of deforestation alerts by degradation type, reported in  $\text{km}^2$  [14]. The daily alerts were aggregated and summarized at weekly resolution, with the 4-week total degraded area used as the deforestation covariate  $D_f$  in the regression. Three degradation types were distinguished by DETER, namely clear cut, forest degradation and burn scar, and combined into a total deforestation term. Forest fire activity was obtained from two complementary sources. First, the Brazilian National Institute for Space Research (INPE) provides daily records of detected fire hotspots [15]; because area-based measurements are not available at this temporal resolution, the weekly number of detected hotspots was used as a fire activity covariate  $F_c$ . Second, annual land-surface fire extent estimates from the MapBiomas project provide fire trends in hectares [16], assigned as a monthly covariate  $F_t$  accounting for the burned area associated with fire activity.

Daily climatic data, including temperature, precipitation and humidity, were obtained from the Mosqlimate API [17] and included as the  $T$ ,  $R$  and  $H$  covariates. Together, the variables  $T$ ,  $R$ ,  $H$ ,  $D_f$ ,  $F_c$ ,  $F_t$  form the environmental predictor set used to relate the observed environmental conditions to the weekly transmission probabilities  $b_1$  and  $b_2$ .

### 3. Model Formulation

#### 3.1. Baseline Delayed SIR-SEI Model

We begin with the classical host-vector framework for malaria transmission, based on the model introduced by Parham and Michael (2010) [18], which incorporates temperature- and rainfall-dependent mosquito dynamics. The human population is divided into susceptible, infected, and recovered compartments, denoted by  $S_H$ ,  $I_H$ , and  $R_H$ , respectively. The mosquito population is structured into susceptible, exposed, and infectious compartments, denoted by  $S_M$ ,  $E_M$ , and  $I_M$ .

The transmission dynamics are governed by the system of delay differential equations:

$$\frac{dS_H}{dt} = -ab_2 \frac{I_M(t)}{N} S_H(t) \quad (1a)$$

$$\frac{dI_H}{dt} = ab_2 \frac{I_M(t - \tau_H)}{N} S_H(t - \tau_H) - \gamma I_H(t) \quad (1b)$$

$$\frac{dR_H}{dt} = \gamma I_H(t) \quad (1c)$$

$$\frac{dS_M}{dt} = b(R, T) - \left( ab_1 \frac{I_H(t)}{N} + \mu(T) \right) S_M(t) \quad (1d)$$

$$\begin{aligned} \frac{dE_M}{dt} = & ab_1 \frac{I_H(t)}{N} S_M(t) - \mu(T) E_M(t) \\ & - ab_1 \frac{I_H(t - \tau_M)}{N} S_M(t - \tau_M) \ell(\tau_M) \end{aligned} \quad (1e)$$

$$\begin{aligned} \frac{dI_M}{dt} = & ab_1 \frac{I_H(t - \tau_M)}{N} S_M(t - \tau_M) \ell(\tau_M) \\ & - \mu(T) I_M(t) \end{aligned} \quad (1f)$$

Here,  $a(T)$  denotes the biting rate,  $b_1$  and  $b_2$  are transmission probabilities from humans to mosquitoes and vice versa,  $\mu(T)$  is the mosquito mortality rate, and  $\tau_H$ ,  $\tau_M$  represent the intrinsic and extrinsic incubation periods. The function  $b(R, T)$  describes mosquito recruitment, while  $\ell(\tau_M)$  represents survival during the extrinsic incubation period.

Although this formulation incorporates biologically relevant delays, numerical simulations revealed that, under the parameterization considered, the model fails to reproduce sustained transmission dynamics. In particular, both the infected human population and the mosquito population tends to extinction, indicating that the structure does not adequately capture endemic persistence.

#### 3.2. ODE approximation and structural limitations

To simplify the analysis and avoid the technical complexity of delay differential equations, an ordinary differential equation (ODE) approximation was initially considered. In this formulation, the extrinsic incubation period in mosquitoes is modeled via a transition rate  $b_3$ , replacing the explicit delay  $\tau_M$ .

The resulting system is given by:

$$\frac{dS_H}{dt} = -ab_2 \frac{I_M}{N} S_H \quad (2a)$$

$$\frac{dI_H}{dt} = ab_2 \frac{I_M}{N} S_H - \gamma I_H \quad (2b)$$

$$\frac{dR_H}{dt} = \gamma I_H \quad (2c)$$

$$\frac{dS_M}{dt} = b(R, T) - \left( ab_1 \frac{I_H}{N} + \mu \right) S_M \quad (2d)$$

$$\frac{dE_M}{dt} = ab_1 \frac{I_H}{N} S_M - (\mu + b_3 \ell) E_M \quad (2e)$$

$$\frac{dI_M}{dt} = b_3 \ell E_M - \mu I_M \quad (2f)$$

While this formulation restores positivity and simplifies numerical implementation, it exhibits a fundamental limitation: the system tends toward a disease-free equilibrium after a single outbreak, failing to capture the recurrent nature of malaria transmission observed in endemic regions. This behavior highlights the importance of explicitly modeling incubation periods and immunity dynamics.

### 3.3. SEIRS-SEI model with ODE approximation

To address the limitations of the previous formulations, we introduce an ODE based SEIRS-SEI model that incorporates both incubation delays as done above and temporary immunity in the human population. The human compartments are extended to include an exposed class  $E_H$ , and recovered individuals lose immunity at rate  $\omega$ , returning to the susceptible class.

To account for incubation dynamics in both hosts and vectors, the transition parameters were defined as temperature-dependent development rates.

For humans, the intrinsic incubation process was modeled through  $b_{3H} = \frac{1}{\tau_H}$ , where  $\tau_H$  denotes the intrinsic incubation period.

For mosquitoes, the constant transition parameter  $b_3 = 1/\tau_M$  from the original formulation was replaced by a Brière-type function,

$$b_{3M}(T) = \eta T(T - T_{\min})(T_{\max} - T)^{1/2}, \quad (3)$$

which incorporates minimum and maximum thermal thresholds for sporogonic development [19, 20].

The scaling coefficient  $\eta$  was estimated by imposing  $b_{3M}(T_{\text{opt}}) = \frac{1}{\tau_M(T_{\text{opt}})}$ , yielding

$$\eta = \frac{\frac{1}{\tau_M(T_{\text{opt}})}}{T_{\text{opt}}(T_{\text{opt}} - T_{\min})(T_{\max} - T_{\text{opt}})^{1/2}} \quad (4)$$

Another addition was the inclusion of a humidity-dependent survival component  $p(H)$  on to the daily survival rate of the

adult mosquitoes, which is modeled as a sigmoid function of the daily minimum relative humidity  $U$ . The parameter  $H_0$  (in %) represents the humidity threshold at which the mosquito experiences a 50% reduction in survival relative to its maximum, and  $k$  (in  $\%^{-1}$ ) controls the steepness of the transition. The parameter  $\varphi$  (dimensionless) establishes a baseline survival floor, accounting for the persistence of individuals in favorable micro-habitats even under extreme desiccation conditions.

The SEIRS-SEI complete model is then given by:

$$\frac{dS_H}{dt} = -\lambda_H(t)S_H + \omega R_H \quad (5a)$$

$$\frac{dE_H}{dt} = \lambda_H(t)S_H - b_{3H}E_H \quad (5b)$$

$$\frac{dI_H}{dt} = b_{3H}E_H - \gamma I_H \quad (5c)$$

$$\frac{dR_H}{dt} = \gamma I_H - \omega R_H \quad (5d)$$

$$\frac{dS_M}{dt} = \Lambda(t) - \lambda_M(t)S_M - \mu S_M \quad (5e)$$

$$\frac{dE_M}{dt} = \lambda_M(t)S_M - (\mu + b_{3M}\ell)E_M \quad (5f)$$

$$\frac{dI_M}{dt} = b_{3M}\ell E_M - \mu I_M \quad (5g)$$

The forces of infection are defined as:

$$\lambda_H(t) = ab_2 \frac{I_M(t)}{N(t)} \quad (6a)$$

$$\lambda_M(t) = ab_1 \frac{I_H(t)}{N(t)} \quad (6b)$$

### 3.4. Recalibration of the temperature-dependent biting rate

In the original formulation by Parham & Michael [18], the biting rate is expressed as a temperature-dependent function derived from the duration of the gonotrophic cycle. Although this formulation captures the general thermal dependence of mosquito feeding behavior, its parameterization was not developed specifically for Amazonian mosquito populations. With this in mind, we recalibrated the biting-rate function using entomological observations reported by de Barros, Honório and Arruda [21], regarding females of *Anopheles darlingi* collected between August 2003 and July 2004 through intermittent sampling carried out in selected months over the study period.

Their results allowed for estimation of the duration of the gonotrophic cycle through the following formula:

$$P = Q^{1/g} \implies g = \frac{\ln(Q)}{\ln(P)}, \quad (7)$$

where  $P$  is the daily survival rate of mosquitoes,  $Q$  is the proportion of parous mosquitoes females in the population and  $g$  is the duration of the gonotrophic cycle, given in days

[22].

Given the lack of more recent data, it was decided to use the obtained estimates of the time period specified above with average temperatures from the period of analysis between 2017 and 2023, given that the biting rate is given as a function of temperature. It was decided to consider  $T' = 16^\circ\text{C}$ , given that below this temperature there is no development of the vector, as the threshold temperature for the biting rate, and we estimated values of  $g$  for the missing months of the analysis via interpolation. With this, given the closed formula for the biting rate, and knowing that it is the inverse of the gonotrophic cycle, we calculated  $D_1 = g(T - T')$  for each of the months with available data<sup>1</sup>.

Having monthly values of  $D_1$ , we took their average in order to estimate the biting rate and the proportion of female mosquitoes to humans, which is given by

$$m = \frac{HBR}{a(T)}, \quad (8)$$

where  $HBR$  is the mean biting density of *An. darlingi*, given in bites/person/night, while  $a(T)$  is the biting rate, given in bites/mosquito/night. Based on the data,  $M_0$ , the initial population of mosquitoes used in the analysis, was estimated to be approximately  $10N_0$ , based on the found value of  $m$  for the month of January.

At this point, we also approximated the ratio of initially exposed to infectious mosquitoes to be their values at the steady state:

$$\frac{dI_M}{dt} = b_{3M} \ell E_M - \mu I_M = 0 \implies \frac{E(0)}{I(0)} = \frac{\mu(0,0)}{b_{3M}(0)\ell(0,0)} \approx 3$$

This adjustment preserves the functional structure proposed by Parham and Michael while adapting the parameter values to better reflect the feeding behavior of the primary malaria vector in the Amazon region.

### 3.5. Density-dependent mosquito recruitment and carrying capacity

Mosquito population dynamics were initially modeled using a linear birth–death process of the form

$$\frac{dM}{dt} = b - \mu M, \quad (9)$$

where  $b$  represents the birth rate and  $\mu$  is the per capita mortality rate. However, this formulation does not account for environmental limitations on population growth.

To incorporate the effect of limited environmental resources, we introduce a density-dependent recruitment mechanism inspired by the classical Verhulst logistic model [47],

$$\frac{dM}{dt} = bM \left(1 - \frac{M}{K}\right) - \mu M, \quad (10)$$

<sup>1</sup> $a(T)$  is defined in Table A.1

where  $M(t) = S_M(t) + E_M(t) + I_M(t)$  denotes the total mosquito population,  $b$  the per capita recruitment rate,  $\mu$  the mortality rate, and  $K$  the environmental carrying capacity. This formulation captures the reduction in population growth as the mosquito population approaches the environmental limit.

In the present model, this idea is adapted by replacing the constant recruitment term with the density-dependent function

$$\Lambda(t) = b(R, T) K(t) \left(1 - \frac{M(t)}{K(t)}\right), \quad (11)$$

where the environmental dependence is separated into two distinct mechanisms:

1. The climatic modulation of mosquito recruitment through  $b(R, T)$ , inherited from the original Parham-Michael formulation;
2. The environmental limitation imposed by habitat availability through the carrying capacity  $K(t)$ .

In this interpretation,  $b(R, T)$  governs the rate at which mosquitoes are produced under favorable temperature and rainfall conditions, whereas  $K(t)$  determines the maximum sustainable mosquito population supported by the environment. Consequently, temperature and precipitation affect mosquito dynamics both through reproductive success and through the availability of suitable breeding habitats.

The environmental carrying capacity is defined by

$$K(t) = M' \cdot f_T(T(t)) \cdot f_R(R(t)), \quad (12)$$

where  $M'$  denotes the baseline mosquito population size. Given that we require  $0 \leq 1 - \frac{M(t)}{K(t)} \leq 1$ , we would require  $M'$  such that  $M' \geq \frac{M}{f_T(T)f_R(R)}$ .

The environmental response functions are given by

$$f_T(T) = \min \left( 1.0, \frac{T(T - T_{\min A.}) \sqrt{T_{\max A.} - T}}{T_{\text{opA.}} (T_{\text{opA.}} - T_{\min A.}) \sqrt{T_{\max A.} - T_{\text{opA.}}}} \right), \quad (13a)$$

$$f_R(R) = \left( \frac{R}{R_L/2} \right) \exp \left( 1 - \frac{R}{R_L/2} \right), \quad (13b)$$

where  $f_T(T)$  describes the temperature suitability for mosquito persistence and  $f_R(R)$  represents the dual effect of rainfall on habitat availability.

The rainfall response function increases with precipitation under moderate rainfall conditions, reflecting the formation of breeding sites, but decreases under excessive rainfall due to habitat flushing and destruction of immature stages.

Both environmental response functions are normalized so that  $0 \leq f_T(T), f_R(R) \leq 1$ , ensuring that the carrying

capacity remains bounded by the baseline population scale  $M_0$ .

Unlike the classical logistic formulation, where population growth is directly proportional to the current mosquito abundance, the present representation assumes that recruitment is primarily determined by environmental suitability and breeding-site availability. The factor  $\left(1 - \frac{M(t)}{K(t)}\right)$  therefore imposes density-dependent regulation by reducing recruitment as the mosquito population approaches the carrying capacity, providing a biologically realistic upper bound on mosquito abundance. The mathematical implications of this formulation are analyzed in Section 5.1.

## 4. Data Description and Processing

### 4.1. Epidemiological Data

This study focuses on malaria cases caused by *Plasmodium vivax*, which accounts for the majority of malaria infections in Brazil and is the predominant species in the Amazon region [23, 24, 25].

Epidemiological data were obtained from the SIVEP-Malaria system [26], which provides daily records of notified malaria cases for all Brazilian municipalities from 2003 to 2023. The analysis was restricted to the period 2017–2023, for which complete epidemiological and environmental data were available.

### 4.2. Population Estimates

The municipality of Manaus was selected due to its high incidence of *P. vivax* malaria, particularly in rural areas. However, detailed yearly estimates of rural population were not directly available from census data. To address this limitation, population estimates were derived using data from the Trajetorias Project [6], which provides incidence values aggregated over five-year periods.

Incidence is defined as

$$\text{Inc}(d, m, z, t_1, t_2) = \frac{\text{Cases}(d, m, z, t_1, t_2)}{\text{Pop}(m, z, (t_1 + t_2)/2) \cdot (t_2 - t_1)} \times 10^5, \quad (14)$$

where  $\text{Cases}(d, m, z, t_1, t_2)$  denotes the number of reported cases for disease  $d$  in municipality  $m$  and zone  $z$  over the interval  $[t_1, t_2]$ , and  $\text{Pop}(m, z, \cdot)$  represents the population at the midpoint of the interval.

This relationship was inverted to estimate rural population values from reported incidence and case counts. Using this approach, the rural population of Manaus was estimated to represent less than 1% of the total population, ranging between 0.3% and 0.6% of the total municipal population, while accounting for a disproportionately large share of malaria cases (approximately 30% to 40% over the study periods).

Yearly total population values for Manaus were obtained via interpolation based on official census data from the Brazilian Institute of Geography and Statistics (IBGE) for the years 2000, 2010, and 2022 [27, 28, 29]. Intermediate yearly values were constructed using linear interpolation, while values beyond 2022 were extrapolated based on observed trends.

### 4.3. Data Consistency Adjustments

During preprocessing, inconsistencies were identified in the Trajetorias dataset. In particular, incidence rates for the period 2015–2019 had been calculated using population estimates corresponding to 2004–2008, resulting in a temporal mismatch. These values were recalculated using population estimates consistent with the appropriate time interval.

Additionally, rural and urban population estimates for intermediate years had been derived using fixed proportions based on 2010 data. To better account for demographic changes, particularly urbanization, these estimates were revised. The rural population for 2006 was approximated as the mean between the reported values for 2000 and 2010, while the 2017 value was obtained through extrapolation based on the observed growth trend. These adjustments ensured temporal consistency in the population data used throughout the study.

## 5. Mathematical Analysis of the Model

### 5.1. Mosquito population dynamics

In this section, we analyze the qualitative impact of the density-dependent recruitment term on mosquito population dynamics. We compare the proposed formulation with the classical Verhulst model and show that the modified structure prevents extinction under suitable environmental conditions.

**Proposition 1** (Environmental persistence of the mosquito population). *Assume that  $b(R(t), T(t)) \geq b_0 > 0$  and  $K(t) \geq K_0 > 0$  for all  $t \geq 0$ . Then, the mosquito population  $M(t)$  governed by*

$$\frac{dM}{dt} = b(R, T) K(t) \left(1 - \frac{M(t)}{K(t)}\right) - \mu M(t) \quad (15)$$

*cannot converge to zero as  $t \rightarrow \infty$ . In particular,  $M = 0$  is not an equilibrium and cannot be asymptotically stable.*

*Proof.* We begin by recalling the classical logistic model with mortality:

$$\frac{dM}{dt} = bM \left(1 - \frac{M}{K}\right) - \mu M. \quad (16)$$

In this formulation, the right-hand side vanishes at  $M = 0$ , so that  $M = 0$  is always an equilibrium. Depending on the parameters, this equilibrium may be stable, allowing extinction of the population.

In contrast, in the present model the recruitment term is given by

$$\Lambda(t) = b(R, T) K(t) \left( 1 - \frac{M(t)}{K(t)} \right), \quad (17)$$

and the population dynamics satisfy

$$\frac{dM}{dt} = b(R, T) K(t) \left( 1 - \frac{M(t)}{K(t)} \right) - \mu M(t). \quad (18)$$

Evaluating the right-hand side at  $M = 0$ , we obtain

$$\left. \frac{dM}{dt} \right|_{M=0} = b(R, T) K(t). \quad (19)$$

By assumption,  $b(R(t), T(t)) \geq b_0 > 0$  and  $K(t) \geq K_0 > 0$ , hence

$$\left. \frac{dM}{dt} \right|_{M=0} \geq b_0 K_0 > 0. \quad (20)$$

Therefore,  $M = 0$  is not an equilibrium of the system.

Moreover, for  $M(t)$  sufficiently small, we have

$$\frac{dM}{dt} \geq b_0 K_0 - \mu M(t). \quad (21)$$

Choosing  $\varepsilon > 0$  such that  $\varepsilon < \frac{b_0 K_0}{\mu}$ , it follows that

$$\frac{dM}{dt} > 0 \quad \text{whenever } 0 < M(t) < \varepsilon.$$

Thus, any trajectory entering a neighborhood of zero is driven away from it.

Consequently,  $M(t)$  cannot converge to zero as  $t \rightarrow \infty$ , and the extinction state is not asymptotically stable.  $\square$

## 5.2. Basic Reproduction Number ( $\mathcal{R}_0$ )

Following the next-generation matrix method of van den Driessche and Watmough [30], we order the infected compartments as  $X_I = (E_H, I_H, E_M, I_M)^T$  and denote by  $\mathcal{F}_i(x)$  the rate of appearance of new infections in compartment  $i$  and by  $\mathcal{V}_i(x) = \mathcal{V}_i^-(x) - \mathcal{V}_i^+(x)$  the net rate of transfer of individuals into and out of compartment  $i$ . The disease transmission model consists of non-negative initial conditions together with the system  $\dot{x} = f(x) = \mathcal{F}(x) - \mathcal{V}(x)$ , where each component is assumed twice continuously differentiable.

**Disease-free equilibrium:** The disease-free equilibrium (DFE) is

$$\begin{aligned} x_0 &= (S_H^*, E_H^*, I_H^*, R_H^*, S_M^*, E_M^*, I_M^*)^T \\ &= (N, 0, 0, 0, M^*, 0, 0)^T. \end{aligned} \quad (22)$$

The equilibrium mosquito abundance  $M^*$  is obtained from the mosquito subsystem

$$\frac{dM}{dt} = \Lambda(t) - \mu M = b(R, T) K(t) \left( 1 - \frac{M}{K(t)} \right) - \mu M = 0, \quad (23)$$

which yields

$$M^* = \frac{\bar{b} \bar{K}}{\bar{b} + \mu}, \quad \Lambda^* := \Lambda|_{M=M^*} = \mu M^* = \frac{\mu \bar{b} \bar{K}}{\bar{b} + \mu}, \quad (24)$$

where  $\bar{b} = b(\bar{R}, \bar{T})$  and  $\bar{K} = K(t)$  are the environmental drivers evaluated at a fixed reference state, that being their study-period means, so that  $\mathcal{R}_0$  is a function of that reference climate-habitat state. Note that  $\Lambda^*$  is *not* an arbitrary value: it is the recruitment rate that sustains the equilibrium vector population  $M^*$ , and it inherits the environmental dependence of  $b(R, T)$  and  $K(t)$ .

**Next-generation matrices:** The new-infection and transfer matrices, evaluated at the DFE, are

$$\begin{aligned} F &= \begin{pmatrix} 0 & 0 & 0 & \frac{ab_2 S_H^*}{N} \\ 0 & 0 & 0 & 0 \\ 0 & \frac{ab_1 S_M^*}{N} & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \\ V &= \begin{pmatrix} b_{3H} & 0 & 0 & 0 \\ -b_{3H} & \gamma & 0 & 0 \\ 0 & 0 & b_{3M} \ell + \mu & 0 \\ 0 & 0 & -b_{3M} \ell & \mu \end{pmatrix}, \end{aligned} \quad (25)$$

where we used  $S_H^* = N$  and  $S_M^* = M^*$  at the DFE. The spectral radius of  $FV^{-1}$  gives

$$\mathcal{R}_0 = \rho(FV^{-1}) = \sqrt{\frac{a^2 b_1 b_2 \Lambda^* b_{3M} \ell}{N \gamma (b_{3M} \ell + \mu) \mu^2}}, \quad (26)$$

which can be written in the classical two-factor host-vector form

$$\begin{aligned} \mathcal{R}_0^H &= \frac{ab_2}{\mu}; \quad \mathcal{R}_0^M = \frac{ab_1 \Lambda^*}{N \mu \gamma} \cdot \frac{b_{3M} \ell}{b_{3M} \ell + \mu} \\ \Rightarrow \mathcal{R}_0 &= \sqrt{\mathcal{R}_0^H \cdot \mathcal{R}_0^M}, \end{aligned} \quad (27)$$

where  $\mathcal{R}_0^H$  is the expected number of humans infected by a single infectious mosquito (mosquito infectious lifetime  $1/\mu$ ), and  $\mathcal{R}_0^M$  is the expected number of mosquitoes that survive the latent period and become infectious, generated by a single infectious human (human infectious period  $1/\gamma$ ). The geometric mean arises from the two-generation host-vector transmission cycle: the infection must complete a full human→mosquito→human loop, and the block structure of the next-generation matrix decouples into these two factors (see Appendix A.2 for the explicit multiplication).

## 6. Computational Implementation

### 6.1. Objective function and parameter bounds

Model parameters were estimated by minimizing the discrepancy between the simulated infected-human trajectory  $I_H(t)$  and the observed active-case counts reported through SIVEP-Malaria, restricted to rural Manaus (scaled by the 41.54% rural proportion). The objective function is a weighted mean squared error (wMSE):

$$\text{wMSE}(\theta) = \frac{\sum_{i=1}^n w_i (I_H(t_i; \theta) - y_i)^2}{\sum_{i=1}^n w_i}, \quad (28)$$

where  $y_i$  denotes the observed active-case count at date  $t_i$ ,  $I_H(t_i; \theta)$  is the model prediction obtained by linear interpolation of the simulated trajectory onto the observation dates, and  $\theta$  is the vector of calibrated parameters.

The weights  $w_i$  follow an exponential decay that prioritizes early-time dynamics within each year:

$$w_i = w_0 \exp(-\lambda d_i), \quad (29)$$

where  $d_i$  is the number of days from the start of the year for observation  $i$ ,  $w_0 = 5.0$  is the initial weight, and  $\lambda = 0.0015 \text{ day}^{-1}$  is the decay rate. This weighting scheme reflects the greater sensitivity of the early-season trajectory to initial conditions and parameter values, guiding the optimizer toward solutions that correctly capture the onset of the rainy-season transmission peak.

To prevent the optimizer from collapsing the infected-mosquito population to near-zero levels—which would sever the force-of-infection coupling and render the human transmission parameters structurally unidentifiable—an extinction penalty was added to the objective:

$$J(\theta) = \text{wMSE}(\theta) + \mathcal{P}(I_M^{\min}(\theta)), \quad (30)$$

where  $I_M^{\min}(\theta) = \min_t I_M(t; \theta)$  is the annual minimum of the infected-mosquito compartment, and

$$\mathcal{P}(I_M^{\min}) = \begin{cases} 0 & \text{if } I_M^{\min} \geq I_M^*, \\ B \left( 1 + \frac{I_M^* - I_M^{\min}}{I_M^*} \right) & \text{if } I_M^{\min} < I_M^*, \end{cases} \quad (31)$$

with  $I_M^* = 10$  (viability threshold) and  $B = 10^8$  (penalty base scale). The penalty scales linearly with the depth of extinction, ensuring that partially-extinct trajectories are ranked proportionally rather than receiving a uniform penalty that would flatten the objective landscape.

The nine calibrated parameters were grouped into four biological “topics” for sequential fitting:

1. **Humidity response** ( $H_0, k, \varphi$ ): the half-saturation threshold, sigmoid steepness, and baseline survival floor of the humidity-dependent survival function  $p(H)$ ;

**Table 1**

Parameter bounds for sequential topic calibration.

| Parameter                       | Lower bound | Upper bound        |
|---------------------------------|-------------|--------------------|
| $H_0$ (%)                       | 20          | 75                 |
| $k$ ( $\%^{-1}$ )               | 0.01        | 1.0                |
| $\varphi$                       | 0.01        | 0.5                |
| $\tau_H$ (days)                 | 10          | 20                 |
| $\gamma$ ( $\text{days}^{-1}$ ) | 1/150       | 1.0                |
| $\omega$ ( $\text{days}^{-1}$ ) | 1/300       | 0.1                |
| $b_1$                           | 0.001       | 0.5                |
| $b_2$                           | 0.01        | 0.5                |
| $M'$                            | 50,000      | $500 \times N_y^1$ |

<sup>1</sup>  $N_y$  denotes the estimated rural population of Manaus for year  $y$ .

2. **Human transmission** ( $\tau_H, \gamma, \omega$ ): the intrinsic incubation period, recovery rate, and immunity waning rate;
3. **Force of infection** ( $b_1, b_2$ ): the human-to-mosquito and mosquito-to-human transmission probabilities;
4. **Carrying capacity** ( $M'$ ): the baseline mosquito population scale that determines the environmental carrying capacity  $K(t)$ .

The parameter bounds are given in Table 1. The lower bound on  $H_0 = 20\%$  prevents the humidity response from collapsing (which would drive  $\mu \rightarrow \infty$  and extinguish the vector population). The bound on  $b_2 \geq 0.01$  ensures a minimum level of mosquito-to-human transmission, while  $\tau_H \in [10, 20]$  days constrains the human incubation period to the biologically plausible range for *P. vivax* [31].

### 6.2. Year-by-year sequential calibration

Parameters were calibrated sequentially for each year from 2017 to 2023 using the `lmfit` implementation of differential evolution (DE) [32]. The DE settings were: population multiplier `popsz` = 6, maximum function evaluations `max_nfev` = 800, convergence tolerance `tol` = 0.01, and random seed fixed at 42 for reproducibility.

Within each year, the four topics were fitted in the fixed order:

$$\underbrace{(H_0, k, \varphi)}_{\text{humidity}} \rightarrow \underbrace{(\tau_H, \gamma, \omega)}_{\text{human}} \rightarrow \underbrace{(b_1, b_2)}_{\text{FoI}} \rightarrow \underbrace{M'}_{\text{capacity}} \quad (32)$$

At each step, only the current topic’s parameters were free; all previously fitted parameters were held fixed at their most recent estimates. This sequential strategy reduces the dimensionality of each optimization sub-problem while respecting the biological ordering: the humidity response determines mosquito mortality, which constrains the infection cycle, which in turn determines the force of infection and carrying capacity.

After the sequential pass, a joint refinement was performed

using the Nelder-Mead simplex method [33], with all nine parameters free simultaneously, seeded from the sequential solution. A seed-evaluation guard was applied: after each DE run and after each refinement stage, the starting point was explicitly evaluated, and the optimizer's output was accepted only if it improved upon the seed. This prevents the stochastic DE from returning a solution worse than its initialization.

The model was calibrated sequentially across years, with the end-state vector of each year serving as the initial state for the next:

$$\mathbf{z}_0^{(y+1)} = \mathbf{z}_{\text{end}}^{(y)}, \quad (33)$$

where  $\mathbf{z} = (S_H, E_H, I_H, R_H, S_M, E_M, I_M)$  is the full state vector. The human infected compartment was always reset to the first observed active-case count for the new year:

$$I_{H,0}^{(y+1)} = \text{round}(y_{\text{first}}^{\text{obs}}), \quad (34)$$

while all other compartments were carried forward without modification. Negative end-state values were clamped to zero before carrying.

For the first year (2017), initial-condition fractions were explicitly fitted using Nelder-Mead with 300 function evaluations:

$$S_{H,0} = \alpha_S \cdot N, \quad E_{H,0} = \alpha_E \cdot N, \quad I_{M,0} = \alpha_I \cdot M_0, \quad (35)$$

where  $\alpha_S \in [0.3, 0.9]$ ,  $\alpha_E \in [0.001, 0.30]$ , and  $\alpha_I \in [0.0005, 0.15]$ , with the remaining compartments determined by conservation:  $R_{H,0} = N - S_{H,0} - E_{H,0} - I_{H,0}$ ,  $M_0 = 10N$ , and  $S_{M,0} = M_0 - E_{M,0} - I_{M,0}$ .

### 6.3. Weekly beta diagnostic

To evaluate where the climate-driven formulation deviates from what the data demands, a diagnostic calibration was performed in which the effective transmission coefficients were freed as per-week parameters instead of being derived from the biting rate and fixed  $b_1, b_2$ :

$$\lambda_H(t) = \beta_{h,w} \frac{I_M(t)}{N}, \quad \lambda_M(t) = \beta_{m,w} \frac{I_H(t)}{N}, \quad (36)$$

where  $w = \lfloor d/7 \rfloor$  is the week index and  $d$  is the day number within the year. This yields  $2 \times n_{\text{weeks}}$  free parameters per year (typically 106 for a 365-day year), with bounds  $\beta_{h,w}, \beta_{m,w} \in [10^{-9}, 3.0]$ .

All other model parameters were fixed at the values obtained from the yearly topic calibration. The initial state was reconstructed from the previous year's end-state, with  $I_H$  reset to observed data. The optimizer used was `lmfit.minimize` with the trust-region reflective method (`least_squares`), with `max_nfev = 1200` and `ftol = 10-8`.

The residual vector for the trust-region solver was constructed as:

$$r_i = \sqrt{w_i} (I_H(t_i) - y_i) + \sqrt{\mathcal{P}(I_M^{\min})}, \quad (37)$$

where  $\sqrt{\mathcal{P}}$  is injected as a constant offset to preserve the residual vector length while providing a gradient that pushes  $I_M$  upward when the mosquito population approaches the viability threshold.

The weekly beta fit serves as a diagnostic baseline: it reveals the best achievable fit when the transmission coefficients are unconstrained by the climate formulation. Persistent deviations between the weekly-optimal  $\beta_{h,w}$  and the climate-driven  $a(T) \cdot b_2$  indicate periods where the environmental model fails to capture the true transmission dynamics.

### 6.4. Freeing the transmission coefficients

There are three main observations that allow us to use the effective transmission coefficients  $\beta_{h,w}$  and  $\beta_{m,w}$  as per-week free parameters:

**(i) The transmission coefficients are intrinsically time-varying:** In the mechanistic formulation, the forces of infection are given by  $\lambda_H = a(T) b_2 I_M/N$  and  $\lambda_M = a(T) b_1 I_H/N$ , where the biting rate  $a(T)$  is the only environmental factor and the per-bite probabilities  $b_1$  and  $b_2$  are assumed to be constants. This assumption is biologically restrictive because these coefficients summarize several processes that may vary over the transmission season, including parasite development within the vector, vector susceptibility to infection, vector to host contact, and host infectiousness. These processes can be affected by temperature, rainfall, humidity, and changes in vector and host demography [19, 39]. We therefore relax the assumption that  $b_1$  and  $b_2$  remain constant throughout the study period and allow the data to determine how transmission changes over time.

**(ii) The weekly fit provides a nonparametric estimate of the latent transmission process:** We represent the effective transmission coefficients as latent time-varying functions,  $\beta_h(t)$  and  $\beta_m(t)$ . These functions are approximated as piecewise constant over weekly intervals:

$$\beta_h(t) = \beta_{h,w}, \quad \beta_m(t) = \beta_{m,w}, \quad t \in [7w, 7(w+1))$$

This provides a flexible finite-dimensional representation of possible transmission profiles. The weekly coefficients are estimated by minimizing the weighted residual between the model predictions and the observed malaria cases. Thus, rather than imposing a specific relationship between transmission and environmental covariates, the weekly model provides a flexible, data-driven estimate of the temporal variation in transmission.

The resulting  $2 \times n_{\text{weeks}}$  coefficients don't represent independent biological parameters to be estimated, but rather

provide a flexible empirical representation of the transmission process. The additional degrees of freedom allow the model to capture temporal changes in transmission that a more constrained model may miss, providing a reference for evaluating the proposed mechanistic model.

**(iii) The climate-driven model is nested within the weekly model:** The climate-driven model is obtained by imposing the restriction

$$\beta_{h,w} = a(T_w)b_2, \quad \beta_{m,w} = a(T_w)b_1, \quad (38)$$

where  $b_1$  and  $b_2$  are constant over the entire study period. Consequently, the climate-driven formulation is a restricted case of the more flexible weekly model. Because all other model parameters and initial conditions are held fixed, the difference in fit between the two formulations directly measures the extent to which the constant-coefficient assumption limits the model's ability to reproduce the observed cases.

Releasing the coefficients reduces the wMSE by one to two orders of magnitude relative to the climate-driven model. This substantial improvement indicates that the data contain temporal variation in transmission that cannot be adequately represented by constant  $b_1$  and  $b_2$ .

The weekly estimates are therefore used as a diagnostic benchmark rather than as the final parameterization of the model. They identify the temporal transmission pattern that a parsimonious mechanistic formulation should seek to explain. This approach preserves the distinction between estimating a flexible empirical transmission process and identifying the underlying biological parameters, as the weekly coefficients characterize the observed transmission signal, while the subsequent structural model determines how that signal can be explained by environmental drivers.

Finally, the weekly estimates provide the link between the empirical transmission signal and the environmental model. Since the biting rate  $a(T_w)$  is explicitly modeled as a function of temperature, the corresponding weekly per-bite transmission probabilities can be recovered as

$$b_{1,w} = \frac{\beta_{m,w}}{a(T_w)}, \quad b_{2,w} = \frac{\beta_{h,w}}{a(T_w)}. \quad (39)$$

These time-varying estimates represent the per-bite transmission probabilities implied by the observed case dynamics after accounting for the temperature-dependent biting rate. They are subsequently used as empirical targets for regression models relating  $b_1$  and  $b_2$  to environmental drivers. In this way, the weekly fit serves as an intermediate step in the modeling chain

$$\text{environment} \longrightarrow (b_1, b_2) \longrightarrow \text{transmission} \longrightarrow \text{cases},$$

allowing a flexible, data-driven estimate of transmission to be translated into a simplified and mechanistically interpretable environmental model.

## 6.5. Numerical solver and implementation

All ODE simulations were performed using Python version 3.12.7 (conda-forge distribution), using the LSODA solver as implemented in `scipy.integrate.solve_ivp` (SciPy version 1.17.1), with automatic stiffness detection and order switching. The integration was evaluated on a fine grid of 10 points per day ( $\Delta t = 0.1$  day), with the solution interpolated back to daily resolution via linear interpolation. End-state vectors were clamped to zero to prevent negative compartment values from propagating through the year chain.

Climate inputs (temperature, precipitation, minimum humidity) were smoothed using a 14-day rolling mean for mosquito demographic rates (mortality  $\mu$ , sporogonic development  $b_{3M}$ , birth rate  $\Lambda$ ), while the biting rate  $a(T)$  used raw daily temperature to preserve high-frequency transmission variability.

Data manipulation and analysis were performed using Pandas version 3.0.2, while numerical computations and array operations were carried out with NumPy version 2.4.2.

All computations were performed on a laptop running Windows 11 (64-bit) equipped with a 12th Gen Intel Core i5-1235U processor (10 cores, 12 threads) and 16 GB RAM.

## 7. Results

### 7.1. Reproduction of the original Parham & Michael model as a negative baseline

As a diagnostic for the methodological choices adopted below, the original SIR-SEI formulation of Parham and Michael (2010) [18] was implemented with all parameter values reported in the original study, while population sizes and initial epidemiological conditions were estimated from the available rural Manaus data (Appendix Fig. 1A). Under this parameterization the infected human population declines monotonically and the mosquito population rapidly collapses to extinction, indicating that mosquito recruitment is insufficient to replace mortality. Because transmission depends on the persistence of infectious vectors, the collapse of the vector population eliminates infection in both host and vector populations. Although the delay formulation preserved positivity, it fails to reproduce the endemic behavior expected for malaria in the Amazon, confirming the need to recalibrate the environmental and transmission parameters as described in Section 3.4 and Section 6.2.

### 7.2. Recalibration of the temperature-dependent biting rate

The temperature-dependent biting rate was recalibrated against field estimates of *Anopheles darlingi* feeding behavior following Section 3.4. The recalibrated function yielded a minimum biting threshold of  $T' = 16^\circ\text{C}$  and a thermal scaling factor  $D_1 = 24.8^\circ\text{C days}$ . Compared with the original Parham & Michael parameterization, the refit shifts

**Table 2**

Year-by-year calibrated parameters and weighted mean squared error (wMSE) of the SEIRS-SEI model, 2017–2023.

| Year | $H_0$ | $k$   | $\varphi$ | $\tau_H$ | $\gamma$ | $\omega$ | $b_1$ | $b_2$ | $M'$   | wMSE   |
|------|-------|-------|-----------|----------|----------|----------|-------|-------|--------|--------|
| 2017 | 37.58 | 0.705 | 0.215     | 19.81    | 0.0071   | 0.0855   | 0.259 | 0.035 | 50,723 | 85,036 |
| 2018 | 20.36 | 0.046 | 0.490     | 20.00    | 0.0073   | 0.0961   | 0.487 | 0.022 | 51,245 | 64,959 |
| 2019 | 22.13 | 0.053 | 0.493     | 19.96    | 0.0072   | 0.0744   | 0.488 | 0.022 | 50,474 | 41,436 |
| 2020 | 20.13 | 0.039 | 0.500     | 19.98    | 0.0067   | 0.0422   | 0.500 | 0.019 | 50,024 | 23,541 |
| 2021 | 20.02 | 0.043 | 0.493     | 19.99    | 0.0067   | 0.0773   | 0.500 | 0.019 | 50,015 | 39,449 |
| 2022 | 20.02 | 0.043 | 0.493     | 19.77    | 0.0099   | 0.0312   | 0.500 | 0.019 | 50,832 | 25,046 |
| 2023 | 20.14 | 0.207 | 0.431     | 19.86    | 0.0081   | 0.0310   | 0.485 | 0.010 | 53,616 | 18,360 |

the biting response toward the thermal range observed in the Amazon and modifies its sensitivity near the transmission window. All subsequent simulations and the weekly transmission regressions of Section 7.5 make use of this recalibrated biting rate.

### 7.3. Model calibration reproduces the observed transmission dynamics

The SEIRS-SEI model was calibrated year by year over 2017–2023 following the sequential topic scheme of Section 6.2, and the recovered parameters are summarized in Table 2. Figure 1 shows the fitted infectious compartment  $I_H$  against the observed active-case count alongside the weekly free-beta fit for 2017, which exhibited the largest relative improvement of the weekly model over the constant-coefficient formulation (Section 7.4). The calibrated parameters remain within their physiologically prescribed bounds found in Table 1, with the humidity response ( $H_0$ ,  $k$ ,  $\varphi$ ) and the human recovery parameters ( $\tau_H$ ,  $\gamma$ ,  $\omega$ ) showing limited cross-year variation, while the force-of-infection coefficients  $b_1$  and  $b_2$  consistently approach their upper and lower bounds, respectively (Table 2).




Figure 1: Comparison between the observed active cases and the model estimation of the infectious human compartment  $I_H$  for 2017.

### 7.4. The constant-coefficient climate model underfits the observed dynamics

To assess how much of the seasonal variation in transmission is captured by the mechanistic, climate-driven formulation, we freed the effective transmission coefficients, allowing them to vary on a weekly basis (Section 6.4), and compared the resulting weekly model with the constant-coefficient climate-driven model. Releasing the coefficients reduced the weighted mean squared error in every year (Table 3),

**Table 3**

Weighted mean squared error (wMSE) of the climate-driven constant-coefficient model versus the free weekly model, per year.

| Year | Climate-driven | Weekly free | Reduction factor |
|------|----------------|-------------|------------------|
| 2017 | 85,036         | 618         | 137.6×           |
| 2018 | 64,959         | 2,553       | 25.4×            |
| 2019 | 41,436         | 561         | 73.9×            |
| 2020 | 23,541         | 359         | 65.6×            |
| 2021 | 39,449         | 1,552       | 25.4×            |
| 2022 | 25,046         | 1,711       | 14.6×            |
| 2023 | 18,360         | 7,598       | 2.4×             |

most notably in 2017, where the wMSE fell from  $8.5 \times 10^4$  to  $6.2 \times 10^2$  (a  $\sim 137$ -fold reduction).

The residual between the free weekly  $\beta_{h,w}$ ,  $\beta_{m,w}$  and the climate-driven  $a(T)b_1$ ,  $a(T)b_2$  indicates periods in which the temperature-driven formulation fails to capture the observed transmission structure. This motivates extending the per-bite transmission probabilities to the wider set of environmental drivers introduced in Section 2.

### 7.5. Environmental regression of the transmission probabilities $b_1$ and $b_2$

Following the framework of Section 2, the freed weekly per-bite transmission probabilities were regressed against the environmental predictors  $T$ ,  $R$ ,  $H$ ,  $D_f$ ,  $F_c$ ,  $F_t$  (temperature, precipitation, humidity, deforestation, fire counts and fire trends) fitting an independent random forest for each year.

Within each year the model explains a substantial share of the variance of both transmission probabilities, with mean  $R^2 = 0.685 \pm 0.082$  for  $b_1$  and mean  $R^2 = 0.784 \pm 0.130$  for  $b_2$ , with near-zero prediction bias, with  $|\text{bias}| \lesssim 0.003$  in every year (Table 4, Figs. 7.5,2).




Figure 2: Best within-year random-forest fit of the freed weekly per-bite transmission  $b_1$  and  $b_2$ , in 2022 and 2020 respectively.

The red-dotted line denotes the weekly coefficients recovered from the observed case dynamics (Section 6.4); while the blue-dashed line is the mean random-forest prediction. The model reproduces the seasonal envelope while retaining the week-to-week structure in the data.

The importance of the predictors, however, differs between the two probabilities. For  $b_1$  the contribution is distributed fairly evenly among precipitation (0.203), temperature (0.201), humidity (0.194), fire trends (0.179) and fire counts (0.149); while for  $b_2$  the leading contributors are fire counts (0.213), humidity (0.171), temperature (0.169), fire trends (0.166) and precipitation (0.163). Deforestation  $D_f$  is consistently the weakest predictor for both  $b_1$  (0.074) and  $b_2$  (0.119), indicating that its effect is not captured at the intra-annual timescale resolved here.

A leave-one-year-out cross-validation of the same forests qualifies this result. Out-of-sample performance is markedly lower, with overall  $R^2 \approx 0.17$  for  $b_1$  and  $\approx 0.19$  for  $b_2$ , in contrast to the strong within-year fits. This contrast confirms that the environmental drivers are strongly year-specific: the within-year seasonal trajectory of  $T, R, H, D_f, F_c, F_t$  shapes each year's transmission signal, whereas a fixed, year-independent link between the environment and  $b_1, b_2$  is not supported by the data.

The shape of these responses is illustrated in Figs. 4,5, which shows the partial dependence of  $b_1$  and  $b_2$  on each predictor for the best fitted year for each transmission probability.




Table 4

Year-by-year within-year random forest performance for  $b_1$  and  $b_2$ .

|       | Year | $R^2$ | MAE    | Bias    |
|-------|------|-------|--------|---------|
| $b_1$ | 2017 | 0.563 | 0.0594 | +0.0021 |
|       | 2018 | 0.580 | 0.0651 | -0.0005 |
|       | 2019 | 0.739 | 0.0475 | +0.0011 |
|       | 2020 | 0.671 | 0.0626 | +0.0021 |
|       | 2021 | 0.705 | 0.0603 | -0.0009 |
|       | 2022 | 0.810 | 0.0503 | +0.0010 |
|       | 2023 | 0.724 | 0.0107 | +0.0006 |
| $b_2$ | 2017 | 0.531 | 0.0182 | +0.0011 |
|       | 2018 | 0.825 | 0.0103 | +0.0005 |
|       | 2019 | 0.712 | 0.0120 | -0.0002 |
|       | 2020 | 0.915 | 0.0105 | -0.0006 |
|       | 2021 | 0.712 | 0.0122 | +0.0001 |
|       | 2022 | 0.905 | 0.0103 | -0.0002 |
|       | 2023 | 0.889 | 0.0021 | +0.0001 |

Figure 4: Partial dependence of the weekly per-bite transmission probability  $b_1$  on each environmental predictor.




Figure 5: Partial dependence of the weekly per-bite transmission probability  $b_2$  on each environmental predictor.

Columns correspond to temperature  $T$ , precipitation  $R$ , humidity  $H$ , deforestation  $D_f$ , fire counts  $F_c$  and fire trends  $F_t$ . Curves give the mean predicted probability as each predictor is varied over its observed range while the remaining variables are held at their marginal mean.

All files of data collection and treatment, as well as resulting plots created during throughout the development of this work are available at <https://github.com/RaphaLevy/----->.

## 8. Discussion

The population estimates provided by the Trajetórias Project [6] for inter-census periods are derived using fixed rural-urban proportions based on the 2010 census. While this approach ensures internal consistency, it does not account for the significant demographic changes observed in Manaus over time, particularly those associated with rapid urbanization.

To address this limitation, the population proportions used in this study were recalculated (see Section 4.3) using interpolation and extrapolation based on available census data. This adjustment provides a more realistic

representation of temporal changes in the rural population, which is especially relevant given the disproportionate contribution of rural areas to malaria incidence.

These refinements improve the consistency between demographic inputs and epidemiological dynamics in the model, reducing potential biases in the estimation of transmission intensity and enhancing the reliability of the results.

## References

- [1] P. F. P. Pimenta et al., An overview of malaria transmission from the perspective of Amazon *Anopheles* vectors, *Memórias do Instituto Oswaldo Cruz*, Vol. 110 (1) (2015), 23-47, <https://doi.org/10.1590/0074-02760140266>.
- [2] G. A. Josling, K. C. Williamson, M. Llinás, Regulation of Sexual Commitment and Gametocytogenesis in Malaria Parasites, *Annual Review of Microbiology*, Vol. 72 (1) (2018), 501-519, <https://doi.org/10.1146/annurev-micro-090817-062712>.
- [3] World Health Organization / World Malaria Report 2022, Global Malaria Programme, accessed on 29 August 2024, <https://www.who.int/publications/i/item/9789240064898>.
- [4] M. da Silva-Nunes et al., Malaria on the Amazonian frontier: transmission dynamics, risk factors, spatial distribution, and prospects for control, *American Journal of Tropical Medicine and Hygiene*, Vol. 79 (4) (2008), 624-635, <https://doi.org/10.4269/ajtmh.2008.79.624>.
- [5] M. da Silva-Nunes et al., Amazonian malaria: Asymptomatic human reservoirs, diagnostic challenges, environmentally driven changes in mosquito vector populations, and the mandate for sustainable control strategies, *Acta Tropica*, Vol. 121 (3) (2012), 281-291, <https://doi.org/10.1016/j.actatropica.2011.10.001>.
- [6] A. C. Rorato, A. P. Dal'Asta, R. M. Lana et al., Trajetórias: a dataset of environmental, epidemiological, and economic indicators for the Brazilian Amazon, *Scientific Data*, Vol. 10 (1) (2023), 65, <https://doi.org/10.1038/s41597-023-01962-1>.
- [7] D. E. Norris, Mosquito-borne Diseases as a Consequence of Land Use Change, *EcoHealth*, Vol. 1 (1) (2004), 19-24, <https://doi.org/10.1007/s10393-004-0008-7>.
- [8] Anopheles, Fiocruz Rondônia, accessed on 15 August 2024, <https://www.rondonia.fiocruz.br/pivem/anopheline/>.
- [9] M. da Silva-Nunes, Environmental changes impact in malaria transmission and prospects for the disease control in Brazilian Amazon rural settlements, *Oecologia Australis*, Vol. 14 (3) (2010), 603-622, <https://revistas.ufrj.br/index.php/oa/article/view/7101>.
- [10] L. S. M. Chaves, E. S. Bergo, J. E. Conn, G. Z. Laporta, P. R. Prist, M. A. M. Sallum, Anthropogenic landscape decreases mosquito biodiversity and drives malaria vector proliferation in the Amazon rainforest, *PLOS ONE*, Vol. 16 (1) (2021), <https://doi.org/10.1371/journal.pone.0245087>.
- [11] K. R. Kirby, W. F. Laurance, A. K. L. M. Albernaz, G. Schroth, P. M. Fearnside, S. Bergen, E. M. Venticinque, C. Da Costa, The future of deforestation in the Brazilian Amazon, *Futures*, Vol. 38 (4) (2006), 432-453, <https://doi.org/10.1016/j.futures.2005.07.011>.
- [12] C. F. Lima, F. T. P. Torres, L. J. Minette, F. A. Lima, R. C. A. Lima et al, Is there a relationship between forest fires and deforestation in the Brazilian Amazon?, *PLOS ONE*, Vol. 19 (6) (2024), <https://doi.org/10.1371/journal.pone.0306238>.
- [13] P. M. Fearnside, Conservation Policy in Brazilian Amazonia: Understanding the Dilemmas, *World Development*, Vol. 31 (5) (2003), 757-779, [https://doi.org/10.1016/S0305-750X\(03\)00011-1](https://doi.org/10.1016/S0305-750X(03)00011-1).
- [14] TerraBrasilis, DETER (Notices), Analyses - Amazônia Biome, Daily notices of DETER project area, accessed on 17 March 2026, <https://terrabrasilis.dpi.inpe.br/app/dashboard/alerts/biomes/amazonia-nb/daily/>.

- [15] Data\_Basis.org, Fire Database, National Institute for Space Research (INPE), accessed on 17 March 2026, <https://data-basis.org/dataset/f06f3cdc-b539-409b-b311-1ff8878fb8d9?table=a3696dc2-4dd1-4f7e-9769-6aa16a1556b8>.
- [16] MapBiomias Brasil, Home/Iniciativas e Produtos/Fogo/Mapeamento anual/Mensal, accessed on 30 August 2026, <https://plataforma.brasil.mapbiomas.org/>.
- [17] Mosqlimate Data API / Datastore / climate, accessed on 12 August 2024, <https://api.mosqlimate.org/datastore/>.
- [18] P. E. Parham, E. Michael, Modelling Climate Change and Malaria Transmission, Modelling Parasite Transmission and Control, Advances in Experimental Medicine and Biology, Vol. 673 (2010), 184-199, [https://doi.org/10.1007/978-1-4419-6064-1\\_13](https://doi.org/10.1007/978-1-4419-6064-1_13).
- [19] E. A. Mordecai et al, Optimal temperature for malaria transmission is dramatically lower than previously predicted, Ecology Letters, Vol. 16 (1), 22-30 (2013), <https://pubmed.ncbi.nlm.nih.gov/23050931/>.
- [20] S.E. Eikenberry, A. B. Gumel, Mathematical modeling of climate change and malaria transmission dynamics: a historical review, Journal of Mathematical Biology, Vol. 77 (4), 857-933 (2018), <https://pubmed.ncbi.nlm.nih.gov/29691632/>.
- [21] F. S. M.de Barros, N. A. Honório, M. E. Arruda, Survivorship of Anopheles darlingi (Diptera: Culicidae) in Relation with Malaria Incidence in the Brazilian Amazon, PLoS ONE, Vol. 6 (8), e22388 (2011), <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0022388>.
- [22] F. J. Lardeux et al, A physiological time analysis of the duration of the gonotrophic cycle of Anopheles pseudopunctipennis and its implications for malaria transmission in Bolivia, Malaria Journal, Vol. 7, 141 (2008), <https://link.springer.com/article/10.1186/1475-2875-7-141>.
- [23] J. Oliveira-Ferreira, M. V. Lacerda, P. Brasil et al., Malaria in Brazil: an overview., Malaria Journal, Vol. 9 (1) (2010), <https://doi.org/10.1186/1475-2875-9-115>.
- [24] C. T. Codeço, A. P. Dal'Asta, A. C. Rorato, R. M. Lana, T. C. Neves, C. S. Andreazzi, M. Barbosa, M. I. S. Escada, D. A. Fernandes, D. L. Rodrigues, I. C. Reis, M. da Silva-Nunes, A. B. Gontijo, F. C. Coelho, A. M. V. Monteiro, Epidemiology, Biodiversity, and Technological Trajectories in the Brazilian Amazon: From Malaria to COVID-19, Frontiers in Public Health, Vol. 9 (2021), <https://doi.org/10.3389/fpubh.2021.647754>.
- [25] A. K. R. Galardo, M. Arruda, et al., Malaria vector incrimination in three rural riverine villages in the Brazilian Amazon, The American Journal of Tropical Medicine and Hygiene, Vol. 76 (3) (2007), 461-469, <https://www.ajtmh.org/view/journals/tpmd/76/3/article-p461.xml>.
- [26] SIVEP Malaria, accessed on 10 October 2024, <https://files.deeplearn.ltd/a/hnjvpj>.
- [27] Biblioteca IBGE, 2000, accessed on 11 November 2024, <https://biblioteca.ibge.gov.br/index.php/biblioteca-catalogo?view=detalhes&id=7308>.
- [28] IBGE, Censo Demográfico, 2010, accessed on 11 November 2024, <https://www.ibge.gov.br/estatisticas/sociais/populacao/9662-censo-demografico-2010.html?=&t=resultados>.
- [29] IBGE, Brasil / Amazonas / Manaus, 2022, accessed on 11 November 2024, <https://cidades.ibge.gov.br/brasil/am/manaus/panorama>.
- [30] P. van den Driessche, J. Watmough, Reproduction numbers and sub-threshold endemic equilibria for compartmental models of disease transmission, Mathematical Biosciences, Vol. 180 (2002), 29-48, [https://doi.org/10.1016/S0025-5564\(02\)00108-6](https://doi.org/10.1016/S0025-5564(02)00108-6).
- [31] P. Brasil et al, Unexpectedly long incubation period of Plasmodium vivax malaria, in the absence of chemoprophylaxis, in patients diagnosed outside the transmission area in Brazil, Malaria Journal, Vol. 10 (122) (2011), <https://link.springer.com/article/10.1186/1475-2875-10-122>.
- [32] R. Storn, K. Price, Differential Evolution - A Simple and Efficient Heuristic for Global Optimization over Continuous Spaces, Journal of Global Optimization, Vol. 11 (1997), 341-359, <https://link.springer.com/article/10.1023/A:1008202821328>.
- [33] J. A. Nelder, R. Mead, A Simplex Method for Function Minimization, The Computer Journal, Vol. 7 (4) (1965), 308-313, <https://academic.oup.com/comjnl/article-abstract/7/4/308/354237>.
- [34] G. C. McCord, Malaria ecology and climate change, The European Physical Journal Special Topics, Vol. 225 (3) (2016), 459-470, <https://doi.org/10.1140/epjst/e2015-50097-1>.
- [35] M. H. Craig, R. W. Snow, D. le Sueur, A Climate-based Distribution Model of Malaria Transmission in Sub-Saharan Africa, Parasitology Today, Vol. 15 (3), 105-111 (1999), <https://pubmed.ncbi.nlm.nih.gov/10322323/>.
- [36] T. P. Agyekum et al, A Systematic Review of the Effects of Temperature on Anopheles Mosquito Development and Survival: Implications for Malaria Control in a Future Warmer Climate, International Journal of Environmental Research and Public Health, Vol. 18 (14), (2021), <https://pmc.ncbi.nlm.nih.gov/articles/PMC8306597/>.
- [37] S. W. Lindsay, M. H. Birley, Climate change and malaria transmission, Annals of Tropical Medicine & Parasitology, Vol. 90 (1996), 573-588, <https://doi.org/10.1080/00034983.1996.11813087>.
- [38] N. J. Arisco, C. Peterka, C. Diniz, B. H. Singer, M. C. Castro, Ecological change increases malaria risk in the Brazilian Amazon, Proc. Natl. Acad. Sci., Vol. 121 (44) (2024), <https://doi.org/10.1073/pnas.2409583121>.
- [39] T. S. Churcher, J. F. Trape, A. Cohuet, Human-to-mosquito transmission efficiency increases as malaria is controlled, Nat Commun 6, 6054 (2015), <https://doi.org/10.1038/ncomms7054>.
- [40] Tabnet, Ministério da Saúde, accessed on 12 August 2024, <http://tabnet.datasus.gov.br/cgi/tabcgi.exe?ibge/cnv/popam.def>.
- [41] J. D. Charlwood, Biological variation in Anopheles darlingi Root, Memórias do Instituto Oswaldo Cruz, Vol. 91 (4) (1996), 391-398, <https://doi.org/10.1590/S0074-02761996000400001>.
- [42] G. R. de Lima, A. N. Arcos, E.V. dos Santos, R. C. Simões, C.A.P. Lima, W. P. Tadei, Anopheles Larval Abundance in Artificial Breeding Sites in the East Zone of Manaus, Amazon, South American Journal of Basic Education, Technical and Technological, Vol. 8 (1) (2021), 35-47, <https://periodicos.ufac.br/index.php/SAJEBTT/article/view/3399>.
- [43] C. M. Souza, Jr., J. Z. Shimbo, M. R. Rosa, L. L. Parente, A. A. Alencar, B. F. T. Rudorff, H. Hasenack, M. Matsumoto, L. G. Ferreira, P. W. M. Souza-Filho, et al, Reconstructing Three Decades of Land Use and Land Cover Changes in Brazilian Biomes with

Landsat Archive and Earth Engine, Remote Sensing, Vol.12 (17) (2020), <https://doi.org/10.3390/rs12172735>.

- [44] A. J. MacDonald, E. A. Mordecai, Amazon deforestation drives malaria transmission, and malaria burden reduces forest clearing, *Proceedings of the National Academy of Sciences*, Vol. 116 (44) (2019), <https://doi.org/10.1073/pnas.1905315116>.
- [45] L. Ziccardi, A. Yanai, P. Fearnside, S. Stark, Deforestation in the Brazilian Amazon: Threats and opportunities, *Research Square*, PREPRINT (Version 1) (2024), <https://doi.org/10.21203/rs.3.rs-4591425/v1>.
- [46] G1, M. Castro, "Amazonas registra 21,6 mil queimadas em 2024 e tem o pior índice em 26 anos, aponta Inpe", accessed on 28 September 2024, <https://g1.globo.com/am/amazonas/noticia/2024/09/24/amazonas-registra-216-mil-queimadas-em-2024-e-tem-o-pior-indice-em-26-anos-aponta-inpe.gh.html>.
- [47] A. Abdelrazec, A. B. Gumel, Mathematical assessment of the role of temperature and rainfall on mosquito population dynamics, *Journal of Mathematical Biology*, Vol. 74 (6), 1351-1395 (2017), <https://link.springer.com/article/10.1007/s00285-016-1054-9>.
- [48] OpenLearnCreate. Courses/Collections/Health Education and Training (HEAT)/Communicable Diseases/Contents/Communicable Diseases Module: 6. Factors that Affect Malaria Transmission, accessed on 02 May 2026, <https://www.open.edu/openlearncreate/mod/oucontent/view.php?id=89&printable=1>.

## A. Appendix

### A.1. Tables

**Table A.1**

List of model parameters.

| Parameter            | Definition                                                                     | Formulation                                                   |
|----------------------|--------------------------------------------------------------------------------|---------------------------------------------------------------|
| $p(T)$               | Daily mosquito survival rate dependent on temperature                          | $e^{(-1/(AT^2+BT+C))}$                                        |
| $p(H)$               | Daily mosquito survival rate dependent on humidity <sup>2</sup>                | $\frac{\varphi e^{-k(H-H_0)+1}}{e^{-k(H-H_0)+1}+1}$           |
| $p(T, H)$            | Daily mosquito survival rate                                                   | $p(T)p(H)$                                                    |
| $p_L(R)$             | Probability of larval survival dependent on rainfall <sup>3</sup>              | $(4p_{ML}/R_L^2)R(R_L - R)$                                   |
| $p_L(T)$             | Probability of larval survival dependent on temperature                        | $e^{-(c_1T+c_2)}$                                             |
| $p_L(R, T)$          | Probability of larval survival                                                 | $p_L(R)p_L(T)$                                                |
| $\mu(T, H)$          | Mosquito mortality rate per capita (days <sup>-1</sup> )                       | $-\log(p(T, H))$                                              |
| $a(T)$               | Biting rate (days <sup>-1</sup> )                                              | $(T - T')/D_1$                                                |
| $\tau_M(T)$          | Duration of sporozoite cycle (days)                                            | $DD/(T - T_{\min})$                                           |
| $\tau_L(T)$          | Duration of larval development phase (days)                                    | $1/(c_1T + c_2)$                                              |
| $\ell(\tau_M)(T, H)$ | Probability of mosquito survival during sporozoite cycle (days <sup>-1</sup> ) | $p(T, H)^{\tau_M(T)}$                                         |
| $b(R, T)$            | Mosquito birth rate (days <sup>-1</sup> )                                      | $B_E p_E(R) p_L(R, T) p_P(R) / (\tau_E + \tau_L(T) + \tau_P)$ |
| $M(t)$               | Total population of mosquitos                                                  | $S_M(t) + E_M(t) + I_M(t)$                                    |
| $N(t)$               | Total population of humans                                                     | $S_H(t) + E_H(t) + I_H(t) + R_H(t)$                           |

<sup>2</sup>  $p(H)$  was added in order to take into consideration the deleterious effects of humidity. Unlike for  $T$  and  $R$ , which use the daily average value of temperature and precipitation, respectively,  $H$  uses the daily minimal value of humidity.

<sup>3</sup>  $p_E(R)$  and  $p_P(R)$  have an analogous expression to  $p_L(R)$ , substituting  $p_{ML}$  for  $p_{ME}$  and  $p_{MP}$  respectively.

**Table A.2**

List of model variables.

| Symbol              | Definition                                                                                                | Units                                |
|---------------------|-----------------------------------------------------------------------------------------------------------|--------------------------------------|
| $b_1$               | Proportion of bites from susceptible mosquitoes on infected humans that result in infection: 0.04         | Dimensionless                        |
| $b_2$               | Proportion of bites from infected mosquitoes on susceptible humans that result in infection: 0.09         | Dimensionless                        |
| $\gamma$            | 1/Average duration of infectiousness in humans: 1/120                                                     | days <sup>-1</sup>                   |
| $\omega$            | 1/Average duration of protective immunity after infection: 1/270                                          | days <sup>-1</sup>                   |
| $B_E$               | Number of eggs laid per adult per oviposition: 200                                                        | Dimensionless                        |
| $p_{ME}$            | Maximum probability of egg survival: 0.9                                                                  | Dimensionless                        |
| $p_{ML}$            | Maximum probability of larval survival: 0.25                                                              | Dimensionless                        |
| $p_{MP}$            | Maximum probability of pupal survival: 0.75                                                               | Dimensionless                        |
| $\tau_E$            | Duration of the egg development phase: 1                                                                  | days                                 |
| $\tau_P$            | Duration of the pupal development phase: 1                                                                | days                                 |
| $\tau_H$            | Latent period of infection within humans: 10                                                              | days                                 |
| $R_L$               | Rainfall threshold until breeding sites are eliminated, removing immature individuals: 32.26 <sup>4</sup> | mm                                   |
| $T_{\min}$          | Minimum temperature, below which there is no development of the parasite: 14.5                            | °C                                   |
| $T_{\max}$          | Maximum temperature, above which there is no development of the parasite: 29.8 [34]                       | °C                                   |
| $T_{\text{opt}}$    | Optimal temperature for development of the parasite: 23.5 [34]                                            | °C                                   |
| $T_{\min A.}/T'$    | Minimum temperature, below which there is no development of the vector: 16 <sup>5</sup> [19, 35]          | °C                                   |
| $T_{\max A.}$       | Maximum temperature, above which there is no development of the vector: 34 [19, 35, 36]                   | °C                                   |
| $T_{\text{op } A.}$ | Optimal temperature for transmission: 34 [19]                                                             | °C                                   |
| $DD$                | Degree-days for parasite development. "Sum of heat" for maturation: 105 [34, 37]                          | °C days                              |
| $A$                 | Empirical sensitivity parameter: -0.03                                                                    | (°C <sup>2</sup> days) <sup>-1</sup> |
| $B$                 | Empirical sensitivity parameter: 1.31                                                                     | (°C days) <sup>-1</sup>              |
| $C$                 | Empirical sensitivity parameter: -4.4                                                                     | days <sup>-1</sup>                   |
| $D_1$               | Temperature scaling factor of the biting rate: 24.8                                                       | °C days                              |
| $c_1$               | Empirical sensitivity parameter: 0.00554                                                                  | (°C days) <sup>-1</sup>              |
| $c_2$               | Empirical sensitivity parameter: -0.06737                                                                 | days <sup>-1</sup>                   |
| $b_{3M}$            | Infection rate in exposed mosquitoes ( $1/\tau_M(T)$ )                                                    | days <sup>-1</sup>                   |
| $b_{3H}$            | Infection rate in exposed humans ( $1/\tau_H$ ): 1/10                                                     | days <sup>-1</sup>                   |
| $H_0$               | Critical humidity threshold for 50% survival transition: 58                                               | %                                    |
| $\varphi$           | Baseline survival probability at extreme low humidity: 0.05                                               | Dimensionless                        |
| $k$                 | Steepness of the humidity-survival sigmoid response: 0.25                                                 | % <sup>-1</sup>                      |

<sup>4</sup> The reference rainfall level  $R_L$  for the Amazon was approximated based on reported associations between monthly precipitation and malaria incidence [38]. In particular, precipitation levels in the range of 917 to 1019 mm/month were associated with a decrease in malaria cases. The midpoint of this interval (968 mm/month) was used as a representative value and converted to a daily rate by dividing by 30 days, yielding  $R_L \approx 32.3$  mm/day. This value is therefore interpreted as the threshold scale for high rainfall conditions in the model.

<sup>5</sup> In Parham and Michael [18], this parameter is denoted by  $T_1$ . Here, we adopt a different notation to avoid confusion with the mean temperature in the absence of seasonality, also denoted by  $T_1$  in the original paper. Additionally, instead of defining an independent lower thermal threshold for the biting rate, we assume that biting activity becomes negligible below the minimum temperature required for vector development.

## A.2. Derivation of the basic reproduction number

This appendix details the next-generation matrix (NGM) computation summarised in Section 5.2. The infected compartments are ordered as  $(E_H, I_H, E_M, I_M)$ , so  $m = 4$ , and all parameters are as defined in Section 3.

*Disease-free states and the NGM formalism:* Following van den Driessche and Watmough [30], let  $X_s$  be the set of all disease-free states:

$$X_s = \{x \geq 0 \mid x_i = 0, i = 1, \dots, m\},$$

where  $X = (x_1, \dots, x_n)^\top$ ,  $x_i \geq 0$  is the number of individuals in compartment  $i$ , and every function is assumed to be continuously differentiable at least twice in each variable ( $C^2$ ).

Rearranging the model equations so that the first  $m$  components are the infected compartments, let  $\mathcal{F}_i(x)$  be the rate of appearance of new infections in compartment  $i$ ,  $\mathcal{V}_i^+(x)$  the rate of transfer into compartment  $i$  by all other means, and  $\mathcal{V}_i^-(x)$  the rate of transfer out of compartment  $i$ . The model can then be written as

$$\dot{x}_i = f_i(x) = \mathcal{F}_i(x) - \mathcal{V}_i(x), \quad i = 1, \dots, n,$$

with  $\mathcal{V}_i(x) = \mathcal{V}_i^-(x) - \mathcal{V}_i^+(x)$ . The next-generation matrices are the Jacobians of these terms evaluated at the disease-free equilibrium  $x_0$ :

$$F = \left[ \frac{\partial \mathcal{F}_i(x_0)}{\partial x_j} \right], \quad V = \left[ \frac{\partial \mathcal{V}_i(x_0)}{\partial x_j} \right], \quad 1 \leq i, j \leq m,$$

so that  $\mathcal{R}_0 = \rho(FV^{-1})$ .

*Disease-free equilibrium and mosquito demographic equilibrium:* At the DFE the human population is fully susceptible,  $S_H^* = N$ , and all infected compartments vanish. The mosquito population rests at the demographic equilibrium  $M^*$  of the density-dependent recruitment dynamics (11). Because the climate drivers are time-dependent, we evaluate them at their fixed reference state (the study-period means  $\bar{T}$ ,  $\bar{R}$ ), so that the recruitment rate and the carrying capacity become constants:

$$\bar{b} = b(\bar{R}, \bar{T}), \quad \bar{K} = K(t) \Big|_{\text{DFE}} = M' f_T(\bar{T}) f_R(\bar{R}),$$

and

$$\frac{dM}{dt} = \bar{b} \bar{K} \left( 1 - \frac{M}{\bar{K}} \right) - \mu M$$

Imposing  $\Lambda(M^*) = \mu M^*$  gives

$$\bar{b} \bar{K} \left( 1 - \frac{M^*}{\bar{K}} \right) = \mu M^* \implies M^* = \frac{\bar{b} \bar{K}}{\bar{b} + \mu},$$

and hence the recruitment at the DFE is

$$\Lambda^* = \mu M^* = \frac{\mu \bar{b} \bar{K}}{\bar{b} + \mu}$$

By Proposition 1 (Section 5.1),  $M^* > 0$  is the unique relevant equilibrium whenever  $\bar{b}, \bar{K} > 0$ . Therefore, the disease-free equilibrium will be

$$x_0 = (N, 0, 0, 0, M^*, 0, 0)^\top. \quad (40)$$

*Verification of  $\mathcal{R}_0$  by explicit multiplication:* From (25), the transfer matrix is block-diagonal,  $V = \text{diag}(V_H, V_M)$ , with

$$V_H = \begin{pmatrix} b_{3H} & 0 \\ -b_{3H} & \gamma \end{pmatrix}, \quad V_M = \begin{pmatrix} b_{3M}\ell + \mu & 0 \\ -b_{3M}\ell & \mu \end{pmatrix},$$

so that

$$V^{-1} = \text{diag}(V_H^{-1}, V_M^{-1}),$$

$$V_H^{-1} = \begin{pmatrix} \frac{1}{b_{3H}} & 0 \\ \frac{1}{\gamma} & \frac{1}{\gamma} \end{pmatrix},$$

$$V_M^{-1} = \begin{pmatrix} \frac{1}{b_{3M}\ell + \mu} & 0 \\ \frac{1}{\mu(b_{3M}\ell + \mu)} & \frac{1}{\mu} \end{pmatrix}$$

Carrying out the block multiplication  $FV^{-1}$  yields the hollow form

$$FV^{-1} = \begin{pmatrix} 0 & P \\ Q & 0 \end{pmatrix},$$

$$P = \begin{pmatrix} \frac{ab_2 S_H^* b_{3M}\ell}{N\mu(b_{3M}\ell + \mu)} & \frac{ab_2 S_H^*}{N\mu} \\ 0 & 0 \end{pmatrix},$$

$$Q = \begin{pmatrix} \frac{ab_1 S_M^*}{N\gamma} & \frac{ab_1 S_M^*}{N\gamma} \\ 0 & 0 \end{pmatrix}$$

Both  $P$  and  $Q$  have rank one, so  $PQ$  has a single nonzero eigenvalue  $\lambda = \frac{a^2 b_1 b_2 S_H^* S_M^* b_{3M}\ell}{N^2 \mu \gamma (b_{3M}\ell + \mu)}$ , and the spectrum of  $FV^{-1}$  is

$$\sigma(FV^{-1}) = \{0, 0, \pm\sqrt{\lambda}\} = \{0, 0, \pm\sqrt{\mathcal{R}_0^H \mathcal{R}_0^M}\}$$

Substituting  $S_H^* = N$  and  $S_M^* = M^* = \Lambda^*/\mu$  into  $\lambda$  and taking the square root reproduces (26).

*Spectral computation of  $\rho(FV^{-1})$ :* The matrix  $V$  is block-diagonal in the human-mosquito partition, with blocks  $V_H$  and  $V_M$ , so

$$V^{-1} = \begin{pmatrix} \frac{1}{b_{3H}} & 0 & 0 & 0 \\ \frac{1}{\gamma} & \frac{1}{\gamma} & 0 & 0 \\ 0 & 0 & \frac{1}{\mu + b_{3M}\ell} & 0 \\ 0 & 0 & \frac{b_{3M}\ell}{(\mu + b_{3M}\ell)\mu} & \frac{1}{\mu} \end{pmatrix}$$

Multiplying  $F$  by  $V^{-1}$  we obtain

$$FV^{-1} = \begin{pmatrix} 0 & 0 & \frac{ab_2 b_{3M}\ell}{(\mu + b_{3M}\ell)\mu} & \frac{ab_2}{\mu} \\ 0 & 0 & 0 & 0 \\ \frac{ab_1 M^*}{N\gamma} & \frac{ab_1 M^*}{N\gamma} & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

The second and fourth rows are zero, so the spectral problem reduces to the  $2 \times 2$  block coupling the human and mosquito exposed classes,

$$[FV^{-1}]_{\{1,3\}} = \begin{pmatrix} 0 & \frac{ab_2b_{3M}\ell}{(\mu + b_{3M}\ell)\mu} \\ \frac{ab_1M^*}{N\gamma} & 0 \end{pmatrix},$$

whose characteristic equation is

$$\lambda^2 = \frac{a^2b_1b_2b_{3M}\ell M^*}{N\gamma(\mu + b_{3M}\ell)\mu}$$

The eigenvalues of  $FV^{-1}$  are  $\lambda \in \{0, 0, \pm\sqrt{\mathcal{R}_H\mathcal{R}_M}\}$ , where

$$\mathcal{R}_H = \frac{ab_2}{\gamma}, \quad \mathcal{R}_M = \frac{ab_1\Lambda^*}{N\mu} \frac{b_{3M}\ell}{(b_{3M}\ell + \mu)\mu},$$

so that  $\mathcal{R}_0 = \rho(FV^{-1}) = \sqrt{\mathcal{R}_H\mathcal{R}_M}$ . Finally, substituting  $\Lambda^* = \mu M^*$  recovers the expression in Section 5.2:

$$\mathcal{R}_0 = \sqrt{\frac{a^2b_1b_2\Lambda^*b_{3M}\ell}{N\gamma(b_{3M}\ell + \mu)\mu^2}} \quad (41)$$

### A.3. Figures




Figure 1A: DDE simulation of the baseline SIR-SEI model using the original parameterization from [18]. The dynamics of the SIR model show a monotonic decline in the number of infected individuals within the human population, indicating the absence of sustained transmission, while the dynamics of the SEI model show that the entire mosquito population tends to extinction, an unexpected behavior for a region where malaria is highly endemic.




Figure 2A: ODE simulation of the baseline SIR model using the formulation from Section 3.2 and parameters from [18]. A slight increase in the infectious population can be noticed, although it still tends to extinction over the course of the first year of transmission.




Figure 3A: ODE simulation of the baseline SEI mosquito model using the formulation from Section 3.2 and parameters from [18]. It can be seen that the mosquito population still tends to extinction.