# Simple Harmonic Motion

Dr Mohammad Abdur Rashid



#### References

University Physics with Modern Physics


- Hugh D. Young, Roger A. Freedman

Chapter 14: Periodic Motion

#### Vertical Simple Harmonic Motion

(a)


- **(b)** A body is suspended from the spring. It is in equilibrium when the upward force exerted by the stretched spring equals the body's weight.
- (c) If the body is displaced from equilibrium, the net force on the body is proportional to its displacement. The oscillations are SHM.




# A system that can have periodic motion


$$f = \frac{1}{T} \qquad T = \frac{1}{f}$$

$$\omega = 2\pi f = \frac{2\pi}{T}$$



# Describing oscillation






$$F_x = -kx$$

(restoring force exerted by an ideal spring)


$$a_x = \frac{d^2x}{dt^2} = -\frac{k}{m}x$$
 (simple harmonic motion)

$$\omega = \sqrt{\frac{k}{m}}$$




The restoring force exerted by an idealized spring is directly proportional to the displacement (Hooke's law,  $F_x = -kx$ ): the graph of  $F_x$  versus x is a straight line.

$$x = A\cos\theta$$




$$x = A\cos(\omega t + \phi)$$
 (displacement in SHM)

$$T = \frac{1}{f} = \frac{2\pi}{\omega} = 2\pi \sqrt{\frac{m}{k}}$$
 (simple harmonic motion)




(a) Increasing m; same A and k

Mass *m* increases from curve 1 to 2 to 3. Increasing *m* alone




**(b)** Increasing *k*; same *A* and *m* 

Force constant *k* increases from curve 1 to 2 to 3. Increasing *k* alone



(c) Increasing A; same k and m

Amplitude *A* increases from curve 1 to 2 to 3. Changing *A* alone has



$$x = A\cos(\omega t + \phi)$$

$$T = 2\pi \sqrt{\frac{m}{k}}$$

$$x = A\cos(\omega t + \phi)$$
 (displacement in SHM)

$$v_x = \frac{dx}{dt} = -\omega A \sin(\omega t + \phi)$$
 (velocity in SHM)

$$a_x = \frac{dv_x}{dt} = \frac{d^2x}{dt^2} = -\omega^2 A \cos(\omega t + \phi)$$
 (acceleration in SHM)


$$v_x = \pm \sqrt{\frac{k}{m}} \sqrt{A^2 - x^2}$$


$$a_x = -\omega^2 x = -\frac{k}{m} x$$

$$a_x = -\omega^2 x = -\frac{k}{m} x$$

(a) Displacement x as a function of time t


(b) Velocity  $v_x$  as a function of time t





(c) Acceleration  $a_x$  as a function of time t

The  $v_x$ -t graph is shifted by  $\frac{1}{4}$  cycle from the x-t graph.



The  $a_x$ -t graph is shifted by  $\frac{1}{4}$  cycle from the  $v_x$ -t graph and by  $\frac{1}{2}$  cycle from the x-t graph.


# Energy in Simple Harmonic Motion

$$E = \frac{1}{2}mv_x^2 + \frac{1}{2}kx^2 = \frac{1}{2}kA^2 = \text{constant}$$

(total mechanical energy in SHM)

$$E = \frac{1}{2}mv_x^2 + \frac{1}{2}kx^2 = \frac{1}{2}m[-\omega A\sin(\omega t + \phi)]^2 + \frac{1}{2}k[A\cos(\omega t + \phi)]^2$$
  
=  $\frac{1}{2}kA^2\sin^2(\omega t + \phi) + \frac{1}{2}kA^2\cos^2(\omega t + \phi)$   
=  $\frac{1}{2}kA^2$ 


#### Energy in Simple Harmonic Motion




### Energy in Simple Harmonic Motion

(a) The potential energy U and total mechanical energy E for a body in SHM as a function of displacement x

The total mechanical energy E is constant.



**(b)** The same graph as in **(a)**, showing kinetic energy *K* as well




At these points the energy is half kinetic and half potential.


# Angular frequency, frequency, and period in SHM

A spring is mounted horizontally, with its left end fixed. A spring balance attached to the free end and pulled toward the right (Fig. 14.8a) indicates that the stretching force is proportional to the displacement, and a force of 6.0 N causes a displacement of 0.030 m. We replace the spring balance with a 0.50-kg glider, pull it 0.020 m to the right along a frictionless air track, and release it from rest (Fig. 14.8b). (a) Find the force constant k of the spring. (b) Find the angular frequency  $\omega$ , frequency f, and period T of the resulting oscillation.

# Angular frequency, frequency, and period in SHM

**14.8** (a) The force exerted *on* the spring (shown by the vector F) has x-component  $F_x = +6.0$  N. The force exerted by the spring has x-component  $F_x = -6.0$  N. (b) A glider is attached to the same spring and allowed to oscillate.





16

# Angular frequency, frequency, and period in SHM

**EXECUTE:** (a) When x = 0.030 m, the force the spring exerts on the spring balance is  $F_x = -6.0$  N. From Eq. (14.3),

$$k = -\frac{F_x}{x} = -\frac{-6.0 \text{ N}}{0.030 \text{ m}} = 200 \text{ N/m} = 200 \text{ kg/s}^2$$

(b) From Eq. (14.10), with m = 0.50 kg,

$$\omega = \sqrt{\frac{k}{m}} = \sqrt{\frac{200 \text{ kg/s}^2}{0.50 \text{ kg}}} = 20 \text{ rad/s}$$

$$f = \frac{\omega}{2\pi} = \frac{20 \text{ rad/s}}{2\pi \text{ rad/cycle}} = 3.2 \text{ cycle/s} = 3.2 \text{ Hz}$$

$$T = \frac{1}{f} = \frac{1}{3.2 \text{ cycle/s}} = 0.31 \text{ s}$$

(a) Find the maximum and minimum velocities attained by the oscillating glider of Example 14.2. (b) Find the maximum and minimum accelerations. (c) Find the velocity  $v_x$  and acceleration  $a_x$  when the glider is halfway from its initial position to the equilibrium position x = 0. (d) Find the total energy, potential energy, and kinetic energy at this position.

**EXECUTE:** (a) From Eq. (14.22), the velocity  $v_x$  at any displacement x is

$$v_x = \pm \sqrt{\frac{k}{m}} \sqrt{A^2 - x^2}$$

The glider's maximum *speed* occurs when it is moving through x = 0:

$$v_{\text{max}} = \sqrt{\frac{k}{m}} A = \sqrt{\frac{200 \text{ N/m}}{0.50 \text{ kg}}} (0.020 \text{ m}) = 0.40 \text{ m/s}$$

Its maximum and minimum (most negative) velocities are +0.40 m/s and -0.40 m/s, which occur when it is moving through x = 0 to the right and left, respectively.

(b) From Eq. (14.4),  $a_x = -(k/m)x$ . The glider's maximum (most positive) acceleration occurs at the most negative value of x, x = -A:

$$a_{\text{max}} = -\frac{k}{m}(-A) = -\frac{200 \text{ N/m}}{0.50 \text{ kg}}(-0.020 \text{ m}) = 8.0 \text{ m/s}^2$$

The minimum (most negative) acceleration is  $a_{min} = -8.0 \text{ m/s}^2$ , which occurs at x = +A = +0.020 m.

(c) The point halfway from  $x = x_0 = A$  to x = 0 is x = A/2 = 0.010 m. From Eq. (14.22), at this point


$$v_x = -\sqrt{\frac{200 \text{ N/m}}{0.50 \text{ kg}}} \sqrt{(0.020 \text{ m})^2 - (0.010 \text{ m})^2} = -0.35 \text{ m/s}$$

We choose the negative square root because the glider is moving from x = A toward x = 0. From Eq. (14.4),

$$a_x = -\frac{200 \text{ N/m}}{0.50 \text{ kg}} (0.010 \text{ m}) = -4.0 \text{ m/s}^2$$

Figure 14.14 shows the conditions at x = 0,  $\pm A/2$ , and  $\pm A$ .





(d) The energies are

$$E = \frac{1}{2}kA^2 = \frac{1}{2}(200 \text{ N/m})(0.020 \text{ m})^2 = 0.040 \text{ J}$$


$$U = \frac{1}{2}kx^2 = \frac{1}{2}(200 \text{ N/m})(0.010 \text{ m})^2 = 0.010 \text{ J}$$

$$K = \frac{1}{2}mv_x^2 = \frac{1}{2}(0.50 \text{ kg})(-0.35 \text{ m/s})^2 = 0.030 \text{ J}$$

### Vertical Simple Harmonic Motion

(a)

- **(b)** A body is suspended from the spring. It is in equilibrium when the upward force exerted by the stretched spring equals the body's weight.
- (c) If the body is displaced from equilibrium, the net force on the body is proportional to its displacement. The oscillations are SHM.



# Readings

University Physics with Modern Physics

- Hugh D. Young, Roger A. Freedman

Chapter 14: Periodic Motion