

Environmental Science and Ecotechnology

Soil Microbiomes Reversely Respond to Heavy Metals and Polycyclic Aromatic Hydrocarbons in Highly Contaminated Industrial Sites

--Manuscript Draft--

Manuscript Number:	ESE-D-21-00113
Full Title:	Soil Microbiomes Reversely Respond to Heavy Metals and Polycyclic Aromatic Hydrocarbons in Highly Contaminated Industrial Sites
Short Title:	Soil Microbiomes Reversely Respond to Heavy Metals and PAHs
Article Type:	Full Length Article
Section/Category:	Environmental Science
Keywords:	soil microbiomes; e-waste; coking plant; heavy metal; polycyclic aromatic hydrocarbons
Corresponding Author:	Zhen-Ni Yang, Ph.D. Chinese Academy of Sciences CHINA
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	Chinese Academy of Sciences
Corresponding Author's Secondary Institution:	
First Author:	Zhen-Ni Yang, Ph.D.
First Author Secondary Information:	
Order of Authors:	Zhen-Ni Yang, Ph.D.
	Ze-Shen Liu
	Ke-Huan Wang
	Zong-Lin Liang
	Rashidin Abdugheni
	Ye Huang
	Run-Hua Wang
	Hong-Lin Ma
	Xiao-Kang Wang
	Mei-Ling Yang
	Bing-Ge Zhang
	De-Feng Li
	Cheng-Ying Jiang
	Philippe F.-X. Corvini
	Shuang-Jiang Liu
Order of Authors Secondary Information:	
Abstract:	Industrial sites from e-waste dismantling plants and coking plants are characterized by severe pollution with heavy metals (HMs) and/or polycyclic aromatic hydrocarbons (PAHs) in soil. Mixed contaminations (HMs+PAHs) complicate land reclamation and influence the microbial diversity and function of soil microbiomes. In this study, we determined HMs and PAHs contaminations of an e-waste dismantling plant and a coking plant, and evaluated the influences of HMs and PAHs on soil microbiomes. The

	<p>microbiomes (diversity and abundance) of all sites were determined using high-throughput sequencing of 16S rRNA genes, and canonical correlation analysis was conducted to investigate the relations between soil microbiomes and contaminants. The e-waste dismantling plant was polluted mainly with HMs (such as $2,379.07 \pm 227.46$ mg/kg of Mn) and slightly with PAHs (10.36 ± 0.74 mg/kg), whereas the coking plant was contaminated severely with PAHs ($12,558.06 \pm 611.19$ mg/kg) and moderately with HMs (such as 672.61 ± 7.13 mg/kg of Mn). The abundances of bacterial taxa such as MND1, Bryobacter and Galiella were positively related to the concentration of HMs, whereas the abundance of genera such as Sulfuritalea, Pseudomonas and Sphingobium were positively related to the concentration of PAHs. To our knowledge, we found for the first time that HMs and PAHs in the severely contaminated industrial sites reversely changed the soil microbiomes. This study contributes to the understanding of the combined effect of HMs and PAHs on soil microbiomes.</p>
Suggested Reviewers:	<p>Xingyu Liu China nonferrous Mining Group Co., Ltd. wellwoodliu@163.com experts in heavy metal pollution control and bioremediation</p> <p>Yujie He Nanjing University heyujie@nju.edu.cn expert in soil bioremediation</p> <p>Jiandong Jiang Nanjing Agriculture University jiang_jjd@njau.edu.cn expert in environmental microbiology</p>
Opposed Reviewers:	
Additional Information:	
Question	Response

November 2nd, 2021

Dear Editors:

With the agreement from all listed authors, I submit a manuscript entitled “*Soil Microbiomes Reversely Respond to Heavy Metals and Polycyclic Aromatic Hydrocarbons in Highly Contaminated Industrial Sites*”(authors: Zhen-Ni Yang, Ze-Shen Liu, Ke-Huan Wang, Zong-Lin Liang, Rashidin Abdugheni, Ye Huang, Run-Hua Wang, Hong-Lin Ma, Xiao-Kang Wang, Mei-Ling Yang, Bing-Ge Zhang, De-Feng Li, Cheng-Ying Jiang, Philippe F.-X. Corvini and Shuang-Jiang Liu) for your consideration as an article in *Environmental Science and Ecotechnology*. We confirm that the manuscript has not been published previously by any of the authors and is not under consideration for publication in another journal at the time of submission.

We believe that this article is of great interest to the readership of *Environmental Science and Ecotechnology*. Soil contaminations from industries have attracted serious concerns. Electronic waste (e-waste) dismantling and coking are important contamination sources of heavy metals (HMs) and polycyclic aromatic hydrocarbons (PAHs). The mixed contaminations (HMs+PAHs) affected the diversity and function of the soil microbiomes. Intensive understanding of how soil microbiomes interact with contaminants is pivotal to develop more effective processes for bioremediation.

In the study, we aimed to determine the contaminations of HMs and PAHs in an e-waste dismantling plant and a coking plant, and investigate the effect of the mixed contamination on the soil microbiomes. We found that the e-waste dismantling plant site was featured by heavy contamination of HMs but minor PAHs, whereas the coking plant site was featured by serious contamination of PAHs and moderate HMs. The abundance of bacterial genera such as *Sulfuritalea* increased in soil contaminated severely with PAHs; however, other bacterial genera such as *Bryobacter* increased in soil polluted with HMs. To our knowledge, we found for the first time that HMs and PAHs in the severely contaminated industrial sites reversely changed the soil microbiomes. This study promotes the understanding of the mixed effect of HMs and PAHs on soil microbiomes.

We believe that the present results will be of immediate interest to scientists who have interest in soil microbiomes and mixed contaminations of HMs and PAHs. For these reasons, we believe that this manuscript warrants publication in Article form in *Environmental Science and Ecotechnology*.

Thank you very much for your kind consideration. We are looking forward to hearing from you soon.

Yours sincerely,



Shuang-Jiang Liu

Professor

Institute of Microbiology, Chinese Academy of Sciences,
Shandong University

Email: liusj@im.ac.cn or liusj@sdu.edu.cn

Declaration of interests

☒ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☐The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

**Soil Microbiomes Reversely Respond to Heavy Metals and Polycyclic Aromatic
Hydrocarbons in Highly Contaminated Industrial Sites**

Zhen-Ni Yang^{a,1}, Ze-Shen Liu^{a,1}, Ke-Huan Wang^{a,b}, Zong-Lin Liang^{a,b}, Rashidin
Abdugheni^{a,b}, Ye Huang^{a,b}, Run-Hua Wang^{a,b}, Hong-Lin Ma^{a,b}, Xiao-Kang Wang^{a,b},
Mei-Ling Yang^d, Bing-Ge Zhang^e, De-Feng Li^a, Cheng-Ying Jiang^a, Philippe F.-X.
Corvini^f and Shuang-Jiang Liu^{a,c,*}

^a State Key Laboratory of Microbial Resources, Institute of Microbiology, Chinese
Academy of Sciences, Beijing 100101, China

^b University of Chinese Academy of Sciences, Beijing 100049, China

^c State Key Laboratory of Microbial Technology, Microbial Technology Institute,
Shandong University, Qingdao 226237, Shandong Province, China

^d School of Life Sciences, Hebei University, Baoding 071002, Hebei Province, China

^e School of Public Health, Xuzhou Medical University, Xuzhou 221004, Jiangsu
Province, China

^f School of Life Sciences, University of Applied Sciences and Arts Northwestern
Switzerland, Muttensz 4132, Switzerland

¹ The two authors contributed equally to this work.

* Correspondence: liusj@im.ac.cn or liusj@sdu.edu.cn (S. J. Liu)

Abstract: Industrial sites from e-waste dismantling plants and coking plants are characterized by severe pollution with heavy metals (HMs) and/or polycyclic aromatic hydrocarbons (PAHs) in soil. Mixed contaminations (HMs+PAHs) complicate land reclamation and influence the microbial diversity and function of soil microbiomes. In this study, we determined HMs and PAHs contaminations of an e-waste dismantling plant and a coking plant, and evaluated the influences of HMs and PAHs on soil microbiomes. The microbiomes (diversity and abundance) of all sites were determined using high-throughput sequencing of 16S rRNA genes, and canonical correlation analysis was conducted to investigate the relations between soil microbiomes and contaminants. The e-waste dismantling plant was polluted mainly with HMs (such as $2,379.07 \pm 227.46$ mg/kg of Mn) and slightly with PAHs (10.36 ± 0.74 mg/kg), whereas the coking plant was contaminated severely with PAHs ($12,558.06 \pm 611.19$ mg/kg) and moderately with HMs (such as 672.61 ± 7.13 mg/kg of Mn). The abundances of bacterial taxa such as MND1, *Bryobacter* and *Galiella* were positively related to the concentration of HMs, whereas the abundance of genera such as *Sulfuritalea*, *Pseudomonas* and *Sphingobium* were positively related to the concentration of PAHs. To our knowledge, we found for the first time that HMs and PAHs in the severely contaminated industrial sites reversely changed the soil microbiomes. This study contributes to the understanding of the combined effect of HMs and PAHs on soil microbiomes.

1 39 **Keywords:** soil microbiomes; e-waste; coking plant; heavy metal; polycyclic
2
3
4 40 aromatic hydrocarbons
5
6 41
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1. Introduction

Soil contaminations resulting from industrial activities have raised serious concerns (Park *et al.* 2021, Zhao *et al.* 2021). Currently, industrial processes such as electronic waste (e-waste) dismantling, metal smelting and coking are important sources of contamination with heavy metals (HMs) and polycyclic aromatic hydrocarbons (PAHs) (Guo *et al.* 2021, Kong *et al.* 2021). Previous studies demonstrated that mixed contaminations with HMs and PAHs have different influences on the diversity and community structure of the soil microbiomes compared to single contamination (Gauthier *et al.* 2014). Intensive understanding of how soil microbiomes responding to mixed pollution is essential to develop more effective bioremediation processes (Lee *et al.* 2018).

HMs widely occur in soil (Li *et al.* 2021b) and human activities such as industrial processes might lead to high concentration of HMs in soil (Zhou *et al.* 2016). HMs such as Cd, Cr and Pb are toxic to organisms even at low concentration (Rehman *et al.* 2018, Seleiman *et al.* 2020, Usman *et al.* 2013). Studies have reported that soil microbiomes respond and adapt to HMs, resulting in microbial metabolic activity, population diversity and abundance changes (Beattie *et al.* 2018, Lin *et al.* 2019b, Zhao *et al.* 2020). For example, some species of *Achromobacter* and *Bacillus* were able to eliminate the toxicity of some HMs (*e.g.*, Cr, Mn and As) via adsorption, oxidation and reduction mechanisms (Bachate *et al.* 2013, Yin *et al.* 2019). PAHs are also relevant soil pollutants (Marsh *et al.* 2005) and are released into the environment

63 as a consequence of industrial processes and human activities, *e.g.*, incomplete
64 combustion and pyrolysis of organic substances, transportation and waste incineration
65 (Song *et al.* 2021, Wheatley and Sadhra 2004, Zhang and Wang 2011). PAHs cause
66 great concerns due to their high toxicity and low bioavailability (Tomczyk *et al.* 2020).
67 Besides, PAHs represent a threat to human health due to their teratogenic,
68 carcinogenic, and mutagenic properties (Wu *et al.* 2021). PAHs, especially those with
69 more than four benzene rings, are recalcitrant to degradation in the environment,
70 owing to their high hydrophobicity and their low bioavailability (Gupta *et al.* 2019).
71 Soil contamination with PAHs resulted in the decrease of soil microbial diversity and
72 abundance and metabolic function. The abundances of genera such as *Rhizobacter*,
73 *Sphingobium*, *Mycobacterium*, *Bacillus* and *Pseudarthrobacter* increased in
74 PAHs-contaminated soil (Li *et al.* 2019).

75 Many contaminated sites are affected with both HMs and PAHs, and mixed
76 contamination has caused increasing concerns (Liu *et al.* 2017). The effects of mixed
77 contamination with HMs and PAHs on microbes and environments are complex and
78 difficult to understand (Haarstad *et al.* 2012). Studies demonstrated that HMs affect
79 PAH degradation (Guo *et al.* 2010, Zhang *et al.* 2011) and conversely, that PAHs
80 altered the transport of HMs (Cao *et al.* 2008, Gorria *et al.* 2006). In addition, the
81 effect of mixed contamination is species-and-dose-dependent. For example, the
82 microbial adsorption and degradation of PAHs were dependent on both the bacterial
83 species and concentration of HMs (Chen *et al.* 2016, Yin *et al.* 2020). The adsorption

1 84 of phenanthrene on cyanobacteria cells decreased with the increase of Cu(II)
2
3 85 concentration below 0.04 mM, but enhanced with increasing Cu(II) concentration in
4
5
6 86 the range from 0.04 mM to 0.2 mM (Tao *et al.* 2013). Bourceret *et al.* (Bourceret *et al.*
7
8
9 87 2016) found that *Proteobacteria*, *Actinobacteria*, and *Bacteroidetes* were the
10
11
12 88 dominant phyla in soil contaminated with HMs and PAHs. Gran-Scheuch *et al.*
13
14
15 89 (Gran-Scheuch *et al.* 2020) reported that the population of PAHs degraders or metal
16
17
18 90 resistant bacteria increased in the soil of King George Island (Antarctica) where HMs
19
20
21 91 and PAHs were detected. Although attention has been paid to mixed contamination
22
23
24 92 with HMs and PAHs (Lin *et al.* 2019a, Picariello *et al.* 2020), the understanding of
25
26
27 93 how microbiomes respond and adapt to mixed contamination is still limited (Liu *et al.*
28
29
30 94 2017).

31
32 95 In this study, we aimed to determine the level of contamination with HMs and
33
34
35 96 PAHs in an e-waste dismantling plant and a coking plant and to investigate the effect
36
37
38 97 of this mixed contamination on the soil microbiomes. We found that the e-waste
39
40
41 98 dismantling plant was polluted mainly with HMs and slightly with PAHs, whereas the
42
43
44 99 coking plant was contaminated severely with PAHs and moderately with HMs. The
45
46
47 100 abundance of bacterial taxa such as MND1, *Bryobacter* and *Galiella* increased with
48
49
50 101 the increasing concentration of HMs, whereas that of *Sulfuritalea*, *Pseudomonas* and
51
52
53 102 *Sphingobium* increased with the increasing concentration of PAHs. To our knowledge,
54
55
56 103 we report for the first time that HMs and PAHs in severely contaminated industrial
57
58
59 104 sites reversely changed the soil microbiomes.

2. Materials and Methods

2.1. Soil sampling

An initial investigation of the level of soil contamination with HMs and PAHs was carried out at an e-waste dismantling plant and a coking plant. The two plants were located 48.8 km far from each other and showed similar climatic features. Based on the degree of contamination with HMs and PAHs, five samples (annotated samples A-E) were selected for further investigation (Table 1).

Table 1. Information of sampling sites.

Samples	Sampling sites	Characteristics ¹	Depth (cm)
A	Paddy field (control)	HMs (I) and PAHs (I)	0–30
B	E-waste dismantling plant	HMs (V) and PAHs (II)	0–30
C	Coking plant	HMs (III) and PAHs (III)	0–30
D	Coking plant	HMs (IV) and PAHs (IV)	0–30
E	Coking plant	HMs (II) and PAHs (V)	0–30

¹ Roman numbers from ‘I’ (low) to ‘V’ (high) refer to the contamination levels of HMs and PAHs according to their total concentration. The concentration of HMs and PAHs in detail were given in Figure 1.

Sample A collected from the paddy field near the e-waste dismantling plant served as the negative control in this study. Sample B was obtained from the e-waste dismantling plant, whereas samples C-E were from the coking plant. In the coking plant, sample C was collected from the field where coal and iron were stacked, sample D was taken near the coking process, and sample E was significantly polluted with tar oil. The five samples were classified in different level categories, *i.e.*, Level ‘I’ to Level ‘V’ (Table 1) on the basis of the concentration of HMs and PAHs. Sampling

was conducted on December 12th, 2020. Five soil subsamples from each site were collected and mixed to obtain one composite soil sample. After drying and sieving (0.15 mm), triplicate soil samples of 100 g for each sampling site were used for soil characterization.

2.2. Chemical analysis

Chemical reagents including hydrochloric acid, nitric acid, acetonitrile, acetone and *n*-hexane of analytical grade were purchased from Sinopharm Chemical Reagent Co., Ltd (China). Decafluorobiphenyl and dichloromethane were from Shanghai Eon Chemical Technology Co., Ltd (China) and Shanghai Macklin Regent Co., Ltd (China), respectively. Acetonitrile of HPLC grade was from Concord Technology (Tianjin) Co., Ltd (China). Standard solution of PAHs was from TMstandard Co., Ltd (China).

2.2.1. Analyses of HMs

Concentration of HMs were determined using an inductively coupled plasma optical emission spectrometer (ICP-OES; Optima 5300 DV, PerkinElmer, USA) after acidic digestion (da Silva *et al.* 2016). Triplicate soil samples (around 1.0 g each) in 7.0 mL of aqua regia (HCl:HNO₃, 3:1, v/v) were digested at 185°C for 40 min using microwave (Topwave, analytikjena, Germany). After digestion, the acid solution kept faint boiling on a hot plate in the fuming cupboard until 2~3 mL of volume left. The residual acid solution was dissolved in ultrapure Milli-Q water to a final volume of 50 mL. The concentration of 15 elements (*i.e.*, Mn, As, Cr, Pb, Cu, Zn, Ni, Ba, Sr, V, La,

Li, Co, Mo and Sc) in the solution was measured using ICP-OES, and the final concentration in the soil samples was calculated for each element and the total concentration (Σ HMs) was also determined.

2.2.2. Analyses of PAHs

PAHs in the soil were extracted using the Soxhlet method and analyzed by means of high-performance liquid chromatography (HPLC; 1260 Infinity, Agilent Technologies, USA) (Wu *et al.* 2018). Triplicate soil samples of 10.0 g each were extracted with 100 mL of acetone and *n*-hexane (1:1, v/v) using a Soxhlet extractor and spiked with 200 ng of decafluorobiphenyl. The extract was then filtered through fiberglass meshes and concentrated and cleaned up using a rotary evaporator and Si SPE Cartridge (CNW, ANPEL Laboratory Technologies (Shanghai) Inc., China), respectively. A mixture of *n*-hexane and dichloromethane (1:1, v/v) was used to dissolve the PAHs. The extract was concentrated by a rotary evaporator and dissolved in acetonitrile to a final volume of 5.0 mL. The concentration of PAHs in the concentrated acetonitrile solution was analyzed using a HPLC with Eclipse plus C18 column (Agilent, USA) and acetonitrile-water as mobile phase according to the gradient (6:4, acetonitrile/water, from the start to 18 min; only acetonitrile from 18 to 28.5 min; finally, 6:4, acetonitrile/water, from 28.5 min to the end). The PAHs analyzed were naphthalene (NAP), acenaphthylene (ACY), fluorene (FLO), phenanthrene (PHE), anthracene (ANT), fluoranthene (FLA), pyrene (PYR), benzo[a]anthracene (BaA), chrysene (CHR), benzo[b]fluoranthene (BbF),

benzo[k]fluoranthene (BkF), benzo[a]pyrene (BaP), dibenzo[a,h]anthracene (DhA), benzo[ghi]perylene (BgP) and indeno[1,2,3-cd]pyrene (IcP). The concentration of the 15 PAHs in the soil samples was calculated for each PAH and the total concentration (Σ PAHs) was also calculated.

2.3. Characteristics of microbial communities

The soil DNA was extracted from triplicate samples using DNeasy[®] PowerSoil[®] Kit (QIAGEN, Germany). The primer set 515F (5'-GTGCCAGCMGCCGCGG-3')/907R (5'-CCGTCAATTCMTTTRAGTTT-3') was used to amplify the V4–V5 region of bacterial 16S ribosomal RNA (16S rRNA) genes (Shan *et al.* 2015). The PCR products were sequenced by Shanghai Majorbio Bio-pharm Technology Co., Ltd (China) using Illumina MiSeq Platform (San Diego, USA). Microbial diversity and composition were analyzed using QIIME2 (v2020.2). Shannon index was calculated using software MOTHUR (v1.30). One-way analysis of variance (ANOVA) was used to establish statistical differences between the groups with $p < 0.05$. Beta diversity on the level of amplicon sequence variants (ASV) was evaluated by principal co-ordinates analysis (PcoA). Linear discriminant analysis (LDA) coupled with effect size measurements (LEfSe) analysis was performed to screen different bacterial species. Canonical correlation analysis (CCA) was conducted by means of the software R (v.3.3.1) using Vegan package to determine the correlation between the soil microbiomes diversity and the level of PAHs/HMs.

3. Results and Discussion

3.1. The e-waste dismantling plant and the coking plant sites are featured with HMs and PAHs

Soil samples from the e-waste dismantling plant and the coking plant were screened according to their contamination with HMs and PAHs. Five soil samples were finally selected to investigate the effect of the co-occurrence of HMs and PAHs on the soil microbiomes, *i.e.* sample A from the paddy field as the negative control, sample B from the e-waste dismantling plant and samples C-D from the coking plant. The concentration of HMs and PAHs in these samples is displayed in Figure 1. Sample A showed the lowest $\sum\text{HMs}$ ($1,609.56 \pm 65.85$ mg/kg) and the lowest $\sum\text{PAHs}$ (4.40 ± 0.84 mg/kg). Mn (679.6 ± 37.29 mg/kg), Zn (307.7 ± 21.4 mg/kg) and Ba (229.62 ± 25.95 mg/kg) were the major HMs (75.61%), and FLO consisting of 3 rings (3.00 ± 0.68 mg/kg) was the major PAH (68.18%). Sample B from the e-waste dismantling plant had $\sum\text{HMs}$ of $16,780.08 \pm 1,278.91$ mg/kg, which was ten times higher than that in sample A and also the highest among all the soil samples. Metals such as Cu, Pb, Zn, Ni and As were widely used in electronic products and their total concentration in sample B accounted for 75.92% of $\sum\text{HMs}$ (Yang *et al.* 2020a, Yang *et al.* 2020b). $\sum\text{PAHs}$ in sample B was twice that of sample A. All the fifteen PAHs were detected, and FLO (2.06 ± 0.11 mg/kg) and BbF (1.65 ± 0.15 mg/kg) were the major PAHs (35.82%).




Figure 1. Analyses of HMs (left) and PAHs (right) in the soil samples. Number in each cell indicates the mean concentration of the corresponding pollutant. Sample A (negative control) was collected from the paddy field, sample B was from the e-waste dismantling plant, and samples C-E were from the coking plant. Roman numbers in superscript refer to the contamination levels based on Σ HMs or Σ PAHs, *i.e.*, ‘I’ (low) to ‘V’ (high).

In the coking plant, Σ HMs in samples C-E were 1.17 to 1.94 times of that in sample A. In samples C-E, Mn showed the highest concentration (672.61 to 1,306.82 mg/kg). The concentration of As (23.03 to 99.51 mg/kg) was the most heterogeneously distributed in samples C-E compared to sample A, followed by Zn, Cr and Ni. The high concentration of Mn, As, Zn, Cr and Ni in soil may be caused by the storage and combustion of coal (Zajusz-Zubek *et al.* 2017). The Σ PAHs in samples C-E ranged from 208.00 to 12,558.06 mg/kg, which were 47.24 to 2,852.29 times higher than that of sample A. In samples C and D, the concentration of PAHs

with more than 4 rings (*e.g.*, PYR of 21.49 to 81.41 mg/kg, CHR of 29.65 to 60.67 mg/kg, and BbF of 22.37 to 95.32 mg/kg) were higher than that with 3 rings and less (*e.g.*, NAP of 0.93 to 16.94 mg/kg and ACY of 2.70 to 11.29 mg/kg). Sample E showed the highest \sum PAHs (12,558.06 \pm 611.19 mg/kg) among the five samples, and 3-ring FLO of 11,740.06 \pm 620.1 mg/kg (93.49%) was the main PAH while the concentration of the other PAH congeners (*i.e.*, ACY, PYR and BbF) was below 211.68 mg/kg. The high \sum PAHs probably resulted from coking activity and the very high concentration of FLO could be originated from tar oil, the main product of the coking activity (Vasilieva *et al.* 2012).

Previous studies showed that e-waste dismantling plants have generally high concentration of HMs but low concentration of PAHs, and coking plant usually have high concentration of PAHs but low concentration of HMs (Rachwal *et al.* 2015, Sun *et al.* 2014, Tang *et al.* 2010). This study showed similar features and the concentration of HMs and PAHs were much higher than those reported in the literature. The high contaminations with HMs and PAHs might help in understanding the response of soil microbiomes to mixed contamination with HMs and PAHs.

3.2. The α - and β -diversities of the microbiomes at the e-waste dismantling and the coking sites

After quality control, 831,803 sequences of 16S rRNA genes were obtained. The measurement of the sequences was sufficient according to the rarefaction curves (Figure S1). These sequences were clustered into 28,746 ASV (amplicon sequence

variants) and assigned into 1,271 genera of 50 bacterial phyla. The α -diversities of the soil microbiomes, *i.e.*, Shannon indexes, were evaluated using those data (Figure 2a; other indexes, *i.e.*, Ace, Chao1 and Simpson indexes, in Table S1).




Figure 2. The α -(a) and β -(b) diversities of soil microbiomes. Symbols of the sampling sites are explained in Figure 1. Roman numbers refer to \sum HMs (the first superscript) and \sum PAHs (the last superscript) in the soil samples increased successively from ‘I’ (low) to ‘V’ (high).

The Shannon indexes of samples A, C and D (7.10~7.33) were not significantly different from each other ($p > 0.05$), but they were significantly higher than those of samples B (6.31 ± 0.15) and E (5.01 ± 0.20) ($p < 0.05$), indicating that the richness of soil microbiomes in samples B and E was apparently lower than that in samples A, C and D. Interestingly, samples B and E had the highest \sum HMs and \sum PAHs, respectively, among all samples. This indicated that soil microbial diversity was decreased by high \sum HMs in sample B ($16,780.08 \pm 1,278.91$ mg/kg) or high \sum PAHs in sample E ($12,558.06 \pm 611.19$ mg/kg). In addition, the Shannon index of sample B was significantly higher than that of sample E ($p < 0.05$). This could indicate that PAHs represent a bigger threat to soil microbiomes than HMs. The β -diversity of the soil

microbiomes was evaluated by means of PCoA (Figure 2b). Samples were distant from each other, indicating that the community structures of the soil microbiomes were affected by the concentration of HMs or PAHs.

3.3. Features of soil microbial community at phylum and genus levels

The microbial communities were then analyzed at phylum and genus levels (Figure 3). There were twelve bacterial phyla with relative abundance more than 1% in the soil samples (Figure 3a). The top ten phyla found in the soil samples were *Proteobacteria*, *Acidobacteriota*, *Actinobacteriota*, *Bacteroidota*, *Chloroflexi*, *Planctomycetota*, *Myxococcota*, *Gemmatimonadota*, *Firmicutes* and *Cyanobacteria*. The *Actinobacteriota* (21.49%) and *Firmicutes* (4.31%) were relatively abundant in sample A but less in the other samples. *Acidobacteriota* (24.55%) and *Myxococcota* (6.90%) showed the highest relative abundances in sample B with the highest \sum HMs. The relative abundance of *Proteobacteria* increased significantly from 28.24% to 78.69% along with the increasing \sum PAHs from sample A (4.4 ± 0.84 mg/kg) to sample E ($12,558.06 \pm 611.19$ mg/kg). These results were similar to those of previous studies reporting that *Acidobacteriota* and *Myxococcota* are more frequently detected in HMs-heavily contaminated environment (Dell'Anno *et al.* 2021) and that *Proteobacteria* were more abundant when \sum PAHs increased (Liu *et al.* 2019).




Figure 3. Bacterial phyla (a) and genera (b) in the soil samples. Unclassified genera in different domains (d), phyla (p), classes (c), orders (o) and families (f) are given. Symbols of the sampling sites are explained in Figure 1. Roman numbers refer to Σ HMs (the first superscript) and Σ PAHs (the last superscript) in the soil samples increased successively from ‘I’ (low) to ‘V’ (high).

Thirty-nine genera had relative abundances above 1.5%, including twenty-two classified genera (56.41%) and seventeen unclassified genera (43.59%) (Figure 3b). Such a high proportion of unclassified genera showed that many bacteria still need to be characterized in the co-contaminated sites. The genera in sample A were quite diverse, and each of them was present with a similar relative abundance. In sample B, MND1 (7.95%), *Bryobacter* (7.54%) and an unclassified genus of *Vicinamibacteriales* order (6.66%) showed high relative abundances. *Flavobacterium*

genus showed the highest abundance in sample C (5.39%) and *Pseudomonas* genus (6.01%) was well represented in sample D. *Sulfuritalea* genus was detected in samples D and E but barely in samples A-C. The relative abundance of *Sulfuritalea* genus in sample E was high, *i.e.*, 21.34%. Besides, *Rhodanobacter* (9.06%), an unclassified genus of the *Comamonadaceae* family (9.35%), *Sphingobium* (4.03%), *Pandora* (3.70%) and *Pseudoxanthomonas* (2.92%) showed higher relative abundances in sample E.

LEfSe (linear discriminant analysis effect size) analysis with LDA (linear discriminant analysis) score >4.0 was useful to identify the main genera which were differently represented in the soil microbiomes of the various places of sampling (Figure 4). There were eight and six typical genera with LDA score >4.0 in samples B and D, respectively, but only three in samples A, C and D. Unclassified genera of *Gaiellales* (3.93%) and *Acidobacteriales* (3.85%) orders and *Massilia* (2.18%) mainly existed in sample A and all of them showed relative abundances <1% in the other samples. The three genera might be sensitive to HMs and PAHs. Genera such as MND1, *Bryobacter* and an unclassified genus of *Vicinamibacteriales* order showed the highest relative abundances in sample B (6~8%). Genera including *Sulfuritalea*, *Rhodanobacter* and an unclassified genus of *Comamonadaceae* family were the highest in sample E (9% to 22%), whereas the three genera showed relative abundances ranging from 0.20% to 4.15% in samples A-D and decreased along with the increase of \sum PAHs. The results suggested that bacteria tolerant to high \sum HMs

existed widely in the soil but failed to colonize soils with high concentration of PAHs. Differently, bacteria tolerant to high Σ PAHs dominated in PAHs-polluted soils but were hardly found in HMs-polluted soils. Besides, genera such as *Flavobacterium* in sample C (5.39%) and *Pseudomonas* in sample D (6.01%) may perform better than other genera in the soil polluted by HMs and PAHs at moderate concentration.




Figure 4. LEfSe results of bacterial genera (a) and relative abundances of typical genera in samples (b). LEfSe was performed with LDA score > 4.0 and $p < 0.05$. Symbols of the sampling sites are explained in Figure 1.

3.4. HMs and PAHs impact differently the soil microbiome composition

The impact of HMs and PAHs on the soil microbiome composition was evaluated with CCA (canonical correlation analysis) (Figure 5a and Table S2). Results showed that microbial community structures of these soil samples were mainly influenced by eight congeners of PAHs (*i.e.*, ACY, FLO, BgP, PYR, DhA, BbF, FLA

and IcP) and four HMs (*i.e.*, Sr, Mn, As and Ni). Sample A with low concentration of HMs and PAHs was far from the others in Figure 5a. The microbiomes of samples B, C and D were mainly influenced by HMs, and the microbiome of sample E was highly impacted by PAHs. The result of CCA coincided with the concentration of HMs and PAHs in the soil samples. Sample B and E were highly contaminated by either HMs ($16,780.08 \pm 1,278.91$ mg/kg) or PAHs ($12,558.06 \pm 611.19$ mg/kg). The microbial community structures of them were therefore determined by HMs and PAHs, respectively. For samples C and D polluted by HMs ($2,124.21$ to $3,118.97$ mg/kg) and PAHs (208.00 to 698.74 mg/kg) at moderate concentration, the microbial community structures were mainly shaped by HMs.




Figure 5. Canonical correlation analysis (CCA) showed reverse influences of HMs and PAHs on the microbial community structure (a) and correlation between the main kinds of contaminants and

bacterial phyla (b) and genera (c). Symbols of the sampling sites and the abbreviations of PAHs (e.g., ACY, FLO and PYR) are explained in Figure 1.

The influences of four HMs (i.e., Mn, Ni, As and Sr) and PAHs (i.e., ACY, FLO, PYR and BgP) on bacterial phyla (Figure 5b) and genera (Figure 5c) were analyzed further. Interestingly, bacteria in soil polluted with high concentration of HMs or PAHs almost separated in two groups. At phylum level, the abundance of *Acidobacteriota*, *Myxococcota* and *Nitrospirota* were positively correlated to the concentration of HMs but negatively to that of PAHs, while the abundance of *Proteobacteria* and *Bacteroidota* were positively related to the concentration of PAHs and indistinctively related with that of HMs. At genus level, the abundance of MND1, *Bryobacter*, *Gaiella*, RB41, *Acidibacter* and *Nitrospira* were positively related to the concentration of HMs and partially negatively to that of PAHs, whereas the abundance of *Sulfuritalea*, *Pseudomonas*, *Rhodanobacter*, *Flavobacterium*, *Sphingobium*, *Pandoraea* and *Pseudoxanthomonas* were positively related to the concentration of PAHs and partially negatively to that of HMs. The correlation analysis results shown in Figure 5 was in accordance with LEfSe results shown in Figure 4. To our knowledge, this is the first study reporting that soil microbiomes in highly HMs- or PAHs- contaminated soils change in opposite directions, and that bacteria tolerant to high concentration of HMs or PAHs were completely different. The different abilities of bacteria to cope with HMs and PAHs may be related to their properties of cell membranes and enzymes. For example, many stains of

1 356 *Cyanobacteria* may produce negatively charged extracellular polymeric substances,
2
3
4 357 which are considered as the first protective barrier against metal stress
5
6 358 (Cassier-Chauvat and Chauvat 2014). In contrast, *Sphingobium* spp. were usually
7
8
9 359 characterized by high cell surface hydrophobicity due to their rich polysaccharidic
10
11
12 360 content and hence tend to adsorb PAHs (Garcia *et al.* 2018).
13
14

15 361 In previous studies, genera such as *Bryobacter*, *Gaiella* and *Nitrospira* were
16
17
18 362 reported to resist to HMs (Hao *et al.* 2018, Kulicheyskaya *et al.* 2014, Li *et al.* 2021a)
19
20
21 363 and *Pseudomonas*, *Sphingobium* and *Pseudoxanthomonas* were largely reported to be
22
23
24 364 able to degrade PAHs (Liang *et al.* 2021, Madueno *et al.* 2018b, Shinde *et al.* 2020).
25
26
27 365 Among these bacteria, many strains of *Pseudomonas* and *Sphingobium* were reported
28
29
30 366 to degrade PAHs and resist to HMs (Afegbua and Batty 2019, Madueno *et al.* 2018a).
31
32
33 367 Few strains of *Sulfuritalea* and *Rhodanobacter* were reported to be able to degrade
34
35
36 368 PAHs (Li *et al.* 2017, Sperfeld *et al.* 2019). Knowledge of genera such as *Bryobacter*,
37
38
39 369 *Flavobacterium* and *Pseudoxanthomonas* is still limited to the bacterial community
40
41
42 370 level, and scarce strains were obtained and studied at laboratory level (Li *et al.* 2021a,
43
44
45 371 Miao *et al.* 2020) (Wang *et al.* 2020). There are still many HMs-tolerant and
46
47
48 372 PAHs-degrading strains to be studied. This study represents a contribution to the
49
50
51 373 application of PAHs-degrading strains for the remediation of sites polluted with HMs
52
53
54 374 and PAHs. For example, *Sphingobium* may be used to remediate soils severely
55
56
57 375 polluted with PAHs ($10^3\sim 10^4$ mg/kg) but only slightly with HMs, while *Pseudomonas*
58
59
60
61
62
63
64
65

could be applied to soil contaminated by both HMs and PAHs at moderate concentration ($10^2 \sim 10^3$ mg/kg).

4. Conclusions

Fifteen HMs and fifteen PAHs were analyzed in contaminated soil samples from an e-waste dismantling plant and a coking plant. Results showed that the microbiomes of these samples were mainly influenced by four HMs (*i.e.*, Sr, Mn, As and Ni) and eight PAHs (*i.e.*, ACY, FLO, BgP, PYR, DhA, BbF, FLA and IcP). The abundance of bacterial taxa such as MND1, *Bryobacter* and *Galiella* were positively related to the concentration of HMs, whereas the abundance of genera such as *Sulfuritalea*, *Pseudomonas* and *Sphingobium* were positively related to the concentration of PAHs. These results demonstrated that the soil microbiomes of the contaminated sites were driven in different way by HMs and PAHs.

Acknowledgements

This study was funded by National Natural Science Foundation of China (Grant Nos. 41991333; 31861133002), European Union's Horizon 2020 Research and Innovation Program Under Grant Agreement (No. 826244), CAS Engineering Laboratory for Advanced Microbial Technology of Agriculture, Chinese Academy of Sciences (KFJ-PTXM-016) and Science and Technology Basic Resources Survey Special Project (2019FY100700).

Author Contributions

Methodology, data analysis and original draft preparation, Z. N. Yang and Z. S. Liu; measurements of PAHs and HMs, K. H. Wang and Z. L. Liang; sample preparation, DNA extraction and bacterial analysis, R. Abdugheni, Y. Huang, R. H. Wang, H. L. Ma, X. K. Wang, M. L. Yang and B. G. Zhang; review and editing, C. Y. Jiang, D. F. Li, P.F.-X. Corvini and S. J. Liu; funding acquisition, S. J. Liu. All authors have read and agreed to the published version of the manuscript.

Abbreviations

E-waste, electronic waste; HMs, heavy metals; PAHs, polycyclic aromatic hydrocarbons; NAP, naphthalene; ACY, acenaphthylene; FLO, fluorene; PHE, phenanthrene; ANT, anthracene; FLA, fluoranthene; PYR, pyrene; BaA, benzo[A]anthracene; CHR, chrysene; BbF, benzo[b]fluoranthene; BkF, benzo[k]fluoranthene; BaP, benzo[a]pyrene; DhA, dibenzo[a,h]anthracene; BgP, benzo[ghi]perylene; IcP, indeno[1,2,3-cd]pyrene; 16S rRNA, 16S ribosomal RNA; ASV, amplicon sequence variants; PcoA, principal co-ordinates analysis; LEfSe, linear discriminant analysis (LDA) coupled with effect size measurements; CCA, canonical correlation analysis.

References

- Afegbua, S.L. and Batty, L.C., 2019. Effect of plant growth promoting bacterium; *Pseudomonas putida* UW4 inoculation on phytoremediation efficacy of monoculture and mixed culture of selected plant species for PAH and lead spiked soils. Int. J. Phytoremediat. 21(3), 200-208. <https://doi.org/10.1080/15226514.2018.1501334>.
- Bachate, S.P., Nandre, V.S., Ghatpande, N.S. and Kodam, K.M., 2013. Simultaneous reduction of Cr(VI) and oxidation of As(III) by *Bacillus firmus* TE7 isolated from tannery effluent. Chemosphere 90(8), 2273-2278. <https://doi.org/10.1016/j.chemosphere.2012.10.081>.
- Beattie, R.E., Henke, W., Campa, M.F., Hazen, T.C., McAliley, L.R. and Campbell, J.H., 2018. Variation in microbial community structure correlates with heavy-metal contamination in soils decades after mining ceased. Soil Biol. Biochem. 126, 57-63. <https://doi.org/10.1016/j.soilbio.2018.08.011>.
- Bourceret, A., Cebon, A., Tisserant, E., Poupin, P., Bauda, P., Beguiristain, T. and Leyval, C., 2016. The bacterial and fungal diversity of an aged PAH- and heavy metal-contaminated soil is affected by plant cover and edaphic parameters. Microb. Ecol. 71(3), 711-724. <https://doi.org/10.1007/s00248-015-0682-8>.
- Cao, L.K., Shen, G.Q. and Lu, Y.T., 2008. Combined effects of heavy metal and polycyclic aromatic hydrocarbon on soil microorganism communities. Environ. Geol. 54(7), 1531-1536. <https://doi.org/10.1007/s00254-007-0934-0>.

437 Cassier-Chauvat, C. and Chauvat, F., 2014. Responses to oxidative and heavy metal
 438 stresses in Cyanobacteria: recent advances. *Int J Mol Sci* 16(1), 871-886.
 439 <https://doi.org/10.3390/ijms16010871>.
 440 Chen, C., Lei, W.R., Lu, M., Zhang, J.N., Zhang, Z., Luo, C.L., Chen, Y.H., Hong, Q.
 441 and Shen, Z.G., 2016. Characterization of Cu(II) and Cd(II) resistance
 442 mechanisms in *Sphingobium* sp. PHE-SPH and *Ochrobactrum* sp. PHE-OCH
 443 and their potential application in the bioremediation of heavy
 444 metal-phenanthrene co-contaminated sites. *Environ. Sci. Pollut. R.* 23(7),
 445 6861-6872. <https://doi.org/10.1007/s11356-015-5926-0>.
 446 da Silva, F.B.V., do Nascimento, C.W.A., Araujo, P.R.M., da Silva, L.H.V. and da
 447 Silva, R.F., 2016. Assessing heavy metal sources in sugarcane Brazilian soils: an
 448 approach using multivariate analysis. *Environ. Monit. Assess.* 188(8).
 449 <https://doi.org/10.1007/s10661-016-5409-x>.
 450 Dell'Anno, F., Rastelli, E., Tangherlini, M., Corinaldesi, C., Sansone, C., Brunet, C.,
 451 Balzano, S., Ianora, A., Musco, L., Montereali, M.R. and Dell'Anno, A., 2021.
 452 Highly contaminated marine sediments can host rare bacterial taxa potentially
 453 useful for bioremediation. *Front. Microbiol.* 12.
 454 <https://doi.org/10.3389/fmicb.2021.584850>.
 455 Garcia, M.C., Carmona, J.A., Santos, J., Alfaro, M.C. and Munoz, J., 2018. Effect of
 456 temperature and shear on the microstructure of a microbial polysaccharide

secreted by *Sphingomonas* species in aqueous solution. Int. J. Biol. Macromol. 118, 2071-2075. <https://doi.org/10.1016/j.ijbiomac.2018.07.070>.

Gauthier, P.T., Norwood, W.P., Prepas, E.E. and Pyle, G.G., 2014. Metal-PAH mixtures in the aquatic environment: a review of co-toxic mechanisms leading to more-than-additive outcomes. Aquat Toxicol 154, 253-269. <https://doi.org/10.1016/j.aquatox.2014.05.026>.

Gorria, M., Tekpli, X., Sergent, O., Huc, L., Gaboriau, F., Rissel, M., Chevanne, M., Dimanche-Boitrel, M.T. and Lagadic-Gossmann, D., 2006. Membrane fluidity changes are associated with benzo[a]pyrene-induced apoptosis in F258 cells - Protection by exogenous cholesterol. Ann. NY Acad. Sci. 1090, 108-112. <https://doi.org/10.1196/annals.1378.011>.

Gran-Scheuch, A., Ramos-Zuniga, J., Fuentes, E., Bravo, D. and Perez-Donoso, J.M., 2020. Effect of co-contamination by PAHs and heavy metals on bacterial communities of diesel contaminated soils of South Shetland Islands, Antarctica. Microorganisms 8(11). <https://doi.org/10.3390/microorganisms8111749>.

Guo, G.H., Zhang, D.G. and Wang, Y., 2021. Source apportionment and source-specific health risk assessment of heavy metals in size-fractionated road dust from a typical mining and smelting area, Gejiu, China. Environ. Sci. Pollut. R. 28(8), 9313-9326. <https://doi.org/10.1007/s11356-020-11312-y>.

Guo, H.J., Luo, S.L., Chen, L.A., Xiao, X., Xi, Q.A., Wei, W.Z., Zeng, G.M., Liu, C.B., Wan, Y., Chen, J.L. and He, Y.J., 2010. Bioremediation of heavy metals by

growing hyperaccumulaor endophytic bacterium *Bacillus* sp. L14. Bioresour. Technol. 101(22), 8599-8605. <https://doi.org/10.1016/j.biortech.2010.06.085>.

Gupta, G., Kumar, V. and Pal, A.K., 2019. Microbial Degradation of High Molecular Weight Polycyclic Aromatic Hydrocarbons with Emphasis on Pyrene. Polycycl. Aromat. Comp. 39(2), 124-138. <https://doi.org/10.1080/10406638.2017.1293696>.

Haarstad, K., Bavor, H.J. and Maehlum, T., 2012. Organic and metallic pollutants in water treatment and natural wetlands: a review. Water Sci. Technol. 65(1), 76-99. <https://doi.org/10.2166/wst.2011.831>.

Hao, L.T., Zhang, B.G., Feng, C.P., Zhang, Z.Y., Lei, Z.F., Shimizu, K., Cao, X.L., Liu, H. and Liu, H.P., 2018. Microbial vanadium (V) reduction in groundwater with different soils from vanadium ore mining areas. Chemosphere 202, 272-279. <https://doi.org/10.1016/j.chemosphere.2018.03.075>.

Kong, D.G., Du, Y.M., Luo, M., Liu, G.W., Su, X.T., Luo, D.G., Huang, X.X., Wei, L.Z., Liu, Y. and Wu, Q.H., 2021. Environmental effects of heavy metals from the e-waste dismantling site, South China. Soil Sediment Contam. <https://doi.org/10.1080/15320383.2021.1887810>.

Kulicheyskaya, I.S., Suzina, N.E., Rijpstra, W.I.C., Damste, J.S.S. and Dedysh, S.N., 2014. *Paludibaculum fermentans* gen. nov., sp nov., a facultative anaerobe capable of dissimilatory iron reduction from subdivision 3 of the *Acidobacteria*. Int. J. Syst. Evol. Micr. 64, 2857-2864. <https://doi.org/10.1099/ij.s.0.066175-0>.

499 Lee, D.W., Lee, H., Lee, A.H., Kwon, B.O., Khim, J.S., Yim, U.H., Kim, B.S. and
 500 Kim, J.J., 2018. Microbial community composition and PAHs removal potential
 501 of indigenous bacteria in oil contaminated sediment of Taean coast, Korea.
 502 Environ. Pollut. 234, 503-512. <https://doi.org/10.1016/j.envpol.2017.11.097>.
 503 Li, B.Q., Xu, R., Sun, X.X., Han, F., Xiao, E.Z., Chen, L., Qiu, L. and Sun, W.M.,
 504 2021a. Microbiome-environment interactions in antimony-contaminated rice
 505 paddies and the correlation of core microbiome with arsenic and antimony
 506 contamination. Chemosphere 263.
 507 <https://doi.org/10.1016/j.chemosphere.2020.128227>.
 508 Li, J.B., Zhang, D.Y., Song, M.K., Jiang, L.F., Wang, Y.J., Luo, C.L. and Zhang, G.,
 509 2017. Novel bacteria capable of degrading phenanthrene in activated sludge
 510 revealed by stable-isotope probing coupled with high-throughput sequencing.
 511 Biodegradation 28(5-6), 423-436. <https://doi.org/10.1007/s10532-017-9806-9>.
 512 Li, L., Yan, X.Y., Li, J., Tian, Y.S. and Ren, P., 2021b. Advances in cotton tolerance to
 513 heavy metal stress and applications to remediate heavy metal-contaminated
 514 farmland soil. Phyton-Int. J. Exp. Bot. 90(1), 35-50.
 515 <https://doi.org/10.32604/phyton.2021.012276>.
 516 Li, X., Qu, C., Bian, Y., Gu, C., Jiang, X. and Song, Y., 2019. New insights into the
 517 responses of soil microorganisms to polycyclic aromatic hydrocarbon stress by
 518 combining enzyme activity and sequencing analysis with metabolomics. Environ
 519 Pollut 255(Pt 2), 113312. <https://doi.org/10.1016/j.envpol.2019.113312>.

- 520 Liang, J.Q., Xu, J.T., Zhao, W.J., Wang, J.F., Chen, K., Li, Y.Q. and Tian, Y., 2021.
- 521 Benzo[a]pyrene might be transported by a TonB-dependent transporter in
- 522 *Novosphingobium pentaromativorans* US6-1. J. Hazard. Mater. 404.
- 523 <https://doi.org/10.1016/j.jhazmat.2020.124037>.
- 524 Lin, Y., Ye, Y., Hu, Y. and Shi, H., 2019a. The variation in microbial community
- 525 structure under different heavy metal contamination levels in paddy soils.
- 526 Ecotoxicol Environ Saf 180, 557-564.
- 527 <https://doi.org/10.1016/j.ecoenv.2019.05.057>.
- 528 Lin, Y.B., Ye, Y.M., Hu, Y.M. and Shi, H.K., 2019b. The variation in microbial
- 529 community structure under different heavy metal contamination levels in paddy
- 530 soils. Ecotoxicol. Environ. Saf. 180, 557-564.
- 531 <https://doi.org/10.1016/j.ecoenv.2019.05.057>.
- 532 Liu, S.H., Zeng, G.M., Niu, Q.Y., Liu, Y., Zhou, L., Jiang, L.H., Tan, X.F., Xu, P.,
- 533 Zhang, C. and Cheng, M., 2017. Bioremediation mechanisms of combined
- 534 pollution of PAHs and heavy metals by bacteria and fungi: A mini review.
- 535 Bioresour. Technol. 224, 25-33. <https://doi.org/10.1016/j.biortech.2016.11.095>.
- 536 Liu, X.R., Liu, M., Chen, X., Yang, Y., Hou, L.J., Wu, S.X. and Zhu, P.K., 2019.
- 537 Indigenous PAH degraders along the gradient of the Yangtze Estuary of China:
- 538 Relationships with pollutants and their bioremediation implications. Mar. Pollut.
- 539 Bull. 142, 419-427. <https://doi.org/10.1016/j.marpolbul.2019.03.064>.

540 Madueno, L., Coppotelli, B.M., Festa, S., Alvarez, H.M. and Morelli, I.S., 2018a.
 541 Insights into the mechanisms of desiccation resistance of the Patagonian
 542 PAH-degrading strain *Sphingobium* sp. 22B. J Appl Microbiol 124(6),
 543 1532-1543. <https://doi.org/10.1111/jam.13742>.
 544 Madueno, L., Coppotelli, B.M., Festa, S., Alvarez, H.M. and Morelli, I.S., 2018b.
 545 Insights into the mechanisms of desiccation resistance of the Patagonian
 546 PAH-degrading strain *Sphingobium* sp. 22B. J. Appl. Microbiol. 124(6),
 547 1532-1543. <https://doi.org/10.1111/jam.13742>.
 548 Marsh, N.D., Ledesma, E.B., Wornat, M.J., Tan, M.P., Zhu, D.L. and Law, C.K., 2005.
 549 Newly identified products of benzene droplet combustion: Polycyclic aromatic
 550 hydrocarbons of three to ten rings. Polycycl. Aromat. Comp. 25(3), 227-244.
 551 <https://doi.org/10.1080/10406630591007143>.
 552 Miao, R., Guo, M., Zhao, X., Gong, Z., Jia, C., Li, X. and Zhuang, J., 2020. Response
 553 of soil bacterial communities to polycyclic aromatic hydrocarbons during the
 554 phyto-microbial remediation of a contaminated soil. Chemosphere 261, 127779.
 555 <https://doi.org/10.1016/j.chemosphere.2020.127779>.
 556 Park, M.K., Cho, H.K., Cho, I.G., Lee, S.E. and Choi, S.D., 2021. Contamination
 557 characteristics of polychlorinated naphthalenes in the agricultural soil of two
 558 industrial cities in South Korea. Chemosphere 273.
 559 <https://doi.org/10.1016/j.chemosphere.2021.129721>.

- 560 Picariello, E., Baldantoni, D. and De Nicola, F., 2020. Acute effects of PAH
561 contamination on microbial community of different forest soils. Environ. Pollut.
562 262, 114378. <https://doi.org/10.1016/j.envpol.2020.114378>.
- 563 Rachwal, M., Magieral, T. and Wawer, M., 2015. Coke industry and steel metallurgy
564 as the source of soil contamination by technogenic magnetic particles, heavy
565 metals and polycyclic aromatic hydrocarbons. Chemosphere 138, 863-873.
566 <https://doi.org/10.1016/j.chemosphere.2014.11.077>.
- 567 Rehman, M.Z.U., Rizwan, M., Hussain, A., Saqib, M., Ali, S., Sohail, M.I., Shafiq, M.
568 and Hafeez, F., 2018. Alleviation of cadmium (Cd) toxicity and minimizing its
569 uptake in wheat (*Triticum aestivum*) by using organic carbon sources in
570 Cd-spiked soil. Environ. Pollut. 241, 557-565.
571 <https://doi.org/10.1016/j.envpol.2018.06.005>.
- 572 Seleiman, M.F., Ali, S., Refay, Y., Rizwan, M., Alhammad, B.A. and El-Hendawy,
573 S.E., 2020. Chromium resistant microbes and melatonin reduced Cr uptake and
574 toxicity, improved physio-biochemical traits and yield of wheat in contaminated
575 soil. Chemosphere 250. <https://doi.org/10.1016/j.chemosphere.2020.126239>.
- 576 Shan, J., Ji, R., Yu, Y.J., Xie, Z.B. and Yan, X.Y., 2015. Biochar, activated carbon, and
577 carbon nanotubes have different effects on fate of C-14-catechol and microbial
578 community in soil. Sci. Rep.-UK 5. <https://doi.org/10.1038/srep16000>.

579 Shinde, V.L., Suneel, V., Rathore, C. and Shenoy, B.D., 2020. Degradation of tarballs
 580 using associated bacterial consortia. 3 Biotech. 10(3).
 581 <https://doi.org/10.1007/s13205-020-2095-8>.
 582 Song, W.H., Cao, F., Lin, Y.C., Haque, M.M., Wu, X., Zhang, Y.X., Zhang, C.Y., Xie,
 583 F. and Zhang, Y.L., 2021. Extremely high abundance of polycyclic aromatic
 584 hydrocarbons in aerosols from a typical coal-combustion rural site in China: Size
 585 distribution, source identification and cancer risk assessment. Atmos. Res. 248.
 586 <https://doi.org/10.1016/j.atmosres>.
 587 Sperfeld, M., Diekert, G. and Studenik, S., 2019. Anaerobic aromatic compound
 588 degradation in *Sulfuritalea hydrogenivorans* sk43H. Fems Microbiol Ecol 95(1).
 589 <https://doi.org/10.1093/femsec/fiy199>.
 590 Sun, L., Liao, X., Yan, X., Zhu, G. and Ma, D., 2014. Evaluation of heavy metal and
 591 polycyclic aromatic hydrocarbons accumulation in plants from typical industrial
 592 sites: potential candidate in phytoremediation for co-contamination. Environ Sci
 593 Pollut Res Int 21(21), 12494-12504. <https://doi.org/10.1007/s11356-014-3171-6>.
 594 Tang, X.J., Shen, C.F., Shi, D.Z., Cheema, S.A., Khan, M.I., Zhang, C.K. and Chen,
 595 Y.X., 2010. Heavy metal and persistent organic compound contamination in soil
 596 from Wenling: An emerging e-waste recycling city in Taizhou area, China. J.
 597 Hazard. Mater. 173(1-3), 653-660.
 598 <https://doi.org/10.1016/j.jhazmat.2009.08.134>.

- 599 Tao, Y., Li, W., Xue, B., Zhong, J., Yao, S. and Wu, Q., 2013. Different effects of
600 copper (II), cadmium (II) and phosphate on the sorption of phenanthrene on the
601 biomass of cyanobacteria. J. Hazard. Mater. 261, 21-28.
602 <https://doi.org/10.1016/j.jhazmat.2013.06.062>.
- 603 Tomczyk, B., Siatecka, A., Jedruchniewicz, K., Sochacka, A., Bogusz, A. and
604 Oleszczuk, P., 2020. Polycyclic aromatic hydrocarbons (PAHs) persistence,
605 bioavailability and toxicity in sewage sludge- or sewage sludge-derived
606 biochar-amended soil. Sci. Total Environ. 747.
607 <https://doi.org/10.1016/j.scitotenv.2020.141123>.
- 608 Usman, A.R.A., Almaroai, Y.A., Ahmad, M., Vithanage, M. and Ok, Y.S., 2013.
609 Toxicity of synthetic chelators and metal availability in poultry manure amended
610 Cd, Pb and As contaminated agricultural soil. J. Hazard. Mater. 262, 1022-1030.
611 <https://doi.org/10.1016/j.jhazmat.2013.04.032>.
- 612 Vasilieva, V., Scherr, K.E., Edelmann, E., Hasinger, M. and Loibner, A.P., 2012.
613 Comprehensive GC(2)/MS for the monitoring of aromatic tar oil constituents
614 during biodegradation in a historically contaminated soil. J. Biotechnol. 157(4),
615 460-466. <https://doi.org/10.1016/j.jbiotec.2011.08.006>.
- 616 Wang, Q., Hou, J., Yuan, J., Wu, Y., Liu, W., Luo, Y. and Christie, P., 2020. Evaluation
617 of fatty acid derivatives in the remediation of aged PAH-contaminated soil and
618 microbial community and degradation gene response. Chemosphere 248, 125983.
619 <https://doi.org/10.1016/j.chemosphere.2020.125983>.

- Wheatley, A.D. and Sadhra, S., 2004. Polycyclic aromatic hydrocarbons in solid residues from waste incineration. *Chemosphere* 55(5), 743-749. <https://doi.org/10.1016/j.chemosphere.2003.10.055>.
- Wu, B., Guo, S.H. and Wang, J.N., 2021. Assessment of the human health risk of polycyclic aromatic hydrocarbons in soils from areas of crude oil exploitation. *Environ. Res.* 193. <https://doi.org/10.1016/j.envres.2020.110617>.
- Wu, S.X., Liu, X.R., Liu, M., Chen, X., Liu, S., Cheng, L., Lin, X.B. and Li, Y., 2018. Sources, influencing factors and environmental indications of PAH pollution in urban soil columns of Shanghai, China. *Ecol. Indic.* 85, 1170-1180. <https://doi.org/10.1016/j.ecolind.2017.11.067>.
- Yang, J., Bertram, J., Schettgen, T., Heitland, P., Fischer, D., Seidu, F., Felten, M., Kraus, T., Fobil, J.N. and Kaifie, A., 2020a. Arsenic burden in e-waste recycling workers - A cross-sectional study at the Agbogbloshie e-waste recycling site, Ghana. *Chemosphere* 261. <https://doi.org/10.1016/j.chemosphere.2020.127712>.
- Yang, S.Y., Gu, S.B., He, M.J., Tang, X.J., Ma, L.N.Q., Xu, J.M. and Liu, X.M., 2020b. Policy adjustment impacts Cd, Cu, Ni, Pb and Zn contamination in soils around e-waste area: Concentrations, sources and health risks. *Sci. Total Environ.* 741. <https://doi.org/10.1016/j.scitotenv.2020.140442>.
- Yin, C., Xiong, W., Qiu, H., Peng, W., Deng, Z., Lin, S. and Liang, R., 2020. Characterization of the phenanthrene-degrading *Sphingobium yanoikuyae* SJTF8

in heavy metal co-existing liquid medium and analysis of its metabolic pathway.

Microorganisms 8(6). <https://doi.org/10.3390/microorganisms8060946>.

Yin, Y.J., Li, D., Wang, Y.Q., Xu, Z.H., Xu, G.J., Zhao, Z.Q., Li, S. and Song, L.Y.,
2019. Concurrent Removal of Mn(II) and Cr(VI) by *Achromobacter* sp. TY3-4.
Geomicrobiol. J. 36(4), 317-325.
<https://doi.org/10.1080/01490451.2018.1544324>.

Zajusz-Zubek, E., Radko, T. and Mainka, A., 2017. Fractionation of trace elements
and human health risk of submicron particulate matter (PM1) collected in the
surroundings of coking plants. Environ. Monit. Assess. 189(8).
<https://doi.org/10.1007/s10661-017-6117-x>.

Zhang, W.H., Zhuang, L.W., Yuan, Y.A., Tong, L.Z. and Tsang, D.C.W., 2011.
Enhancement of phenanthrene adsorption on a clayey soil and clay minerals by
coexisting lead or cadmium. Chemosphere 83(3), 302-310.
<https://doi.org/10.1016/j.chemosphere.2010.12.056>.

Zhang, Y.J. and Wang, J., 2011. Distribution and source of polycyclic aromatic
hydrocarbons (PAHs) in the surface soil along main transportation routes in
Jiaxing City, China. Environ. Monit. Assess. 182(1-4), 535-543.
<https://doi.org/10.1007/s10661-011-1895-z>.

Zhao, Q.Q., Bai, J.H., Gao, Y.C., Zhang, G.L., Lu, Q.Q. and Jia, J., 2021. Heavy metal
contamination in soils from freshwater wetlands to salt marshes in the Yellow

660 River Estuary, China. Sci. Total Environ. 774.
 661 <https://doi.org/10.1016/j.scitotenv.2021.145072>.
 662 Zhao, X.Q., Huang, J., Zhu, X.Y., Chai, J.C. and Ji, X.L., 2020. Ecological effects of
 663 heavy metal pollution on soil microbial community structure and diversity on
 664 both sides of a river around a mining Area. Int. J. Env. Res. Pub. He. 17(16).
 665 <https://doi.org/10.3390/ijerph17165680>.
 666 Zhou, Y.Y., Tang, L., Zeng, G.M., Zhang, C., Zhang, Y. and Xie, X., 2016. Current
 667 progress in biosensors for heavy metal ions based on DNazymes/DNA
 668 molecules functionalized nanostructures: A review. Sensor. Actuat. B-Chem. 223,
 669 280-294. <https://doi.org/10.1016/j.snb.2015.09.090>.

Figure 1







Figure 2a






[Click here to access/download;Figure;Figure 3a.tif](#)



[Click here to access/download;Figure;Figure 3b.tif](#)







Figure 4b

[Click here to access/download;Figure;Figure 4b.tif](#)






Figure 5b





Highlights

1. Mixed contamination (HMs+PAHs) was detected in soil from the e-waste dismantling plant and the coking plant.
2. The e-waste dismantling plant site was featured by heavy contamination of HMs but minor PAHs.
3. The coking plant was featured by severe contamination of PAHs and moderate HMs.
4. The abundance of bacterial genera such as *Sulfuritalea*, *Pseudomonas* and *Sphingobium* increased with severe pollution of PAHs.
5. The abundance of bacterial genera such as *Bryobacter*, *Galiella* and *Nitrospira* increased in soil contaminated with HMs.

