

DXXXXXXXXXXXXX  
XIXXXXXXXXXXXXX  
XXAXXXXXXXXXXXXX  
XXXGXXXXXXXXXXXX  
XXXOXXXXXXXXXXXX  
XXXXNXXXXXXXXXXX  
XXXXXXAXXXXXXXXX  
XXXXXXLXXXXXXXXX  
XXXXXXXXIXXXXXXX  
XXXXXXXXXSXXXXXX  
XXXXXXXXXXAXXXX  
XXXXXXXXXXXTXXX  
XXXXXXXXXXIXX  
XXXXXXXXXXOX  
XXXXXXXXXXNX

# Diagonalisation

## The Road to Infinities, Truth and Gödel

Rasmus Bakken — Mahin Hossain

University of Oxford

Michaelmas 2025

Summary: We introduce provability inside a formal system, prove Gödel's diagonal lemma, and use it to show the undefinability of truth in a formal logical setting.

*“All Cretans are liars.”*

— EPIMENIDES, A CRETAN

## Lecture 3 Summary

- What is a proof: define Hilbert-style system and axioms; example derivations.
- The Diagonal Lemma: construction of self-referential sentences via substitution and the diagonal operator.
- Undefinability of truth (Tarski): apply diagonal lemma to rule out a truth-defining formula in  $\mathcal{L}$ .

# Roadmap

1 What is a Proof?

2 The Diagonal Lemma

3 The Undefinability of Truth

## Motivating the concept

- From last time: a proof of  $\varphi_k$  is finite sequence of  $\mathcal{L}$ -sentences  $\varphi_1, \dots, \varphi_k$  with some condition.

# Motivating the concept

- From last time: a proof of  $\varphi_k$  is finite sequence of  $\mathcal{L}$ -sentences  $\varphi_1, \dots, \varphi_k$  with some condition.
- The relevant condition: for all  $1 \leq i \leq k$ 
  - $\varphi_i$  is an axiom, or

# Motivating the concept

- From last time: a proof of  $\varphi_k$  is finite sequence of  $\mathcal{L}$ -sentences  $\varphi_1, \dots, \varphi_k$  with some condition.
- The relevant condition: for all  $1 \leq i \leq k$ 
  - $\varphi_i$  is an axiom, or
  - $\varphi_i$  follows from  $\varphi_j$  and  $\varphi_m$  by some rule of inference where  $j \leq m < i$ .
- We use a Hilbert-style calculus.



# Motivating the concept

- From last time: a proof of  $\varphi_k$  is finite sequence of  $\mathcal{L}$ -sentences  $\varphi_1, \dots, \varphi_k$  with some condition.
- The relevant condition: for all  $1 \leq i \leq k$ 
  - $\varphi_i$  is an axiom, or
  - $\varphi_i$  follows from  $\varphi_j$  and  $\varphi_m$  by some rule of inference where  $j \leq m < i$ .
- We use a Hilbert-style calculus.
- Logical vs non-logical axioms, and only two rules of inference.

## Definition (First-order Predicate Logic)

Let  $\varphi$ ,  $\psi$  and  $\xi$  be  $\mathcal{L}$  formula, then the following are axioms of first-order logic:

**C1**  $\varphi \rightarrow (\psi \rightarrow \varphi).$

**C2**  $(\varphi \rightarrow (\psi \rightarrow \xi)) \rightarrow ((\varphi \rightarrow \psi) \rightarrow (\varphi \rightarrow \xi)).$

**C3**  $(\varphi \rightarrow \psi) \rightarrow (\neg\psi \rightarrow \neg\varphi).$

## Definition (First-order Predicate Logic)

Let  $\varphi$ ,  $\psi$  and  $\xi$  be  $\mathcal{L}$  formula, then the following are axioms of first-order logic:

C1  $\varphi \rightarrow (\psi \rightarrow \varphi)$ .

C2  $(\varphi \rightarrow (\psi \rightarrow \xi)) \rightarrow ((\varphi \rightarrow \psi) \rightarrow (\varphi \rightarrow \xi))$ .

C3  $(\varphi \rightarrow \psi) \rightarrow (\neg\psi \rightarrow \neg\varphi)$ .

C4  $\forall v_i \varphi(v_i) \rightarrow \varphi(t)$ , where  $t$  is a term of  $\mathcal{L}$  such that it does not contain a variable  $v_j$  where  $v_i$  occurs free in the scope of  $\forall v_j$  in  $\varphi$ .

C5  $\forall v_i(\varphi \rightarrow \psi) \rightarrow (\varphi \rightarrow \forall v_i \psi)$ , provided  $v_i$  does not occur free in  $\psi$ .

C6  $\forall v_i(v_i = v_i)$ .

C7 If  $F$  and  $G$  are atomic formulae, where  $G$  results from replacing some but not necessarily all of  $v_i$  in  $F$  by  $v_j$ , then  $\forall v_i \forall v_j(v_i = v_j \rightarrow (F \rightarrow G))$  is an axiom.

# Rules of Inference

## Definition

The rules of inference are the following:

**MP** From  $\varphi$  and  $\varphi \rightarrow \psi$  you may infer  $\psi$ .

**Gen** From  $\varphi$  you may infer  $\forall v_i \varphi$ .

Example: proving  $\varphi \rightarrow \varphi$

$$\mathbf{1} \quad (\varphi \rightarrow ((\varphi \rightarrow \varphi) \rightarrow \varphi)) \rightarrow ((\varphi \rightarrow (\varphi \rightarrow \varphi)) \rightarrow (\varphi \rightarrow \varphi)) \quad \text{C2}$$

## Example: proving $\varphi \rightarrow \varphi$

**1**  $(\varphi \rightarrow ((\varphi \rightarrow \varphi) \rightarrow \varphi)) \rightarrow ((\varphi \rightarrow (\varphi \rightarrow \varphi)) \rightarrow (\varphi \rightarrow \varphi))$  C2

**2**  $\varphi \rightarrow ((\varphi \rightarrow \varphi) \rightarrow \varphi)$  C1

## Example: proving $\varphi \rightarrow \varphi$

- 1  $(\varphi \rightarrow ((\varphi \rightarrow \varphi) \rightarrow \varphi)) \rightarrow ((\varphi \rightarrow (\varphi \rightarrow \varphi)) \rightarrow (\varphi \rightarrow \varphi))$  C2
- 2  $\varphi \rightarrow ((\varphi \rightarrow \varphi) \rightarrow \varphi)$  C1
- 3  $((\varphi \rightarrow (\varphi \rightarrow \varphi)) \rightarrow (\varphi \rightarrow \varphi))$  MP line 1 and line 2

## Example: proving $\varphi \rightarrow \varphi$

- 1  $(\varphi \rightarrow ((\varphi \rightarrow \varphi) \rightarrow \varphi)) \rightarrow ((\varphi \rightarrow (\varphi \rightarrow \varphi)) \rightarrow (\varphi \rightarrow \varphi))$  C2
- 2  $\varphi \rightarrow ((\varphi \rightarrow \varphi) \rightarrow \varphi)$  C1
- 3  $((\varphi \rightarrow (\varphi \rightarrow \varphi)) \rightarrow (\varphi \rightarrow \varphi))$  MP line 1 and line 2
- 4  $\varphi \rightarrow (\varphi \rightarrow \varphi)$  C1



## Example: proving $\varphi \rightarrow \varphi$

- 1  $(\varphi \rightarrow ((\varphi \rightarrow \varphi) \rightarrow \varphi)) \rightarrow ((\varphi \rightarrow (\varphi \rightarrow \varphi)) \rightarrow (\varphi \rightarrow \varphi))$  C2
- 2  $\varphi \rightarrow ((\varphi \rightarrow \varphi) \rightarrow \varphi)$  C1
- 3  $((\varphi \rightarrow (\varphi \rightarrow \varphi)) \rightarrow (\varphi \rightarrow \varphi))$  MP line 1 and line 2
- 4  $\varphi \rightarrow (\varphi \rightarrow \varphi)$  C1
- 5  $\varphi \rightarrow \varphi$  MP line 3 and 4

# Peano Arithmetic

## Definition (Peano Arithmetic)

Peano Arithmetic, or *PA* for short, is the theory given by the following set of axioms:

**PA1**  $\forall v_i (S(v_i) \neq 0)$ .

**PA2**  $\forall v_i \forall v_j (S(v_i) = S(v_j) \rightarrow v_i = v_j)$ .

# Peano Arithmetic

## Definition (Peano Arithmetic)

Peano Arithmetic, or *PA* for short, is the theory given by the following set of axioms:

**PA1**  $\forall v_i (S(v_i) \neq 0).$

**PA2**  $\forall v_i \forall v_j (S(v_i) = S(v_j) \rightarrow v_i = v_j).$

**PA3**  $\forall v_i (v_i + 0 = v_i).$

**PA4**  $\forall v_i \forall v_j (v_i + S(v_j) = S(v_i + v_j)).$

# Peano Arithmetic

## Definition (Peano Arithmetic)

Peano Arithmetic, or *PA* for short, is the theory given by the following set of axioms:

**PA1**  $\forall v_i (S(v_i) \neq 0).$

**PA2**  $\forall v_i \forall v_j (S(v_i) = S(v_j) \rightarrow v_i = v_j).$

**PA3**  $\forall v_i (v_i + 0 = v_i).$

**PA4**  $\forall v_i \forall v_j (v_i + S(v_j) = S(v_i + v_j)).$

**PA5**  $\forall v_i (v_i * 0 = 0).$

**PA6**  $\forall v_i \forall v_j (v_i * S(v_j) = v_i * v_j + v_i).$

# Peano Arithmetic

## Definition (Peano Arithmetic)

Peano Arithmetic, or *PA* for short, is the theory given by the following set of axioms:

PA1  $\forall v_i (S(v_i) \neq 0).$

PA2  $\forall v_i \forall v_j (S(v_i) = S(v_j) \rightarrow v_i = v_j).$

PA3  $\forall v_i (v_i + 0 = v_i).$

PA4  $\forall v_i \forall v_j (v_i + S(v_j) = S(v_i + v_j)).$

PA5  $\forall v_i (v_i * 0 = 0).$

PA6  $\forall v_i \forall v_j (v_i * S(v_j) = v_i * v_j + v_i).$

PA7  $\forall v_i (\exp(v_i, 0) = \bar{1}).$

PA8  $\forall v_i \forall v_j (\exp(v_i, S(v_j)) = \exp(v_i, v_j) * v_i).$

# Peano Arithmetic

## Definition (Peano Arithmetic)

Peano Arithmetic, or *PA* for short, is the theory given by the following set of axioms:

**PA1**  $\forall v_i (S(v_i) \neq 0).$

**PA2**  $\forall v_i \forall v_j (S(v_i) = S(v_j) \rightarrow v_i = v_j).$

**PA3**  $\forall v_i (v_i + 0 = v_i).$

**PA4**  $\forall v_i \forall v_j (v_i + S(v_j) = S(v_i + v_j)).$

**PA5**  $\forall v_i (v_i * 0 = 0).$

**PA6**  $\forall v_i \forall v_j (v_i * S(v_j) = v_i * v_j + v_i).$

**PA7**  $\forall v_i (\exp(v_i, 0) = \bar{1}).$

**PA8**  $\forall v_i \forall v_j (\exp(v_i, S(v_j)) = \exp(v_i, v_j) * v_i).$

**Ind $\mathcal{L}$**  For any  $\mathcal{L}$  formula with exactly one free variable  $\varphi(v_i)$ ,  
 $(\varphi(0) \wedge \forall v_i (\varphi(v_i) \rightarrow \varphi(S(v_i)))) \rightarrow \forall v_i \varphi(v_i)$  is an axiom.

## Example in PA: $\bar{1} + \bar{1} = \bar{2}$ (pt.1)

### Theorem

$PA \vdash \bar{1} + \bar{1} = \bar{2}$ .

### Proof.

**1**  $\forall v_i \forall v_j (v_i + S(v_j) = S(v_i + v_j))$  PA4

## Example in PA: $\bar{1} + \bar{1} = \bar{2}$ (pt.1)

### Theorem

$$PA \vdash \bar{1} + \bar{1} = \bar{2}.$$

### Proof.

- 1  $\forall v_i \forall v_j (v_i + S(v_j) = S(v_i + v_j))$  PA4
- 2  $\forall v_i \forall v_j (v_i + S(v_j) = S(v_i + v_j)) \rightarrow \forall v_j (S(0) + S(v_j) = S(S(0) + v_j))$   
C4
- 3  $\forall v_j (S(0) + S(v_j) = S(S(0) + v_j))$  MP with line 1 and line 2



## Example in PA: $\bar{1} + \bar{1} = \bar{2}$ (pt.1)

### Theorem

$$PA \vdash \bar{1} + \bar{1} = \bar{2}.$$

### Proof.

- 1  $\forall v_i \forall v_j (v_i + S(v_j) = S(v_i + v_j))$  PA4
- 2  $\forall v_i \forall v_j (v_i + S(v_j) = S(v_i + v_j)) \rightarrow \forall v_j (S(0) + S(v_j) = S(S(0) + v_j))$   
C4
- 3  $\forall v_j (S(0) + S(v_j) = S(S(0) + v_j))$  MP with line 1 and line  
2
- 4  $\forall v_j (S(0) + S(v_j) = S(S(0) + v_j)) \rightarrow S(0) + S(0) = S(S(0) + 0)$  C4
- 5  $S(0) + S(0) = S(S(0) + 0)$  MP with line 3 and 4



## Example in PA: $\bar{1} + \bar{1} = \bar{2}$ (pt.2)

### Theorem

$PA \vdash \bar{1} + \bar{1} = \bar{2}$ .

### Proof.

**5**  $S(0) + S(0) = S(S(0) + 0)$  MP with line 3 and 4

**6**  $\forall v_i (v_i + 0 = v_i)$  PA3

## Example in PA: $\bar{1} + \bar{1} = \bar{2}$ (pt.2)

### Theorem

$PA \vdash \bar{1} + \bar{1} = \bar{2}$ .

### Proof.

**5**  $S(0) + S(0) = S(S(0) + 0)$  MP with line 3 and 4

**6**  $\forall v_i (v_i + 0 = v_i)$  PA3

**7**  $\forall v_i (v_i + 0 = v_i) \rightarrow S(0) + 0 = S(0)$  C4

**8**  $S(0) + 0 = S(0)$  MP with line 6 and line  
7



## Example in PA: $\overline{1} + \overline{1} = \overline{2}$ (pt.3)

### Theorem

$PA \vdash \overline{1} + \overline{1} = \overline{2}$ .

### Proof.

**5**  $S(0) + S(0) = S(S(0) + 0)$  MP with line 3 and 4

**8**  $S(0) + 0 = S(0)$  MP with line 6 and line  
7

**9**  $\forall v_i \forall v_j (v_i = v_j \rightarrow (S(0) + S(0) = S(v_i) \rightarrow S(0) + S(0) = S(v_j)))$  C7

## Example in PA: $\bar{1} + \bar{1} = \bar{2}$ (pt.3)

### Theorem

$$PA \vdash \bar{1} + \bar{1} = \bar{2}.$$

### Proof.

**5**  $S(0) + S(0) = S(S(0) + 0)$  MP with line 3 and 4

**8**  $S(0) + 0 = S(0)$  MP with line 6 and line  
7

**9**  $\forall v_i \forall v_j (v_i = v_j \rightarrow (S(0) + S(0) = S(v_i) \rightarrow S(0) + S(0) = S(v_j)))$  C7

**10**  $\forall v_i \forall v_j (v_i = v_j \rightarrow (S(0) + S(0) = S(v_i) \rightarrow S(0) + S(0) = S(v_j))) \rightarrow \forall v_j (S(0) + 0 = v_j \rightarrow (S(0) + S(0) = S(S(0) + 0) \rightarrow S(0) + S(0) = S(v_j)))$  C4

**11**  $\forall v_j (S(0) + 0 = v_j \rightarrow (S(0) + S(0) = S(S(0) + 0) \rightarrow S(0) + S(0) = S(v_j)))$   
MP line 9 and line 10



## Example in PA: $\bar{1} + \bar{1} = \bar{2}$ (pt.4)

### Theorem

$$PA \vdash \bar{1} + \bar{1} = \bar{2}.$$

### Proof.

**5**  $S(0) + S(0) = S(S(0) + 0)$  MP with line 3 and 4

**8**  $S(0) + 0 = S(0)$  MP with line 6 and line  
7

**11**  $\forall v_j (S(0) + 0 = v_j \rightarrow (S(0) + S(0) = S(S(0) + 0) \rightarrow S(0) + S(0) = S(v_j)))$   
MP line 9 and line 10

## Example in PA: $\bar{1} + \bar{1} = \bar{2}$ (pt.4)

### Theorem

$$PA \vdash \bar{1} + \bar{1} = \bar{2}.$$

### Proof.

**5**  $S(0) + S(0) = S(S(0) + 0)$  MP with line 3 and 4

**8**  $S(0) + 0 = S(0)$  MP with line 6 and line  
7

**11**  $\forall v_j (S(0) + 0 = v_j \rightarrow (S(0) + S(0) = S(S(0) + 0) \rightarrow S(0) + S(0) = S(v_j)))$   
MP line 9 and line 10

**12**  $\forall v_j (S(0) + 0 = v_j \rightarrow (S(0) + S(0) = S(S(0) + 0) \rightarrow S(0) + S(0) = S(v_j))) \rightarrow S(0) + 0 = S(0) \rightarrow (S(0) + S(0) = S(S(0))) \rightarrow S(0) + S(0) = S(S(0)))$  C4

**13**  $S(0) + 0 = S(0) \rightarrow (S(0) + S(0) = S(S(0) + 0) \rightarrow S(0) + S(0) = S(S(0)))$   
MP line 11 and line 12

## Example in PA: $\bar{1} + \bar{1} = \bar{2}$ (pt.4)

### Theorem

$$PA \vdash \bar{1} + \bar{1} = \bar{2}.$$

### Proof.

**5**  $S(0) + S(0) = S(S(0) + 0)$  MP with line 3 and 4

**8**  $S(0) + 0 = S(0)$  MP with line 6 and line  
7

**11**  $\forall v_j (S(0) + 0 = v_j \rightarrow (S(0) + S(0) = S(S(0) + 0) \rightarrow S(0) + S(0) = S(v_j)))$   
MP line 9 and line 10

**12**  $\forall v_j (S(0) + 0 = v_j \rightarrow (S(0) + S(0) = S(S(0) + 0) \rightarrow S(0) + S(0) = S(v_j))) \rightarrow S(0) + 0 = S(0) \rightarrow (S(0) + S(0) = S(S(0))) \rightarrow S(0) + S(0) = S(S(0)))$  C4

**13**  $S(0) + 0 = S(0) \rightarrow (S(0) + S(0) = S(S(0) + 0) \rightarrow S(0) + S(0) = S(S(0)))$   
MP line 11 and line 12

**14**  $(S(0) + S(0) = S(S(0) + 0) \rightarrow S(0) + S(0) = S(S(0)))$  MP line 8 and line 13



## Example in PA: $\bar{1} + \bar{1} = \bar{2}$ (pt.5)

### Theorem

$PA \vdash \bar{1} + \bar{1} = \bar{2}$ .

### Proof.

**5**  $S(0) + S(0) = S(S(0) + 0)$  MP with line 3 and 4

**14**  $(S(0) + S(0) = S(S(0) + 0) \rightarrow S(0) + S(0) = S(S(0)))$  MP line 8 and line 13

**15**  $S(0) + S(0) = S(S(0))$  MP line 5 and line 14



# Roadmap

1 What is a Proof?

2 The Diagonal Lemma

3 The Undefinability of Truth

# Gödel's Diagonal Lemma

## Diagonal Lemma

If  $\varphi(v_n)$  is an  $\mathcal{L}$ -formula with one free variable, then there is an  $\mathcal{L}$ -sentence  $\gamma$  such that

$$PA \vdash \gamma \leftrightarrow \varphi(\ulcorner \gamma \urcorner).$$

- For any one-place formula, we can manufacture a sentence that says of itself that it has property  $\varphi$ .
- This turns Gödel codes + substitution into genuine *self-reference*.

# Substitution on Formulae

- Given a formula  $\varphi$  and a term  $t$ ,  $\varphi(t/v_i)$  means:
  - uniformly replace all *free* occurrences of  $v_i$  in  $\varphi$  by  $t$ .
- Example:  $\varphi$  is  $S(v_5) = S(S(0))$ .
  - $\varphi(S(0)/v_5)$  is  $S(S(0)) = S(S(0))$ .
  - $\varphi(0/v_5)$  is  $S(0) = S(S(0))$ .
  - $\varphi(v_3/v_5)$  is  $S(v_3) = S(S(0))$ .
- Beware **variable capture**:
  - quantifiers can accidentally bind new free variables,
  - we avoid this by renaming bound variables before substituting.

# Internalising Substitution: The Function *sub*

## Lemma (informal)

There is an  $\mathcal{L}$ -definable function  $sub(v_1, v_2, v_3)$  such that

$$PA \vdash \overline{\varphi} = sub(\overline{\psi}, \overline{n}, \overline{t})$$

iff  $\varphi = \psi(t/v_n)$ .

- So substitution on formulas can be mirrored by a calculation on Gödel numbers.
- We take the existence of such a *sub*-function for granted (construction is tedious but standard).

# The Diagonal Operator

## Definition (Diagonal Operator)

Let  $\varphi(v_n)$  be a formula with  $v_n$  its only free variable. Define

$$\text{dia}(\overline{\ulcorner \varphi(v_n) \urcorner}) := \text{sub}(\overline{\ulcorner \varphi \urcorner}, \overline{n}, \overline{\overline{\ulcorner \varphi \urcorner}}).$$

- Intuition:

$$\text{dia}(\overline{\ulcorner \varphi \urcorner}) = \overline{\ulcorner \varphi(\overline{\ulcorner \varphi \urcorner} / v_n) \urcorner}.$$

- It takes the code of  $\varphi$  and returns the code of the result of plugging that very code back into  $\varphi$ .
- This is the core mechanism behind self-reference.

# Proof Idea of the Diagonal Lemma

- Fix a formula  $\varphi(v_n)$ .
- Consider the auxiliary formula

$$\theta(v_n) := \exists x (x = \text{dia}(v_n) \wedge \varphi(x)).$$

- Now look at the sentence

$$\gamma := \exists x (x = \text{dia}(\overline{\ulcorner \theta \urcorner}) \wedge \varphi(x)).$$

- Using the properties of *sub* and *dia*, one checks inside *PA* that

$$\text{dia}(\overline{\ulcorner \theta \urcorner}) = \overline{\ulcorner \gamma \urcorner}.$$

- Hence *PA* proves

$$\gamma \leftrightarrow \varphi(\overline{\ulcorner \gamma \urcorner}),$$

which is exactly the conclusion of the Diagonal Lemma.

# Roadmap

1 What is a Proof?

2 The Diagonal Lemma

3 The Undefinability of Truth



# What Should a Truth Predicate Do?

- Suppose  $\varphi(v_i)$  is meant to say:

$$\varphi(\ulcorner \psi \urcorner) = \text{“}\psi \text{ is true”}.$$

- We would like  $\varphi$  to respect logical structure, e.g. disjunction:

$$PA \vdash \varphi(\overline{\ulcorner \psi_1 \vee \psi_2 \urcorner}) \leftrightarrow \varphi(\overline{\ulcorner \psi_1 \urcorner}) \vee \varphi(\overline{\ulcorner \psi_2 \urcorner}).$$

- More strongly: there is a coding function  $x \dot{\vee} y$  with

$$x = \ulcorner \psi_1 \urcorner, y = \ulcorner \psi_2 \urcorner \Rightarrow x \dot{\vee} y = \ulcorner \psi_1 \vee \psi_2 \urcorner,$$

and we might want

$$PA \vdash \forall x \forall y (\varphi(x \dot{\vee} y) \leftrightarrow \varphi(x) \vee \varphi(y)).$$

## Tarski Biconditionals (for $\varphi$ )

For every  $\mathcal{L}$ -sentence  $\psi$ ,

$$PA \vdash \varphi(\overline{\ulcorner \psi \urcorner}) \leftrightarrow \psi.$$

- This is a very natural *minimal* requirement for a truth definition.
- Read: “ $\varphi$  holds of the code of  $\psi$  iff  $\psi$ .”
- Tarski’s theorem: no such  $\varphi$  exists in the language of  $PA$ .

# Tarski's Undefinability Theorem

## Theorem (Tarski)

There is no  $\mathcal{L}$ -formula  $\varphi(v_i)$  such that for all  $\mathcal{L}$ -sentences  $\psi$ ,

$$PA \vdash \varphi(\overline{\ulcorner \psi \urcorner}) \leftrightarrow \psi.$$

- So: the intuitive notion of *truth in the standard model of PA* is not arithmetically definable.
- This contrasts with many other notions (e.g. “even”, “provable in *PA*”) which *are* definable in  $\mathcal{L}$ .

# Proof Idea: The Liar Sentence

## Proof sketch

Assume, for contradiction, that such a  $\varphi(v_i)$  exists.

- Apply the Diagonal Lemma to  $\neg\varphi(v_i)$ .
- Obtain a sentence  $\lambda$  such that

$$PA \vdash \lambda \leftrightarrow \neg\varphi(\ulcorner \lambda \urcorner).$$

- But by the Tarski biconditionals we also have

$$PA \vdash \lambda \leftrightarrow \varphi(\ulcorner \lambda \urcorner).$$

- Combining:

$$PA \vdash \lambda \leftrightarrow \neg\lambda,$$

contradicting the consistency of  $PA$ .

# The Liar and Beyond

- The sentence  $\lambda$  above is a formal **liar sentence**:

$\lambda$  says “I am not true (w.r.t.  $\varphi$ )”.

- Tarski's theorem shows:
  - No  $\mathcal{L}$ -formula can define a fully adequate truth predicate for arithmetic.
  - Truth is strictly more complex than many other arithmetical properties.
- One response: extend the language with a new primitive truth predicate  $T$ .
- Example theory:

$$TB := PA \cup \{ T^{\overline{\ulcorner \psi \urcorner}} \leftrightarrow \psi \mid \psi \text{ an } \mathcal{L}\text{-sentence} \}.$$

- The liar reappears, but now at the *extended* level, and must be handled with care.