

# Econometrics

## Properties of OLS and the FWL Theorem

J. Eduardo Vera-Valdés  
eduardo@math.aau.dk

Department of Mathematical Sciences  
Aalborg University  
Denmark



**AALBORG UNIVERSITY**  
DENMARK

# Outline



## Introduction

Formalities

## Geometrical Properties of OLS

## The Frisch-Waugh-Lovell Theorem

Applications

Goodness of Fit

## OLS is BLUE

Bias and Consistency

Precision

## Summing Up

Econometrics

J. Eduardo

Vera-Valdés

eduardo@math.aau.

Introduction

Formalities

Geometrical

Properties of OLS

The Frisch-

Waugh-Lovell

Theorem

Applications

Goodness of Fit

OLS is BLUE

Bias and Consistency

Precision

Summing Up

# Outline



## Introduction

### Formalities

## Geometrical Properties of OLS

## The Frisch-Waugh-Lovell Theorem

### Applications

### Goodness of Fit

## OLS is BLUE

### Bias and Consistency

### Precision

## Summing Up

### Econometrics

J. Eduardo

Vera-Valdés

eduardo@math.aau.

### 2 Introduction

Formalities

Geometrical  
Properties of OLS

The Frisch-  
Waugh-Lovell  
Theorem

Applications

Goodness of Fit

OLS is BLUE

Bias and Consistency

Precision

Summing Up

Department of  
Mathematical Sciences  
Aalborg University  
Denmark

# Formalities



**Course:** 10 lectures followed by exercise sessions and 2 self-studies. (1 extra “in case” lecture).

**Evaluation:** Oral exam without preparation.

**Literature:** “Econometric Theory and Methods” (ETM) by Russell Davidson and James G. MacKinnon.

Alternatively, “Econometric Analysis” (EA) by William H. Greene for a more analytical perspective, and “A Guide to Modern Econometrics” (AGME) by Marno Verbeek for a more intuitive approach.

**Software:** R via RStudio.

**Data:** The “Ecdat” package in R, but more importantly, data from your projects if appropriate.

Econometrics

J. Eduardo  
Vera-Valdés

eduardo@math.aau.

Introduction

3

Formalities

Geometrical

Properties of OLS

The Frisch-  
Waugh-Lovell  
Theorem

Applications

Goodness of Fit

OLS is BLUE

Bias and Consistency

Precision

Summing Up

## Knowledge

- ▶ understanding of the most common estimation methods in econometrics.
- ▶ knowledge about finite sample and asymptotic properties of the ordinary least squares estimator.
- ▶ knowledge about method of moments estimators, e.g., the instrumental variables estimator.
- ▶ knowledge about models for panel data, e.g., random effects and fixed effects estimators.
- ▶ knowledge about models for discrete and limited dependent variables, e.g., censored data and sample selection.
- ▶ knowledge about estimation of multivariate models and systems of equations.
- ▶ knowledge about simulation methods for inference, e.g., Monte Carlo and bootstrapped tests.

Econometrics

J. Eduardo  
Vera-Valdés  
eduardo@math.aau.

Introduction

4 Formalities

Geometrical  
Properties of OLS

The Frisch-  
Waugh-Lovell  
Theorem

Applications

Goodness of Fit

OLS is BLUE

Bias and Consistency

Precision

Summing Up

## Skills

- ▶ are able to argue for the importance of using econometric/statistical methods in the analysis of a given economic problem.
- ▶ are able to build econometric models and judge their applicability.

Econometrics

J. Eduardo  
Vera-Valdés

eduardo@math.aau.

Introduction

5

Formalities

Geometrical  
Properties of OLS

The Frisch-  
Waugh-Lovell  
Theorem

Applications

Goodness of Fit

OLS is BLUE

Bias and Consistency

Precision

Summing Up

## Competencies

- ▶ are able to demonstrate deep understanding of the theory of econometric models and know how to reason within the models.
- ▶ are able to communicate the results of an econometric analysis to non-specialists.
- ▶ are able to analyse economic data using the available software.

Econometrics

J. Eduardo  
Vera-Valdés

eduardo@math.aau.

Introduction

6

Formalities

Geometrical  
Properties of OLS

The Frisch-  
Waugh-Lovell  
Theorem

Applications

Goodness of Fit

OLS is BLUE

Bias and Consistency

Precision

Summing Up

# Schedule (subject to change)



Lectures on Wednesdays from 8.15 to 12.00 (two exceptions noted below)

Week	Subject
1	Properties of OLS and FWL Theorem
2	Misspecification
3	GLS and FGLS
4	Instrumental Variables
5	Panel Data Models
6	Self-study
7	Seemingly Unrelated Regressions
8	Simultaneous Equations Models
9	GARCH Models*
10	Cointegration*
11	Self-study
12	Censored and Truncated Data

Econometrics

J. Eduardo  
Vera-Valdés

eduardo@math.aau.

Introduction

Formalities

Geometrical  
Properties of OLS

The Frisch-  
Waugh-Lovell  
Theorem

Applications

Goodness of Fit

OLS is BLUE

Bias and Consistency

Precision

Summing Up

Department of  
Mathematical Sciences  
Aalborg University  
Denmark



# Outline



Introduction

Formalities

Geometrical Properties of OLS

The Frisch-Waugh-Lovell Theorem

Applications

Goodness of Fit

OLS is BLUE

Bias and Consistency

Precision

Summing Up

Econometrics

J. Eduardo

Vera-Valdés

eduardo@math.aau.

Introduction

Formalities

8

Geometrical  
Properties of OLS

The Frisch-  
Waugh-Lovell  
Theorem

Applications

Goodness of Fit

OLS is BLUE

Bias and Consistency

Precision

Summing Up

Department of  
Mathematical Sciences  
Aalborg University  
Denmark

40

# Ordinary Least Squares



## Assumptions for OLS:

- ▶ Correct specification
- ▶ Linear independence
- ▶ Exogeneity
- ▶ Homoskedasticity
- ▶ No autocorrelation
- ▶ Normality

Econometrics

J. Eduardo  
Vera-Valdés  
eduardo@math.aau.

Introduction

Formalities

9 Geometrical  
Properties of OLS

The Frisch-  
Waugh-Lovell  
Theorem

Applications

Goodness of Fit

OLS is BLUE

Bias and Consistency

Precision

Summing Up

Department of  
Mathematical Sciences  
Aalborg University  
Denmark

# Orthogonal Projections



When we estimate a regression model, we map  $Y$  into a vector of fitted values  $X\hat{\beta}$  and a vector of residuals  $\hat{U}$ .

OLS uses the two projection matrices:

$$P_X = X(X^T X)^{-1} X^T, \quad \text{and} \quad M_X = I - P_X.$$

And make,

$$X\hat{\beta} = P_X Y, \quad \text{and} \quad \hat{U} = M_X Y.$$

It is easy to show that  $P_X$  and  $M_X$  are orthogonal projection matrices, and that  $P_X X = X$ , and  $M_X X = 0$ .

Furthermore, they are **complementary projections** since  $P_X M_X = 0$  (which you will show).

Econometrics

J. Eduardo  
Vera-Valdés

eduardo@math.aau.

Introduction

Formalities

10 Geometrical  
Properties of OLS

The Frisch-  
Waugh-Lovell  
Theorem

Applications

Goodness of Fit

OLS is BLUE

Bias and Consistency

Precision

Summing Up

# Orthogonal Projections




OLS decomposes  $Y$  in

$$Y = P_X Y + M_X Y,$$

where  $X\hat{\beta} = P_X Y$  are the fitted values, and  $\hat{U} = M_X Y$  are the residuals.

This decomposition can be represented by a right-angled triangle.



Econometrics

J. Eduardo  
Vera-Valdés  
eduardo@math.aau.

Introduction

Formalities

11 Geometrical  
Properties of OLS

The Frisch-  
Waugh-Lovell  
Theorem

Applications

Goodness of Fit

OLS is BLUE

Bias and Consistency

Precision

Summing Up

Department of  
Mathematical Sciences  
Aalborg University  
Denmark

# Orthogonal Projections



By Pythagora's Theorem,

$$||Y||^2 = ||P_X Y||^2 + ||M_X Y||^2,$$

so that

$$||P_X Y||^2 \leq ||Y||^2, \text{ and } ||\hat{U}||^2 \leq ||U||^2.$$

Moreover, note that

$$X^T \hat{U} = X^T M_X Y = 0,$$

so that **OLS residuals are orthogonal to all the regressors.**

In particular, if the regressors include a constant, then the residuals sum to zero,  $\sum_{t=1}^N \hat{U}_t = 0$ .

Econometrics

J. Eduardo  
Vera-Valdés  
eduardo@math.aau.

Introduction

Formalities

12 Geometrical  
Properties of OLS

The Frisch-  
Waugh-Lovell  
Theorem

Applications

Goodness of Fit

OLS is BLUE

Bias and Consistency

Precision

Summing Up

Department of  
Mathematical Sciences  
Aalborg University  
Denmark

# Outline



Introduction

Formalities

Geometrical Properties of OLS

The Frisch-Waugh-Lovell Theorem

Applications

Goodness of Fit

OLS is BLUE

Bias and Consistency

Precision

Summing Up

Econometrics

J. Eduardo

Vera-Valdés

eduardo@math.aau.

Introduction

Formalities

Geometrical

Properties of OLS

13

The Frisch-Waugh-Lovell Theorem

Applications

Goodness of Fit

OLS is BLUE

Bias and Consistency

Precision

Summing Up

Department of  
Mathematical Sciences  
Aalborg University  
Denmark

40

# The Frisch-Waugh-Lovell Theorem

## Groups of Regressors



Assume we broke up the regressors into two groups  $X = [X_1 \ X_2]$  so that

$$Y = X_1\beta_1 + X_2\beta_2 + U,$$

we are interested in analysing the effect of this partition on the estimators.

First, suppose that  $X_1$  is orthogonal to  $X_2$ , then we obtain the same estimate for  $\beta_1$  above than the one using the regression

$$Y = X_1\beta_1 + V.$$

Analogously, we obtain the same estimate for  $\beta_2$  if we remove  $X_1$  from the regression.

Econometrics

J. Eduardo  
Vera-Valdés  
eduardo@math.aau.

Introduction

Formalities

Geometrical

Properties of OLS

14

**The Frisch-  
Waugh-Lovell  
Theorem**

Applications

Goodness of Fit

OLS is BLUE

Bias and Consistency

Precision

Summing Up

# The Frisch-Waugh-Lovell Theorem

## Groups of Regressors



In the general case, the two regressions

$$Y = X_1\beta_1 + M_{X_1}X_2\beta_2 + U,$$

and

$$Y = M_{X_1}X_2\beta_2 + V,$$

yield identical estimates for  $\beta_2$ .

Nonetheless, they do not yield the same residuals.

Econometrics

J. Eduardo

Vera-Valdés

eduardo@math.aau.

Introduction

Formalities

Geometrical

Properties of OLS

15

**The Frisch-Waugh-Lovell Theorem**

Applications

Goodness of Fit

OLS is BLUE

Bias and Consistency

Precision

Summing Up



# The Frisch-Waugh-Lovell Theorem



## The FWL Theorem

The OLS estimates of  $\beta_2$  in the regressions

$$Y = X_1\beta_1 + X_2\beta_2 + U,$$

and

$$M_{X_1}Y = M_{X_1}X_2\beta_2 + U,$$

are numerically identical.

Moreover, the residuals in both regressions are numerically identical.

Econometrics

J. Eduardo

Vera-Valdés

eduardo@math.aau.

Introduction

Formalities

Geometrical

Properties of OLS

16

**The Frisch-Waugh-Lovell Theorem**

Applications

Goodness of Fit

OLS is BLUE

Bias and Consistency

Precision

Summing Up

# The Frisch-Waugh-Lovell Theorem

## Applications



Assume the regression includes a constant

$$Y = \iota\beta_0 + X\beta_1 + U,$$

where  $\iota$  is a vector of ones.

Then the FWL theorem states that the estimator from  $\beta_1$  is the same if we instead run the regression

$$M_\iota Y = M_\iota X\beta_1 + U.$$

Econometrics

J. Eduardo  
Vera-Valdés

eduardo@math.aau.

Introduction

Formalities

Geometrical

Properties of OLS

The Frisch-  
Waugh-Lovell  
Theorem

17

Applications

Goodness of Fit

OLS is BLUE

Bias and Consistency

Precision

Summing Up

40

Department of  
Mathematical Sciences  
Aalborg University  
Denmark

# The Frisch-Waugh-Lovell Theorem

## Applications



Many economic activities are strongly affected by the season, for obvious reasons like Christmas shopping, or summer holidays.

It is thus common to model this seasonality using seasonal dummy variables.

$$Y = \alpha_1 s_1 + \alpha_2 s_2 + \alpha_3 s_3 + \alpha_4 s_4 + X\beta + U,$$

where are the  $s_i$  seasonal dummy variables.

The FWL theorem tells us that we can estimate  $\beta$  using the raw data and the above regression or by a simpler regression without the dummy variables using deseasonalised (by regression) data.

Econometrics

J. Eduardo  
Vera-Valdés  
eduardo@math.aau.

Introduction

Formalities

Geometrical  
Properties of OLS

The Frisch-  
Waugh-Lovell  
Theorem

18 Applications

Goodness of Fit

OLS is BLUE

Bias and Consistency

Precision

Summing Up

Department of  
Mathematical Sciences  
Aalborg University  
Denmark

# The Frisch-Waugh-Lovell Theorem

## Applications



Another set of variables that are often encountered are time trends.

We consider a regression like

$$Y = \alpha_0 \iota + \alpha_1 T + X\beta + U,$$

where  $T^T = [1, 2, \dots, N]$ .

An increasing time trend might be appropriate, for instance, in a model of a production function where technical progress is taking place.

The FWL Theorem shows that it is possible to use detrended variables.

Econometrics

J. Eduardo  
Vera-Valdés

eduardo@math.aau.

Introduction

Formalities

Geometrical  
Properties of OLS

The Frisch-  
Waugh-Lovell  
Theorem

19 Applications

Goodness of Fit

OLS is BLUE

Bias and Consistency

Precision

Summing Up

# The Frisch-Waugh-Lovell Theorem

## Goodness of Fit



Recall that OLS decomposes  $Y$  in

$$Y = P_X Y + M_X Y,$$

where  $P_X Y$  is the part that  $X$  is able to explain from  $Y$ .

We define the coefficient of determination or (uncentered)  $R^2$  by

$$R^2 = \frac{||P_X Y||^2}{||Y||^2},$$

which fall between 0 and 1 given what we previously discussed.

It is invariant under nonsingular linear transformations of the regressors and, because it is defined as a ratio, it is invariant to changes in the scale of  $Y$ .

Econometrics

J. Eduardo

Vera-Valdés

eduardo@math.aau.

Introduction

Formalities

Geometrical

Properties of OLS

The Frisch-

Waugh-Lovell

Theorem

Applications

20

Goodness of Fit

OLS is BLUE

Bias and Consistency

Precision

Summing Up

40

# The Frisch-Waugh-Lovell Theorem

## Goodness of Fit



Nonetheless, the (uncentered)  $R^2$  is not invariant to translations

$$\tilde{Y} := Y + \alpha \iota = X\beta + U.$$

If  $X$  includes a constant, we have that

$$\tilde{Y} = P_X(\tilde{Y}) + M_X(\tilde{Y}) = P_X Y + \alpha \iota + M_X Y,$$

so that

$$R^2 = \frac{\|P_X Y + \alpha \iota\|^2}{\|Y + \alpha \iota\|^2}.$$

By choosing  $\alpha$ , we can make the  $R^2$  as close to 1 as we want.

Econometrics

J. Eduardo  
Vera-Valdés

eduardo@math.aau.

Introduction

Formalities

Geometrical

Properties of OLS

The Frisch-  
Waugh-Lovell  
Theorem

Applications

21

Goodness of Fit

OLS is BLUE

Bias and Consistency

Precision

Summing Up

40

Department of  
Mathematical Sciences  
Aalborg University  
Denmark

# The Frisch-Waugh-Lovell Theorem

## Goodness of Fit



The last result is misleading because the fit is accounted almost exclusively by the constant.

To avoid this problem, for regressions that **include a constant**, the FWL theorem tells us that we can demean the series without changing the estimates or residuals.

This gives rise to the (centered)  $R^2$  defined as

$$R^2 = \frac{||P_X M_\iota Y||^2}{||M_\iota Y||^2},$$

which is unaffected by translations.

Econometrics

J. Eduardo  
Vera-Valdés  
eduardo@math.aau.

Introduction

Formalities

Geometrical  
Properties of OLS

The Frisch-  
Waugh-Lovell  
Theorem

Applications

22 Goodness of Fit

OLS is BLUE

Bias and Consistency

Precision

Summing Up

Department of  
Mathematical Sciences  
Aalborg University  
Denmark

# Outline



Introduction

Formalities

Geometrical Properties of OLS

The Frisch-Waugh-Lovell Theorem

Applications

Goodness of Fit

OLS is BLUE

Bias and Consistency

Precision

Summing Up

Econometrics

J. Eduardo

Vera-Valdés

eduardo@math.aau.

Introduction

Formalities

Geometrical

Properties of OLS

The Frisch-

Waugh-Lovell

Theorem

Applications

Goodness of Fit

23

OLS is BLUE

Bias and Consistency

Precision

Summing Up

Department of  
Mathematical Sciences  
Aalborg University  
Denmark

40



# OLS is BLUE



## The Gauss-Markov Theorem

In a regression under correct specification, exogenous regressors, homoskedastic and no-autocorrelated errors, then the OLS estimator is more efficient than any other linear unbiased estimator.

In this sense, the OLS estimator is the Best Linear Unbiased Estimator, or BLUE.

Econometrics

J. Eduardo

Vera-Valdés

eduardo@math.aau.

Introduction

Formalities

Geometrical

Properties of OLS

The Frisch-  
Waugh-Lovell  
Theorem

Applications

Goodness of Fit

24 OLS is BLUE

Bias and Consistency

Precision

Summing Up

To show that  $E[(X^T X)^{-1} X^T U] = 0$ , and thus that  $\hat{\beta}$  is unbiased, there are two possibilities.

(a)  $X$  is **nonstochastic** and  $U$  has mean zero.

(b)  $X$  is **exogenous**, i.e.,  $E[U|X] = 0$ .

The nonstochastic assumption may be sensible when we are able to control the inputs, like in an experimental setting.

Nonetheless, it is often not a reasonable assumption to make in applied econometric work.

The exogeneity assumption is reasonable in the context of some types of data like cross-sectional data in which each observation might correspond to an individual household or person.

On the other hand, for time-series data, in which each observation might correspond to a year, quarter, or day, the exogeneity assumption is a very strong one.

Econometrics

J. Eduardo  
Vera-Valdés  
eduardo@math.aau.

Introduction

Formalities

Geometrical  
Properties of OLS

The Frisch-  
Waugh-Lovell  
Theorem

Applications

Goodness of Fit

OLS is BLUE

26

Bias and Consistency  
Precision



Summing Up

# Bias



The exogeneity assumption imposes that all errors are not related to past and future values of the regressors.

For instance,  $\hat{\beta}$  is biased under the weaker assumption of **predetermined** regressors,  $E(U_t|X_t) = 0$ .



Econometrics

J. Eduardo  
Vera-Valdés

eduardo@math.aau.

Introduction

Formalities

Geometrical  
Properties of OLS

The Frisch-  
Waugh-Lovell  
Theorem

Applications

Goodness of Fit

OLS is BLUE

27

Bias and Consistency

Precision

Summing Up

40

Department of  
Mathematical Sciences  
Aalborg University  
Denmark

# Consistency



Another statistical property we may want from our estimators is consistency; that is, we want the estimator to asymptotically achieve the true value.

Thus, we are interested in whether

$$\text{plim}_{n \rightarrow \infty} \hat{\beta} = \beta$$

Recalling that we have assumed a correct specification, and the more general stochastic regressors assumptions, consistency requires that

$$\text{plim}_{n \rightarrow \infty} (X^T X)^{-1} X^T U = 0$$

Econometrics

J. Eduardo  
Vera-Valdés  
eduardo@math.aau.

Introduction

Formalities

Geometrical  
Properties of OLS

The Frisch-  
Waugh-Lovell  
Theorem

Applications

Goodness of Fit

OLS is BLUE

28

Bias and Consistency  
Precision

Summing Up

40

Department of  
Mathematical Sciences  
Aalborg University  
Denmark

Hence, consistency requires:

(a)  $\text{plim}_{n \rightarrow \infty} (X^T X)^{-1}$  exists, in the sense of having a finite limit, possibly after some standardization.

(b)  $\text{plim}_{n \rightarrow \infty} X^T U = 0$ , possibly after some standardization.

# Consistency



Given that we are interested in the limit as the sample size  $n$  increases, we would expect the standardization terms to depend on  $n$ .

Note that

$$(X^T X)^{-1} X^T U = \left( \frac{1}{n} X^T X \right)^{-1} \left( \frac{1}{n} X^T U \right).$$

So that we look for conditions so that  $\text{plim}_{n \rightarrow \infty} \left( \frac{1}{n} X^T X \right)^{-1} = S_{XX} < \infty$  and  $\text{plim}_{n \rightarrow \infty} \frac{1}{n} X^T U = 0$ .

Econometrics

J. Eduardo

Vera-Valdés

eduardo@math.aau.

Introduction

Formalities

Geometrical

Properties of OLS

The Frisch-

Waugh-Lovell

Theorem

Applications

Goodness of Fit

OLS is BLUE

30

Bias and Consistency

Precision

Summing Up

40

# Consistency



Using the law of large numbers, we can show that **predeterminedness**,  $E(U_t|X_t)$ , implies

$$\text{plim}_{n \rightarrow \infty} \frac{1}{n} X^T U = 0.$$

Formally, the proof requires the conditions for a law of large numbers to apply.

Thus, the errors cannot be too correlated or have unbounded variances.

Econometrics

J. Eduardo

Vera-Valdés

eduardo@math.aau.

Introduction

Formalities

Geometrical  
Properties of OLS

The Frisch-  
Waugh-Lovell  
Theorem

Applications

Goodness of Fit

OLS is BLUE

31

Bias and Consistency

Precision

Summing Up

40

Department of  
Mathematical Sciences  
Aalborg University  
Denmark



# Consistency



Another use of the law of large numbers can be used to show that

$$\text{plim}_{n \rightarrow \infty} \left( \frac{1}{n} X^T X \right)^{-1} = S_{XX} < \infty.$$

So that we are implicitly assuming that the regressors are not too correlated, and the variances do not increase without bound.

Typically, these conditions are satisfied by most type of regressors.

Yet, these are some regressors where this is not satisfied.

Econometrics

J. Eduardo

Vera-Valdés

eduardo@math.aau.

Introduction

Formalities

Geometrical

Properties of OLS

The Frisch-

Waugh-Lovell

Theorem

Applications

Goodness of Fit

OLS is BLUE



32

Bias and Consistency

Precision

Summing Up

# Consistency



Econometrics

J. Eduardo

Vera-Valdés

eduardo@math.aau.

Introduction

Formalities

Geometrical  
Properties of OLS

The Frisch-  
Waugh-Lovell  
Theorem

Applications

Goodness of Fit

OLS is BLUE

33

Bias and Consistency

Precision

Summing Up

40

Department of  
Mathematical Sciences  
Aalborg University  
Denmark

# Precision



Turning our attention now to analyse the second moment of the OLS estimator, we measure the precision of the OLS estimator by its covariance matrix.

In order to derive some properties of the covariance matrix of the OLS estimator, we must make assumptions regarding the second moments of the error term.

The usual assumptions about the errors' second moment of **homoskedasticity** and **no autocorrelation** can be written as

$$\text{Var}(U|X) = E(UU^T|X) = \sigma^2\mathbb{I},$$

where  $\mathbb{I}$  is a  $n \times n$  identity matrix.

Econometrics

J. Eduardo  
Vera-Valdés

eduardo@math.aau.

Introduction

Formalities

Geometrical

Properties of OLS

The Frisch-  
Waugh-Lovell  
Theorem

Applications

Goodness of Fit

OLS is BLUE

Bias and Consistency

Precision

Summing Up

34

# Precision



Under the assumptions on the error term, and **exogeneity** of the regressors, the variance of the OLS estimator is

$$\text{Var}(\hat{\beta}) = \sigma^2(X^T X)^{-1}.$$

Thus, the precision of the OLS estimator can be shown to depend on three things:

- ▶ The variance of the error term
- ▶ The sample size
- ▶ The relation between the regressors

Econometrics

J. Eduardo  
Vera-Valdés  
eduardo@math.aau.

Introduction

Formalities

Geometrical  
Properties of OLS

The Frisch-  
Waugh-Lovell  
Theorem

Applications

Goodness of Fit

OLS is BLUE

Bias and Consistency

Precision

Summing Up

35

40

The dependence of the variance of the OLS estimator on the variance of the **homoskedastic** error term is straightforward.

The dependence on the sample size can be seen if we write

$$\text{Var}(\hat{\beta}) = \sigma^2(X^T X)^{-1} = \left(\frac{1}{n}\sigma^2\right) \left(\frac{1}{n}X^T X\right)^{-1},$$

and assuming, as before, that  $\text{plim}_{n \rightarrow \infty} \left(\frac{1}{n}X^T X\right)^{-1} = S_{XX}$ .

Econometrics

J. Eduardo  
Vera-Valdés  
eduardo@math.aau.

Introduction

Formalities

Geometrical  
Properties of OLS

The Frisch-  
Waugh-Lovell  
Theorem

Applications

Goodness of Fit

OLS is BLUE

Bias and Consistency

Precision

Summing Up

36

To show the dependence on the relation between regressors consider

$$y = x_1\beta_1 + X_2\beta_2 + U,$$

where  $X = [x_1, X_2]$ , and  $x_1$  is a column vector.

From the FWL theorem,  $\hat{\beta}_1$  is the same as the one from

$$M_{X_2}y = M_{X_2}x_1\beta_1 + V.$$

Thus,

$$\text{Var}(\hat{\beta}_1) = \sigma^2 / (x_1^T M_{X_2} x_1),$$

which shows its dependence on the relation between the regressors.

Econometrics

J. Eduardo  
Vera-Valdés  
eduardo@math.aau.

Introduction

Formalities

Geometrical  
Properties of OLS

The Frisch-  
Waugh-Lovell  
Theorem

Applications

Goodness of Fit

OLS is BLUE

Bias and Consistency


Precision

Summing Up

37

40

# Precision



Econometrics

J. Eduardo  
Vera-Valdés  
eduardo@math.aau.

Introduction

Formalities

Geometrical  
Properties of OLS

The Frisch-  
Waugh-Lovell  
Theorem

Applications

Goodness of Fit

OLS is BLUE

Bias and Consistency

38

Precision

Summing Up

40

Department of  
Mathematical Sciences  
Aalborg University  
Denmark

# Outline



Introduction

Formalities

Geometrical Properties of OLS

The Frisch-Waugh-Lovell Theorem

Applications

Goodness of Fit

OLS is BLUE

Bias and Consistency

Precision

Summing Up

Econometrics

J. Eduardo

Vera-Valdés

eduardo@math.aau.

Introduction

Formalities

Geometrical

Properties of OLS

The Frisch-

Waugh-Lovell

Theorem

Applications

Goodness of Fit

OLS is BLUE

Bias and Consistency

Precision

39

Summing Up

40

Department of  
Mathematical Sciences  
Aalborg University  
Denmark



# Summing Up



- ▶ We have seen that OLS uses orthogonal projections to decompose the regressand in fitted values and residuals.
- ▶ The residuals are orthogonal to all the regressors.
- ▶ If the regression includes a constant, the residuals have mean zero.
- ▶ We study the FWL theorem and applications to demeaned series, seasonality, and trends.
- ▶ Showed numerical properties of the  $R^2$ .
- ▶ Under correct specification and exogenous regressors, OLS is an unbiased estimator.
- ▶ Under just predetermined regressors, OLS are consistent (with “well-behaved” regressors).

Econometrics

J. Eduardo  
Vera-Valdés  
eduardo@math.aau.

Introduction

Formalities

Geometrical  
Properties of OLS

The Frisch-  
Waugh-Lovell  
Theorem

Applications

Goodness of Fit

OLS is BLUE

Bias and Consistency

Precision

40 Summing Up