

# Generalized Method of Moments

## Econometrics

J. Eduardo Vera-Valdés  
eduardo@math.aau.dk

Department of Mathematical Sciences  
Aalborg University  
Denmark



**AALBORG UNIVERSITY**  
DENMARK

# Outline



Introduction

Example

GMM

Tests

Summing Up

Generalized  
Method of  
Moments

J. Eduardo  
Vera-Valdés  
eduardo@math.aau.

Introduction

Example

GMM

Tests

Summing Up

# Outline



Introduction

Example

GMM

Tests

Summing Up

Generalized  
Method of  
Moments

J. Eduardo  
Vera-Valdés  
eduardo@math.aau.

2 Introduction

Example

GMM

Tests

Summing Up

# Introduction



Generalized method of moments (GMM) provides a unifying framework for the analysis of many familiar estimators.

It includes OLS, IV, and GLS as special cases.

It offers a convenient method of estimation in certain models which were computationally very burdensome to estimate by traditional methods.

The estimation relies on population moments whose expectation is zero when evaluated at the true parameters.

GMM estimate is then obtained by minimizing a quadratic form in the analogous sample moments.

Generalized  
Method of  
Moments

J. Eduardo  
Vera-Valdés  
eduardo@math.aau.

3 Introduction

Example

GMM

Tests

Summing Up

# Outline



Introduction

Example

GMM

Tests

Summing Up

Generalized  
Method of  
Moments

J. Eduardo  
Vera-Valdés  
eduardo@math.aau.

Introduction

4 Example

GMM

Tests

Summing Up

# Example



Instrumental variables can be obtained as a GMM estimator.

We obtain a moment condition per each instrument.

If the number of instruments is the same as the number of regressors we recover IV.

For more instruments than regressors, we obtain GIV.

Generalized  
Method of  
Moments

J. Eduardo  
Vera-Valdés  
eduardo@math.aau.

Introduction

5 Example

GMM

Tests

Summing Up

# Outline



Introduction

Example

GMM

Tests

Summing Up

Generalized  
Method of  
Moments

J. Eduardo  
Vera-Valdés  
eduardo@math.aau.

Introduction

Example

6 GMM

Tests

Summing Up

17

Department of  
Mathematical Sciences  
Aalborg University  
Denmark

Suppose we have  $r$  moment conditions given by

$$E[f(x_t, \beta_0)] = 0,$$

where  $x$  is a stationary ergodic random vector,  $f$  is a  $r \times 1$  vector of (possibly nonlinear) continuous functions of  $\beta$  a  $q \times 1$  parameter vector, and  $\beta_0$  is the true value of  $\beta$ .

We further assume that a law of large numbers can be applied to  $f$  so that the sample mean converges a.s. to the population mean,

$$\lim_{n \rightarrow \infty} n^{-1} \sum_{t=1}^n f(x_t, \beta) = E[f(x_t, \beta)].$$

Generalized  
Method of  
Moments

J. Eduardo  
Vera-Valdés  
eduardo@math.aau.

Introduction

Example

7 GMM

Tests

Summing Up



The core principle of GMM is to estimate the parameters by using a quadratic form on the sample moments.

Let  $g_n(\beta) = n^{-1} \sum_{t=1}^T f(x_t, \beta)$ , and define the quadratic form

$$Q_n(\beta) = g_n'(\beta) W_n g_n(\beta),$$

where  $W_n$  is a positive definite symmetric matrix which converges in probability to a positive definite symmetric matrix  $W$ .

The GMM estimator for  $\beta$  is the value that minimizes  $Q_n(\beta)$ .

Generalized  
Method of  
Moments

J. Eduardo  
Vera-Valdés  
eduardo@math.aau.

Introduction

Example

8 GMM

Tests

Summing Up

The first order conditions are given by

$$G'_n(\hat{\beta})W_ng_n(\hat{\beta}) = 0,$$

where  $G_n(\hat{\beta})$  is the  $r \times q$  matrix given by  $[G_n(\hat{\beta})]_{ij} = \partial g_{ni}(\beta)/\partial \beta_j$ .

If the system is just identified,  $r = q$ , the matrices  $G_n(\hat{\beta})$  and  $W_n$  are nonsingular, so that the solution does not depend on  $W_n$ .

Unlike the linear case, the f.o.c. does not imply an analytical solution, so they must be solved numerically.

Although we cannot obtain an explicit solution, it can be shown that GMM is a consistent estimator.

From the population moment condition and the LLN, we have that  $g_n(\beta_0) \rightarrow 0$ .

$W_n$  positive definite implies that  $Q_n(\beta) \geq 0 \forall \beta$ . Moreover, we assume that the model is asymptotically identified; that is,  $E[f(x_t, \beta)] \neq 0 \forall \beta \neq \beta_0$ .

Thus, if  $\hat{\beta}$  minimizes  $Q_n()$ , then  $\hat{\beta} \rightarrow \beta_0$ .

Note that the estimator is consistent regardless of the  $W_n$  matrix used.

Generalized  
Method of  
Moments

J. Eduardo  
Vera-Valdés  
eduardo@math.aau.

Introduction

Example

10 GMM

Tests

Summing Up

To show that GMM is asymptotically normal, we use a first-order Taylor expansion approximation of  $g_n(\beta)$  around  $\beta_0$ ,

$$g_n(\hat{\beta}) = g_n(\beta_0) + G_n(\bar{\beta})(\hat{\beta} - \beta_0) + o_p(n^{-1}),$$

where  $\|\hat{\beta} - \bar{\beta}\| \leq \|\hat{\beta} - \beta_0\|$ .

Premultiplying both sides by  $G'_n(\hat{\beta})W_n$ , the left-hand side is zero by construction, so asymptotically,

$$n^{1/2}(\hat{\beta} - \beta_0) = -[G'_n(\beta_0)W_nG_n(\beta_0)]^{-1}G'_n(\beta_0)W_n n^{1/2}g_n(\beta_0).$$

Hence, under regularity conditions that guarantee that the CLT for  $g_n(\beta_0)$  holds,  $n^{1/2}(\hat{\beta} - \beta_0)$  follows a normal distribution.

From the previous derivation, note that the variance of  $n^{1/2}(\hat{\beta} - \beta_0)$  is given by

$$[G'_n(\beta_0)W_nG_n(\beta_0)]^{-1}G'_n(\beta_0)W_nS_WW_nG_n(\beta_0)[G'_n(\beta_0)W_nG_n(\beta_0)]^{-1},$$

where  $S_W = E[f(x_t, \beta_0)f'(x_t, \beta_0)]$  is the variance-covariance matrix for  $g_n(\beta_0)$ .

Note that it has a sandwich form, which suggests that it is an inefficient estimator.

We obtain the efficient estimator by taking  $W_n = S_w^{-1}$ .

Generalized  
Method of  
Moments

J. Eduardo  
Vera-Valdés  
eduardo@math.aau.

Introduction

Example

12 GMM

Tests

Summing Up

We do not know  $S_w$  in general.

Hence, we can use a multi-step estimation methodology:

- ▶ We use an initial (most likely inefficient)  $W_0$  matrix to estimate  $\hat{\beta}$  and construct an estimate for  $\hat{S}_w$ . A typical implementation will make  $W_0 = I$ .
- ▶ We use our estimate for  $\hat{S}_w$  in a second iteration of GMM.
- ▶ Keep iterating if necessary.

Generalized  
Method of  
Moments

J. Eduardo  
Vera-Valdés  
eduardo@math.aau.

Introduction

Example

13 GMM

Tests

Summing Up

# Outline



Introduction

Example

GMM

Tests

Summing Up

Generalized  
Method of  
Moments

J. Eduardo  
Vera-Valdés  
eduardo@math.aau.

Introduction

Example

GMM

14 Tests

Summing Up

17

Department of  
Mathematical Sciences  
Aalborg University  
Denmark

# Test for Overidentification



Generalized  
Method of  
Moments

J. Eduardo  
Vera-Valdés  
eduardo@math.aau.

Introduction

Example

GMM

15 Tests

Summing Up

Similar to the GIV case, we can/should test the model specification using the overidentifying restrictions.

The test-statistic is given by

$$\tau_n = n g_n'(\hat{\beta}) \hat{S}_w^{-1} g_n(\hat{\beta}),$$

which follows a chi-square distribution with  $r - q$  degrees of freedom.



# Outline



Introduction

Example

GMM

Tests

Summing Up

Generalized  
Method of  
Moments

J. Eduardo  
Vera-Valdés  
eduardo@math.aau.

Introduction

Example

GMM

Tests

16 Summing Up

17

Department of  
Mathematical Sciences  
Aalborg University  
Denmark

# Summing Up



- ▶ We have studied a method to obtain consistent estimates based on population moments.
- ▶ The method relies on minimizing a quadratic form in the sample moments.
- ▶ There must be at least as many moment conditions as parameters to estimate.
- ▶ GMM can estimate nonlinear models.
- ▶ Weak instruments can have perverse finite sample effects on the estimation.
- ▶ The test for overidentification allows us to check the validity of the moment conditions.

Generalized  
Method of  
Moments

J. Eduardo  
Vera-Valdés  
eduardo@math.aau.

Introduction

Example

GMM

Tests

17 Summing Up