

15. Error analysis for ODE IVP solvers.

Numerical Analysis E2021

Institute of Mathematics
Aalborg University



AALBORG UNIVERSITY
DENMARK

Errors in ODE problems

Numerical Analysis
E2021

Errors In ODE
Problems

Order Of Euler's
Method

1

When solving ODEs numerically, we get errors from two sources:

- ▶ *Discretisation error*: a property of the differential equation and the applied method.
 - ▶ *Local*: the error that would be made in one step if the previous values were exact and if there were no roundoff error.
 - ▶ *Global*: the difference between the computed solution, still ignoring roundoff, and the true solution.
- ▶ *Roundoff error*: a property of the computer hardware and the program used.

4

Error in ODE problems

Discretisation error

Numerical Analysis
E2021

Errors In ODE
Problems

Order Of Euler's
Method

2

Let u be a function given by

$$\begin{aligned}u'_n &= f(t, y_n), \\ u_n(t_n) &= y_n.\end{aligned}$$

Then the local resp. global discretisation errors are given by

$$d_n = y_{n+1} - u_n(t_{n+1}), \quad e_n = y_n - y(t_n).$$

We say that an ODE solver is of order p if $d_n = \mathcal{O}(h_n^{p+1})$.

4

Order of Euler's method

Numerical Analysis
E2021

Errors In ODE
Problems

Order Of Euler's
Method

3

Recall Euler's method

$$y_{n+1} = y_n + h_n f(t_n, y_n).$$

Assume the local solution $u_n(t)$ has a continuous second derivative. Then, using Taylor series near the point t_n ,

$$u_n(t) = u_n(t_n) + (t - t_n)u'_n(t_n) + \mathcal{O}((t - t_n)^2).$$

Now, by using the differential equation and the initial condition defining $u_n(t)$, we obtain

$$u_n(t_{n+1}) = y_n + h_n f(t_n, y_n) + \mathcal{O}(h_n^2).$$

Consequently, we have

$$d_n = y_{n+1} - u_n(t_{n+1}) = \mathcal{O}(h_n^2),$$

thus Euler's method is of order 1.

4

Order and global error

Numerical Analysis
E2021

Errors In ODE
Problems

Order Of Euler's
Method

4

Consider the global discretisation error at a fixed point t_f . As accuracy requirements increase, the step size h_n decreases, thus the total number of steps N required to reach t_f increases. Rough estimate:

$$N \approx (t_f - t_0)/h,$$

where h is the average step size. We roughly have that if the local error is $\mathcal{O}(h^{p+1})$, then the global error is $N \cdot \mathcal{O}(h^{p+1}) = \mathcal{O}(h^p)$.

This explains the use of $p + 1$ as the exponent in the definition of order.

MATLAB demo of exercise 7.7.