

8. Zero finding for real functions of one variable. Present the three methods, bisection, secant and IQI, and discuss advantages and disadvantages of the three methods.

Numerical Analysis E2021

Institute of Mathematics
Aalborg University



AALBORG UNIVERSITY
DENMARK

For function of one variable we are often interested in solving for x in the following equation

$$F(x) = y \quad (1)$$

which can be transformed into a root-finding problem by defining $f(x) := F(x) - y$, resulting in

$$f(x) = 0 \Leftrightarrow F(x) = y \quad (2)$$

Thus, we wish to have methods for determining zeroes.

Bisection

Numerical Analysis
E2021

Motivation

Bisection

Newton's Method


Secant Method

Inverse Quadratic
Interpolation

Summary

2

If we have an interval $[a, b]$ where the function changes sign, then bisection is a slow but sure method, where we in every iteration bisect the search interval.



In IEEE double-precision it is guaranteed to find two successive floating-point numbers in 52 steps.

Newton's Method

Numerical Analysis
E2021

Motivation

Bisection

Newton's Method

Secant Method

Inverse Quadratic
Interpolation


Summary

3

Newton's Method is based on the following iterative process

$$x_{n+1} := x_n - \frac{f(x_n)}{f'(x_n)} \quad (3)$$

the idea is we draw the tangent to the graph, and determine this tangents intersection with the x -axis to get our next point.



However, f must be smooth, f' should be nice to compute, and starting guess must be close to root.

However, we do have nice local convergence properties, **if** f' and f'' exist, and are continuous, and our initial point is close to our computed solution. Furthermore,

$$e_{n+1} = \mathcal{O}(e_n^2) \quad (4)$$

Secant Method

Numerical Analysis
E2021

Motivation

Bisection

Newton's Method

Secant Method

Inverse Quadratic
Interpolation

Summary

4

The secant method is Newton's method where we replace the derivative with an approximation using the secant. That is

$$x_{n+1} := x_n - \frac{f(x_n)}{s_n}, \quad \text{where } s_n := \frac{f(x_n) - f(x_{n-1})}{x_n - x_{n-1}} \quad (5)$$

The advantage over Newton's is of course then that we do not have to compute the derivative at every step. It also has similar convergence properties, and assuming continuity of f' and f'' , one can prove that

$$e_{n+1} = \mathcal{O}(e_n^\phi) \quad (6)$$

6

Inverse Quadratic Interpolation

Numerical Analysis
E2021

Motivation

Bisection

Newton's Method

Secant Method

Inverse Quadratic
Interpolation

Summary

5

IQI uses the three previous points in order to determine the next one. For the point

$$(a, f(a)), \quad (b, f(b)), \quad (c, f(c)) \quad (7)$$

we interpolate the three points into a quadratic function of y , $P(y)$, such that

$$a = P(f(a)), \quad b = P(f(b)), \quad c = P(f(c)) \quad (8)$$

our next iterative step will then be $x = P(0)$. However, we require that our three points $f(a), f(b), f(c)$ are all distinct, which we cant guarantee. Furthermore, it can be shown that

$$e_{n+1} = \mathcal{O}(e_n^{1.84}) \quad (9)$$

6

Summary

Numerical Analysis
E2021

Motivation

Bisection

Newton's Method

Secant Method

Inverse Quadratic
Interpolation

Summary

6

We have several methods for polynomial interpolation:

- ▶ Bisection
 - ▶ Always has a solution, but slow
 - ▶ Linear convergence
- ▶ Newton's Method
 - ▶ Quick convergence guaranteed under assumptions
 - ▶ Quadratic convergence
- ▶ Secant Method
 - ▶ No convergence-guarantee.
 - ▶ Doesn't require derivative, but may have issues if root is a critical point
 - ▶ Superlinear convergence
- ▶ IQI
 - ▶ Converges quickly when close to a root
 - ▶ Can fail entirely
 - ▶ Poor if initial point is far from solution

MATLAB demo of exercise 4.3.

6