

Test your knowledge : Class Quiz

Quiz . 20 Total Points

😊 Great job! Keep pushing for that perfect score! 🚀🌟

Your marks : 19 / 20

1. What is statistics?

- ☐ A branch of mathematics
- ☐ A collection of random numbers
- ☒ A discipline that concerns the collection, organization, analysis, interpretation, and presentation of data.
- ☐ A type of computer programming language

✓ Correct!

2. Why is statistics important in everyday life?

- ☐ It's not important in everyday life
- ☒ It helps make informed decisions based on data
- ☐ It's only used by scientists and researchers
- ☐ It's mainly used for entertainment purposes

✓ Correct!

3. What are descriptive statistics?

- ☐ A type of statistical test
- ☒ A way to summarize and present data
- ☐ A mathematical constant
- ☐ A type of survey

✓ Correct!


4. Define inferential statistics

- ☐ A type of descriptive statistic
- ☒ A method to make predictions based on data
- ☐ A way to organize data
- ☐ A type of data visualization

✓ Correct!

5. Explain the difference between a population and a sample

- ☐ A population is a type of sample
- ☒ A sample is a subset of a population



5. Explain the difference between a population and a sample

- ☐ A population is a type of sample
- ☒ A sample is a subset of a population
- ☐ They are the same thing
- ☐ A population is always larger than a sample

✓ Correct!

6. What is categorical data?

- ☒ Data that consists of categories or labels
- ☐ Data measured on a continuous scale
- ☐ Data with high variability
- ☐ Data collected from experiments

✓ Correct!

7. Provide an example of ordinal data

- ☐ Temperature in degrees Celsius

☐ A ... in ...



7. Provide an example of ordinal data

- ☐ Temperature in degrees Celsius
- ☐ Age in years
- ☒ Likert scale ratings (e.g., "strongly agree," "agree," "neutral," "disagree," "strongly disagree")
- ☐ The number of cars in a parking lot

✓ Correct!

8. What is the difference between discrete and continuous data?


- ☐ Discrete data can take on any value, while continuous data can only take on whole numbers
- ☐ Discrete data is measured with intervals, while continuous data is counted
- ☒ Discrete data consists of distinct values, while continuous data can take on an infinite number of values within a range.
- ☐ Discrete data is always qualitative, while continuous data is always quantitative.

✓ Correct!

9. What is the measure of central tendency?

- ☐ A way to measure how spread out data points are
- ☐ A way to identify outliers in a dataset





9. What is the measure of central tendency?

- ☐ A way to measure how spread out data points are
- ☐ A way to identify outliers in a dataset
- ☒ A way to summarize the center or typical value of a dataset
- ☐ A method to calculate the range of data

✓ Correct!

10. Which of the following is a measure of central tendency?

- ☐ Variance
- ☐ Interquartile range
- ☒ Mode
- ☐ Standard deviation

✓ Correct!

11. If you have a dataset with extreme outliers, which measure of central tendency is least affected?

- ☐ Mean
- ☒ Median



11. If you have a dataset with extreme outliers, which measure of central tendency is least affected?

- ☐ Mean
- ☒ Median
- ☐ Mode
- ☐ Range

✓ Correct!

12. What is the mean of the numbers 3, 5, 7, and 9?

- ☐ 3
- ☐ 5
- ☐ 7
- ☒ 6

✓ Correct!

13. What does the range of a dataset measure?

- ☐ The average value of the dataset
- ☒ The spread or variability of the dataset



13. What does the range of a dataset measure?

- ☐ The average value of the dataset
- ☒ The spread or variability of the dataset
- ☐ The number of data points in the dataset
- ☐ The center of the dataset

✓ Correct!

14. Which of the following measures the average deviation from the mean?


- ☒ Variance
- ☐ Range
- ☐ Median
- ☐ Mode

✓ Correct!

15. The standard deviation is a measure of:

- ☐ The average value in a dataset
- ☐ The range of a dataset





15. The standard deviation is a measure of:

- ☐ The average value in a dataset
- ☐ The range of a dataset
- ☒ The spread or dispersion of a dataset
- ☐ The center of a dataset

✓ Correct!

16. What is the interquartile range (IQR)?


- ☐ The difference between the maximum and minimum values in a dataset
- ☐ The range of values within one standard deviation of the mean
- ☒ The range between the first and third quartiles of a dataset
- ☐ The square root of the variance

✓ Correct!

17. If the variance of a dataset is zero, what can you conclude about the data?

- ☐ The data is highly variable
- ☐ The data is normally distributed





17. If the variance of a dataset is zero, what can you conclude about the data?

- ☐ The data is highly variable
- ☐ The data is normally distributed
- ☒ All the data points are identical
- ☐ The data is negatively skewed

✓ Correct!

18. What is an outlier in a dataset?

- ☒ A data point that is extremely high or low compared to other data points
- ☐ The mean value of the dataset
- ☐ The most frequently occurring value in the dataset
- ☐ A data point that is exactly in the center of the dataset

✓ Correct!

19. Which of the following is not a common method for detecting outliers?

- ☐ Z-score
- ☐ Box plot



19. Which of the following is not a common method for detecting outliers?

- ☐ Z-score
- ☐ Box plot
- ☒ Mode
- ☐ Tukey's method

✓ Correct!

20. Which measure of dispersion is most affected by outliers?

- ☒ Range
- ☐ Variance
- ☐ Standard deviation
- ☐ Interquartile range

✗ Incorrect!

Variance is the correct answer

Submit

