Numerical verification of the refined Mazur–Tate conjecture in ranks 2 and 3, with an elementary formula for the Mazur–Tate pairing

Oct 21, 2026 · @Rauf Alencar

Abstract

Let E/Q be an elliptic curve of Mordell–Weil rank r and, for a prime m of good reduction, let θ_m ∈ Z[(Z/m)^x/±1] be the Mazur–Tate theta element built from plus modular symbols. The refined conjecture of Mazur and Tate (1987), in the formulation of Burns–Kurihara–Sano and Ota, predicts that θ_m lies in the r-th power of the augmentation ideal I and prescribes its image in I^r/I^{r+1} as a Tamagawa-type factor times the discriminant of the Mazur–Tate height pairing. We report a numerical verification of this leading-term prediction, with both sides computed independently, for nine elliptic curves of rank 2 (389a1, 433a1, 446d1, 571b1, 655a1, 709a1, 916c1, 1088j1, 1132a1) and one of rank 3 (5077a1), in about 700 non-trivial checks at prime levels m, with no failures; the curves were chosen to separate the roles of the Tamagawa numbers, the real components and the torsion, and an independent re-implementation with the constant predicted in advance confirmed the identity on seven further curves (237 non-trivial levels), and once the normalizations (period, sign of the modular symbol, orientation of the pairing, torsion) are made explicit the identity holds exactly in Ota's form, including his sign (−1)^{ν(m)}, assuming the standard normalization of the Artin map (uniformizer ↦ arithmetic Frobenius) used by Tan, which we have not checked in Ota's text; an independent re-implementation with the constant predicted in advance also confirms the identity in rank 1 on eight curves (533 levels). At composite level $S = m_1m_2$ we verify Ota's inclusion–exclusion form of the conjecture for 389a1, including the mixed coefficient and the sign (−1)^{ν(S)}. The pairing side is computed by an elementary recipe: for R = c_Tam·#E(F_m)·P in the formal group at m, the diagonal value ⟨R,R⟩_m is the class of t(R)/d(R) in $F_m^x/\pm1$, where t = −x/y and d^2 is the denominator of x(R). We derive this recipe from the biextension definition of the pairing, assuming the Néron-model description of the canonical local splittings (as in Tan 1995; we have not had access to Mazur–Tate 1983), obtaining the closed formula ⟨R,S⟩ = [γ/A(R+S)] with γ the constant term of the primitive line through R and S and A the numerator of the x-coordinate; in particular $\langle R,R\rangle = [A(R)]^{-1}$. One elementary integrality step is left unproved at $\ell = 2$ (verified at 349 points on six curves and at 3631 further points on 453 curves). We know of no earlier numerical verification of the refined conjecture in rank 3, nor of one in rank 2 with a closed formula for the pairing in terms of the coordinates of the points themselves; Bley–Macías Castillo (2022) verified the equivalent 3-adic eTNC statement for 389a1, 433a1 and 446d1 at a handful of primes via Selmer groups and Hilbert symbols. As a secondary result we verify, in 20 cases over three curves and p ∈ {3,5}, that the Kolyvagin derived class κ_ℓ attached to a rank-2 curve and a Heegner field spans exactly the line $\ker(E(Q)/p \to E(F_\ell)/p)$, with an explicit nonvanishing criterion — a statement implicit in Howard and Mazur–Rubin and stated in an unpublished draft hosted on W. Stein's website; the nonvanishing part was tested in two cases by Jetchev–Lauter–Stein (2009), with which we agree, and the direction had not been tested. All computations are in PARI/GP 2.17 and are reproducible from the accompanying code.

1. Setup and notation

Throughout, E/Q is an elliptic curve of conductor N given by a minimal Weierstrass model, with Mordell–Weil group E(Q) of rank r and a fixed basis $P_1, \dots, P_r$ of E(Q)/E(Q)tors (saturated). We write $c_{Tam} = \prod_{\ell|N} c_\ell$ for the product of Tamagawa numbers and $c_\infty \in \{1, 2\}$ for the number of connected components of E(R).

Modular symbols. Let x^+ be the plus modular symbol of E, normalized as in PARI's `msfromell1`: for a rational cusp a/m, $[a/m]^+ := x^+(\{a/m\} - \{\infty\}) \in Q$ is the real part of $2\pi i\int$ from a/m to i∞ of f(z)dz, divided by ω_1, the least positive real period of the Néron lattice (the real period over the connected component of E(R)). For the curves below, $[a/m]^+ \in Z$ when $c_\infty = 2$ and $[a/m]^+ \in \tfrac{1}{2}Z$ when $c_\infty = 1$.

Theta elements. For a prime m ∤ N let $G_m = (Z/m)^x$ and $G_m^+ = G_m/\pm1$, cyclic of order n := (m−1)/2. Since $[-a/m]^+ = [a/m]^+$, the Mazur–Tate element

$$\theta_m \;=\; \sum_{a\in(\mathbb{Z}/m)^\times} [a/m]^+\,\sigma_a \;\in\; \mathbb{Q}[G_m^+]$$

is well defined. Fixing a primitive root g mod m and writing σ for the image of σ_g, we identify $Z[G_m^+] = Z[\sigma]/(\sigma^n - 1)$ and set x = σ − 1, so the augmentation ideal is I = (x). We expand $\theta_m = \Sigma_k a_k x^k$ with $a_k = \Sigma_t c_t C(t, k)$, where c_t is the coefficient of σ^t. The relation $(1+x)^n = 1$ gives $n x \equiv -C(n,2) x^2 - C(n,3) x^3 - \dots \pmod{I^n}$, so for $k \geq 1$ one has $I^k/I^{k+1} \cong Z/n$, and the class of θ_m in I^r/I^{r+1}, assuming $\theta_m \in I^r$, is obtained from $(a_0, \dots, a_r)$ by r successive reductions: if $a_j = n\cdot k_j$, replace a_{j-1+i} by $a_{j-1+i} - k_j C(n,i)$ for $i \geq 2$. We write $c_r(m) \in Z/n$ for the resulting class. The condition "$\theta_m \in I^r$" is exactly that $a_0 \equiv 0$ and that each intermediate reduced coefficient is $\equiv 0 \pmod n$; that $\theta_m \in I^r$ (the "order of vanishing" half of the conjecture) is a theorem of Ota (2018, Thm. 1.2) for the Z_p-parts with $p \geq 5$ (2 and 3 inverted), E without complex multiplication and further hypotheses; our verification is over Z, including the 2- and 3-parts; we verify it in every case below.

The refined conjecture at prime level. In the form of Burns–Kurihara–Sano (2021, eqs. (8)–(9)) and Ota (2018, Conj. 2.4) for S = {m}:

$$\theta_m \;\equiv\; (-1)^{\nu(m)}\,\#Ш(E/\mathbb{Q})\,J_m\,R_F \pmod{I^{r+1}}, \qquad J_m = \#\operatorname{coker}\Big(E(\mathbb{Q}) \to E(\mathbb{F}_m) \oplus \bigoplus_{\ell|N} E(\mathbb{Q}_\ell)/E_0(\mathbb{Q}_\ell)\Big)$$

where $R_F = \det(\langle x_i, y_j\rangle)$ is the discriminant of the Mazur–Tate pairing $\langle\cdot,\cdot\rangle$: E(Q) × E_S(Q) → $G_m^+ \cong I/I^2$, with x_i a basis of E(Q) and y_j a basis of the "S-Selmer" sublattice E_S(Q) = kernel of the map defining J_m. For the curves considered, #Ш = 1 (analytically) and E(Q)_tors = 0, except 1088j1 with $E(Q)_{tors} \cong Z/2$.

2. An elementary formula for the Mazur–Tate pairing at a prime level

The Mazur–Tate pairing is defined idèlically via a biextension (Mazur–Tate 1983, 1987; see also Tan 1995, Def. 2.6–2.8). Its definition does not yield a closed formula; earlier numerical work computed it by other means (Portillo-Bobadilla 2019 in rank 1; Séguin 2014 in Darmon's Heegner-point setting; Bley–Macías Castillo 2022 via Selmer groups). We use the following recipe, suggested by the analogy with the Mazur–Stein–Tate formula h_p(P) = (1/p)·log_p(σ_p(P)/d(P)) for cyclotomic p-adic heights:

Recipe. Let R ∈ E(Q) lie in the formal group E_1(Q_m), write x(R) = a/d^2 in lowest terms with d > 0, and let t(R) = −x(R)/y(R) be the formal parameter. Then v_m(t) = v_m(d) ≥ 1, the ratio t(R)/d(R) is an m-adic unit, and we set

$$h(R) \;:=\; \log_g\Big[\,t(R)/d(R) \bmod m\,\Big] \;\in\; \mathbb{Z}/n, \qquad \langle R,R\rangle_m \;:=\; 2\,h(R) \;=\; \log_g\big[(t/d)^2\big] \;=\; -\log_g\big[A(R)\big],$$

where x(R) = A(R)/d^2 and the last equality uses $B^2 \equiv A^3$ (mod m) for y(R) = B/d^3, since t/d = −A/B exactly. We call h the *half-pairing*: it is a quadratic form on E_S(Q) and the pairing is its polarization, ⟨R_1,R_2⟩ := h(R_1+R_2) − h(R_1) − h(R_2), which gives back 2h(R) on the diagonal; no division by 2 is ever needed. The pairing matrix of a basis R_1, …, R_r of a sublattice is H := (⟨R_i,R_j⟩), with H_ii = 2h(R_i). The m-adic sigma function does not appear because σ_m(t) = t·(1 + O(t)) and t ≡ 0 (mod m), so σ_m(R)/d(R) ≡ t(R)/d(R) (mod m). The justification of these formulas is in the paragraph "The local computation" below.

Points in the formal group. For an arbitrary basis P_i of E(Q) we take R_i = M·P_i with M = c_Tam·#E(F_m). Multiplication by #E(F_m) kills the reduction mod m, and multiplication by c_Tam puts the point into E_0(Q_ℓ) at every bad ℓ, so R_i lies in the lattice E_S(Q) of §1 (this is the only place where the Tamagawa numbers enter, and §4 shows that omitting c_Tam breaks the identity for 446d1, the one curve on which the omission was tested). The coordinates of R_i are large rationals (roughly M^2·ĥ(P_i)/log 10 digits) but PARI handles them exactly for m up to a few thousand.

Status of the recipe: a consequence of Tan's definition, up to one local computation. We have not had access to Mazur–Tate (1983, 1987), but Tan (1995, §2) gives the definition in a form that can be compared with the recipe. Tan's pairing E(K) × E_S(K) → G is the composite of a pairing with values in C_S = ∏ C_v (Def. 2.6) with the Artin map C_S → G (Def. 2.8), where the local groups are: at a tamely ramified place v ∈ S, A_v = E(K_v), B_v = E_1'(K_v) (points with trivial reduction) and C_v = K_v^x/(1 + π_v O_v) ≅ Z × k_v^x; at an unramified non-archimedean v, B_v = E_0(K_v) and C_v = K_v^x/O_v^x ≅ Z, the pairing being the Néron local symbol; at a real place unramified in L/K, C_v = 1. For K = Q, S = {m}, L = Q(μ_m)$^+$ this says three things. (i) The domain of the second argument is exactly the lattice E_S(Q) = {R : R ∈ E_1(Q_m), R ∈ E_0(Q_ℓ) for ℓ | N} that the recipe requires — the hypotheses of the recipe are the hypotheses of the definition. (ii) The Artin map kills the uniformizer m at the place m and the subgroup 1 + mZ_m, so only the unit part of the local value survives, as a class in F_m^x/±1 = G_m^+; and at ℓ ≠ m it sends ℓ^k to Frob_l^k = [ℓ^k]. Hence ⟨R,R⟩ = Art(ψ_m(p)) · ∏_l [ℓ]^ψ_l(p) for any point p of the biextension over (R,R), the product being independent of p because the global units are killed (φ(Q^x) = 0). (iii) The splitting ψ_m is the unique one on the formal group (Hom(E_1, Hom(E_1, C_m)) = 0 since E_1 is pro-m and C_m has no m-torsion), so it coincides with the reduction mod (1 + mZ_m) of the Mazur–Tate m-adic σ-function, which is t·(1 + O(t)); and the Néron symbols at ℓ | d sum to ord_l of the denominator. The recipe t/d is precisely this product, read as the quadratic form whose polarization is the pairing; t/d is the half-pairing h, whose polarization is the pairing; the local computation below shows that the pairing itself is [(t/d)^2] on the diagonal. What remains to make this a proof is the local computation at m and ℓ for one explicit representative p over (R,R), with the Q^x-ambiguity removed; we have not carried it out. The orientation of the Artin map on local units (u ↦ σ(u) or σ(u^{-1})) decides whether Tan's pairing is ours or its inverse; we fix the orientation that makes the formulas below hold, and §4(b) shows from the data that Tan's is the inverse.

The local computation. We then carried out the computation that the previous paragraph left open; it is short. Realize the Poincaré biextension of E × E by G_m in the usual way: a point of P over (a, b) is a non-zero rational function f on E with divisor (a+b) + (O) − (a) − (b), and the G_m-action is scaling. For a = R, b = S in E_S(Q) take

f = (x − x(R+S)) / ℓ(x, y), ℓ = the line through R and S (the tangent line if R = S),

which has the right divisor, and scale f by a rational number so that both the numerator d(R+S)^{2}x − A(R+S) and the denominator ℓ_0 = αy + βx + γ (α, β, γ ∈ Z, gcd = 1) are primitive integral forms. Here x(R+S) = A(R+S)/d(R+S)^2 in lowest terms. The canonical local splittings of Mazur–Tate, in the description through the Néron model, are: at an unramified place ℓ (B_ℓ = E_0(Q_ℓ), C_ℓ = Z) the order of f along the identity component of the special fibre — which is 0 for a primitive f — and at the tamely ramified place m (B_m = E_1(Q_m), C_m = Q_m^x/(1+mZ_m)) the m-adic order together with the reduction of f to the special fibre: since R, S, R+S all reduce to O, the reduction $\bar{f}$ has trivial divisor and is a non-zero constant in F_m^x, and this constant is ψ_m(f). The reduction is immediate from the primitive forms: d(R+S) ≡ 0, so the numerator reduces to −A(R+S); and since R, S ∈ E_1(Q_m) the coefficients α, β of the primitive line are divisible by m (the line through two points of E_1 has slope and intercept with m-adic poles of order ≥ 1 and ≥ 3, so after clearing denominators the constant term is the unit), and the denominator reduces to the constant γ. Hence ψ_m(f) = −A(R+S)/γ ∈ F_m^x, all other components are trivial, and by Artin reciprocity

⟨R, S⟩ = [γ / A(R+S)] ∈ F_m^x/±1 = G_m^+,

where we fix the orientation of the Artin map on local units so that this holds (the other orientation inverts every value). This is a closed formula for the Mazur–Tate pairing, valid for all R, S ∈ E_S(Q) with R, S, R+S ≠ O. It remains to compare it with the recipe. For R = S, writing x(R) = A/d^2, y(R) = B/d^3, the tangent-line constant is γ = (BW − ΛA)/c_L with W = 2B + a_1Ad + a_3d^3, Λ = 3A^2 + 2a_2Ad^2 + a_4d^4 − a_1Bd and c_L the content of the line; modulo m (d ≡ 0, and $B^2 \equiv A^3$ from the Weierstrass equation) this is γ ≡ −A^3/c_L, while A(2R) ≡ A^4/δ^2, where δ^2 := N_4/A(2R) is the cancellation in the duplication formula x(2R) = N_4/(d^2W^2), N_4 := A^4 − $b_4A^2d^4$ − 2b_6Ad^6 − b_8d^8 (δ is an integer). So ⟨R,R⟩ = [−δ^2/(c_L·A)], and the recipe gives ⟨R,R⟩ = 2h(R) = [(t/d)^2] = [A/B]^2 = [A]$^{-1}$ (because t/d = −A/B exactly and $B^2 \equiv A^3$). The two agree if and only if $\delta^2 \equiv \pm$c_L (mod m). The content c_L is always 1: since BW − ΛA ≡ −A^3 (mod d) is coprime to d, c_L = gcd(W, Λ), and a prime ℓ divides both W = d^3·∂F/∂y(R) and Λ = −d^4·∂F/∂x(R) (F the Weierstrass

equation) only if $\bar{R}$ is a singular point of the reduction, which $R \in E_0(Q_l)$ excludes. For δ at an odd prime ℓ (good or bad) with $R \in E_0(Q_l)$: $\ell \mid W$ means $\bar{R}$ is a non-singular 2-torsion point of the reduction, where $2y + a_1x + a_3$ is a uniformizer, so a point at distance j from the 2-torsion has double at distance exactly j from O (the formal group law is $[2]s = 2s + \dots$ with 2 a unit), which forces $v_l(N_4) = 0$, i.e. $\ell \nmid \delta$. At $\ell = 2$ the same argument does not apply: it uses that doubling preserves the distance to O, i.e. $[2]s = 2s + \dots$ with 2 a unit, and 2 is not a unit in the formal group at $\ell = 2$; we have not written out another argument. Numerically (chkg2.gp), $\delta = c_L = 1$ and $\gamma/A(2R) \equiv \pm(t/d)^2$ at all 349 points $R_i = M{\cdot}P_i$ with $7 \le m \le 113$ on six curves (389a1, 446d1, 1088j1, 916c1, 655a1, 5077a1), and in a separate sweep (gsweep.py) $\delta = c_L = 1$ at all 3631 points $c_{Tam}{\cdot}k{\cdot}P$, $1 \le k \le 8$, P running over the generators, on the 453 curves of positive rank and conductor ≤ 400, which include every reduction type at 2. For points outside $E_0(Q_2)$ on 446d1 and 1088j1 one finds $\delta = c_L = 2$ and 8, so the hypothesis $R \in E_0$ is used. For $R \ne S$ the same computation with the chord gives $\langle R,S\rangle = [\gamma_{R,S}/A(R+S)]$, and at 129 pairs (R_1, R_2) over five curves this equals $[(t/d)(R+S)/((t/d)(R)(t/d)(S))]$, i.e. exactly the polarization $h(R+S) - h(R) - h(S)$ of the recipe — with no 2-torsion ambiguity, which the polarization of a quadratic form alone would leave when n is even. Scripts: `biext.gp`, `chkg2.gp`, `chord.gp`, `gsweep.py`.

In summary: granting the Néron-model description of the canonical local splittings (which is how Mazur–Tate 1983 construct them and how Tan 1995 uses them), the Mazur–Tate pairing on $E_S(Q) \times E_S(Q)$ at prime level is given exactly by the recipe of this section, the only unproved step being the elementary statement $\delta = 1$ at $\ell = 2$ for points of $E_0(Q_2)$, verified in all cases at hand. The diagonal takes the memorable form $\langle R,R\rangle = [A(R)]^{\mp 1}$: the Mazur–Tate self-pairing of a point of $E_S(Q)$ is the class of the numerator of its x-coordinate.

3. The identity tested

Let $I_m := [E(Q) : E_S(Q)]$ be the size of the image of $E(Q)$ in $E(F_m) \oplus \bigoplus_{\ell \mid N} E(Q_\ell)/E_0(Q_\ell)$, so that $J_m = M/I_m$ with $M = c_{Tam}{\cdot}\#E(F_m)$ as above. Since the pairing is bilinear with $E_S(Q)$ in one argument, $\langle x_i, y_j\rangle = M^{-1}\langle M x_i, y_j\rangle$, and since $M{\cdot}E(Q) \subset E_S(Q)$ has index M^r/I_m, a change of basis gives

$$J_m\, R_F \;=\; \frac{M}{I_m}\cdot\frac{I_m}{M^{2r}}\cdot\det\big(\langle R_i,R_j\rangle_m\big) \;=\; \frac{\det\big(\langle R_i,R_j\rangle_m\big)}{M^{2r-1}}, \qquad R_i = MP_i.$$

The index I_m therefore cancels, and the prediction of §1 becomes, for $Ш = 1$ and up to a constant u_E depending only on normalizations,

$$\boxed{M^{2r-1}\,c_r(m) \;\equiv\; u_E\cdot\det_{r\times r}\big(\langle R_i,R_j\rangle_m\big) \pmod n, \qquad n=(m-1)/2}$$

In practice we test $2^r\, M^{2r-1}\, c_r(m) \equiv u_E{\cdot}\det H \pmod n$ with $H = (\langle R_i,R_j\rangle)$ the pairing matrix of §2; the factor 2^r on the left is a historical artefact of an earlier normalization and is simply absorbed into u_E (so $u_E = 2^r \times$ the constant of the boxed identity). Both sides depend on the primitive root g: replacing g by g^k multiplies c_r by k^{-r} and $\det H$ by k^{-r}, so the identity is independent of g. The constant u_E must be the same for all m; we determine it from the first few primes and then test it on all others. The test is non-trivial at a prime m when (lhs, rhs) $\ne (0, 0)$; when $\gcd(M, n) > 1$ the identity constrains $c_r(m)$ only modulo $n/\gcd$, and we flag these as weaker checks.

A first version of this computation used the factor N_m/I_m^2 ($N_m := \#E(F_m)$) instead of M^{-3} for $r = 2$; the two agree modulo 3 whenever $3 \nmid N_m$ (because $N_m^4 \equiv 1 \bmod 3$) but not modulo n, and the mod-3 test passed without revealing the error. We mention this because it illustrates why the full-level test is the one that carries information.

4. Results

All ten curves satisfy $\theta_m \in I^r$ at every prime level tested (the successive reductions of §1 never failed), and the boxed identity of §3 holds at every level with a constant u_E depending only on the curve.

Curve (Cremona)	r	N	c_{Tam}	c_∞	Basis P_i	u_E	prime levels m	non-trivial checks	of which gcd(M,n) > 1	failures
389a1	2	389	1	2	(−1,1), (0,0)	8	$5 \le m \le 1500$	221	106	**0**
709a1	2	709	1	2	saturated `ellrank` basis	8	$m \le 700$	111	58	**0**
446d1	2	$446 = 2{\cdot}223$	2 ($c_2 = 2$)	2	(2,0), (1,0)	8	$23 \le m \le 400$	61	—	**0**
571b1	2	571	1	1	(1,−1), (0,−2)	4	$23 \le m \le 330$	54	—	**0**
5077a1	3	5077	1	2	(1,−1), (2,−1), (0,−3)	−16	$23 \le m \le 330$	49	—	**0**
655a1	2	$655 = 5{\cdot}131$	2	1	(1,2), (3,2)	4	$23 \le m \le 330$	51	—	**0**
1132a1	2	$1132 = 2^2{\cdot}283$	3	1	(1,1), (−1,−3)	4	$23 \le m \le 330$	45	—	**0**
916c1	2	$916 = 2^2{\cdot}229$	3	2	(0,1), (2,1)	8	$23 \le m \le 330$	46	—	**0**
1088j1 (torsion Z/2)	2	$1088 = 2^6{\cdot}17$	4	2	(−1,8), (7,16)	2	$23 \le m \le 300$	44	—	**0**
433a1	2	433	1	1	(0,1), (−1,0)	4	$7 \le m \le 113$	21	—	**0**

"Non-trivial" excludes levels where both sides vanish (15 of the 236 levels for 389a1, for instance). Levels dividing N are excluded. The curves 655a1, 1132a1, 916c1 and 1088j1 were chosen, after a first draft of this note, to separate the three invariants that could enter the constant: $c_\infty = 1$ with $c_{Tam} = 2$ and 3 (655a1, 1132a1), $c_\infty = 2$ with odd c_{Tam} (916c1), and non-trivial torsion (1088j1). The result is that u_E depends on c_∞ and on the torsion, and not at all on the Tamagawa numbers (which are fully absorbed by the factor c_{Tam} in M):

$u_E = (-2)^r \cdot c_\infty$ / $\#E(Q)_{tors}^2$ in all ten cases (8, 8, 8, 4, −16, 4, 4, 8, 2, 4).

Every factor, including the sign, is accounted for. (a) 2^r is the factor we put on the left-hand side of the test (§3) and cancels; in terms of the pairing matrix H the identity is M^{2r−1} c_r(m) ≡ $(-1)^r \cdot (c_\infty / \#E(\mathbb{Q})\mathrm{tors}^2) \cdot \det H$. (b) The sign $(-1)^r$ — + for the rank-2 curves, − for the rank-3 curve, and − for rank 1 (eight curves, 533 levels, in the independent re-implementation of §4 below) — is the product of two conventions. First, the pairing: Tan's Artin map (his §1, Weil group W_D = $K^x\backslash A^x/U_D$) sends a uniformizer π_v to the Frobenius [v]; by global reciprocity a local unit u at m then corresponds to σ_u^{-1} (choose a ∈ Z with a ≡ u mod m; the principal idèle a gives $\prod_{\ell|a}$ Frob_ℓ^{v_ℓ(a)} = σ_a, so [u at m] = σ_a^{-1}), whereas our recipe uses u ↦ σ_u. So Tan's pairing is the inverse of ours, P := −H, i.e. ⟨R,R⟩ = [A(R)] rather than $[A(R)]^{-1}$, and $\det P = (-1)^r \det H$, which removes the $(-1)^r$. Second, the theta element: PARI's plus symbol satisfies $x^+(\{0\} - \{\infty\}) = -L(E,1)/\omega_1$ (checked on seven rank-0 curves of both discriminant signs), whereas Mazur–Tate normalize $[0]^+ = +L(E,1)/\Omega^+$; so θ_m of §1 is $-c_\infty$ times the Mazur–Tate element θ_m^+, and this global −1 is exactly Ota's $(-1)^{\nu(m)} = -1$ at prime level — provided Ota's Artin map is normalized as Tan's (uniformizer ↦ arithmetic Frobenius), which is the standard choice but which we have not checked in his text; this is question 3 of our letter to him. An earlier draft of this paragraph said the discrepancy could not be removed by an orientation change; that was wrong because it took the sign of θ_m for granted, and the rank-1 data (which force a factor $(-1)^{r+1} = -(-1)^r$, not $(-1)^{r(r-1)/2}$) exposed it. The composite-level test below confirms the relative sign $(-1)^{\nu(S)-\nu(m)}$ as well. (c) The symbols x^+ of PARI are normalized by the least real period ω_1 of the Néron lattice, which is the full real period when $c_\infty = 1$ but only half of it when $c_\infty = 2$; and θ_m as defined in §1 is $2\cdot\Sigma\,[a/m]^+\,\sigma_a$ over a ∈ G_m^+, because both a and −a are summed. If instead the symbols are normalized by Ω^+_int := the generator of the real projection of the lattice — which is ω_1 when $c_\infty = 2$ and $\omega_1/2$ when $c_\infty = 1$, and is the normalization that makes $[a/m]^+$ an integer in all cases — and $\theta_m^+ := -\Sigma\,[a/m]^+\,\sigma_a$ over a ∈ G_m^+ (the sign from (b)), then $\theta_m = -c_\infty \cdot \theta_m^+$ and the factor c_∞ disappears. (d) The factor $1/\#E(\mathbb{Q})\mathrm{tors}^2$ is the torsion normalization of the discriminant in Tan's Def. 2.13 ($\mathrm{disc}_S = |A_{\mathrm{tors}}|^{-1}|B_{\mathrm{tors}}|^{-1}$ det) combined with the torsion contribution to the index $[E(\mathbb{Q}) : E_S(\mathbb{Q})]$ in J_m; it is the same tors^2 as in the Birch–Swinnerton-Dyer formula and in Burns–Kurihara–Sano's eq. (9). With these normalizations the identity reads, for all curves and ranks 0–3 (rank 0: eight curves and 249 levels in the re-implementation, including #Ш = 4 and 9; ranks 1–3: this note and the re-implementation),

$$\theta_m^+ \equiv (-1)^{\nu(m)} \cdot \#Ш \cdot J_m \cdot \mathrm{disc}_m(P) = - \#Ш \cdot \det P \,/\, (M^{2r-1} \cdot \#E(\mathbb{Q})\mathrm{tors}^2) \pmod{I^{r+1}},$$

with P = (⟨R_i,R_j⟩) in Tan's orientation, ⟨R,R⟩ = [A(R)] (and #Ш = 1 for every curve of positive rank here),

which is Ota's Conj. 2.4 at prime level with no undetermined constant. The value u_E = 4 for 571b1 supersedes the value 2 quoted in an earlier draft, which we could not reproduce.

The Tamagawa test (446d1). With R_i = #E(F_m)·P_i, i.e. omitting c_Tam = 2, the identity fails at 28 levels of a run over 5 ≤ m ≤ 400. With R_i = 2·#E(F_m)·P_i, so that R_i ∈ $E_0(\mathbb{Q}_2)$ as the definition of E_S(Q) requires, it holds at 61 of 61 non-trivial levels with the same u = 8 as for the other $c_\infty = 2$ curves. This is the definition of J_m through the joint cokernel in Burns–Kurihara–Sano being confirmed in detail.

The vanishing levels. For 389a1 at m ≡ 1 (mod 9) the mod-3 reduction of the identity was examined separately: the levels at which ord_I θ_m jumps from 2 to 4 (m = 37, 163, 613, 757, 829, 937 among m ≤ 1200) are exactly the levels at which $\det\langle R_i,R_j\rangle \equiv 0 \pmod 3$. The frequency of such jumps (6 of the 20 levels m ≡ 1 (mod 9) in the mod-3 run, m ≤ 1200) matches the probability that a random symmetric 2×2 matrix over F_3 is singular (9/27). Jumps are always by an even amount (ord ∈ {2, 4, 6}), consistent with the alternating character of the second derived pairing (Bertolini–Darmon 1994, Thm. 3.18). At levels with 3 | #E(F_m), where the mod-3 identity is not directly applicable, the implication "leading term ≢ 0 mod p ⇒ p ∤ J_m" of Ota's theorem — which is proved only for p ≥ 5 and under further hypotheses, so that at p = 3 it is a consistency check and not a theorem — would force $c_2 \equiv 0 \pmod 3$ at m = 127, 487, 919, which is what occurs. The re-implementation checked the same implication at p = 5 and 7, which lie in the range of Ota's theorem (his remaining hypotheses were not checked case by case), at the 12 levels in eight curves (m ≤ 3000; m ≤ 10000 for 389a1 and 709a1) where p | J_m, with no violations; no level with p ∈ {7, 11, 13} dividing J_m occurs in that range beyond the four at p = 7, which is expected, since such a level needs p | (m−1)/2, p | #E(F_m) and a non-surjective reduction of E(Q) at once.

Rank 3. For 5077a1 the class $c_3(m) \in I^3/I^4$ and the 3×3 determinant agree with u = −16 at all 49 non-trivial levels tested. The order-of-vanishing statement $\theta_m \in I^3$ alone (tested mod 3 on 8 further levels m ≡ 1 mod 9) showed the leading coefficient vanishing mod 3 more often than in rank 2 (6 of 8), which the full-level computation explains as degeneracy of the 3×3 pairing mod 3 rather than any defect in the conjecture.

Composite levels (Ota's inclusion–exclusion). For $S = m_1m_2$ the conjecture (Ota, Conj. 2.4, which refers to Mazur–Tate 1987, (2.5)–(2.6), (3.1) for d_T and μ_{S,T}; we reconstructed μ_{S,T} from the norm relation rather than from that paper) reads, for $\tilde\theta_S$ the image of θ_S in Z[G_S^+] (Ota's notation), $\tilde\theta_S \equiv \#Ш \cdot \Sigma_{T \mid S} (-1)^{\nu(T)}\, J_T\, \eta_r(\mu_{S,T}(d_T))$ in $I_S{}^r/I_S{}^{r+1}$, where d_T is the discriminant of the level-T pairing and μ_{S,T} lifts $\mathrm{Sym}^r(G_T{}^+)$ to $\mathrm{Sym}^r(G_S{}^+)$ and multiplies by φ(S/T). We push everything to the quotient $\bar G = Z/n_1 \times Z/n_2$ of $G_S{}^+$ ($n_i = (m_i - 1)/2$), where $I^2/I^3 \cong Z/n_1{\cdot}x_1{}^2 \oplus Z/g{\cdot}x_1x_2 \oplus Z/n_2{\cdot}x_2{}^2$ with $g = \gcd(n_1, n_2)$ (Passi's theorem $I^2/I^3 \cong \mathrm{Sym}^2(\bar G)$), and the level-S pairing of R ∈ E_S(Q) is the pair of level-m_i recipes $(h^1(R), h^2(R))$. Writing H^k for the polarized matrix at m_k and $R_i = c_{Tam}\cdot\#E(F_{m_1})\cdot\#E(F_{m_2})\cdot P_i$, the discriminant is $\det(H^1x_1 + H^2x_2) = D_1x_1{}^2 + D_{12}x_1x_2 + D_2x_2{}^2$ with $D_k = \det H^k$ and $D_{12} = H^1{}_{11}H^2{}_{22} + H^2{}_{11}H^1{}_{22} - 2H^1{}_{12}H^2{}_{12}$. Two consequences, in rank 2: the terms $T = m_1, m_2$ contribute only to $x_i{}^2$, where the sum over T collapses (by the norm relation $\pi_{S/m_2}\theta_S = (a_{m_2} - \sigma - \sigma^{-1})\theta_{m_1}$, which is a theorem) to the statement that the $x_1{}^2$-coefficient of θ_S equals $(a_{m_2} - 2)\cdot c_2(m_1) \bmod n_1$; and the mixed coefficient receives only the term T = S, giving the one genuinely new prediction (a_{11} denotes the coefficient of x_1x_2 in the expansion of θ_S, after the reductions of §1 in each variable)

$$4\,M^3 \cdot a_{11}(\theta_S) \equiv -u_E \cdot D_{12} \pmod g, \quad M = c_{Tam}\cdot\#E(F_{m_1})\cdot\#E(F_{m_2}),$$

with the same u_E = 8 as at prime level and the sign reversed, because ν(S) = 2. For 389a1 we tested all 28 pairs of primes $7 \le m_1 < m_2 \le 61$ with g ≥ 3 (code `mt_ota.gp`, `mt_ota_run.py`): $\theta_S \in I_S{}^2$ in every case; the $x_1{}^2$ and $x_2{}^2$ norm-relation checks hold in every case; and the mixed identity holds in every case, non-trivially (both sides ≠ 0) at 10 pairs with g ∈ {3, 5, 6, 7}, including (11, 31) mod 5 and (29, 43) mod 7, where the alternatives u = −8 (no sign change) and u = ±4, ±16 are excluded. The remaining pairs have $D_{12} \equiv 0 \equiv a_{11} \pmod g$ and are consistent but uninformative. The cost is in the points, not the symbols: R_i has about $(m_1m_2)^2\cdot\hat h(P_i)$ nats, i.e. 10^5–10^6 digits for $m_i \le 61$, so m_i of a few hundred are out of reach of exact arithmetic

without a p-adic shortcut. An earlier draft of this paragraph, written before the formula was implemented, said that the three second-order coefficients "do not follow a single-constant extrapolation of the prime-level identity"; that was correct and is explained by the T = m_i terms above.

Comparison with Bley–Macías Castillo at their levels. Bley and Macías Castillo (2022, §4.4) verify the 3-part of the equivariant Tamagawa number conjecture for E over the cubic subfield F of Q(ζ_ℓ) for 389a1 (ℓ = 7, 13, 43), 433a1 (ℓ = 7, 13, 31, 37) and 446d1 (ℓ = 19). Their statement (Thm. 2.1, condition (11)) is that L*/Reg lies in $Z_3[G]^\times$, where Reg is built from the Néron–Tate regulator and a 2×2 matrix Ψ of Mazur–Tate pairing values ⟨P_i, P_j⟩ ∈ I(G)/I(G)² ≅ Z/3 (minus the Bertolini–Darmon pairing, computed from generalized 3-Selmer groups and Hilbert symbols in Magma); their published output (file ETNC_V.out on W. Bley's page) records only the verdict "ETNC is valid" per pair, not the values of Ψ or of L*, and their code requires det Ψ ≠ 0 (mod 3) as part of the verdict. The comparison we can make without Magma is therefore at the level of that determinant: in our data, det⟨R_i, R_j⟩_m (mod 3) is non-zero at all eight of their levels (389a1: 2, 1, 2 at m = 7, 13, 43; 433a1: 2, 2, 1, 1 at m = 7, 13, 31, 37; 446d1: 1 at m = 19), in agreement with the nonvanishing their verification certifies, and our full identity mod n = (m−1)/2 holds at each of these levels (433a1, c_∞ = 1, is the tenth curve in the table, added for this comparison, with u_E = 4 at 21 levels). The two pairings are related by the projection $G_m^+ \to \mathrm{Gal}(F/Q) \cong Z/3$, so this is the expected agreement rather than an independent confirmation of the values; matching Ψ entry by entry would require re-running their code.

Independent re-implementation and blind test. After this note reached its present form, the prime-level identity was re-implemented from the text alone (without access to our scripts) in a separate session, and the constant $u_E = (-2)^r c_\infty / \#E(Q)\mathrm{tors}^2$ was recorded in advance for seven curves not in the table above. The identity then held on all of them — 563a1, 643a1, 664a1 (c_Tam = 4), 681c1 (c_Tam = 2), 707a1 (c_Tam = 2), 997b1, and the rank-3 curve 11197a1 — at 237 non-trivial levels (7 ≤ m ≤ 200 in rank 2, m ≤ 150 in rank 3) with no failures, u_E as predicted in every case, including the sign −16 for the second rank-3 curve. The same re-implementation, with the hypothesis u_E = −2c∞ recorded in advance, verified the rank-1 case on 37a1, 43a1, 53a1, 57a1, 58a1, 61a1, 77a1 and 79a1 at 533 levels m ≤ 400 with no failures (the opposite sign agrees at 1–3 levels per curve, by chance); all eight curves have trivial torsion, so the general law reduces there to u_E = −2c∞. In rank 0 the identity is the statement about the augmentation class, $c_0(m) = \Sigma c_t$, and the law takes the form $u_E \cdot \#E(Q)\mathrm{tors}^2 = c_\infty \cdot \#Ш$ (hypothesis recorded in advance); it held at all 249 levels m ≤ 400 on eight curves, including the two with non-trivial Ш: Negative controls fail as they should: a wrong u_E, omitting c_Tam, pairing the points of one curve with the theta element of another, or the wrong sign in rank 3 each give at most 2 accidental agreements out of 26–39. The re-implementation also reproduces the Appendix exactly. Both implementations use PARI/GP 2.17.2, so this is independence of code and of the choice of constant, not of the library.

Curve	c∞	c_Tam	#E(Q)tors	#Ш	levels	failures
11a1	1	5	5	1	29	0
14a1	1	6	6	1	18	0
15a1	2	8	8	1	30	0
32a1	1	4	4	1	39	0
67a1	1	1	1	1	39	0
75a1	1	1	1	1	27	0
571a1	1	1	1	4	37	0
681b1	2	4	4	9	30	0

This is the only place where the factor #Ш of the conjecture is tested with #Ш > 1; in all curves of positive rank considered here #Ш = 1. With #Ш included, the constant is $u_E = (-2)^r \cdot c_\infty \cdot \#Ш / \#E(Q)\mathrm{tors}^2$ in ranks 0–3.

5. Direction of Kolyvagin classes in rank 2

Let E/Q have rank 2 and root number +1, K an imaginary quadratic field satisfying the Heegner hypothesis for N, and p an odd prime with E[p] irreducible, p ∤ N·c_Tam·#E(K)tors. For a Kolyvagin prime ℓ (inert in K, p | a_l, p | ℓ+1) let $y_l \in E(K[\ell])$ be the Heegner point of conductor ℓ and $\kappa_l \in H^1(K, E[p])$ the Kolyvagin derived class of $\Sigma\, \sigma\, D_l\, y_l$, the sum over coset representatives σ of G = Gal(K[ℓ]/K[1]) ≅ Z/(ℓ+1) in Gal(K[ℓ]/K). Since the Heegner point y_K ∈ E(K) is torsion (here literally 0), $\kappa_l \in \mathrm{Sel}_p(E/K)^+ = E(Q)/pE(Q) \cong F_p^2$ when Ш(E/Q)[p] = 0. Let Q generate E^D(Q) ⊗ Q, D = disc K; in every case below rank E^D(Q) = 1, checked with ellrank.

Statement tested. (a) κ_ℓ ≠ 0 if and only if the reduction map E(Q)/p → E(F_ℓ)/pE(F_ℓ) is nonzero and Q is not p-divisible in E^D(F_ℓ). (b) When κ_ℓ ≠ 0, F_p·κ_ℓ = ker(E(Q)/p → E(F_ℓ)/pE(F_ℓ)); in particular the line is independent of K.

Both follow from the transverse local condition at ℓ in Kolyvagin systems (Howard 2004, Def. 1.2.2; Mazur–Rubin 2004) together with the vanishing of y_K and the core-rank-one structure theory; the statement appears as Prop. 0.2 of an unpublished and incomplete draft hosted on W. Stein's website (no author or date given; its proof refers to a missing lemma). We are not aware of any numerical test of (b).

Method. The h = ℓ+1 (or 2(ℓ+1) when h(K) = 2) Heegner points of conductor ℓ are computed to 150–450 digits from the modular parametrization; x(y_ℓ) has a rational minimal polynomial f of degree h, recognized exactly (denominators up to 10^{113}). A reduced primitive element θ = B(x) (`polredbest`) gives L = Q(θ); the Galois action of a generator σ of G on the roots is recovered, without interpolation, from the integer resolvents ∏(T − (θ_k + c(θ_{σk} + θ_{σ⁻¹k}))), c = 1,2,3, and (for orientation) the O_K-resolvents ∏(T − (θ_k + cθ_{σk})), c = 2,3. The y-coordinate is recovered from the exact square root of $(4x^3 + b_2x^2 + 2b_4x + b_6)/(4D)$ in L (`nfroots`), whose existence encodes the action of complex conjugation. At each prime q ≡ 1 (mod p) totally split in K[ℓ], all points reduce to E(F_q), the derivative $D_\ell = \Sigma\, i\sigma^i$ is applied along the σ-cycle of roots mod q, and the class of the result in E(F_q)/p is read off by the order-p Tate pairing (checked against direct enumeration of pE(F_q)) and compared with the reductions of P_1, P_2 and Q. Twenty such primes identify the class uniquely; the trace relation Tr y_ℓ = a_ℓ y_1 holds at every prime as a check. A control case with a theorem-level prediction (43a1, rank 1, K = Q(√−8), ℓ = 29, where κ = 0 is forced) gave 0 at 15/15 primes.

Curve	p	K	ℓ	predicted	computed
389a1	3	$\mathbb{Q}(\sqrt{-5})$ (h = 2)	17	$\langle P_1 - P_2 \rangle$	$-(P_1 - P_2)$
389a1	3	$\mathbb{Q}(\sqrt{-7})$	5 / 17 / 41 / 59	$\langle P_1{+}P_2 \rangle$ / 0 / $\langle P_1{-}P_2 \rangle$ / $\langle P_2 \rangle$	all as predicted
389a1	3	$\mathbb{Q}(\sqrt{-11})$	17 / 29 / 41	$\langle P_1{-}P_2 \rangle$ / 0 / $\langle P_1{-}P_2 \rangle$	all as predicted
389a1	3	$\mathbb{Q}(\sqrt{-19})$	29 / 41 / 53 / 59	$\langle P_1{-}P_2 \rangle$ / $\langle P_1{-}P_2 \rangle$ / 0 / $\langle P_2 \rangle$	all as predicted
389a1	**5**	$\mathbb{Q}(\sqrt{-7})$, $\mathbb{Q}(\sqrt{-11})$	19	$\langle P_1{+}P_2 \rangle$	as predicted (both)
571b1	3	$\mathbb{Q}(\sqrt{-7})$ / $\mathbb{Q}(\sqrt{-8})$	47, 59 / 47	0, $\langle P_1{+}P_2 \rangle$ / $\langle P_1{+}P_2 \rangle$	all as predicted
709a1	3	$\mathbb{Q}(\sqrt{-7})$	5 / 47 / 59	$\langle P_1 \rangle$ / 0 / $\langle P_1{-}P_2 \rangle$	all as predicted

Twenty cases, three curves, two primes p, five zero predictions, four distinct lines of F_p^2, no failures (the 43a1 control is not counted). Basis vectors P_1, P_2 are the saturated `ellrank` generators of each curve (for 389a1 in §5: (−1,1), (0,0) in the first row, `ellrank`'s (0,−1), (1,0) elsewhere; since (0,−1) = $-P_2$ and (1,0) = $P_1 + P_2$ in the first basis, $P_1 - P_2 \equiv P_2' - P_1'$ (mod 3) and the label $\langle P_1{-}P_2 \rangle$ denotes the same line in both). For h(K) = 2 the class must be summed over the two cosets of G with consistent orientation; summing one coset gives a well-defined but wrong class, which is how the h = 2 computation first failed.

Comparison with Jetchev–Lauter–Stein. Two of the cases above were computed before, by a different method: JLS (2009, Prop. 3.10 and Rem. 3.11) show $\kappa_5 \neq 0$ for 389a1 and for 709a1 with $K = \mathbb{Q}(\sqrt{-7})$, p = 3, by proving that the Heegner point P_5 is not 3-divisible in E(K[5]) (the polynomial $\varphi_3(x) - x(P_5)\psi_3(x)^2$ has no root in K[5]). Our table gives $\kappa_5 \neq 0$ in both cases, with the additional information of the line: $\langle P_1{+}P_2 \rangle$ for 389a1 and $\langle P_1 \rangle$ for 709a1. JLS did not compute the direction of the class, and the two methods share no code or intermediate data beyond the Heegner points themselves; their nonvanishing results and ours agree.

6. Relation to the literature and what is new

Theory. Mazur–Tate (1987) formulated the refined conjectures; we cite it only through Tan, Ota and Burns–Kurihara–Sano, not having had access to it. Ota (2018) proved $\theta_S \in I^{r}$ (the order part) for the p-parts under hypotheses, and the implication "leading term $\neq 0$ mod $p \Rightarrow Ш[p] = 0$ and $p \nmid J$", both for $p \geq 5$ (his §§3, 5). Burns–Kurihara–Sano (2021, published IMRN 2025) recast the leading-term conjecture in the form used here and proved cases in rank 1 (Cor. 1.2: supersingular reduction, squarefree conductor and further conditions, the leading term up to an automorphism of Gal(F/Q)); they also remark (§1) that the only supporting numerical evidence then available was for r = 1 (Portillo-Bobadilla); Bullach–Honnor (2025) prove substantial parts of Mazur–Tate's Conjectures 3, 4 and 6 for $p > 3$ under a big-image hypothesis (e.g. E semistable and $p \geq 11$) and mild local conditions; their leading-term result is trivial in positive rank (their Rem. 1.8(b)). Bertolini–Darmon (1994, 1995) introduced derived heights for the degenerate case; the parity phenomenon in §4 is consistent with their Thm. 3.18.

Numerics. Portillo-Bobadilla (2019) treats rank 1. Séguin (McGill M.Sc. thesis, 2014/15) tests Darmon's Heegner-point analogue of the refined conjecture, in conductor 37 over real quadratic fields, computing Mazur–Tate pairings in that setting; it is not a test of the Mazur–Tate conjectures over Q. Llerena-Córdova (2024) tests three of Mazur–Tate's conjectures on 425,713 pairs (E, p) with split multiplicative reduction at p, finding failures that disappear when a factor $1/\#E(\mathbb{Q})_{\mathrm{tors}}$ is included; it does not treat the rank ≥ 2 leading-term statement with a Mazur–Tate regulator (its Conj. 2.12 asserts only the order of vanishing in positive rank); its conclusion that $1/\#E(\mathbb{Q})_{\mathrm{tors}}$ must be adjoined is consistent with the factor $1/\#E(\mathbb{Q})_{\mathrm{tors}}^2$ in u_E found here. The closest work is Bley–Macías Castillo (2022): a numerical verification, assuming classical BSD for E/F and $Ш[3] = 0$, of the 3-adic equivariant Tamagawa number conjecture — equivalent to congruences in the augmentation filtration involving the Mazur–Tate pairing — for 389a1 (ℓ = 7, 13, 43), 433a1 and 446d1 (ℓ = 19), in the cubic subfield of $\mathbb{Q}(\zeta_\ell)$, with the pairing computed via generalized Selmer groups and Hilbert symbols in Magma.

What is new here, to the best of our knowledge: (i) a verification of the leading-term conjecture in rank 3; (ii) in rank 2, a verification with θ_m computed from modular symbols at the full level G_m^+ (moduli (m−1)/2 up to 749), with both sides independent, at 21–221 non-trivial levels per curve; (iii) the elementary coordinate formula for the pairing of §2 — $\langle R,S \rangle = [\gamma/A(R+S)]$, $\langle R,R \rangle = [A(R)]^{-1}$ — with its derivation from the biextension, which we have not found stated anywhere; (iv) the explicit confirmation that the Tamagawa factor enters through the joint kernel defining $E_S(\mathbb{Q})$; (v) the direction test for Kolyvagin classes of §5. Items (i)–(iv) are verifications of an existing conjecture, not new mathematics; item (iii) is a formula whose proof is complete except for the two points named in §7.1.

7. Caveats, open points and next steps

1. **Status of the pairing formula.** §2 derives the recipe from the biextension definition, granting the Néron-model description of the canonical local splittings (vertical order at unramified places; m-adic order and reduction to the special fibre at m). What is still not written out is an elementary integrality statement: that the cancellation δ in the duplication formula is 1 at $\ell = 2$ for points of $E_0(\mathbb{Q}_2)$. It is proved for all odd ℓ (and the content of the line is proved to be 1 at every ℓ); the argument fails at $\ell = 2$ because it uses that 2 is a unit in the formal group. It holds at the 349 + 3631 points checked, which cover every reduction type at 2 systematically: 166 curves of good reduction at 2, multiplicative I_n with $n \leq 19$, and all additive types II, III, IV, I_0^*, I_n^*, II*, III*, IV* (count in the companion repository, kodaira2.py). We do not attempt to force an argument at $\ell = 2$ here. Someone with Mazur–Tate (1983) in hand should also confirm that the splittings we used are theirs; Tan's description of (L3)/(L4) and the uniqueness argument of §2 (iii) leave little room for a different choice. With these two points settled the formula $\langle R,S \rangle = [\gamma_{R,S}/A(R+S)]$, and hence $\langle R,R \rangle = [A(R)]^{-1}$, is a theorem.
2. **Powers of 2 and sign — explained.** The constant $u_E = (-2)^r \cdot c_\infty / \#E(\mathbb{Q})_{\mathrm{tors}}^2$ of §4 is explained factor by factor, sign included, and the stress tests on 655a1, 1132a1, 916c1 and 1088j1 confirm that the Tamagawa numbers do not enter it. Our conventions are: $x = \sigma - 1$ (so $\eta_r(g_1 \cdots g_r) = \prod(\sigma(g_i) - 1)$, as in Ota and Burns–Kurihara–Sano), the primitive root g fixed by PARI's znprimroot, PARI's plus symbol (which has $x^+(\{0\}-\{\infty\})$

= −L(E,1)/ω_1, the opposite sign to Mazur–Tate's [0]$^+$ = L(E,1)/Ω^+), and the pairing ⟨R,S⟩ = [γ/A(R+S)] of §2, whose inverse is the pairing in Tan's orientation. Undoing these two sign conventions turns our $(-1)^r$ into Ota's (−1)^{v(m)} = −1 for every r, as §4(b) explains, assuming that Ota normalizes the Artin map as Tan does (uniformizer ↦ arithmetic Frobenius; the standard choice, not checked in his text and asked of him directly); the rank-1 data of the independent re-implementation (533 levels, u_E = −2c∞) were what fixed the dependence on r. The v0.1 text of this item said the sign was open and suggested a factor (−1)^{r(r−1)/2}; both statements were wrong.

3. **Weak levels.** When gcd(M, n) > 1 the identity determines c_r(m) only modulo n/gcd(M,n). About half the levels for 389a1 are of this kind. They are still non-trivial checks, but the headline counts should be read with this in mind.
4. **One analytic input.** #Ш = 1 is taken from the analytic order (strong BSD), and every curve of rank ≥ 1 tested here (ours and the re-implementation's) has #Ш = 1, so the factor #Ш in the conjecture is tested only in rank 0 (by the re-implementation, on 571a1 with Ш = 4 and 681b1 with Ш = 9; Llerena-Córdova's rank-0 tests cover it on a much larger scale at primes of split multiplicative reduction). The identity tested in positive rank is therefore a joint consistency check of Mazur–Tate and of BSD's Ш-term; a curve with Ш ≠ 1 in rank 2 would be a sharper test (none exists in Cremona's tables up to conductor 500,000: the analytic-Ш column of the allbsd files is 1 for all 279,056 optimal rank-2 curves, checked with an awk one-liner recorded in the companion document).
5. **Composite levels** are now tested through Ota's inclusion–exclusion formula for S = m_1m_2 (§4); the mixed coefficient x_1x_2 is the only genuinely new information at composite level and is checked only modulo gcd(n_1, n_2), which limits the strength of each individual check; extending to three prime factors and to larger gcd is straightforward but costly (the points R_i have \∼10^6 digits).
6. **Rank 4** (234446a1, [1,−1,0,−79,289], c_Tam = 2, c_∞ = 2). The code runs unchanged for r = 4 (the identity is in I^4/I^5 with a 4×4 determinant), but PARI's `msfromell` for conductor 234446 needs more than the 7 GB available to us: the attempt was killed by the out-of-memory handler before the first level. On a machine with \∼32 GB, or with a modular-symbol implementation that works in the plus quotient only, this is a few hours of computation.
7. **Bley–Macías Castillo.** The comparison in §4 is at the level of det Ψ ≠ 0 (mod 3), which is all their published output allows. An entry-by-entry match of the 2×2 matrix Ψ against the mod-3 reduction of our pairing matrix would be a sharper test; it requires running their Magma code (available on W. Bley's page), to which we have no access.

None of this touches the archimedean Birch–Swinnerton-Dyer conjecture: the "derivative" in the Mazur–Tate conjecture is in the group-ring direction σ − 1, not in the complex variable s, and the identification of the two is the open problem.

Appendix. A worked example: 389a1 at m = 23

E: $y^2 + y = x^3 + x^2 - 2x$, $P_1 = (-1, 1)$, $P_2 = (0, 0)$, c_Tam = 1, c_∞ = 2, E(Q)_tors = 0. Take m = 23, so n = 11, g = 5 (PARI's primitive root), $G_{23}^+ = \langle\sigma\rangle \cong Z/11$.

Theta element. The plus symbols [a/23]$^+$ for a = 1, …, 22 are

0, −1, 0, 0, −1, −1, −1, 1, 2, 1, 0, 0, 1, 2, 1, −1, −1, −1, 0, 0, −1, 0

(symmetric under a ↔ 23 − a, as they must be). Collecting a and −a by t = $\log_5$(a) mod 11, the coefficient vector of $\theta_{23} = \Sigma\, c_t\, \sigma^t$ is

$(c_0, \ldots, c_{10}) = (0, -2, -2, 2, 0, 0, 2, -2, -2, 0, 4)$.

Expanding $\sigma^t = (1 + x)^t$: $a_0 = \Sigma\, c_t = 0$, $a_1 = \Sigma\, t\, c_t = 22 = 2\cdot 11$, $a_2 = \Sigma\, C(t,2)\, c_t = 116$. So $\theta_{23} \in I$, and since $a_1 = 2n$ the relation $n\cdot x \equiv -C(11,2)x^2$ gives $\theta_{23} \in I^2$ with leading class $c_2(23) = 116 - 2\cdot 55 = 6 \in Z/11$.

Points. #E(F_{23}) = 28 ($a_{23} = -4$), so M = 28 and R_i = 28·P_i. The x-coordinate of R_1 has a 234-digit numerator and denominator $d(R_1)^2$ with

$d(R_1)$ = 607852338707617298241916768285023402187465773080649173756581131907387799193727318114990310685196156742789291848564509 (117 digits),

$d(R_2)$ = 161961604235457625196184711884753920723064530210946525 77 (56 digits), and $d(R_1 + R_2)$ has 81 digits. In all three cases $v_{23}(d) = v_{23}(t) = 1$, as expected for points in $E_1(Q_{23}) \setminus E_2(Q_{23})$. The unit parts give

$(t/d)(R_1) \equiv 14$, $(t/d)(R_2) \equiv 14$, $(t/d)(R_1 + R_2) \equiv 20 \pmod{23}$,

and with $5^{10} \equiv 9 \equiv -14$, $5^5 \equiv 20 \pmod{23}$ the logarithms in Z/11 are $h(R_1) = 10$, $h(R_2) = 10$, $h(R_1+R_2) = 5$. (That $h(R_1) = h(R_2)$ is a coincidence of this level.) The polarized matrix is

$H = (2h(R_1),\ h(R_1+R_2) - h(R_1) - h(R_2);\ \cdot,\ 2h(R_2)) = (20, -15; -15, 20) \equiv (9, 7; 7, 9) \pmod{11}$, $\det H \equiv 81 - 49 \equiv 10 \pmod{11}$.

The identity. Left side: $2^2\cdot M^3\cdot c_2 = 4\cdot 28^3\cdot 6$; with $28 \equiv 6$ and $6^3 \equiv 7 \pmod{11}$ this is $4\cdot 7\cdot 6 = 168 \equiv 3$. Right side: u_E·det H = 8·10 = 80 ≡ 3 (mod 11). The two sides agree, with the same u_E = 8 = $(-2)^2$·c_∞ as at the other 220 non-trivial levels for this curve. In the normalization of §4 ($\theta^+ = \theta/c_\infty$ has leading class 6/2 = 3; torsion trivial) this reads $M^{-3}\cdot\det H = 8\cdot 10 \equiv 3 \equiv \theta_{23}/c_\infty$; with the Mazur–Tate sign of the symbol ($\theta^+ = -\theta/c_\infty$, class $-3 \equiv 8$) and Tan's orientation of the pairing (P = −H, det P = det H), this is $\theta_{23}^+ \equiv -J_{23}\cdot \mathrm{disc}_{23}(P)$ in $I^2/I^3 \cong Z/11$, i.e. Ota's statement with $(-1)^{v} = -1$.

The whole computation is `thetaclass(E,23,2)` and `pairdet(E,23,[P1,P2],28,11,Mod(5,23))` in `mt_rank.gp`, and takes well under a second.

8. Reproducibility, acknowledgements, references

Code and data. All computations use PARI/GP 2.17.2 through cypari2 2.2.4, on Cremona's tables (`allbsd`, `allcurves`). The scripts `mt_full.gp` / `mt_full_run.py` (prime-level identity, rank 2), `mt_rank.gp` / `mt_rank_run.py` (general rank), `mt_lead.gp` (mod-3 version), `mt_comp.gp` (exploratory composite levels), mt_ota.gp / mt_ota_run.py (Ota's composite-level formula), biext.gp / chkg2.gp / chord.gp / gsweep.py (the biextension computation of §2 and the integrality sweep), `heeg.gp` + `kgen3.py` (Kolyvagin classes, any curve and p) and the raw logs are in the accompanying archive. A typical prime-level check for 389a1 up to m = 1500 takes about 40 minutes on one core; the Kolyvagin computation for a field of degree 60 takes 15–20 minutes, dominated by `polredbest`. Two PARI conventions matter: `elltatepairing(E, P, Q, m)` takes the m-torsion point as **P**, and its value lies in $\mathbb{F}_{q^x}/(\mathbb{F}_{q^x})^m$ and must be raised to (q−1)/m; `ellpadicheightmatrix` returns [A, B] with the cyclotomic height matrix equal to $A - s_2 \cdot B$.

Acknowledgements. The computations and the drafting of this note were carried out with the assistance of an AI system (Claude, Anthropic), which also surveyed the literature; all numerical claims were checked by running the code, and the author takes responsibility for the statements. We are grateful to the authors of PARI/GP and to J. Cremona for the tables.

References.

- B. Mazur, J. Tate, *Refined conjectures of the "Birch and Swinnerton-Dyer type"*, Duke Math. J. 54 (1987), no. 2, 711–750.
- B. Mazur, J. Tate, *Canonical height pairings via biextensions*, in Arithmetic and Geometry, Vol. I, Progr. Math. 35, Birkhäuser (1983), 195–237.
- M. Bertolini, H. Darmon, *Derived heights and generalized Mazur–Tate regulators*, Duke Math. J. 76 (1994), no. 1, 75–111; *Derived p-adic heights*, Amer. J. Math. 117 (1995).
- K.-S. Tan, *Refined theorems of the Birch and Swinnerton-Dyer type*, Ann. Inst. Fourier 45 (1995), no. 2, 317–374.
- K. Ota, *Kato's Euler system and the Mazur–Tate refined conjecture of BSD type*, Amer. J. Math. 140 (2018), no. 2, 495–542. arXiv:1509.00682.
- D. Burns, M. Kurihara, T. Sano, *On derivatives of Kato's Euler system and the Mazur–Tate conjecture*, Int. Math. Res. Not. IMRN 2025, no. 4, rnaf012, doi:10.1093/imrn/rnaf012. arXiv:2103.11535.
- W. Bley, D. Macías Castillo, *Congruences for critical values of higher derivatives of twisted Hasse–Weil L-functions, III*, Math. Proc. Cambridge Philos. Soc. 173 (2022), no. 2, 431–456 (online 2021), doi:10.1017/S0305004121000657. arXiv:1912.11260.
- D. Bullach, M. H. L. Honnor, On the refined "Birch–Swinnerton-Dyer type" conjectures of Mazur and Tate, arXiv:2511.07203 (2025).
- B. Mazur, W. Stein, J. Tate, *Computation of p-adic heights and log convergence*, Doc. Math. 2006, Extra Vol. (John H. Coates' Sixtieth Birthday), 577–614.
- B. Howard, *The Heegner point Kolyvagin system*, Compos. Math. 140 (2004), no. 6, 1439–1472. B. Mazur, K. Rubin, *Kolyvagin systems*, Mem. Amer. Math. Soc. 168 (2004), no. 799.
- B. Gross, *Kolyvagin's work on modular elliptic curves*, in L-functions and Arithmetic, LMS Lecture Note Ser. 153 (1991), 235–256.
- D. Jetchev, K. Lauter, W. Stein, *Explicit Heegner points: Kolyvagin's conjecture and non-trivial elements in the Shafarevich–Tate group*, J. Number Theory 129 (2009), no. 2, 284–302. Kolyvagin classes for higher rank elliptic curves, unpublished and incomplete draft, no author or date given, hosted at wstein.org/papers/current/kolydensity/ (Prop. 0.2).
- F. Séguin, Numerical verification of a "Birch and Swinnerton-Dyer type" conjecture, M.Sc. thesis, McGill University (2014/2015). F. X. Portillo-Bobadilla, Experimental evidence on a refined conjecture of the BSD type, Bol. Soc. Mat. Mex. (3) 25 (2019), no. 3, 529–541. arXiv:1709.01204. J.-P. Llerena-Córdova, Numerical study of refined conjectures of the BSD type, arXiv:2412.17703.
- J. Cremona, *Elliptic curve data*, github.com/JohnCremona/ecdata.