

Homework #7

April 11, 2024

Name: Ravi Raghavan

Extension: No

Problem 1. *Voronoi description of halfspace.* Let a and b be distinct points in \mathbb{R}^n . Show that the set of all points that are closer (in Euclidean norm) to a than b , i.e., $\{x : \|x - a\|_2 \leq \|x - b\|_2\}$, is a halfspace. Describe it explicitly as an inequality of the form $c^T x \leq d$. Draw a picture.

Solution. We can see that the following two sets are equivalent:

$$S_1 = S_2 \text{ where } S_1 = \{x : \|x - a\|_2 \leq \|x - b\|_2\} \text{ and } S_2 = \{x : \|x - a\|_2^2 \leq \|x - b\|_2^2\}$$

Let's work with S_2 since it will be a lot easier

$$\begin{aligned} S_2 &= \{x : \|x - a\|_2^2 \leq \|x - b\|_2^2\} \\ S_2 &= \{x : \|x\|_2^2 - 2 \langle x, a \rangle + \|a\|_2^2 \leq \|x\|_2^2 - 2 \langle x, b \rangle + \|b\|_2^2\} \\ S_2 &= \{x : -2 \langle x, a \rangle + \|a\|_2^2 \leq -2 \langle x, b \rangle + \|b\|_2^2\} \\ S_2 &= \{x : 2 \langle x, b - a \rangle \leq \|b\|_2^2 - \|a\|_2^2\} \\ S_2 &= \{x : 2(b - a)^T x \leq \|b\|_2^2 - \|a\|_2^2\} \\ S_2 &= \{x : (b - a)^T x \leq 0.5(\|b\|_2^2 - \|a\|_2^2)\} \end{aligned}$$

This is a closed half-space

Problem 2. Which of the following sets S are polyhedra? If possible, express S in the form $S = \{x | Ax \preceq b, Fx = g\}$

(b) Yes S is a polyhedra.

Let $M_1 = [a_1, a_2, \dots, a_n] \in \mathbb{R}^{1 \times n}$ and let $M_2 = [a_1^2, a_2^2, \dots, a_n^2] \in \mathbb{R}^{1 \times n}$

Let F be the vertical concatenation of 1^T , M_1 , and M_2 . Let g be $[1, b_1, b_2]^T$

Let $A = -I$

We can express S , via compact notation, as $S = \{x | Ax \preceq 0, Fx = g\}$

(c) S is NOT a Polyhedra

Problem 3. *Hyperbolic sets.* Show that the *hyperbolic set* is $\{x \in \mathbb{R}_+^2 : x_1 x_2 \geq 1\}$ is convex. As a generalization, show that $\{x \in \mathbb{R}_+^n : \prod_{i=1}^n x_i \geq 1\}$ is convex. *Hint.* If $a, b \geq 0$ and $0 \leq \theta \leq 1$, then $a^\theta b^{1-\theta} \leq \theta a + (1 - \theta)b$

Solution. Let S be $\{x \in \mathbb{R}_+^n : \prod_{i=1}^n x_i \geq 1\}$. Let's have two vectors j and k that are in S . Let the elements of j be j_1, j_2, \dots, j_n . Let the elements of k be k_1, k_2, \dots, k_n

We want to prove that $\theta j + (1 - \theta)k \in S$.

Since we know that $j \in S$ and $k \in S$, we can state the following:

- $j_1 \geq 0, j_2 \geq 0, \dots, j_n \geq 0$
- $j_1 j_2 j_3 \dots j_n \geq 1$
- $k_1 \geq 0, k_2 \geq 0, \dots, k_n \geq 0$
- $k_1 k_2 k_3 \dots k_n \geq 1$

For $i \in [1, n]$, $j_i, k_i \geq 0$ and $0 \leq \theta \leq 1$, we can see that $0 \leq j_i^\theta k_i^{1-\theta} \leq \theta j_i + (1 - \theta)k_i$.

We can also see that:

$$\prod_{i=1}^n j_i^\theta k_i^{1-\theta} = (\prod_{i=1}^n j_i)^\theta (\prod_{i=1}^n k_i)^{1-\theta}$$

Since $j_1 j_2 j_3 \dots j_n \geq 1$ and $k_1 k_2 k_3 \dots k_n \geq 1$, we can say that:

$$\prod_{i=1}^n j_i^\theta k_i^{1-\theta} = (\prod_{i=1}^n j_i)^\theta (\prod_{i=1}^n k_i)^{1-\theta} \geq 1$$

Since $j_i^\theta k_i^{1-\theta} \leq \theta j_i + (1 - \theta)k_i$,

$$1 \leq \prod_{i=1}^n j_i^\theta k_i^{1-\theta} \leq \prod_{i=1}^n (\theta j_i + (1 - \theta)k_i)$$

We have shown that $\theta j_i + (1 - \theta)k_i \in S$ and that S is a convex set!

Since we have proved the generalized case, we can say that $\{x \in \mathbb{R}_+^2 : x_1 x_2 \geq 1\}$ is convex as well

Problem 4. Problem 2.16:

Show that if S_1 and S_2 are convex sets in \mathbb{R}^{m+n} , then so is their partial sum
 $S = \{(x, y_1 + y_2) : x \in \mathbb{R}^m, y_1, y_2 \in \mathbb{R}^n, (x, y_1) \in S_1, (x, y_2) \in S_2\}$

Solution. Let's say that we have two points in S , namely $(x_1, y_{11} + y_{12})$ and $(x_2, y_{21} + y_{22})$. To prove that S is convex, we need to show that $\theta(x_1, y_{11} + y_{12}) + (1 - \theta)(x_2, y_{21} + y_{22})$ is in S .

Based on the definition of S , we can see the following:

- $(x_1, y_{11}) \in S_1$
- $(x_1, y_{12}) \in S_2$
- $(x_2, y_{21}) \in S_1$
- $(x_2, y_{22}) \in S_2$

Since S_1 and S_2 are convex, based on the definition of convex sets, we can see that:

- $\theta(x_1, y_{11}) + (1 - \theta)(x_2, y_{21}) \in S_1$
 $(\theta x_1 + (1 - \theta)x_2, \theta y_{11} + (1 - \theta)y_{21}) \in S_1$
- $\theta(x_1, y_{12}) + (1 - \theta)(x_2, y_{22}) \in S_2$
 $(\theta x_1 + (1 - \theta)x_2, \theta y_{12} + (1 - \theta)y_{22}) \in S_2$

By definition of Set S , we can see that:

$$(\theta x_1 + (1 - \theta)x_2, \theta y_{11} + (1 - \theta)y_{21} + \theta y_{12} + (1 - \theta)y_{22}) \in S$$

$$(\theta x_1 + (1 - \theta)x_2, \theta(y_{11} + y_{12}) + (1 - \theta)(y_{21} + y_{22})) \in S$$

$$\theta(x_1, y_{11} + y_{12}) + (1 - \theta)(x_2, y_{21} + y_{22}) \text{ is in } S.$$

Problem 5 (Problem 2.19(a)). *Linear-fractional functions and convex sets.* Let $f : \mathbb{R}^m \rightarrow \mathbb{R}^n$ be the linear-fractional function

$$f(x) = (Ax + b)/(c^T x + d), \quad \text{dom } f = \{x | c^T x + d > 0\} \quad (1)$$

In this problem, we study the inverse image of a convex set C under f , i.e.,

$$f^{-1}(C) = \{x \in \text{dom } f : f(x) \in C\} \quad (2)$$

For each of the following sets $C \subseteq \mathbb{R}^n$, give a simple description of $f^{-1}(C)$

Solution. Let's look at the halfspace $C = \{y : g^T y \leq h\}$ (with $g \neq 0$).

$$f^{-1}(C) = \{x : g^T((Ax + b)/(c^T x + d)) \leq h, c^T x + d > 0\}$$

Since $c^T x + d > 0$

$$f^{-1}(C) = \{x : g^T(Ax + b) \leq h(c^T x + d), c^T x + d > 0\}$$

$$f^{-1}(C) = \{x : g^T Ax + g^T b \leq h c^T x + h d, c^T x + d > 0\}$$

$$f^{-1}(C) = \{x : (g^T A - h c^T)x \leq h d - g^T b, c^T x + d > 0\}$$

$$f^{-1}(C) = \{x : (A^T g - h c^T)^T x \leq h d - g^T b, c^T x + d > 0\}$$

Let's call a new vector $p^T = (A^T g - h c^T)^T$ and $q = h d - g^T b$

$$f^{-1}(C) = \{x : p^T x \leq q, c^T x + d > 0\}$$

We can see that $f^{-1}(C)$ is just the intersection of a halfspace and the domain of f !
