

MATH 4630 / 6632 3.0 - Fall 2022

Assignment 3

(Due Date: December 2, 2022)

Question 1: Consider the data given in the EXCEL file tab “q1”.

- a. State the multivariable linear regression model with all the necessary assumptions.
- b. Find the predicted model.
- c. Test the significance of the model.
- d. Regardless of your result in part (c), test if X_1 is significant? How about X_2 ?
- e. Find a 95% working Hotelling confidence region for the mean response when $X_1 = 192$ and $X_2 = 152$.
- f. Find a 95% working Hotelling prediction region for a new response when $X_1 = 192$ and $X_2 = 152$.

Question 2: Timm (1975) reported the results of an experiment in which subjects' respond time to “probe words” at five position (Y_1 is at the beginning of the sentence, Y_2 is in the first quartile of the sentence, Y_3 is in the middle of the sentence, Y_4 is in the third quartile of the sentence, and Y_5 is at end of the sentence). The data are recorded in the EXCEL file tab “q2”.

- a. Use the sample variance and obtain all the principle components.
- b. Timm specifically required the reduction in dimension should cover at least 90% of the total variance. How many principle components are needed? Why?
- c. Repeat parts (a) and (b) using the sample correlation matrix.

Question 3: Use the data in the EXCEL file tab “q2”.

- Find the canonical correlation between (x_1, x_2, x_3) and (x_4, x_5) .
- Test the significance canonical correlations.
- Regardless of your answer in part (b), is each canonical correlation individually significant?

Question 4:

- Show that

$$-\frac{1}{2}(\underline{x} - \underline{\mu}_1)' \Sigma^{-1}(\underline{x} - \underline{\mu}_1) + \frac{1}{2}(\underline{x} - \underline{\mu}_2)' \Sigma^{-1}(\underline{x} - \underline{\mu}_2) \\ (\underline{\mu}_1 - \underline{\mu}_2)' \Sigma^{-1} \underline{x} - \frac{1}{2}(\underline{\mu}_1 - \underline{\mu}_2)' \Sigma^{-1}(\underline{\mu}_1 + \underline{\mu}_2)$$

- Let

$$f_1(x) = (1 - |x|) \quad \text{for } |x| \leq 1 \\ f_2(x) = (1 - |x - 0.5|) \quad \text{for } -0.5 \leq x \leq 1.5$$

- Sketch the two densities in the same graph.
- Identify the classification rule for the case $p_1 = p_2$ and $c(1|2) = c(2|1)$.
- Identify the classification rule for the case $p_1 = 0.2$ and $c(1|2) = c(2|1)$.

Question 5: Use the data in the EXCEL file tab “q5”. Assume the data is a sample from two multivariate normal distributions.

- Identify the classification rule for the case $p_1 = p_2$ and $c(1|2) = c(2|1)$.
- Identify the classification rule for the case $p_1 = 0.25$ and $c(1|2)$ is half of $c(2|1)$.
- Identify the Bayesian rule using $p_1 = 0.6$.
- Based on the rules given in part (a), which population will a new data point $(50, 48, 47, 49)$ be classified into? How about using the rules in part (b)? Using the rule in part (c)?