

MATH 501: REAL ANALYSIS

RUIZHE QIAN

CONTENTS

1. August 25, 2026	1
1.1. Measures	1
1.2. Algebras and σ -algebras	3
1.3. Borel σ -algebras	5
2. August 27, 2026	6
2.1. Product σ -algebras	6
2.2. Products of Borel σ -algebras	8
2.3. Measures	9
2.4. Examples of measures	10
2.5. Basic properties of measures	11

1. AUGUST 25, 2026

Real analysis will be developed through measure theory, Lebesgue integration, and the study of function spaces such as L^p spaces.

1.1. Measures. The goal of a measure is to assign a length, area, or volume to subsets of $\mathbb{R}$, $\mathbb{R}^2$, or, more generally, $\mathbb{R}^n$. Ideally, we would like a function

$$m: 2^{\mathbb{R}^n} \longrightarrow [0, \infty]$$

with the following properties.

- (i) *Countable additivity:* if $E_1, E_2, \dots \subseteq \mathbb{R}^n$ are pairwise disjoint, then

$$m\left(\bigcup_{j=1}^{\infty} E_j\right) = \sum_{j=1}^{\infty} m(E_j).$$

1

A weaker requirement is finite additivity, in which the identity is required only for finite pairwise disjoint families.

(ii) *Invariance under rigid motions:* if

$$\tilde{E} = AE + b, \quad A \in O(n), \quad b \in \mathbb{R}^n,$$

then $m(\tilde{E}) = m(E)$. Here A represents a rotation or reflection, and b a translation.

(iii) *Normalization:*

$$m([0, 1]^n) = 1.$$

Unfortunately, a measure with all these properties cannot be defined on every subset of $\mathbb{R}^n$.

Theorem 1 (Vitali). *Assuming the axiom of choice, there is no countably additive, rigid-motion-invariant map*

$$m: 2^{\mathbb{R}^n} \longrightarrow [0, \infty]$$

satisfying $m([0, 1]^n) = 1$.

Proof. It is enough to consider $n = 1$: if such a map existed on $\mathbb{R}^n$, then

$$\mu(E) = m(E \times [0, 1]^{n-1}), \quad E \subseteq \mathbb{R},$$

would have the required properties on $\mathbb{R}$. We therefore write m for this one-dimensional map.

First, $m(\{x\}) = 0$ for every $x \in \mathbb{R}$. Otherwise, translation invariance and countable additivity applied to a countably infinite set in $[0, 1]$ would contradict $m([0, 1]) = 1$. Consequently, $m([0, 1)) = 1$.

Define an equivalence relation on $[0, 1)$ by

$$x \sim y \iff x - y \in \mathbb{Q}.$$

Using the axiom of choice, select one representative from each equivalence class and let V be the set of representatives. Thus, for every $x \in [0, 1)$, there is a unique $y \in V$ such that $x - y \in \mathbb{Q}$.

For $t \in [0, 1)$, define the translate of V modulo 1 by

$$\begin{aligned} V_t &= V + t \pmod{1} \\ &= \{x + t : x \in V, x + t < 1\} \\ &\quad \cup \{x + t - 1 : x \in V, x + t \geq 1\}. \end{aligned}$$

Finite additivity and translation invariance give

$$m(V_t) = m(V).$$

The family indexed by $t \in [0, 1) \cap \mathbb{Q}$ is a partition of $[0, 1)$. Indeed, if $x \in [0, 1)$ and $y \in V$ represents the equivalence class of x , then either $t = x - y$ or $t = x - y + 1$ lies in $[0, 1) \cap \mathbb{Q}$ and $x \in V_t$. On the other hand, if $V_t \cap V_{t'} \neq \emptyset$, the uniqueness of representatives implies $t = t'$.

Countable additivity now yields

$$1 = m([0, 1)) = \sum_{t \in [0, 1) \cap \mathbb{Q}} m(V_t) = \sum_{t \in [0, 1) \cap \mathbb{Q}} m(V).$$

If $m(V) = 0$, the last sum is 0; if $m(V) > 0$, it is infinite. Either possibility is a contradiction. $\square$

One might try to rescue the construction by requiring only finite additivity together with invariance and normalization. In dimensions at least three, even this is impossible.

Theorem 2 (Banach–Tarski). *If $n \geq 3$, there is no finitely additive, rigid-motion-invariant map*

$$m: 2^{\mathbb{R}^n} \longrightarrow [0, \infty]$$

satisfying $m([0, 1]^n) = 1$.

The resolution is to give up the requirement that a measure be defined on *all* subsets of $\mathbb{R}^n$. In fact, measure theory can be developed on a general underlying set X once an appropriate collection of measurable subsets has been chosen.

1.2. Algebras and σ -algebras.

Definition 1 (Algebra). *An algebra on a set X is a nonempty collection $\mathcal{A} \subseteq 2^X$ such that*

- (a) if $E_1, \dots, E_N \in \mathcal{A}$, then $\bigcup_{j=1}^N E_j \in \mathcal{A}$;
- (b) if $E \in \mathcal{A}$, then $E^c = X \setminus E \in \mathcal{A}$.

Definition 2 (σ -algebra). A σ -algebra on a set X is a nonempty collection $\mathcal{A} \subseteq 2^X$ such that

- (a) if $E_n \in \mathcal{A}$ for every $n \in \mathbb{N}$, then $\bigcup_{n=1}^{\infty} E_n \in \mathcal{A}$;
- (b) if $E \in \mathcal{A}$, then $E^c \in \mathcal{A}$.

Because the collections in these definitions are nonempty and closed under complements, they contain both X and $\emptyset$. By De Morgan's laws, an algebra is closed under finite intersections and a σ -algebra is closed under countable intersections. For example,

$$\bigcap_{n=1}^{\infty} E_n = \left(\bigcup_{n=1}^{\infty} E_n^c \right)^c.$$

When an algebra is already known to be closed under finite set operations, it is enough to check closure under countable *disjoint* unions in order to prove that it is a σ -algebra. Given $E_n \in \mathcal{A}$, set

$$\tilde{E}_1 = E_1, \quad \tilde{E}_n = E_n \setminus \bigcup_{j=1}^{n-1} E_j \quad (n \geq 2).$$

Then the sets $\tilde{E}_n$ are pairwise disjoint and

$$\bigcup_{n=1}^{\infty} \tilde{E}_n = \bigcup_{n=1}^{\infty} E_n.$$

Example 1. The collection $\{\emptyset, X\}$ is the smallest σ -algebra on X .

Example 2. The power set 2^X is the largest σ -algebra on X .

Example 3 (Countable-cocountable σ -algebra). The collection

$$\mathcal{A} = \{E \subseteq X : E \text{ is countable or } E^c \text{ is countable}\}$$

is a σ -algebra on X .

A convenient way to specify a σ -algebra is to give a family of generators.

Definition 3 (Generated σ -algebra). *Let $\mathcal{E} \subseteq 2^X$. The σ -algebra generated by $\mathcal{E}$, denoted $\mathcal{M}(\mathcal{E})$, is the smallest σ -algebra containing $\mathcal{E}$.*

This object exists because 2^X is a σ -algebra containing $\mathcal{E}$, and an arbitrary intersection of σ -algebras is again a σ -algebra. Explicitly,

$$\mathcal{M}(\mathcal{E}) = \bigcap \{ \mathcal{A} : \mathcal{A} \text{ is a } \sigma\text{-algebra on } X \text{ and } \mathcal{E} \subseteq \mathcal{A} \}.$$

1.3. Borel σ -algebras.

Definition 4 (Borel σ -algebra). *Let X be a metric space, or more generally a topological space. The Borel σ -algebra $\mathcal{B}_X$ is the σ -algebra generated by the open subsets of X .*

For example, every open subset of $\mathbb{R}$ is a union of open intervals. The Borel σ -algebra contains every open and closed set, every G_δ set (a countable intersection of open sets), every F_σ set (a countable union of closed sets), and all sets obtained by continuing such countable operations. These classes form the Borel hierarchy.

Proposition 1. *The Borel σ -algebra $\mathcal{B}_{\mathbb{R}}$ is generated by any one of the following collections:*

- (a) $\{(a, b) : a < b\};$
- (b) $\{[a, b] : a < b\};$
- (c) $\{(-\infty, a] : a \in \mathbb{R}\}.$

Proof. Let the three collections above be $\mathcal{E}_1$, $\mathcal{E}_2$, and $\mathcal{E}_3$, respectively.

Every member of $\mathcal{E}_1$ is open, so $\mathcal{M}(\mathcal{E}_1) \subseteq \mathcal{B}_{\mathbb{R}}$. For the reverse inclusion, let $U \subseteq \mathbb{R}$ be open. For every $x \in U$, there is an $\varepsilon > 0$ such that

$$(x - \varepsilon, x + \varepsilon) \subseteq U.$$

By the density of $\mathbb{Q}$ in $\mathbb{R}$, we can choose $p, q \in \mathbb{Q}$ such that

$$x - \varepsilon < p < x < q < x + \varepsilon.$$

Thus every $x \in U$ lies in an interval (p, q) with rational endpoints that is contained in U . It follows that

$$U = \bigcup \{ (p, q) : p, q \in \mathbb{Q}, p < q, (p, q) \subseteq U \}.$$

The indexing collection is a subset of $\mathbb{Q}^2$, so it is countable. Therefore every open subset of $\mathbb{R}$ is a countable union of open intervals. In particular, every open set belongs to $\mathcal{M}(\mathcal{E}_1)$, proving

$$\mathcal{M}(\mathcal{E}_1) = \mathcal{B}_{\mathbb{R}}.$$

Every closed interval is Borel. In the other direction, for $a < b$,

$$(a, b) = \bigcup_{\substack{n \in \mathbb{N} \\ 2/n < b-a}} \left[a + \frac{1}{n}, b - \frac{1}{n} \right].$$

Thus $\mathcal{E}_1 \subseteq \mathcal{M}(\mathcal{E}_2)$, and consequently $\mathcal{M}(\mathcal{E}_2) = \mathcal{B}_{\mathbb{R}}$. Equivalently, the reverse containment can also be seen from

$$[a, b] = \bigcap_{n=1}^{\infty} \left(a - \frac{1}{n}, b + \frac{1}{n} \right).$$

Finally, every left half-line in $\mathcal{E}_3$ is closed and hence Borel. Conversely,

$$(-\infty, b) = \bigcup_{n=1}^{\infty} \left(-\infty, b - \frac{1}{n} \right]$$

and therefore

$$(a, b) = (-\infty, b) \cap (-\infty, a]^c \in \mathcal{M}(\mathcal{E}_3).$$

It follows that $\mathcal{E}_1 \subseteq \mathcal{M}(\mathcal{E}_3)$, so $\mathcal{M}(\mathcal{E}_3) = \mathcal{B}_{\mathbb{R}}$ as well. $\square$

2. AUGUST 27, 2026

We continue our discussion of σ -algebras. Recall that a σ -algebra $\mathcal{M}$ on a set X contains $\emptyset$ and X , is closed under complements, and is closed under countable unions. If (X, d) is a metric space, then its Borel σ -algebra $\mathcal{B}_X$ is generated by the open subsets of X .

2.1. Product σ -algebras. Let I be an index set, let X_i be a nonempty set for each $i \in I$, and set

$$X = \prod_{i \in I} X_i.$$

For each $i \in I$, let $\mathcal{M}_i$ be a σ -algebra on X_i , and let

$$\pi_i: X \longrightarrow X_i, \quad \pi_i((x_j)_{j \in I}) = x_i,$$

be the coordinate projection.

Definition 5 (Product σ -algebra). *The product σ -algebra on X is*

$$\bigotimes_{i \in I} \mathcal{M}_i = \mathcal{M}(\{\pi_i^{-1}(E) : i \in I, E \in \mathcal{M}_i\}).$$

Thus it is the smallest σ -algebra that makes every coordinate projection measurable.

The generating sets $\pi_i^{-1}(E)$ are called *cylinder sets*. They restrict the i th coordinate to E and leave all other coordinates unrestricted.

Lemma 1. *If I is countable, then $\bigotimes_{i \in I} \mathcal{M}_i$ is generated by the measurable rectangles*

$$\left\{ \prod_{i \in I} E_i : E_i \in \mathcal{M}_i \right\}.$$

Proof. Since I is countable, every measurable rectangle satisfies

$$\prod_{i \in I} E_i = \bigcap_{i \in I} \pi_i^{-1}(E_i) \in \bigotimes_{i \in I} \mathcal{M}_i.$$

Conversely, each cylinder set is a measurable rectangle. Indeed,

$$\pi_j^{-1}(E_j) = \prod_{i \in I} F_i, \quad F_i = \begin{cases} E_j, & i = j, \\ X_i, & i \neq j. \end{cases}$$

The two collections therefore generate the same σ -algebra. $\square$

The generators of the coordinate σ -algebras can be used directly to generate the product.

Proposition 2. *Suppose*

$$\mathcal{M}_i = \mathcal{M}(\mathcal{G}_i), \quad \mathcal{G}_i \subseteq 2^{X_i},$$

for every $i \in I$.

(a) *The product σ -algebra is generated by*

$$\mathcal{F} = \{\pi_i^{-1}(G) : i \in I, G \in \mathcal{G}_i\}.$$

(b) If I is countable and $\mathcal{G}_i^* = \mathcal{G}_i \cup \{X_i\}$, then the product σ -algebra is generated by

$$\left\{ \prod_{i \in I} G_i : G_i \in \mathcal{G}_i^* \right\}.$$

Proof. For part (a), every member of $\mathcal{F}$ belongs to $\bigotimes_{i \in I} \mathcal{M}_i$, so

$$\mathcal{M}(\mathcal{F}) \subseteq \bigotimes_{i \in I} \mathcal{M}_i.$$

For the reverse inclusion, fix $i \in I$ and consider

$$\mathcal{A}_i = \{E \subseteq X_i : \pi_i^{-1}(E) \in \mathcal{M}(\mathcal{F})\}.$$

Preimages preserve complements and countable unions, so $\mathcal{A}_i$ is a σ -algebra. It contains $\mathcal{G}_i$ and hence contains $\mathcal{M}(\mathcal{G}_i) = \mathcal{M}_i$. Therefore every cylinder $\pi_i^{-1}(E)$ with $E \in \mathcal{M}_i$ belongs to $\mathcal{M}(\mathcal{F})$, which proves the reverse inclusion.

For part (b), each cylinder from part (a) is a rectangle whose other coordinates are X_j . Conversely, since I is countable, every rectangle in the displayed collection is the countable intersection of the sets $\pi_i^{-1}(G_i)$. When $G_i = X_i$ this set is all of X ; otherwise it is a cylinder from part (a). The conclusion follows. $\square$

2.2. Products of Borel σ -algebras. Let (X_i, d_i) , $i = 1, \dots, n$, be metric spaces. On

$$X = \prod_{i=1}^n X_i$$

define the maximum metric

$$d(x, y) = \max_{1 \leq i \leq n} d_i(x_i, y_i).$$

Proposition 3. *With the notation above,*

$$\bigotimes_{i=1}^n \mathcal{B}_{X_i} \subseteq \mathcal{B}_X.$$

If every (X_i, d_i) is separable, then equality holds:

$$\bigotimes_{i=1}^n \mathcal{B}_{X_i} = \mathcal{B}_X.$$

Proof. Each coordinate projection $\pi_i: X \rightarrow X_i$ is continuous. If $U_i \subseteq X_i$ is open, then $\pi_i^{-1}(U_i)$ is open in X . Since $\mathcal{B}_{X_i}$ is generated by the open subsets of X_i , Proposition 2 gives

$$\bigotimes_{i=1}^n \mathcal{B}_{X_i} \subseteq \mathcal{B}_X.$$

Now suppose that each X_i is separable, and choose a countable dense subset $D_i \subseteq X_i$. The collection

$$\mathcal{G}_i = \{B_{d_i}(p, q) : p \in D_i, q \in \mathbb{Q}, q > 0\}$$

is a countable basis for X_i and hence generates $\mathcal{B}_{X_i}$. The finite product $D = \prod_{i=1}^n D_i$ is countable and dense in X . Moreover,

$$B_d(x, q) = \prod_{i=1}^n B_{d_i}(x_i, q).$$

Thus every ball in X whose center lies in D and whose radius is a positive rational belongs to $\bigotimes_{i=1}^n \mathcal{B}_{X_i}$. These balls form a countable basis for X , so every open subset of X is a countable union of such balls. Hence

$$\mathcal{B}_X \subseteq \bigotimes_{i=1}^n \mathcal{B}_{X_i},$$

which completes the proof. $\square$

Corollary 1. For every $n \in \mathbb{N}$,

$$\mathcal{B}_{\mathbb{R}^n} = \bigotimes_{i=1}^n \mathcal{B}_{\mathbb{R}}.$$

For example, $\mathcal{B}_{\mathbb{R}^n}$ is generated by each of the following collections:

- open rectangles $\prod_{i=1}^n (a_i, b_i)$;
- closed rectangles $\prod_{i=1}^n [a_i, b_i]$; and
- lower orthants $\prod_{i=1}^n (-\infty, a_i]$.

2.3. Measures.

Definition 6 (Measure). *Let $\mathcal{M}$ be a σ -algebra on a set X . A function*

$$\mu: \mathcal{M} \longrightarrow [0, \infty]$$

is a measure if

- (i) $\mu(\emptyset) = 0$;
- (ii) *whenever $E_1, E_2, \dots \in \mathcal{M}$ are pairwise disjoint,*

$$\mu \left(\bigcup_{i=1}^{\infty} E_i \right) = \sum_{i=1}^{\infty} \mu(E_i).$$

The second property is called countable additivity. If it is required only for finite pairwise disjoint families, then μ is called finitely additive.

The sets in $\mathcal{M}$ are called *measurable sets*, and the triple $(X, \mathcal{M}, \mu)$ is called a *measure space*. A measure μ is

- *finite* if $\mu(X) < \infty$;
- *σ -finite* if $X = \bigcup_{i=1}^{\infty} E_i$ for some $E_i \in \mathcal{M}$ with $\mu(E_i) < \infty$ for every i .

A set $N \in \mathcal{M}$ with $\mu(N) = 0$ is called a *null set*. A statement is said to hold *almost everywhere*, abbreviated a.e., if it holds outside some null set.

2.4. Examples of measures.

Example 4 (Weighted counting measure). *Let X be nonempty, let $\mathcal{M} = 2^X$, and let $f: X \rightarrow [0, \infty]$. Define*

$$\mu(E) = \sum_{x \in E} f(x) := \sup \left\{ \sum_{x \in F} f(x) : F \subseteq E, F \text{ finite} \right\}.$$

Then μ is a measure on 2^X .

If $f(x) = 1$ for every $x \in X$, then μ is the counting measure,

$$\mu(E) = \#E.$$

If $x_0 \in X$ and

$$f(x) = \begin{cases} 1, & x = x_0, \\ 0, & x \neq x_0, \end{cases}$$

then

$$\mu(E) = \begin{cases} 1, & x_0 \in E, \\ 0, & x_0 \notin E. \end{cases}$$

This is the point mass, or Dirac measure, at x_0 .

Example 5 (Countable-cocountable measure). Let X be uncountable, and let $\mathcal{M}$ be the countable-cocountable σ -algebra. Then

$$\mu(E) = \begin{cases} 0, & E \text{ is countable,} \\ 1, & E^c \text{ is countable} \end{cases}$$

defines a measure on $(X, \mathcal{M})$.

Example 6 (Finite additivity is not enough). Let $X = \mathbb{N}$ and $\mathcal{M} = 2^{\mathbb{N}}$. Define

$$\nu(E) = \begin{cases} 0, & E \text{ is finite,} \\ \infty, & E \text{ is infinite.} \end{cases}$$

Then ν is finitely additive. It is not a measure, because

$$\nu\left(\bigcup_{n=1}^{\infty} \{n\}\right) = \infty \quad \text{but} \quad \sum_{n=1}^{\infty} \nu(\{n\}) = 0.$$

2.5. Basic properties of measures.

Theorem 3. Let $(X, \mathcal{M}, \mu)$ be a measure space.

(a) Monotonicity: if $E, F \in \mathcal{M}$ and $E \subseteq F$, then

$$\mu(E) \leq \mu(F).$$

(b) Countable subadditivity: if $E_1, E_2, \dots \in \mathcal{M}$, then

$$\mu\left(\bigcup_{i=1}^{\infty} E_i\right) \leq \sum_{i=1}^{\infty} \mu(E_i).$$

(c) Continuity from below: if

$$E_1 \subseteq E_2 \subseteq \dots, \quad E_i \in \mathcal{M},$$

then

$$\mu\left(\bigcup_{i=1}^{\infty} E_i\right) = \lim_{i \rightarrow \infty} \mu(E_i).$$

(d) Continuity from above: if

$$E_1 \supseteq E_2 \supseteq \cdots, \quad E_i \in \mathcal{M}, \quad \mu(E_1) < \infty,$$

then

$$\mu\left(\bigcap_{i=1}^{\infty} E_i\right) = \lim_{i \rightarrow \infty} \mu(E_i).$$

Proof. For part (a), write F as the disjoint union

$$F = E \cup (F \setminus E).$$

Countable additivity gives

$$\mu(F) = \mu(E) + \mu(F \setminus E) \geq \mu(E).$$

For part (b), define

$$F_1 = E_1, \quad F_k = E_k \setminus \bigcup_{j=1}^{k-1} E_j \quad (k \geq 2).$$

The sets F_k are pairwise disjoint, $F_k \subseteq E_k$, and

$$\bigcup_{k=1}^{\infty} F_k = \bigcup_{k=1}^{\infty} E_k.$$

Therefore countable additivity and part (a) imply

$$\mu\left(\bigcup_{k=1}^{\infty} E_k\right) = \sum_{k=1}^{\infty} \mu(F_k) \leq \sum_{k=1}^{\infty} \mu(E_k).$$

For part (c), set

$$F_1 = E_1, \quad F_k = E_k \setminus E_{k-1} \quad (k \geq 2).$$

The sets F_k are pairwise disjoint, and for each N ,

$$E_N = \bigcup_{k=1}^N F_k, \quad \bigcup_{i=1}^{\infty} E_i = \bigcup_{k=1}^{\infty} F_k.$$

It follows that

$$\mu\left(\bigcup_{i=1}^{\infty} E_i\right) = \sum_{k=1}^{\infty} \mu(F_k)$$

$$\begin{aligned}
&= \lim_{N \rightarrow \infty} \sum_{k=1}^N \mu(F_k) \\
&= \lim_{N \rightarrow \infty} \mu(E_N).
\end{aligned}$$

For part (d), define

$$F_i = E_1 \setminus E_i.$$

Since E_i decreases, F_i increases, and

$$\bigcup_{i=1}^{\infty} F_i = E_1 \setminus \bigcap_{i=1}^{\infty} E_i.$$

By part (c),

$$\mu \left(E_1 \setminus \bigcap_{i=1}^{\infty} E_i \right) = \lim_{i \rightarrow \infty} \mu(E_1 \setminus E_i).$$

Because $E_i \subseteq E_1$ and $\mu(E_1) < \infty$, finite additivity gives

$$\mu(E_1 \setminus E_i) = \mu(E_1) - \mu(E_i)$$

and

$$\mu \left(E_1 \setminus \bigcap_{i=1}^{\infty} E_i \right) = \mu(E_1) - \mu \left(\bigcap_{i=1}^{\infty} E_i \right).$$

Substituting these identities into the preceding limit and cancelling the finite quantity $\mu(E_1)$ yields

$$\mu \left(\bigcap_{i=1}^{\infty} E_i \right) = \lim_{i \rightarrow \infty} \mu(E_i).$$

□

Remark 1 (Necessity of the finite-measure hypothesis). The assumption $\mu(E_1) < \infty$ in Theorem 3(d) cannot be omitted. Let

$$X = \mathbb{N}, \quad \mathcal{M} = 2^{\mathbb{N}},$$

let μ be counting measure, and define

$$E_i = \{i, i+1, i+2, \dots\}.$$

Then

$$E_1 \supseteq E_2 \supseteq E_3 \supseteq \cdots, \quad \bigcap_{i=1}^{\infty} E_i = \emptyset.$$

Indeed, for any $n \in \mathbb{N}$, we have $n \notin E_i$ whenever $i > n$. Consequently,

$$\mu \left(\bigcap_{i=1}^{\infty} E_i \right) = \mu(\emptyset) = 0.$$

On the other hand, every E_i is infinite, so

$$\mu(E_i) = \infty \quad \text{for every } i, \quad \lim_{i \rightarrow \infty} \mu(E_i) = \infty.$$

Thus

$$\mu \left(\bigcap_{i=1}^{\infty} E_i \right) = 0 \neq \infty = \lim_{i \rightarrow \infty} \mu(E_i).$$

This does not contradict part (d), because $\mu(E_1) = \mu(\mathbb{N}) = \infty$.