

MATH 551: NUMERICAL ORDINARY DIFFERENTIAL EQUATIONS

RUIZHE QIAN

CONTENTS

1. August 24, 2026	1
1.1. Ordinary differential equations	1
1.2. Examples	2
1.3. Solutions and the Lipschitz condition	3
2. August 26, 2026	4
2.1. Stability of solutions	4
2.2. The solution map	6
2.3. One-step methods	8

1. AUGUST 24, 2026

1.1. **Ordinary differential equations.** An ODE has the form

$$x'(t) = f(t, x(t)).$$

Given an initial condition $x(t_0) = x_0$, the corresponding *initial-value problem* (IVP) is to find $x(t)$ for $t \geq t_0$ such that

$$(1) \quad x'(t) = f(t, x(t)), \quad x(t_0) = x_0.$$

Topics arising in the numerical solution of differential equations include

- non-stiff problems and stiff solvers;
- geometric solvers;
- parameter estimation and Bayesian approaches, where the vector field may depend on a model parameter θ , as in $f(t, x, \theta)$;
- boundary-value problems; and

- stochastic differential equations (SDEs).

We generally consider vector-valued solutions and vector fields,

$$x: \mathbb{R} \longrightarrow \mathbb{R}^n, \quad f: \mathbb{R} \times \mathbb{R}^n \longrightarrow \mathbb{R}^n,$$

which can be written componentwise as

$$x = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}, \quad f(t, x) = \begin{bmatrix} f_1(t, x_1, \dots, x_n) \\ \vdots \\ f_n(t, x_1, \dots, x_n) \end{bmatrix}.$$

1.2. Examples.

Example 1. *The scalar equation*

$$x' = x$$

is an ODE with $n = 1$.

Example 2 (Damped oscillator). *Consider the second-order equation*

$$x'' = -\omega^2 x - x'.$$

Setting $x_1 = x$ and $x_2 = x'$ converts it into the first-order system

$$\begin{cases} x_1' = x_2, \\ x_2' = -\omega^2 x_1 - x_2. \end{cases}$$

Thus, with $\mathbf{x} = (x_1, x_2)^\top$, we have $\mathbf{x}' = \mathbf{f}(x_1, x_2)$.

Example 3 (Chemical kinetics). *A three-species kinetic model is*

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}, \quad \begin{cases} x_1' = -k_1 x_1 + k_2 x_2 x_3, \\ x_2' = k_1 x_1 - k_2 x_2 x_3 - k_3 x_2^2, \\ x_3' = k_3 x_2^2, \end{cases}$$

where $k_1, k_2, k_3 > 0$ are reaction rates. The reaction rates may occur on very different scales, which can lead to stiffness.

Example 4 (Molecular model). *Let q_i be the position of particle i , let m_i be its mass, and let $V(q_1, \dots, q_N)$ be the potential energy. Newton's*


equation is

$$m_i q_i'' = -\frac{\partial V(q_1, \dots, q_N)}{\partial q_i}.$$

Introducing the momentum $p_i = m_i q_i'$ gives

$$\begin{cases} q_i' = \frac{p_i}{m_i}, \\ p_i' = -\frac{\partial V}{\partial q_i}. \end{cases}$$

The positions and momenta can therefore be collected into the state vector

$$\mathbf{x} = \begin{bmatrix} q_1 \\ p_1 \\ q_2 \\ p_2 \\ \vdots \\ q_N \\ p_N \end{bmatrix}.$$


The diagram shows a vector labeled q_i originating from a point labeled i and pointing to a point labeled q_i . There are several other points scattered around, representing a discrete set of points in a space.

1.3. Solutions and the Lipschitz condition. Consider the IVP (1) on an interval containing t_0 .

Definition 1. A function $x(t)$ is a solution of the IVP if it satisfies the integral equation

$$x(t) = x_0 + \int_{t_0}^t f(s, x(s)) ds.$$

Definition 2 (Lipschitz condition). The vector field f is Lipschitz continuous in x on a domain D if there is a constant $L \geq 0$ such that

$$\|f(t, x) - f(t, y)\| \leq L\|x - y\|$$

for all $x, y \in D$ and all relevant t .


Under the usual continuity assumption in t , the Lipschitz condition in x implies that the solution of the IVP exists and is unique on an appropriate interval.

The same condition also controls the sensitivity to the initial data. Let $x(t)$ and $y(t)$ be solutions with initial values x_0 and y_0 , respectively. Their integral equations give

$$\begin{aligned}\|x(t) - y(t)\| &\leq \|x_0 - y_0\| + \int_{t_0}^t \|f(s, x(s)) - f(s, y(s))\| ds \\ &\leq \|x_0 - y_0\| + L \int_{t_0}^t \|x(s) - y(s)\| ds.\end{aligned}$$

By Grönwall's inequality,

$$(2) \quad \|x(t) - y(t)\| \leq \|x_0 - y_0\| e^{L(t-t_0)}.$$



Thus nearby initial values produce solutions whose separation is bounded by (2).

2. AUGUST 26, 2026

The main topics are properties of solutions, time discretization, and consistency error. We continue to consider the IVP

$$(3) \quad x'(t) = f(t, x(t)), \quad x(t_0) = x_0,$$

where $x: \mathbb{R} \rightarrow \mathbb{R}^n$.

2.1. Stability of solutions. The integral formulation of (3) is

$$x(t) = x_0 + \int_{t_0}^t f(s, x(s)) ds.$$

If $y(t)$ is another solution with initial value y_0 , then

$$y(t) = y_0 + \int_{t_0}^t f(s, y(s)) ds,$$

and hence

$$y(t) - x(t) = y_0 - x_0 + \int_{t_0}^t (f(s, y(s)) - f(s, x(s))) ds.$$

Suppose f is Lipschitz continuous in x with Lipschitz constant L , so that

$$\|f(t, y) - f(t, x)\| \leq L\|y - x\|.$$

Taking norms gives

$$(4) \quad \|y(t) - x(t)\| \leq \|y_0 - x_0\| + \int_{t_0}^t L\|y(s) - x(s)\| ds.$$

The following inequality controls this accumulation of error.

Lemma 1 (Grönwall's inequality). *Let $u, v \in C([t_0, t_1], \mathbb{R})$ be nonnegative functions and let $\rho \geq 0$. If*

$$(5) \quad u(t) \leq \rho + \int_{t_0}^t v(s)u(s) ds, \quad t \in [t_0, t_1],$$

then

$$(6) \quad u(t) \leq \rho \exp \left(\int_{t_0}^t v(s) ds \right), \quad t \in [t_0, t_1].$$

In particular, $u \equiv 0$ when $\rho = 0$.

Proof. First suppose that $\rho > 0$ and define

$$U(t) = \rho + \int_{t_0}^t v(s)u(s) ds.$$

Then $U \in C^1([t_0, t_1], \mathbb{R})$, and

$$u(t) \leq U(t), \quad U(t) \geq \rho > 0.$$

Since u and v are nonnegative,

$$U'(t) = v(t)u(t) \leq v(t)U(t).$$

Thus the logarithmic derivative of U satisfies

$$(\log U(t))' = \frac{U'(t)}{U(t)} \leq v(t).$$

Integrating from t_0 to t yields

$$\log U(t) - \log U(t_0) \leq \int_{t_0}^t v(s) ds.$$

Because $U(t_0) = \rho$ and $u(t) \leq U(t)$, applying the exponential function proves (6) when $\rho > 0$.

Now suppose that $\rho = 0$. For any $\varepsilon > 0$, (5) implies

$$u(t) \leq \varepsilon + \int_{t_0}^t v(s)u(s) ds.$$

The first part of the proof, with $\rho = \varepsilon$, gives

$$0 \leq u(t) \leq \varepsilon \exp \left(\int_{t_0}^t v(s) ds \right).$$

Letting $\varepsilon \downarrow 0$ shows that $u(t) = 0$ for every $t \in [t_0, t_1]$. $\square$

Apply Lemma 1 to (4) with

$$u(t) = \|y(t) - x(t)\|, \quad v(t) = L, \quad \rho = \|y_0 - x_0\|.$$

We obtain

$$(7) \quad \|y(t) - x(t)\| \leq \|y_0 - x_0\| e^{L(t-t_0)}.$$

This estimate can be pessimistic, but it establishes continuous dependence on the initial data.

2.2. The solution map. The *solution map*, or *flow map*, is denoted by

$$x(t) = \varphi^{t,t_0}(x_0).$$

For example, consider the scalar logistic equation

$$x' = x(1 - x).$$

For $x_0 \neq 0$, its solution is

$$\varphi^{t,t_0}(x_0) = \frac{1}{1 + \frac{1-x_0}{x_0}e^{-(t-t_0)}}.$$

When $x_0 = 0$, the solution is $x(t) \equiv 0$.

The solution map has the identity and composition properties

$$\begin{aligned}\varphi^{t_0,t_0}(x_0) &= x_0, & \varphi^{t_0,t_0} &= I, \\ \varphi^{t_2,t_1}(\varphi^{t_1,t_0}(x_0)) &= \varphi^{t_2,t_0}(x_0).\end{aligned}$$

The second property is the basis of time marching.

Differentiating the solution map with respect to time gives

$$\begin{aligned}\frac{d}{dt}\varphi^{t,t_0}(x_0) &= f(t, x(t)), \\ \frac{d^2}{dt^2}\varphi^{t,t_0}(x_0) &= f_t(t, x(t)) + f_x(t, x(t))x'(t) \\ &= (f_t + f_x f)(t, x(t)),\end{aligned}$$

where f_x is the Jacobian matrix of f with respect to x . Higher time derivatives can be found by repeated differentiation, for example with automatic differentiation; they can also be viewed through the Koopman operator.

To measure sensitivity with respect to the initial value, define the Wronskian matrix

$$W(t) = \frac{\partial}{\partial x_0}\varphi^{t,t_0}(x_0) = \frac{\partial x(t)}{\partial x_0}.$$

Differentiating with respect to t gives

$$\begin{aligned}W'(t) &= \frac{\partial}{\partial x_0}f(t, x(t)) \\ &= f_x(t, x(t))\frac{\partial x(t)}{\partial x_0}.\end{aligned}$$

Thus, if

$$A(t) = f_x(t, x(t)),$$

then W satisfies the *variational equation*

$$(8) \quad W'(t) = A(t)W(t), \quad W(t_0) = I.$$

2.3. One-step methods. Choose a time grid

$$t_0 < t_1 < \cdots < t_N = t, \quad \tau_j = t_{j+1} - t_j,$$

where the step sizes τ_j are small. Repeated use of the composition property gives

$$\varphi^{t_N, t_0} = \varphi^{t_N, t_{N-1}} \circ \cdots \circ \varphi^{t_2, t_1} \circ \varphi^{t_1, t_0}.$$

It is therefore enough to approximate a single short-time solution map. Writing $t_j = t$ and $t_{j+1} = t + \tau$, Taylor expansion gives

$$(9) \quad \varphi^{t+\tau, t}(x) = x + \tau f(t, x) + \frac{\tau^2}{2} (f_t + f_x f)(t, x) + O(\tau^3).$$

Dropping the terms of order τ^2 and higher defines the Euler one-step map

$$\Psi^{t+\tau, t}(x) = x + \tau f(t, x).$$

Consequently, Euler's method is

$$(10) \quad x_{j+1} = x_j + \tau_j f(t_j, x_j).$$

Its one-step consistency error follows from (9):

$$\varphi^{t+\tau, t}(x) - \Psi^{t+\tau, t}(x) = \frac{\tau^2}{2} (f_t + f_x f)(t, x) + O(\tau^3) = O(\tau^2).$$