

MATH 501: REAL ANALYSIS

RUIZHE QIAN

CONTENTS

1. August 25, 2026	2
1.1. Measures	2
1.2. Algebras and σ -algebras	4
1.3. Borel σ -algebras	5
2. August 27, 2026	7
2.1. Product σ -algebras	7
2.2. Products of Borel σ -algebras	9
2.3. Measures	10
2.4. Examples of measures	11
2.5. Basic properties of measures	12
3. September 1, 2026	15
3.1. Complete measure spaces	15
3.2. Outer measures	17
4. September 3, 2026	20
4.1. The Carathéodory construction	20
4.2. Premeasures	22
5. September 8, 2026	25
5.1. Uniqueness of the extension	25
5.2. Lebesgue–Stieltjes measures on $\mathbb{R}$	27
6. September 10, 2026	32
6.1. Regularity of Lebesgue–Stieltjes measures	33
6.2. Lebesgue measure	36
7. September 15, 2026	37
7.1. Measure and Baire category	37
7.2. The Cantor set	38
7.3. The idea of Lebesgue integration	41

7.4. Measurable functions	42
8. September 17, 2026	43
8.1. Operations on measurable functions	44
8.2. Simple functions	46
8.3. Nonnegative integration	49

1. AUGUST 25, 2026

Real analysis will be developed through measure theory, Lebesgue integration, and the study of function spaces such as L^p spaces.

1.1. Measures. The goal of a measure is to assign a length, area, or volume to subsets of $\mathbb{R}$, $\mathbb{R}^2$, or, more generally, $\mathbb{R}^n$. Ideally, we would like a function

$$m: 2^{\mathbb{R}^n} \longrightarrow [0, \infty]$$

with the following properties.

- (i) *Countable additivity:* if $E_1, E_2, \dots \subseteq \mathbb{R}^n$ are pairwise disjoint, then

$$m\left(\bigcup_{j=1}^{\infty} E_j\right) = \sum_{j=1}^{\infty} m(E_j).$$

A weaker requirement is finite additivity, in which the identity is required only for finite pairwise disjoint families.

- (ii) *Invariance under rigid motions:* if

$$\tilde{E} = AE + b, \quad A \in O(n), \quad b \in \mathbb{R}^n,$$

then $m(\tilde{E}) = m(E)$. Here A represents a rotation or reflection, and b a translation.

- (iii) *Normalization:*

$$m([0, 1]^n) = 1.$$

Unfortunately, a measure with all these properties cannot be defined on every subset of $\mathbb{R}^n$.

Theorem 1 (Vitali). *Assuming the axiom of choice, there is no countably additive, rigid-motion-invariant map*

$$m: 2^{\mathbb{R}^n} \longrightarrow [0, \infty]$$

satisfying $m([0, 1]^n) = 1$.

Proof. It is enough to consider $n = 1$: if such a map existed on $\mathbb{R}^n$, then

$$\mu(E) = m(E \times [0, 1]^{n-1}), \quad E \subseteq \mathbb{R},$$

would have the required properties on $\mathbb{R}$. We therefore write m for this one-dimensional map.

First, $m(\{x\}) = 0$ for every $x \in \mathbb{R}$. Otherwise, translation invariance and countable additivity applied to a countably infinite set in $[0, 1]$ would contradict $m([0, 1]) = 1$. Consequently, $m([0, 1)) = 1$.

Define an equivalence relation on $[0, 1)$ by

$$x \sim y \iff x - y \in \mathbb{Q}.$$

Using the axiom of choice, select one representative from each equivalence class and let V be the set of representatives. Thus, for every $x \in [0, 1)$, there is a unique $y \in V$ such that $x - y \in \mathbb{Q}$.

For $t \in [0, 1)$, define the translate of V modulo 1 by

$$\begin{aligned} V_t &= V + t \pmod{1} \\ &= \{x + t : x \in V, x + t < 1\} \\ &\quad \cup \{x + t - 1 : x \in V, x + t \geq 1\}. \end{aligned}$$

Finite additivity and translation invariance give

$$m(V_t) = m(V).$$

The family indexed by $t \in [0, 1) \cap \mathbb{Q}$ is a partition of $[0, 1)$. Indeed, if $x \in [0, 1)$ and $y \in V$ represents the equivalence class of x , then either $t = x - y$ or $t = x - y + 1$ lies in $[0, 1) \cap \mathbb{Q}$ and $x \in V_t$. On the other hand, if $V_t \cap V_{t'} \neq \emptyset$, the uniqueness of representatives implies $t = t'$.

Countable additivity now yields

$$1 = m([0, 1]) = \sum_{t \in [0, 1) \cap \mathbb{Q}} m(V_t) = \sum_{t \in [0, 1) \cap \mathbb{Q}} m(V).$$

If $m(V) = 0$, the last sum is 0; if $m(V) > 0$, it is infinite. Either possibility is a contradiction. $\square$

One might try to rescue the construction by requiring only finite additivity together with invariance and normalization. In dimensions at least three, even this is impossible.

Theorem 2 (Banach–Tarski). *If $n \geq 3$, there is no finitely additive, rigid-motion-invariant map*

$$m: 2^{\mathbb{R}^n} \longrightarrow [0, \infty]$$

satisfying $m([0, 1]^n) = 1$.

The resolution is to give up the requirement that a measure be defined on *all* subsets of $\mathbb{R}^n$. In fact, measure theory can be developed on a general underlying set X once an appropriate collection of measurable subsets has been chosen.

1.2. Algebras and σ -algebras.

Definition 1 (Algebra). *An algebra on a set X is a nonempty collection $\mathcal{A} \subseteq 2^X$ such that*

- (a) *if $E_1, \dots, E_N \in \mathcal{A}$, then $\bigcup_{j=1}^N E_j \in \mathcal{A}$;*
- (b) *if $E \in \mathcal{A}$, then $E^c = X \setminus E \in \mathcal{A}$.*

Definition 2 (σ -algebra). *A σ -algebra on a set X is a nonempty collection $\mathcal{A} \subseteq 2^X$ such that*

- (a) *if $E_n \in \mathcal{A}$ for every $n \in \mathbb{N}$, then $\bigcup_{n=1}^{\infty} E_n \in \mathcal{A}$;*
- (b) *if $E \in \mathcal{A}$, then $E^c \in \mathcal{A}$.*

Because the collections in these definitions are nonempty and closed under complements, they contain both X and $\emptyset$. By De Morgan's laws, an algebra is closed under finite intersections and a σ -algebra is

closed under countable intersections. For example,

$$\bigcap_{n=1}^{\infty} E_n = \left(\bigcup_{n=1}^{\infty} E_n^c \right)^c.$$

When an algebra is already known to be closed under finite set operations, it is enough to check closure under countable *disjoint* unions in order to prove that it is a σ -algebra. Given $E_n \in \mathcal{A}$, set

$$\tilde{E}_1 = E_1, \quad \tilde{E}_n = E_n \setminus \bigcup_{j=1}^{n-1} E_j \quad (n \geq 2).$$

Then the sets $\tilde{E}_n$ are pairwise disjoint and

$$\bigcup_{n=1}^{\infty} \tilde{E}_n = \bigcup_{n=1}^{\infty} E_n.$$

Example 1. The collection $\{\emptyset, X\}$ is the smallest σ -algebra on X .

Example 2. The power set 2^X is the largest σ -algebra on X .

Example 3 (Countable-cocountable σ -algebra). The collection

$$\mathcal{A} = \{E \subseteq X : E \text{ is countable or } E^c \text{ is countable}\}$$

is a σ -algebra on X .

A convenient way to specify a σ -algebra is to give a family of generators.

Definition 3 (Generated σ -algebra). Let $\mathcal{E} \subseteq 2^X$. The σ -algebra generated by $\mathcal{E}$, denoted $\mathcal{M}(\mathcal{E})$, is the smallest σ -algebra containing $\mathcal{E}$.

This object exists because 2^X is a σ -algebra containing $\mathcal{E}$, and an arbitrary intersection of σ -algebras is again a σ -algebra. Explicitly,

$$\mathcal{M}(\mathcal{E}) = \bigcap \{ \mathcal{A} : \mathcal{A} \text{ is a } \sigma\text{-algebra on } X \text{ and } \mathcal{E} \subseteq \mathcal{A} \}.$$

1.3. Borel σ -algebras.

Definition 4 (Borel σ -algebra). Let X be a metric space, or more generally a topological space. The Borel σ -algebra $\mathcal{B}_X$ is the σ -algebra generated by the open subsets of X .

For example, every open subset of $\mathbb{R}$ is a union of open intervals. The Borel σ -algebra contains every open and closed set, every G_δ set (a countable intersection of open sets), every F_σ set (a countable union of closed sets), and all sets obtained by continuing such countable operations. These classes form the Borel hierarchy.

Proposition 1. *The Borel σ -algebra $\mathcal{B}_\mathbb{R}$ is generated by any one of the following collections:*

- (a) $\{(a, b) : a < b\};$
- (b) $\{[a, b] : a < b\};$
- (c) $\{(-\infty, a] : a \in \mathbb{R}\}.$

Proof. Let the three collections above be $\mathcal{E}_1$, $\mathcal{E}_2$, and $\mathcal{E}_3$, respectively.

Every member of $\mathcal{E}_1$ is open, so $\mathcal{M}(\mathcal{E}_1) \subseteq \mathcal{B}_\mathbb{R}$. For the reverse inclusion, let $U \subseteq \mathbb{R}$ be open. For every $x \in U$, there is an $\varepsilon > 0$ such that

$$(x - \varepsilon, x + \varepsilon) \subseteq U.$$

By the density of $\mathbb{Q}$ in $\mathbb{R}$, we can choose $p, q \in \mathbb{Q}$ such that

$$x - \varepsilon < p < x < q < x + \varepsilon.$$

Thus every $x \in U$ lies in an interval (p, q) with rational endpoints that is contained in U . It follows that

$$U = \bigcup \{(p, q) : p, q \in \mathbb{Q}, p < q, (p, q) \subseteq U\}.$$

The indexing collection is a subset of $\mathbb{Q}^2$, so it is countable. Therefore every open subset of $\mathbb{R}$ is a countable union of open intervals. In particular, every open set belongs to $\mathcal{M}(\mathcal{E}_1)$, proving

$$\mathcal{M}(\mathcal{E}_1) = \mathcal{B}_\mathbb{R}.$$

Every closed interval is Borel. In the other direction, for $a < b$,

$$(a, b) = \bigcup_{\substack{n \in \mathbb{N} \\ 2/n < b-a}} \left[a + \frac{1}{n}, b - \frac{1}{n} \right].$$

Thus $\mathcal{E}_1 \subseteq \mathcal{M}(\mathcal{E}_2)$, and consequently $\mathcal{M}(\mathcal{E}_2) = \mathcal{B}_{\mathbb{R}}$. Equivalently, the reverse containment can also be seen from

$$[a, b] = \bigcap_{n=1}^{\infty} \left(a - \frac{1}{n}, b + \frac{1}{n} \right).$$

Finally, every left half-line in $\mathcal{E}_3$ is closed and hence Borel. Conversely,

$$(-\infty, b) = \bigcup_{n=1}^{\infty} \left(-\infty, b - \frac{1}{n} \right]$$

and therefore

$$(a, b) = (-\infty, b) \cap (-\infty, a]^c \in \mathcal{M}(\mathcal{E}_3).$$

It follows that $\mathcal{E}_1 \subseteq \mathcal{M}(\mathcal{E}_3)$, so $\mathcal{M}(\mathcal{E}_3) = \mathcal{B}_{\mathbb{R}}$ as well. $\square$

2. AUGUST 27, 2026

We continue our discussion of σ -algebras. Recall that a σ -algebra $\mathcal{M}$ on a set X contains $\emptyset$ and X , is closed under complements, and is closed under countable unions. If (X, d) is a metric space, then its Borel σ -algebra $\mathcal{B}_X$ is generated by the open subsets of X .

2.1. Product σ -algebras. Let I be an index set, let X_i be a nonempty set for each $i \in I$, and set

$$X = \prod_{i \in I} X_i.$$

For each $i \in I$, let $\mathcal{M}_i$ be a σ -algebra on X_i , and let

$$\pi_i: X \longrightarrow X_i, \quad \pi_i((x_j)_{j \in I}) = x_i,$$

be the coordinate projection.

Definition 5 (Product σ -algebra). *The product σ -algebra on X is*

$$\bigotimes_{i \in I} \mathcal{M}_i = \mathcal{M} \left(\{ \pi_i^{-1}(E) : i \in I, E \in \mathcal{M}_i \} \right).$$

Thus it is the smallest σ -algebra that makes every coordinate projection measurable.

The generating sets $\pi_i^{-1}(E)$ are called *cylinder sets*. They restrict the i th coordinate to E and leave all other coordinates unrestricted.

Lemma 1. *If I is countable, then $\bigotimes_{i \in I} \mathcal{M}_i$ is generated by the measurable rectangles*

$$\left\{ \prod_{i \in I} E_i : E_i \in \mathcal{M}_i \right\}.$$

Proof. Since I is countable, every measurable rectangle satisfies

$$\prod_{i \in I} E_i = \bigcap_{i \in I} \pi_i^{-1}(E_i) \in \bigotimes_{i \in I} \mathcal{M}_i.$$

Conversely, each cylinder set is a measurable rectangle. Indeed,

$$\pi_j^{-1}(E_j) = \prod_{i \in I} F_i, \quad F_i = \begin{cases} E_j, & i = j, \\ X_i, & i \neq j. \end{cases}$$

The two collections therefore generate the same σ -algebra. $\square$

The generators of the coordinate σ -algebras can be used directly to generate the product.

Proposition 2. *Suppose*

$$\mathcal{M}_i = \mathcal{M}(\mathcal{G}_i), \quad \mathcal{G}_i \subseteq 2^{X_i},$$

for every $i \in I$.

(a) *The product σ -algebra is generated by*

$$\mathcal{F} = \{\pi_i^{-1}(G) : i \in I, G \in \mathcal{G}_i\}.$$

(b) *If I is countable and $\mathcal{G}_i^* = \mathcal{G}_i \cup \{X_i\}$, then the product σ -algebra is generated by*

$$\left\{ \prod_{i \in I} G_i : G_i \in \mathcal{G}_i^* \right\}.$$

Proof. For part (a), every member of $\mathcal{F}$ belongs to $\bigotimes_{i \in I} \mathcal{M}_i$, so

$$\mathcal{M}(\mathcal{F}) \subseteq \bigotimes_{i \in I} \mathcal{M}_i.$$

For the reverse inclusion, fix $i \in I$ and consider

$$\mathcal{A}_i = \{E \subseteq X_i : \pi_i^{-1}(E) \in \mathcal{M}(\mathcal{F})\}.$$

Preimages preserve complements and countable unions, so $\mathcal{A}_i$ is a σ -algebra. It contains $\mathcal{G}_i$ and hence contains $\mathcal{M}(\mathcal{G}_i) = \mathcal{M}_i$. Therefore every cylinder $\pi_i^{-1}(E)$ with $E \in \mathcal{M}_i$ belongs to $\mathcal{M}(\mathcal{F})$, which proves the reverse inclusion.

For part (b), each cylinder from part (a) is a rectangle whose other coordinates are X_j . Conversely, since I is countable, every rectangle in the displayed collection is the countable intersection of the sets $\pi_i^{-1}(G_i)$. When $G_i = X_i$ this set is all of X ; otherwise it is a cylinder from part (a). The conclusion follows. $\square$

2.2. Products of Borel σ -algebras. Let (X_i, d_i) , $i = 1, \dots, n$, be metric spaces. On

$$X = \prod_{i=1}^n X_i$$

define the maximum metric

$$d(x, y) = \max_{1 \leq i \leq n} d_i(x_i, y_i).$$

Proposition 3. *With the notation above,*

$$\bigotimes_{i=1}^n \mathcal{B}_{X_i} \subseteq \mathcal{B}_X.$$

If every (X_i, d_i) is separable, then equality holds:

$$\bigotimes_{i=1}^n \mathcal{B}_{X_i} = \mathcal{B}_X.$$

Proof. Each coordinate projection $\pi_i : X \rightarrow X_i$ is continuous. If $U_i \subseteq X_i$ is open, then $\pi_i^{-1}(U_i)$ is open in X . Since $\mathcal{B}_{X_i}$ is generated by the open subsets of X_i , Proposition 2 gives

$$\bigotimes_{i=1}^n \mathcal{B}_{X_i} \subseteq \mathcal{B}_X.$$

Now suppose that each X_i is separable, and choose a countable dense subset $D_i \subseteq X_i$. The collection

$$\mathcal{G}_i = \{B_{d_i}(p, q) : p \in D_i, q \in \mathbb{Q}, q > 0\}$$

is a countable basis for X_i and hence generates $\mathcal{B}_{X_i}$. The finite product $D = \prod_{i=1}^n D_i$ is countable and dense in X . Moreover,

$$B_d(x, q) = \prod_{i=1}^n B_{d_i}(x_i, q).$$

Thus every ball in X whose center lies in D and whose radius is a positive rational belongs to $\bigotimes_{i=1}^n \mathcal{B}_{X_i}$. These balls form a countable basis for X , so every open subset of X is a countable union of such balls. Hence

$$\mathcal{B}_X \subseteq \bigotimes_{i=1}^n \mathcal{B}_{X_i},$$

which completes the proof. $\square$

Corollary 1. *For every $n \in \mathbb{N}$,*

$$\mathcal{B}_{\mathbb{R}^n} = \bigotimes_{i=1}^n \mathcal{B}_{\mathbb{R}}.$$

For example, $\mathcal{B}_{\mathbb{R}^n}$ is generated by each of the following collections:

- open rectangles $\prod_{i=1}^n (a_i, b_i)$;
- closed rectangles $\prod_{i=1}^n [a_i, b_i]$; and
- lower orthants $\prod_{i=1}^n (-\infty, a_i]$.

2.3. Measures.

Definition 6 (Measure). *Let $\mathcal{M}$ be a σ -algebra on a set X . A function*

$$\mu: \mathcal{M} \longrightarrow [0, \infty]$$

is a measure if

- (i) $\mu(\emptyset) = 0$;
- (ii) *whenever $E_1, E_2, \dots \in \mathcal{M}$ are pairwise disjoint,*

$$\mu\left(\bigcup_{i=1}^{\infty} E_i\right) = \sum_{i=1}^{\infty} \mu(E_i).$$

The second property is called countable additivity. If it is required only for finite pairwise disjoint families, then μ is called finitely additive.

The sets in $\mathcal{M}$ are called *measurable sets*, and the triple $(X, \mathcal{M}, \mu)$ is called a *measure space*. A measure μ is

- *finite* if $\mu(X) < \infty$;
- *σ -finite* if $X = \bigcup_{i=1}^{\infty} E_i$ for some $E_i \in \mathcal{M}$ with $\mu(E_i) < \infty$ for every i .

A set $N \in \mathcal{M}$ with $\mu(N) = 0$ is called a *null set*. A statement is said to hold *almost everywhere*, abbreviated a.e., if it holds outside some null set.

2.4. Examples of measures.

Example 4 (Weighted counting measure). Let X be nonempty, let $\mathcal{M} = 2^X$, and let $f: X \rightarrow [0, \infty]$. Define

$$\mu(E) = \sum_{x \in E} f(x) := \sup \left\{ \sum_{x \in F} f(x) : F \subseteq E, F \text{ finite} \right\}.$$

Then μ is a measure on 2^X .

If $f(x) = 1$ for every $x \in X$, then μ is the counting measure,

$$\mu(E) = \#E.$$

If $x_0 \in X$ and

$$f(x) = \begin{cases} 1, & x = x_0, \\ 0, & x \neq x_0, \end{cases}$$

then

$$\mu(E) = \begin{cases} 1, & x_0 \in E, \\ 0, & x_0 \notin E. \end{cases}$$

This is the point mass, or Dirac measure, at x_0 .

Example 5 (Countable-cocountable measure). Let X be uncountable, and let $\mathcal{M}$ be the countable-cocountable σ -algebra. Then

$$\mu(E) = \begin{cases} 0, & E \text{ is countable,} \\ 1, & E^c \text{ is countable} \end{cases}$$

defines a measure on $(X, \mathcal{M})$.

Example 6 (Finite additivity is not enough). Let $X = \mathbb{N}$ and $\mathcal{M} = 2^{\mathbb{N}}$.

Define

$$\nu(E) = \begin{cases} 0, & E \text{ is finite,} \\ \infty, & E \text{ is infinite.} \end{cases}$$

Then ν is finitely additive. It is not a measure, because

$$\nu\left(\bigcup_{n=1}^{\infty} \{n\}\right) = \infty \quad \text{but} \quad \sum_{n=1}^{\infty} \nu(\{n\}) = 0.$$

2.5. Basic properties of measures.

Theorem 3. Let $(X, \mathcal{M}, \mu)$ be a measure space.

(a) Monotonicity: if $E, F \in \mathcal{M}$ and $E \subseteq F$, then

$$\mu(E) \leq \mu(F).$$

(b) Countable subadditivity: if $E_1, E_2, \dots \in \mathcal{M}$, then

$$\mu\left(\bigcup_{i=1}^{\infty} E_i\right) \leq \sum_{i=1}^{\infty} \mu(E_i).$$

(c) Continuity from below: if

$$E_1 \subseteq E_2 \subseteq \dots, \quad E_i \in \mathcal{M},$$

then

$$\mu\left(\bigcup_{i=1}^{\infty} E_i\right) = \lim_{i \rightarrow \infty} \mu(E_i).$$

(d) Continuity from above: if

$$E_1 \supseteq E_2 \supseteq \dots, \quad E_i \in \mathcal{M}, \quad \mu(E_1) < \infty,$$

then

$$\mu\left(\bigcap_{i=1}^{\infty} E_i\right) = \lim_{i \rightarrow \infty} \mu(E_i).$$

Proof. For part (a), write F as the disjoint union

$$F = E \cup (F \setminus E).$$

Countable additivity gives

$$\mu(F) = \mu(E) + \mu(F \setminus E) \geq \mu(E).$$

For part (b), define

$$F_1 = E_1, \quad F_k = E_k \setminus \bigcup_{j=1}^{k-1} E_j \quad (k \geq 2).$$

The sets F_k are pairwise disjoint, $F_k \subseteq E_k$, and

$$\bigcup_{k=1}^{\infty} F_k = \bigcup_{k=1}^{\infty} E_k.$$

Therefore countable additivity and part (a) imply

$$\mu\left(\bigcup_{k=1}^{\infty} E_k\right) = \sum_{k=1}^{\infty} \mu(F_k) \leq \sum_{k=1}^{\infty} \mu(E_k).$$

For part (c), set

$$F_1 = E_1, \quad F_k = E_k \setminus E_{k-1} \quad (k \geq 2).$$

The sets F_k are pairwise disjoint, and for each N ,

$$E_N = \bigcup_{k=1}^N F_k, \quad \bigcup_{i=1}^{\infty} E_i = \bigcup_{k=1}^{\infty} F_k.$$

It follows that

$$\begin{aligned} \mu\left(\bigcup_{i=1}^{\infty} E_i\right) &= \sum_{k=1}^{\infty} \mu(F_k) \\ &= \lim_{N \rightarrow \infty} \sum_{k=1}^N \mu(F_k) \\ &= \lim_{N \rightarrow \infty} \mu(E_N). \end{aligned}$$

For part (d), define

$$F_i = E_1 \setminus E_i.$$

Since E_i decreases, F_i increases, and

$$\bigcup_{i=1}^{\infty} F_i = E_1 \setminus \bigcap_{i=1}^{\infty} E_i.$$

By part (c),

$$\mu \left(E_1 \setminus \bigcap_{i=1}^{\infty} E_i \right) = \lim_{i \rightarrow \infty} \mu(E_1 \setminus E_i).$$

Because $E_i \subseteq E_1$ and $\mu(E_1) < \infty$, finite additivity gives

$$\mu(E_1 \setminus E_i) = \mu(E_1) - \mu(E_i)$$

and

$$\mu \left(E_1 \setminus \bigcap_{i=1}^{\infty} E_i \right) = \mu(E_1) - \mu \left(\bigcap_{i=1}^{\infty} E_i \right).$$

Substituting these identities into the preceding limit and cancelling the finite quantity $\mu(E_1)$ yields

$$\mu \left(\bigcap_{i=1}^{\infty} E_i \right) = \lim_{i \rightarrow \infty} \mu(E_i).$$

□

Remark 1 (Necessity of the finite-measure hypothesis). The assumption $\mu(E_1) < \infty$ in Theorem 3(d) cannot be omitted. Let

$$X = \mathbb{N}, \quad \mathcal{M} = 2^{\mathbb{N}},$$

let μ be counting measure, and define

$$E_i = \{i, i+1, i+2, \dots\}.$$

Then

$$E_1 \supseteq E_2 \supseteq E_3 \supseteq \dots, \quad \bigcap_{i=1}^{\infty} E_i = \emptyset.$$

Indeed, for any $n \in \mathbb{N}$, we have $n \notin E_i$ whenever $i > n$. Consequently,

$$\mu \left(\bigcap_{i=1}^{\infty} E_i \right) = \mu(\emptyset) = 0.$$

On the other hand, every E_i is infinite, so

$$\mu(E_i) = \infty \quad \text{for every } i, \quad \lim_{i \rightarrow \infty} \mu(E_i) = \infty.$$

Thus

$$\mu \left(\bigcap_{i=1}^{\infty} E_i \right) = 0 \neq \infty = \lim_{i \rightarrow \infty} \mu(E_i).$$

This does not contradict part (d), because $\mu(E_1) = \mu(\mathbb{N}) = \infty$.

3. SEPTEMBER 1, 2026

Recall that a measure on a σ -algebra $\mathcal{M}$ over a set X is a map

$$\mu: \mathcal{M} \longrightarrow [0, \infty]$$

such that $\mu(\emptyset) = 0$ and

$$\mu\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} \mu(A_i)$$

whenever $A_1, A_2, \dots \in \mathcal{M}$ are pairwise disjoint. For Lebesgue measure m on $\mathbb{R}$, the normalization is $m([0, 1]) = 1$.

3.1. Complete measure spaces.

Definition 7 (Complete measure). *A measure μ on a σ -algebra $\mathcal{M}$ over X is complete if every subset of a null set is measurable. That is, if $E \subseteq N$ for some $N \in \mathcal{M}$ with $\mu(N) = 0$, then $E \in \mathcal{M}$.*

By monotonicity, every such set E also satisfies

$$0 \leq \mu(E) \leq \mu(N) = 0,$$

so $\mu(E) = 0$.

Every measure space has a canonical completion obtained by adjoining all subsets of its null sets.

Proposition 4 (Completion of a measure). *Let $(X, \mathcal{M}, \mu)$ be a measure space, and let*

$$\mathcal{N} = \{N \in \mathcal{M} : \mu(N) = 0\}.$$

Define

$$\overline{\mathcal{M}} = \{E \cup F : E \in \mathcal{M}, F \subseteq N \text{ for some } N \in \mathcal{N}\}.$$

Then the following statements hold.

- (i) *The collection $\overline{\mathcal{M}}$ is a σ -algebra on X containing $\mathcal{M}$.*
- (ii) *There is a unique measure $\overline{\mu}$ on $\overline{\mathcal{M}}$ extending μ . It is given by*

$$\overline{\mu}(E \cup F) = \mu(E),$$

where $E \in \mathcal{M}$ and F is contained in a member of $\mathcal{N}$. Moreover, $\bar{\mu}$ is complete.

Proof. The collections $\mathcal{M}$ and $\mathcal{N}$ are closed under countable unions. For $\mathcal{N}$, this follows from countable subadditivity: if $N_i \in \mathcal{N}$, then

$$0 \leq \mu \left(\bigcup_{i=1}^{\infty} N_i \right) \leq \sum_{i=1}^{\infty} \mu(N_i) = 0.$$

Hence $\bar{\mathcal{M}}$ is closed under countable unions. Indeed, if $E_i \in \mathcal{M}$ and $F_i \subseteq N_i \in \mathcal{N}$, then

$$\bigcup_{i=1}^{\infty} (E_i \cup F_i) = \left(\bigcup_{i=1}^{\infty} E_i \right) \cup \left(\bigcup_{i=1}^{\infty} F_i \right),$$

and $\bigcup_i F_i \subseteq \bigcup_i N_i \in \mathcal{N}$.

To check closure under complements, let $E \in \mathcal{M}$ and $F \subseteq N \in \mathcal{N}$. Replacing N by $N \setminus E$ and F by $F \setminus E$ does not change $E \cup F$, so we may assume that $E \cap N = \emptyset$. In that case,

$$(E \cup F)^c = (E \cup N)^c \cup (N \setminus F).$$

The first set on the right belongs to $\mathcal{M}$, and the second is contained in the null set N . Thus $(E \cup F)^c$ belongs to $\bar{\mathcal{M}}$. Since $\emptyset \in \bar{\mathcal{M}}$, this proves that $\bar{\mathcal{M}}$ is a σ -algebra. Taking $F = \emptyset$ shows that $\mathcal{M} \subseteq \bar{\mathcal{M}}$.

We next show that the formula for $\bar{\mu}$ is well defined. Suppose

$$E_1 \cup F_1 = E_2 \cup F_2, \quad E_i \in \mathcal{M}, \quad F_i \subseteq N_i \in \mathcal{N}.$$

Then $E_1 \subseteq E_2 \cup N_2$, and therefore

$$\mu(E_1) \leq \mu(E_2 \cup N_2) \leq \mu(E_2) + \mu(N_2) = \mu(E_2).$$

By symmetry, $\mu(E_2) \leq \mu(E_1)$, so $\mu(E_1) = \mu(E_2)$.

Clearly $\bar{\mu}(\emptyset) = 0$. Suppose that $A_i = E_i \cup F_i$ are pairwise disjoint members of $\bar{\mathcal{M}}$, where $E_i \in \mathcal{M}$ and $F_i \subseteq N_i \in \mathcal{N}$. The sets E_i are pairwise disjoint, so

$$\bar{\mu} \left(\bigcup_{i=1}^{\infty} A_i \right) = \mu \left(\bigcup_{i=1}^{\infty} E_i \right)$$

$$\begin{aligned}
 &= \sum_{i=1}^{\infty} \mu(E_i) \\
 &= \sum_{i=1}^{\infty} \bar{\mu}(A_i).
 \end{aligned}$$

Thus $\bar{\mu}$ is a measure, and it extends μ .

If $A = E \cup F \in \overline{\mathcal{M}}$ and $\bar{\mu}(A) = 0$, then $\mu(E) = 0$. Since $A \subseteq E \cup N$ and $E \cup N$ is a measurable null set, every subset of A belongs to $\overline{\mathcal{M}}$. Hence $\bar{\mu}$ is complete.

Finally, let ν be any measure on $\overline{\mathcal{M}}$ that extends μ . If $A = E \cup F$, with $E \in \mathcal{M}$ and $F \subseteq N \in \mathcal{N}$, then

$$E \subseteq A \subseteq E \cup N.$$

Monotonicity and the fact that ν extends μ give

$$\mu(E) = \nu(E) \leq \nu(A) \leq \nu(E \cup N) = \mu(E \cup N) = \mu(E).$$

Thus $\nu(A) = \mu(E) = \bar{\mu}(A)$, which proves uniqueness. $\square$

The measure $\bar{\mu}$ is called the *completion* of μ .

3.2. Outer measures. The goal is to construct measures from simple geometric pieces. For example, to estimate the area of a set $E \subseteq \mathbb{R}^2$, one can cover E by squares of side length s and consider

$$\inf_{s>0} (\text{number of squares in the cover}) s^2.$$

More generally, one covers a set by countably many simple pieces and takes the infimum of the sums of their assigned sizes.

A related guiding principle is that the “reasonable” subsets $E \subseteq [0, 1]^2$ should be those for which outer and inner area agree. If the unit square has area 1, its inner estimate for E is

$$1 - m^*([0, 1]^2 \setminus E).$$

Definition 8 (Outer measure). *An outer measure on a set X is a map*

$$\mu^*: 2^X \longrightarrow [0, \infty]$$

satisfying the following properties.

- (i) $\mu^*(\emptyset) = 0$.
- (ii) If $A \subseteq B \subseteq X$, then $\mu^*(A) \leq \mu^*(B)$.
- (iii) For every sequence $A_1, A_2, \dots \subseteq X$,

$$\mu^*\left(\bigcup_{j=1}^{\infty} A_j\right) \leq \sum_{j=1}^{\infty} \mu^*(A_j).$$

Unlike a measure, an outer measure is defined on every subset of X ; its countable subadditivity therefore applies without a measurability assumption.

Proposition 5 (Outer measure from coverings). *Let $\mathcal{E} \subseteq 2^X$ contain $\emptyset$ and X , and let*

$$\rho: \mathcal{E} \longrightarrow [0, \infty]$$

satisfy $\rho(\emptyset) = 0$. For $A \subseteq X$, define

$$\mu^*(A) = \inf \left\{ \sum_{j=1}^{\infty} \rho(E_j) : E_j \in \mathcal{E}, \quad A \subseteq \bigcup_{j=1}^{\infty} E_j \right\}.$$

Then μ^ is an outer measure on X .*

Remark 2. Even when $E \in \mathcal{E}$, it need not be true that $\mu^*(E) = \rho(E)$. A more efficient countable cover may have smaller total ρ -size than the single-set cover by E .

Example 7. *Let $X = \mathbb{R}$ and take*

$$\mathcal{E} = \{\emptyset, \mathbb{R}\} \cup \{[a, b] : a, b \in \mathbb{R}, a \leq b\}.$$

Define

$$\rho(\emptyset) = 0, \quad \rho(\mathbb{R}) = \infty, \quad \rho([a, b]) = (b - a)^2.$$

Although $\rho([0, 1]) = 1$, the induced outer measure satisfies $\mu^([0, 1]) = 0$. Indeed, for every $n \in \mathbb{N}$,*

$$[0, 1] = \bigcup_{j=0}^{n-1} \left[\frac{j}{n}, \frac{j+1}{n} \right],$$

and the total ρ -size of this cover is

$$\sum_{j=0}^{n-1} \left(\frac{1}{n}\right)^2 = \frac{1}{n}.$$

Hence $0 \leq \mu^*([0, 1]) \leq 1/n$ for every n , which gives the claim.

Proof of the proposition. The defining collection of sums is nonempty because any set $A \subseteq X$ is covered by the sequence $X, \emptyset, \emptyset, \dots$. Thus $\mu^*(A)$ is well defined as an element of $[0, \infty]$.

The sequence $\emptyset, \emptyset, \dots$ covers $\emptyset$, so

$$0 \leq \mu^*(\emptyset) \leq \sum_{j=1}^{\infty} \rho(\emptyset) = 0.$$

If $A \subseteq B$, then every countable cover of B by members of $\mathcal{E}$ is also a cover of A . Taking infima over the respective collections of covers gives $\mu^*(A) \leq \mu^*(B)$.

It remains to prove countable subadditivity. Let $A_j \subseteq X$ for $j \in \mathbb{N}$. If $\sum_j \mu^*(A_j) = \infty$, there is nothing to prove. Otherwise, fix $\varepsilon > 0$. For every j , choose $E_{j,k} \in \mathcal{E}$, $k \in \mathbb{N}$, such that

$$A_j \subseteq \bigcup_{k=1}^{\infty} E_{j,k} \quad \text{and} \quad \sum_{k=1}^{\infty} \rho(E_{j,k}) \leq \mu^*(A_j) + \varepsilon 2^{-j}.$$

The doubly indexed collection $\{E_{j,k} : j, k \in \mathbb{N}\}$ is countable and covers $\bigcup_j A_j$. Consequently,

$$\begin{aligned} \mu^*\left(\bigcup_{j=1}^{\infty} A_j\right) &\leq \sum_{j=1}^{\infty} \sum_{k=1}^{\infty} \rho(E_{j,k}) \\ &\leq \sum_{j=1}^{\infty} (\mu^*(A_j) + \varepsilon 2^{-j}) \\ &= \sum_{j=1}^{\infty} \mu^*(A_j) + \varepsilon. \end{aligned}$$

Letting $\varepsilon \downarrow 0$ proves the result. $\square$

Definition 9 (Measurability with respect to an outer measure). *Let μ^* be an outer measure on X . A set $A \subseteq X$ is μ^* -measurable if, for*

every $E \subseteq X$,

$$\mu^*(E) = \mu^*(E \cap A) + \mu^*(E \cap A^c).$$

Countable subadditivity always gives

$$\mu^*(E) \leq \mu^*(E \cap A) + \mu^*(E \cap A^c),$$

so the content of the measurability criterion is the reverse inequality.

When $\mu^*(E) < \infty$, the quantity

$$\mu^*(E) - \mu^*(E \cap A^c)$$

can be viewed as the inner measure of $E \cap A$ relative to E .

4. SEPTEMBER 3, 2026

Recall that if μ^* is an outer measure on X , a set $A \subseteq X$ is μ^* -measurable if

$$\mu^*(E) = \mu^*(E \cap A) + \mu^*(E \cap A^c)$$

for every $E \subseteq X$. By countable subadditivity, it is enough to prove the inequality

$$\mu^*(E) \geq \mu^*(E \cap A) + \mu^*(E \cap A^c).$$

4.1. The Carathéodory construction.

Theorem 4 (Carathéodory). *Let μ^* be an outer measure on X , and set*

$$\mathcal{M}^* = \{A \subseteq X : A \text{ is } \mu^*\text{-measurable}\}.$$

Then $\mathcal{M}^$ is a σ -algebra, and*

$$\mu = \mu^*|_{\mathcal{M}^*}$$

is a complete measure on $\mathcal{M}^$.*

Proof. We first show that $\mathcal{M}^*$ is an algebra. The empty set is measurable because

$$\mu^*(E) = \mu^*(E \cap \emptyset) + \mu^*(E \cap X)$$

for every $E \subseteq X$. The defining condition is symmetric in A and A^c , so

$$A \in \mathcal{M}^* \implies A^c \in \mathcal{M}^*.$$

Now let $A, B \in \mathcal{M}^*$ and fix $E \subseteq X$. Applying the measurability criterion first to A and then to B gives

$$\begin{aligned}\mu^*(E) &= \mu^*(E \cap A) + \mu^*(E \cap A^c) \\ &= \mu^*(E \cap A \cap B) + \mu^*(E \cap A \cap B^c) \\ &\quad + \mu^*(E \cap A^c \cap B) + \mu^*(E \cap A^c \cap B^c).\end{aligned}$$

The first three sets on the last two lines have union $E \cap (A \cup B)$. Therefore countable subadditivity implies

$$\mu^*(E) \geq \mu^*(E \cap (A \cup B)) + \mu^*(E \cap (A \cup B)^c).$$

Thus $A \cup B \in \mathcal{M}^*$, and $\mathcal{M}^*$ is an algebra.

This argument also establishes finite additivity on disjoint measurable sets. Indeed, if $A, B \in \mathcal{M}^*$ are disjoint, apply the measurability of A with $E = A \cup B$ to obtain

$$\mu^*(A \cup B) = \mu^*((A \cup B) \cap A) + \mu^*((A \cup B) \cap A^c) = \mu^*(A) + \mu^*(B).$$

To prove closure under countable unions, first let $A_1, A_2, \dots \in \mathcal{M}^*$ be pairwise disjoint, and set

$$B = \bigcup_{i=1}^{\infty} A_i, \quad B_n = \bigcup_{i=1}^n A_i.$$

Since $\mathcal{M}^*$ is an algebra, $B_n \in \mathcal{M}^*$. Repeatedly applying the measurability criterion and using the disjointness of the A_i gives, for every $E \subseteq X$,

$$\mu^*(E) = \sum_{i=1}^n \mu^*(E \cap A_i) + \mu^*(E \cap B_n^c).$$

Because $B^c \subseteq B_n^c$, monotonicity yields

$$\mu^*(E) \geq \sum_{i=1}^n \mu^*(E \cap A_i) + \mu^*(E \cap B^c).$$

Letting $n \rightarrow \infty$ and then using countable subadditivity, we find

$$\mu^*(E) \geq \sum_{i=1}^{\infty} \mu^*(E \cap A_i) + \mu^*(E \cap B^c)$$

$$\begin{aligned}
&\geq \mu^* \left(\bigcup_{i=1}^{\infty} (E \cap A_i) \right) + \mu^*(E \cap B^c) \\
&= \mu^*(E \cap B) + \mu^*(E \cap B^c).
\end{aligned}$$

Hence $B \in \mathcal{M}^*$. An arbitrary countable union can be disjointified by replacing A_i with

$$A_i \setminus \bigcup_{j=1}^{i-1} A_j.$$

It follows that $\mathcal{M}^*$ is a σ -algebra.

The same computation proves countable additivity. If the measurable sets A_i are pairwise disjoint and $B = \bigcup_i A_i$, then countable subadditivity gives

$$\mu^*(B) \leq \sum_{i=1}^{\infty} \mu^*(A_i).$$

On the other hand, for every n ,

$$\mu^*(B) = \sum_{i=1}^n \mu^*(A_i) + \mu^*(B \setminus B_n) \geq \sum_{i=1}^n \mu^*(A_i).$$

Letting $n \rightarrow \infty$ proves

$$\mu^*(B) = \sum_{i=1}^{\infty} \mu^*(A_i).$$

Thus $\mu = \mu^*|_{\mathcal{M}^*}$ is a measure.

Finally, suppose that $A \subseteq X$ and $\mu^*(A) = 0$. For every $E \subseteq X$, subadditivity and monotonicity give

$$\mu^*(E \cap A) + \mu^*(E \cap A^c) \leq 0 + \mu^*(E) = \mu^*(E).$$

Hence $A \in \mathcal{M}^*$. In particular, if $N \in \mathcal{M}^*$ satisfies $\mu(N) = 0$, then every subset $A \subseteq N$ has outer measure zero and therefore belongs to $\mathcal{M}^*$. This proves that μ is complete. $\square$

4.2. Premeasures. The covering construction from the preceding lecture begins with a set function ρ defined on a family of elementary sets.

The example $\rho([a, b]) = (b - a)^2$ showed that the resulting outer measure may fail to agree with ρ , even on elementary sets. To prevent this collapse, we require the initial set function to behave like a measure.

Definition 10 (Premeasure). *Let $\mathcal{A}$ be an algebra on X . A map*

$$\mu_0: \mathcal{A} \longrightarrow [0, \infty]$$

is a premeasure if

- (i) $\mu_0(\emptyset) = 0$;
- (ii) if $A_1, A_2, \dots \in \mathcal{A}$ are pairwise disjoint and $\bigcup_{i=1}^{\infty} A_i \in \mathcal{A}$, then

$$\mu_0\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} \mu_0(A_i).$$

Thus a premeasure has the same additivity property as a measure, but its domain is only required to be an algebra, rather than a σ -algebra.

Example 8 (Half-open intervals). *Let $X = \mathbb{R}$. Consider intervals of the forms*

$$(a, b], \quad (-\infty, b], \quad (a, \infty),$$

together with $\emptyset$, where $a, b \in \mathbb{R}$ and $a < b$. The collection $\mathcal{A}$ of finite unions of such intervals is an algebra. Moreover,

$$\mathcal{M}(\mathcal{A}) = \mathcal{B}_{\mathbb{R}}.$$

This algebra will later be used to construct Lebesgue measure.

The next proposition identifies the hypotheses under which the outer measure construction preserves the initial set function.

Proposition 6 (Extension of a premeasure). *Let μ_0 be a premeasure on an algebra $\mathcal{A}$ over X . For every $E \subseteq X$, define*

$$\mu^*(E) = \inf \left\{ \sum_{i=1}^{\infty} \mu_0(A_i) : A_i \in \mathcal{A}, \quad E \subseteq \bigcup_{i=1}^{\infty} A_i \right\}.$$

Then the following statements hold.

- (i) $\mu^*|_{\mathcal{A}} = \mu_0$.

(ii) Every member of $\mathcal{A}$ is μ^* -measurable.

Consequently, $\mathcal{M}(\mathcal{A}) \subseteq \mathcal{M}^*$, and the measure $\mu^*|_{\mathcal{M}(\mathcal{A})}$ extends μ_0 .

Proof. Let $A \in \mathcal{A}$. The sequence $A, \emptyset, \emptyset, \dots$ is a cover of A , so

$$\mu^*(A) \leq \mu_0(A).$$

Conversely, suppose that $A_i \in \mathcal{A}$ and $A \subseteq \bigcup_{i=1}^{\infty} A_i$. Define

$$B_1 = A \cap A_1, \quad B_k = A \cap \left(A_k \setminus \bigcup_{i=1}^{k-1} A_i \right) \quad (k \geq 2).$$

Then the sets B_k are pairwise disjoint members of $\mathcal{A}$, and

$$A = \bigcup_{k=1}^{\infty} B_k.$$

Since $B_k \subseteq A_k$, monotonicity of the premeasure gives

$$\mu_0(A) = \sum_{k=1}^{\infty} \mu_0(B_k) \leq \sum_{k=1}^{\infty} \mu_0(A_k).$$

Taking the infimum over all such covers yields $\mu_0(A) \leq \mu^*(A)$. Therefore $\mu^*(A) = \mu_0(A)$, proving part (i).

For part (ii), fix $A \in \mathcal{A}$ and $E \subseteq X$. If $\mu^*(E) = \infty$, the desired measurability identity follows at once from countable subadditivity. Suppose instead that $\mu^*(E) < \infty$, and let $\varepsilon > 0$. Choose $A_1, A_2, \dots \in \mathcal{A}$ such that

$$E \subseteq \bigcup_{i=1}^{\infty} A_i \quad \text{and} \quad \mu^*(E) + \varepsilon \geq \sum_{i=1}^{\infty} \mu_0(A_i).$$

Since μ_0 is finitely additive,

$$\begin{aligned} \sum_{i=1}^{\infty} \mu_0(A_i) &= \sum_{i=1}^{\infty} (\mu_0(A_i \cap A) + \mu_0(A_i \cap A^c)) \\ &\geq \mu^*(E \cap A) + \mu^*(E \cap A^c). \end{aligned}$$

Hence

$$\mu^*(E) + \varepsilon \geq \mu^*(E \cap A) + \mu^*(E \cap A^c).$$

Letting $\varepsilon \downarrow 0$ proves that A is μ^* -measurable.

By Theorem 4, $\mathcal{M}^*$ is a σ -algebra. Since it contains $\mathcal{A}$, it also contains $\mathcal{M}(\mathcal{A})$. Part (i) then shows that the restriction of μ^* to $\mathcal{M}(\mathcal{A})$ extends μ_0 . $\square$

5. SEPTEMBER 8, 2026

We continue the construction of measures from premeasures. Recall that if μ_0 is a premeasure on an algebra $\mathcal{A}$, then

$$\mu^*(E) = \inf \left\{ \sum_{j=1}^{\infty} \mu_0(A_j) : A_j \in \mathcal{A}, \quad E \subseteq \bigcup_{j=1}^{\infty} A_j \right\}$$

defines an outer measure. Every member of $\mathcal{A}$ is μ^* -measurable, and

$$\mu = \mu^*|_{\mathcal{M}(\mathcal{A})}$$

is a measure extending μ_0 .

5.1. Uniqueness of the extension. The extension is unique on sets of finite measure. Under a σ -finiteness assumption, it is unique everywhere.

Theorem 5 (Uniqueness of the premeasure extension). *Let μ_0 be a premeasure on an algebra $\mathcal{A}$, and let μ be the measure on $\mathcal{M}(\mathcal{A})$ obtained from the induced outer measure. Suppose that ν is another measure on $\mathcal{M}(\mathcal{A})$ extending μ_0 .*

- (a) *If $E \in \mathcal{M}(\mathcal{A})$ and $\mu(E) < \infty$, then $\nu(E) = \mu(E)$.*
- (b) *If μ_0 is σ -finite, then $\nu = \mu$ on $\mathcal{M}(\mathcal{A})$.*

Proof. We divide the proof of part (a) into three steps.

First, let $E \in \mathcal{M}(\mathcal{A})$ and suppose that $E \subseteq \bigcup_{j=1}^{\infty} A_j$, where $A_j \in \mathcal{A}$. Since ν extends μ_0 , countable subadditivity gives

$$\nu(E) \leq \nu \left(\bigcup_{j=1}^{\infty} A_j \right) \leq \sum_{j=1}^{\infty} \nu(A_j) = \sum_{j=1}^{\infty} \mu_0(A_j).$$

Taking the infimum over all such covers yields

$$(1) \quad \nu(E) \leq \mu^*(E) = \mu(E).$$

Second, suppose that

$$A = \bigcup_{j=1}^{\infty} A_j, \quad A_j \in \mathcal{A}.$$

Set $B_n = \bigcup_{j=1}^n A_j$. Then $B_n \in \mathcal{A}$ and $B_n \uparrow A$. Continuity from below, together with the fact that μ and ν both extend μ_0 , gives

$$\nu(A) = \lim_{n \rightarrow \infty} \nu(B_n) = \lim_{n \rightarrow \infty} \mu(B_n) = \mu(A).$$

Third, let $E \in \mathcal{M}(\mathcal{A})$ satisfy $\mu(E) < \infty$, and fix $\varepsilon > 0$. By the definition of μ^* , there are $A_j \in \mathcal{A}$ such that, with $A = \bigcup_{j=1}^{\infty} A_j$,

$$E \subseteq A, \quad \mu(A) \leq \sum_{j=1}^{\infty} \mu_0(A_j) < \mu(E) + \varepsilon.$$

In particular, $\mu(A \setminus E) < \varepsilon$. By the second step and (1),

$$\begin{aligned} \mu(E) &\leq \mu(A) = \nu(A) \\ &= \nu(E) + \nu(A \setminus E) \\ &\leq \nu(E) + \mu(A \setminus E) < \nu(E) + \varepsilon. \end{aligned}$$

Letting $\varepsilon \downarrow 0$ shows that $\mu(E) \leq \nu(E)$. Together with (1), this proves part (a).

For part (b), σ -finiteness provides sets $A_j \in \mathcal{A}$ such that

$$X = \bigcup_{j=1}^{\infty} A_j, \quad \mu_0(A_j) < \infty.$$

By replacing A_j with $A_j \setminus \bigcup_{i=1}^{j-1} A_i$, we may assume that the sets are pairwise disjoint. If $E \in \mathcal{M}(\mathcal{A})$, then $\mu(E \cap A_j) \leq \mu_0(A_j) < \infty$, so part (a) applies to each $E \cap A_j$. Therefore

$$\mu(E) = \sum_{j=1}^{\infty} \mu(E \cap A_j) = \sum_{j=1}^{\infty} \nu(E \cap A_j) = \nu(E).$$

□

5.2. Lebesgue–Stieltjes measures on $\mathbb{R}$. Let μ be a finite Borel measure on $\mathbb{R}$. Its *distribution function* is

$$(2) \quad F(x) = \mu((-\infty, x]), \quad x \in \mathbb{R}.$$

This function records the measure of every left half-line.

Proposition 7. *The distribution function F has the following properties.*

- (i) *It is increasing.*
- (ii) *It is right-continuous.*
- (iii) *For every $a < b$,*

$$\mu((a, b]) = F(b) - F(a).$$

Proof. If $x \leq y$, then $(-\infty, x] \subseteq (-\infty, y]$, so monotonicity of μ gives $F(x) \leq F(y)$.

If $x_n \downarrow x$, then

$$(-\infty, x_n] \downarrow (-\infty, x].$$

Since μ is finite, continuity from above gives

$$\lim_{n \rightarrow \infty} F(x_n) = \lim_{n \rightarrow \infty} \mu((-\infty, x_n]) = \mu((-\infty, x]) = F(x).$$

Hence F is right-continuous. Finally,

$$(-\infty, b] = (-\infty, a] \dot{\cup} (a, b],$$

and finite additivity gives

$$\mu((a, b]) = \mu((-\infty, b]) - \mu((-\infty, a]) = F(b) - F(a).$$

□

Example 9 (Distribution function of a Dirac measure). *Let $\mu = \delta_{x_0}$ be the Dirac measure at $x_0 \in \mathbb{R}$. Then*

$$F(x) = \delta_{x_0}((-\infty, x]) = \begin{cases} 0, & x < x_0, \\ 1, & x \geq x_0. \end{cases}$$

Thus an atom of a measure appears as a jump in its distribution function.

Remark 3 (Measures finite on bounded sets). A Borel measure need not be finite on all of $\mathbb{R}$. If μ is finite on every bounded Borel set, define the normalized distribution function

$$\tilde{F}(x) = \begin{cases} \mu((0, x]), & x > 0, \\ 0, & x = 0, \\ -\mu((x, 0]), & x < 0. \end{cases}$$

Then $\tilde{F}$ is increasing and right-continuous, and

$$\mu((a, b]) = \tilde{F}(b) - \tilde{F}(a)$$

for all $a < b$. Adding a constant to $\tilde{F}$ does not change these interval increments.

We now reverse the preceding construction at the level of the algebra of half-open intervals. Let $F: \mathbb{R} \rightarrow \mathbb{R}$ be increasing and right-continuous, and set

$$F(-\infty) = \lim_{x \rightarrow -\infty} F(x), \quad F(\infty) = \lim_{x \rightarrow \infty} F(x),$$

where these limits are allowed to be infinite. Let $\mathcal{A}$ be the algebra of finite unions of intervals of the forms

$$(a, b], \quad (-\infty, b], \quad (a, \infty),$$

together with $\emptyset$. Equivalently, its members are finite unions of intervals $(a, b] \cap \mathbb{R}$ with $-\infty \leq a < b \leq \infty$.

Proposition 8 (The Lebesgue–Stieltjes premeasure). *Suppose that*

$$E = \bigcup_{j=1}^N (a_j, b_j]$$

is a representation of $E \in \mathcal{A}$ as a finite union of pairwise disjoint half-open intervals, with extended endpoints allowed as above. Define

$$(3) \quad \mu_0(E) = \sum_{j=1}^N (F(b_j) - F(a_j)), \quad \mu_0(\emptyset) = 0.$$

Then $\mu_0: \mathcal{A} \rightarrow [0, \infty]$ is well defined and is a premeasure.

Proof. We first show that the value in (3) is independent of the chosen representation. Suppose that a half-open interval is partitioned into finitely many pairwise disjoint half-open intervals:

$$(a, b] = \bigcup_{j=1}^N (a_j, b_j].$$

After relabeling, their endpoints satisfy

$$a = a_1 < b_1 = a_2 < b_2 = \cdots < b_{N-1} = a_N < b_N = b.$$

Hence the sum telescopes:

$$\sum_{j=1}^N (F(b_j) - F(a_j)) = F(b) - F(a).$$

The same argument applies when an endpoint is infinite, using the conventions for $F(-\infty)$ and $F(\infty)$.

Now suppose that

$$\bigcup_{i=1}^N I_i = \bigcup_{j=1}^M J_j,$$

where both families consist of pairwise disjoint half-open intervals. The nonempty intersections $I_i \cap J_j$ partition both sides. By the preceding telescoping identity,

$$\begin{aligned} \sum_{i=1}^N \mu_0(I_i) &= \sum_{i=1}^N \sum_{j=1}^M \mu_0(I_i \cap J_j) \\ &= \sum_{j=1}^M \sum_{i=1}^N \mu_0(I_i \cap J_j) = \sum_{j=1}^M \mu_0(J_j). \end{aligned}$$

Thus μ_0 is well defined; the same calculation also gives finite additivity.

It remains to prove countable additivity. After decomposing all sets involved into their interval components and intersecting with each component of the union, it is enough to consider pairwise disjoint half-open intervals I_j such that

$$(a, b] = \bigcup_{j=1}^{\infty} I_j.$$

Finite additivity and monotonicity give, for every n ,

$$\mu_0((a, b]) \geq \mu_0\left(\bigcup_{j=1}^n I_j\right) = \sum_{j=1}^n \mu_0(I_j).$$

Letting $n \rightarrow \infty$ yields

$$(4) \quad \mu_0((a, b]) \geq \sum_{j=1}^{\infty} \mu_0(I_j).$$

For the reverse inequality, first suppose that $a, b \in \mathbb{R}$ and write $I_j = (a_j, b_j]$. Fix $\varepsilon > 0$. Right-continuity provides $0 < \delta < b - a$ and $\delta_j > 0$ such that

$$F(a + \delta) - F(a) < \varepsilon, \quad F(b_j + \delta_j) - F(b_j) < \varepsilon 2^{-j}.$$

The open intervals $(a_j, b_j + \delta_j)$ cover the compact interval $[a + \delta, b]$, so finitely many of them suffice. After discarding redundant intervals, relabel this finite subcover from left to right with indices $1, \dots, r$. Its total enlargement still satisfies

$$\sum_{j=1}^r (F(b_j + \delta_j) - F(b_j)) < \varepsilon,$$

and the ordering may be chosen so that

$$a_1 < a + \delta, \quad a_{j+1} < b_j + \delta_j \ (1 \leq j < r), \quad b < b_r + \delta_r.$$

Monotonicity of F then gives

$$\begin{aligned} F(b) - F(a) &< F(b) - F(a + \delta) + \varepsilon \\ &\leq F(b_r + \delta_r) - F(a_1) + \varepsilon \\ &= F(b_r + \delta_r) - F(a_r) + \sum_{j=1}^{r-1} (F(a_{j+1}) - F(a_j)) + \varepsilon \\ &\leq \sum_{j=1}^r (F(b_j + \delta_j) - F(a_j)) + \varepsilon \\ &< \sum_{j=1}^r (F(b_j) - F(a_j)) + 2\varepsilon \end{aligned}$$

$$\leq \sum_{j=1}^{\infty} \mu_0(I_j) + 2\varepsilon.$$

Letting $\varepsilon \downarrow 0$ proves the reverse of (4) for bounded intervals.

If the interval on the left is unbounded, apply the bounded case to its truncations. For example,

$$(-m, b] \uparrow (-\infty, b], \quad \mu_0((-m, b]) = F(b) - F(-m) \uparrow F(b) - F(-\infty),$$

while the bounded case gives

$$\mu_0((-m, b]) = \sum_{j=1}^{\infty} \mu_0(I_j \cap (-m, b]) \leq \sum_{j=1}^{\infty} \mu_0(I_j).$$

Letting $m \rightarrow \infty$ proves the desired inequality for $(-\infty, b]$. The same argument uses $(a, m] \uparrow (a, \infty)$ or $(-m, m] \uparrow \mathbb{R}$ for the other unbounded cases. Therefore

$$\mu_0\left(\bigcup_{j=1}^{\infty} I_j\right) = \sum_{j=1}^{\infty} \mu_0(I_j),$$

whenever the union belongs to $\mathcal{A}$, so μ_0 is a premeasure. $\square$

The premeasure extension theorem from the preceding lecture therefore applies to μ_0 on $\mathcal{A}$.

Remark 4 (Two Lebesgue–Stieltjes examples). The Borel measure determined by the increments $F(b) - F(a)$ is denoted by μ_F , and its completion is called the *Lebesgue–Stieltjes measure* associated with F .

(i) For $F(x) = x$,

$$\mu_F((a, b]) = b - a.$$

The completed measure is Lebesgue measure on $\mathbb{R}$.

(ii) For the right-continuous step function

$$F(x) = \mathbf{1}_{[x_0, \infty)}(x),$$

the associated measure is the Dirac measure $\mu_F = \delta_{x_0}$.

6. SEPTEMBER 10, 2026

We finish the construction begun in the preceding lecture. Recall that an increasing, right-continuous function $F: \mathbb{R} \rightarrow \mathbb{R}$ defines a pre-measure on the algebra of finite unions of half-open intervals by

$$\mu_0((a, b]) = F(b) - F(a).$$

The extension theorem and its uniqueness statement give the following result.

Theorem 6 (Lebesgue–Stieltjes measure). *Let $F: \mathbb{R} \rightarrow \mathbb{R}$ be increasing and right-continuous. There is a unique Borel measure μ_F on $\mathbb{R}$, finite on bounded sets, such that*

$$\mu_F((a, b]) = F(b) - F(a)$$

for every $a < b$. Its completion $\bar{\mu}_F$ is called the Lebesgue–Stieltjes measure associated with F .

Proof. Proposition 8 shows that the interval increments of F define a premeasure on the algebra $\mathcal{A}$ of finite unions of half-open intervals. The premeasure extension theorem therefore produces a measure on

$$\mathcal{M}(\mathcal{A}) = \mathcal{B}_{\mathbb{R}}.$$

It is finite on bounded intervals because F is real-valued. In particular, the premeasure is σ -finite, so uniqueness follows from Theorem 5. $\square$

The measure depends only on the increments of F . Thus

$$\mu_F = \mu_G \iff F - G \text{ is constant.}$$

Conversely, every Borel measure μ on $\mathbb{R}$ that is finite on bounded sets arises in this way. One convenient normalization is

$$F(x) = \begin{cases} \mu((0, x]), & x > 0, \\ 0, & x = 0, \\ -\mu((x, 0]), & x < 0. \end{cases}$$

Then F is increasing and right-continuous, and uniqueness of the extension gives $\mu = \mu_F$ on $\mathcal{B}_{\mathbb{R}}$.

One could instead work with an increasing, left-continuous function G . With that convention the natural intervals are left-closed and right-open, and

$$\mu_G([a, b)) = G(b) - G(a).$$

Example 10. If $F(x) = x$, then

$$\mu_F((a, b]) = b - a.$$

The completion of μ_F is Lebesgue measure.

Example 11. If $F(x) = \mathbf{1}_{[x_0, \infty)}(x)$, then $\mu_F = \delta_{x_0}$.

6.1. Regularity of Lebesgue–Stieltjes measures. Fix an increasing, right-continuous F , and write μ for the completed measure $\bar{\mu}_F$. Its domain will be denoted by $\mathcal{M}_F$. The outer-measure construction gives, for every $E \in \mathcal{M}_F$,

$$(5) \quad \mu(E) = \inf \left\{ \sum_{j=1}^{\infty} (F(b_j) - F(a_j)) : E \subseteq \bigcup_{j=1}^{\infty} (a_j, b_j] \right\}.$$

There is an equivalent formula involving open intervals.

Lemma 2 (Open covering formula). *For every $E \in \mathcal{M}_F$,*

$$\mu(E) = \inf \left\{ \sum_{j=1}^{\infty} \mu((a_j, b_j)) : E \subseteq \bigcup_{j=1}^{\infty} (a_j, b_j) \right\}.$$

Proof. Let $V(E)$ denote the infimum on the right. If $E \subseteq \bigcup_j (a_j, b_j)$, then countable subadditivity gives

$$\mu(E) \leq \mu \left(\bigcup_{j=1}^{\infty} (a_j, b_j) \right) \leq \sum_{j=1}^{\infty} \mu((a_j, b_j)).$$

Taking the infimum shows that $\mu(E) \leq V(E)$.

For the reverse inequality, suppose first that $\mu(E) < \infty$, and fix $\varepsilon > 0$. By (5), there are intervals $(a_j, b_j]$ covering E such that

$$\sum_{j=1}^{\infty} \mu((a_j, b_j]) < \mu(E) + \varepsilon.$$

Right-continuity of F allows us to choose $\delta_j > 0$ with

$$F(b_j + \delta_j) - F(b_j) < \varepsilon 2^{-j}.$$

The open intervals $(a_j, b_j + \delta_j)$ still cover E , and

$$\begin{aligned} \sum_{j=1}^{\infty} \mu((a_j, b_j + \delta_j)) &\leq \sum_{j=1}^{\infty} (F(b_j + \delta_j) - F(a_j)) \\ &< \sum_{j=1}^{\infty} \mu((a_j, b_j]) + \varepsilon \\ &< \mu(E) + 2\varepsilon. \end{aligned}$$

Hence $V(E) \leq \mu(E)$. If $\mu(E) = \infty$, the already proved inequality $\mu(E) \leq V(E)$ forces $V(E) = \infty$. $\square$

Theorem 7 (Regularity). *Let $E \in \mathcal{M}_F$.*

(i) Outer regularity:

$$\mu(E) = \inf\{\mu(U) : U \supseteq E, U \text{ open}\}.$$

(ii) Inner regularity:

$$\mu(E) = \sup\{\mu(K) : K \subseteq E, K \text{ compact}\}.$$

Proof. If $E \subseteq U$, then $\mu(E) \leq \mu(U)$. Conversely, the open covering formula provides open intervals (a_j, b_j) such that $E \subseteq U = \bigcup_j (a_j, b_j)$ and

$$\mu(U) \leq \sum_{j=1}^{\infty} \mu((a_j, b_j)) < \mu(E) + \varepsilon$$

whenever $\mu(E) < \infty$. The infinite-measure case is immediate. This proves outer regularity.

We first prove inner regularity when E is bounded. Then $\overline{E}$ has finite measure. By outer regularity, there is an open set U containing $\overline{E} \setminus E$ such that

$$\mu(U) < \mu(\overline{E} \setminus E) + \varepsilon.$$

The set

$$K = \overline{E} \setminus U$$

is closed and bounded, hence compact, and $K \subseteq E$. Since all relevant measures are finite,

$$\begin{aligned}\mu(K) &= \mu(\overline{E}) - \mu(\overline{E} \cap U) \\ &\geq \mu(\overline{E}) - \mu(U) \\ &> \mu(\overline{E}) - \mu(\overline{E} \setminus E) - \varepsilon = \mu(E) - \varepsilon.\end{aligned}$$

For arbitrary E , apply the bounded case to $E \cap [-n, n]$. These sets increase to E , so continuity from below gives

$$\mu(E) = \lim_{n \rightarrow \infty} \mu(E \cap [-n, n]).$$

Compact subsets approximating the bounded truncations therefore give the desired supremum, whether $\mu(E)$ is finite or infinite. $\square$

Regularity gives useful Borel representatives for every set in the completion.

Corollary 2. *If $E \in \mathcal{M}_F$, then there are a G_δ set G , an F_σ set H , and null sets N_1, N_2 such that*

$$H \subseteq E \subseteq G, \quad E = G \setminus N_1 = H \cup N_2.$$

In particular, every Lebesgue–Stieltjes measurable set differs from a Borel set by a null set.

Proof. Decompose E into the bounded pieces $E_k = E \cap [k, k+1)$, $k \in \mathbb{Z}$. For every n and k , outer regularity gives an open set $U_{n,k} \supseteq E_k$ for which

$$\mu(U_{n,k} \setminus E_k) < 2^{-n-|k|-2}.$$

Set

$$U_n = \bigcup_{k \in \mathbb{Z}} U_{n,k}, \quad G = \bigcap_{n=1}^{\infty} U_n.$$

Then G is a G_δ set containing E , and $\mu(G \setminus E) = 0$. Thus $E = G \setminus N_1$, where $N_1 = G \setminus E$ is null. Apply the same argument to E^c and take complements to find an F_σ set $H \subseteq E$ such that $E \setminus H$ is null. Taking $N_2 = E \setminus H$ completes the proof. $\square$

6.2. Lebesgue measure. Let m denote Lebesgue measure on the Lebesgue σ -algebra $\mathcal{L}$. Thus m is the completion of the Borel measure generated by $F(x) = x$.

Theorem 8 (Translation and dilation invariance). *If $E \in \mathcal{L}$ and $s \in \mathbb{R}$, then $E + s \in \mathcal{L}$ and*

$$m(E + s) = m(E).$$

If $\lambda \in \mathbb{R} \setminus \{0\}$, then $\lambda E \in \mathcal{L}$ and

$$m(\lambda E) = |\lambda| m(E).$$

For $\lambda = 0$, the set λE is either empty or a singleton and therefore has measure zero. Thus the dilation formula also holds for $\lambda = 0$ when $m(E) < \infty$ (or with the convention $0 \cdot \infty = 0$).

Proof. Translation by s is a homeomorphism, so it preserves Borel sets. On $\mathcal{B}_{\mathbb{R}}$ define

$$m_s(E) = m(E + s).$$

This is a Borel measure, and for every half-open interval,

$$m_s((a, b]) = m((a + s, b + s]) = b - a = m((a, b]).$$

Uniqueness of the premeasure extension gives $m_s = m$ on $\mathcal{B}_{\mathbb{R}}$.

If N is a Borel null set, then $N + s$ is again a Borel null set. Every $E \in \mathcal{L}$ has the form

$$E = B \cup N,$$

where B is Borel and N is contained in a Borel null set. It follows that $E + s$ is Lebesgue measurable and $m(E + s) = m(B + s) = m(B) = m(E)$.

Now fix $\lambda \neq 0$. The map $x \mapsto \lambda x$ is also a homeomorphism. Singletons have measure zero because

$$m(\{x\}) \leq m((x - \varepsilon, x]) = \varepsilon$$

for every $\varepsilon > 0$. Consequently every interval with endpoints $c < d$, regardless of the choice of open or closed endpoints, has measure $d - c$.

Hence

$$m_\lambda(E) = \frac{1}{|\lambda|} m(\lambda E), \quad E \in \mathcal{B}_{\mathbb{R}},$$

is a Borel measure satisfying $m_\lambda((a, b]) = b - a$. Uniqueness gives $m_\lambda = m$ on Borel sets. As above, dilation carries Borel null sets to Borel null sets, so the identity extends to every $E \in \mathcal{L}$. $\square$

Example 12 (Countable sets). *Every singleton is null, and therefore every countable set is null. In particular,*

$$m(\mathbb{Q}) = 0.$$

More concretely, if $\mathbb{Q} \cap (0, 1) = \{q_1, q_2, \dots\}$ and $\varepsilon > 0$, then

$$U_\varepsilon = \bigcup_{j=1}^{\infty} (q_j - \varepsilon 2^{-j}, q_j + \varepsilon 2^{-j})$$

is an open dense set containing $\mathbb{Q} \cap (0, 1)$, while

$$m(U_\varepsilon) \leq \sum_{j=1}^{\infty} 2\varepsilon 2^{-j} = 2\varepsilon.$$

Thus an open dense set can have arbitrarily small measure.

7. SEPTEMBER 15, 2026

We begin with several examples illustrating the difference between measure-theoretic size and topological size. Throughout, m denotes Lebesgue measure and $\mathcal{L}$ denotes the Lebesgue σ -algebra.

7.1. Measure and Baire category. Enumerate the rationals as $\mathbb{Q} = \{q_1, q_2, \dots\}$. For $n \in \mathbb{N}$, define

$$U_n = \bigcup_{k=1}^{\infty} (q_k - 2^{-n-k-2}, q_k + 2^{-n-k-2}).$$

Each U_n is open and dense because it contains $\mathbb{Q}$, but

$$m(U_n) \leq \sum_{k=1}^{\infty} 2^{-n-k-1} = 2^{-n-1}.$$

The set

$$U = \bigcap_{n=1}^{\infty} U_n$$

is a dense G_δ set by the Baire category theorem, yet $m(U) = 0$. It is necessarily uncountable: a countable subset of $\mathbb{R}$ is meager, whereas a dense G_δ subset of $\mathbb{R}$ is comeager.

This example has a complementary form on $[0, 1]$. Let

$$F_n = [0, 1] \setminus U_n, \quad F = \bigcup_{n=1}^{\infty} F_n = [0, 1] \setminus U.$$

Because U_n is open and dense, F_n is closed and nowhere dense. Thus F is meager, being a countable union of nowhere dense sets, but

$$m(F) = 1.$$

Thus the topological and measure-theoretic notions of size need not agree.

7.2. The Cantor set. The middle-thirds Cantor set $C \subseteq [0, 1]$ can be described in two equivalent ways. First, set $C_0 = [0, 1]$ and obtain C_n from C_{n-1} by deleting the open middle third of each interval component. Then

$$C = \bigcap_{n=0}^{\infty} C_n.$$

Equivalently,

$$C = \left\{ \sum_{j=1}^{\infty} a_j 3^{-j} : a_j \in \{0, 2\} \right\}.$$

Proposition 9. *The Cantor set has the following properties.*

- (a) $m(C) = 0$.
- (b) C is compact, has empty interior, and is totally disconnected.
- (c) $|C| = |\mathbb{R}|$.

Proof. The set C_n consists of 2^n closed intervals, each of length 3^{-n} . By continuity from above,

$$m(C) = \lim_{n \rightarrow \infty} m(C_n) = \lim_{n \rightarrow \infty} \left(\frac{2}{3} \right)^n = 0.$$

Each C_n is compact, so their nested intersection C is compact. Since every nonempty interval has positive measure, the equality $m(C) = 0$ implies that C has empty interior. Every connected subset of $\mathbb{R}$ is an interval; since C contains no nontrivial interval, each connected component of C is a singleton. Thus C is totally disconnected.

Finally, the map

$$(b_j)_{j=1}^{\infty} \mapsto \sum_{j=1}^{\infty} 2b_j 3^{-j}, \quad b_j \in \{0, 1\},$$

is a bijection from $\{0, 1\}^{\mathbb{N}}$ onto C . Since the set of binary sequences has the cardinality of $\mathbb{R}$, so does C . $\square$

The Cantor set supports a function that is continuous and increasing but whose derivative vanishes almost everywhere.

Definition 11 (Cantor function). *If $x \in C$ has ternary expansion*

$$x = \sum_{j=1}^{\infty} 2b_j 3^{-j}, \quad b_j \in \{0, 1\},$$

define

$$\Phi(x) = \sum_{j=1}^{\infty} b_j 2^{-j}.$$

Every component of $[0, 1] \setminus C$ is an open interval whose endpoints lie in C and have the same value under the preceding formula. Extend Φ to be constant on each such interval. The resulting map $\Phi: [0, 1] \rightarrow [0, 1]$ is the Cantor function.

The Cantor function is continuous and increasing, with $\Phi(0) = 0$ and $\Phi(1) = 1$. On the open set $[0, 1] \setminus C$ it is locally constant. Since $m(C) = 0$, it follows that

$$\Phi'(x) = 0 \quad \text{for almost every } x \in [0, 1].$$

Nevertheless,

$$\Phi(1) - \Phi(0) = 1 \neq 0 = \int_0^1 \Phi'(x) dx.$$

Thus continuity alone is not sufficient for the strongest form of the fundamental theorem of calculus.




FIGURE 1. The third-stage polygonal approximation Φ_3 to the Cantor function. At every subsequent stage, each sloping segment is replaced by a smaller copy of the first-stage pattern; the limit is constant on every component of $[0, 1] \setminus C$.

Example 13 (A fat Cantor set). Set $K_0 = [0, 1]$. At stage $n \geq 1$, remove from the center of each interval component of K_{n-1} an open interval of length 4^{-n} , and call the remaining set K_n . Thus the first step removes the middle interval $(3/8, 5/8)$ and leaves

$$K_1 = \left[0, \frac{3}{8}\right] \cup \left[\frac{5}{8}, 1\right].$$

There are 2^{n-1} intervals removed at stage n , each of length 4^{-n} . Hence, for the limit set $K = \bigcap_{n=0}^{\infty} K_n$,

$$m(K) = 1 - \sum_{n=1}^{\infty} 2^{n-1} 4^{-n} = 1 - \sum_{n=1}^{\infty} 2^{-n-1} = \frac{1}{2}.$$

The lengths of the interval components tend to zero, while every component splits into two components at the next stage. Consequently, K is compact, perfect, has empty interior, and is totally disconnected. Thus

K has all the topological features of the usual Cantor set but has positive measure. It is called the fat Cantor set (or the Smith–Volterra–Cantor set).




FIGURE 2. The first stages of the fat Cantor construction. The deleted intervals become shorter quickly enough that the limiting set retains measure $1/2$.

7.3. The idea of Lebesgue integration. Let $(X, \mathcal{M}, \mu)$ be a measure space. For $E \in \mathcal{M}$, the *characteristic function* of E is

$$\mathbf{1}_E(x) = \begin{cases} 1, & x \in E, \\ 0, & x \notin E. \end{cases}$$

The integral is required to satisfy

$$\int_X \mathbf{1}_E d\mu = \mu(E).$$

This suggests how to integrate a more general function. Rather than partitioning the domain into short intervals, as in Riemann integration, Lebesgue integration partitions the range. For a mesh size $\varepsilon > 0$ and $k \in \mathbb{Z}$, consider the level sets

$$E_k = \{x \in X : k\varepsilon \leq f(x) < (k+1)\varepsilon\}.$$

When these sets belong to $\mathcal{M}$, the function can be approximated by a simple function, and formally

$$\int_X f d\mu \approx \sum_{k \in \mathbb{Z}} k\varepsilon \mu(E_k).$$

The requirement that the level sets be measurable leads to the notion of a measurable function.

7.4. Measurable functions.

Definition 12. Let $(X, \mathcal{M})$ and $(Y, \mathcal{N})$ be measurable spaces. A map $f: X \rightarrow Y$ is $(\mathcal{M}, \mathcal{N})$ -measurable if

$$f^{-1}(E) \in \mathcal{M}$$

for every $E \in \mathcal{N}$.

For a real- or complex-valued function, the codomain is equipped with its Borel σ -algebra. Thus $f: X \rightarrow \mathbb{R}$ is $\mathcal{M}$ -measurable when it is $(\mathcal{M}, \mathcal{B}_{\mathbb{R}})$ -measurable, and similarly for $f: X \rightarrow \mathbb{C}$.

The extended real line

$$\overline{\mathbb{R}} = \mathbb{R} \cup \{-\infty, +\infty\}$$

is equipped with its Borel σ -algebra. Its topology may be described by the metric

$$d(x, y) = |\arctan x - \arctan y|, \quad \arctan(\pm\infty) = \pm\frac{\pi}{2}.$$

For functions on $\mathbb{R}$, the terminology is worth distinguishing:

- f is *Borel measurable* if it is $(\mathcal{B}_{\mathbb{R}}, \mathcal{B}_{\mathbb{R}})$ -measurable;
- f is *Lebesgue measurable* if it is $(\mathcal{L}, \mathcal{B}_{\mathbb{R}})$ -measurable.

The codomain still carries the Borel σ -algebra in the second definition. In particular, although a composition of appropriately measurable maps is measurable, the composition of two Lebesgue measurable functions $\mathbb{R} \rightarrow \mathbb{R}$ need not be Lebesgue measurable: the inverse image under the outer function is only known to be a Lebesgue set, while the inner function is only required to pull back Borel sets.

Lemma 3 (Checking a generating family). *Suppose $\mathcal{N} = \mathcal{M}(\mathcal{E})$ for some $\mathcal{E} \subseteq 2^Y$. A map $f: X \rightarrow Y$ is $(\mathcal{M}, \mathcal{N})$ -measurable if and only if*

$$f^{-1}(E) \in \mathcal{M}$$

for every $E \in \mathcal{E}$.

Proof. Only the reverse implication requires proof. Set

$$\mathcal{A} = \{E \subseteq Y : f^{-1}(E) \in \mathcal{M}\}.$$

Since preimages preserve complements and countable unions, $\mathcal{A}$ is a σ -algebra. By hypothesis it contains $\mathcal{E}$, so it contains $\mathcal{M}(\mathcal{E}) = \mathcal{N}$. $\square$

Corollary 3. *If X and Y are topological spaces and $f: X \rightarrow Y$ is continuous, then f is $(\mathcal{B}_X, \mathcal{B}_Y)$ -measurable.*

Proof. The Borel σ -algebra $\mathcal{B}_Y$ is generated by the open subsets of Y , and the inverse image of an open set under a continuous map is open. $\square$

Proposition 10 (Half-line criterion). *Let $f: X \rightarrow \overline{\mathbb{R}}$. The following are equivalent.*

- (i) f is measurable.
- (ii) $\{x : f(x) > a\} \in \mathcal{M}$ for every $a \in \mathbb{R}$.
- (iii) $\{x : f(x) \geq a\} \in \mathcal{M}$ for every $a \in \mathbb{R}$.
- (iv) $\{x : f(x) < a\} \in \mathcal{M}$ for every $a \in \mathbb{R}$.
- (v) $\{x : f(x) \leq a\} \in \mathcal{M}$ for every $a \in \mathbb{R}$.

Proof. Each of the four corresponding families of half-lines generates the Borel σ -algebra of $\overline{\mathbb{R}}$. The conclusion follows from Lemma 3. $\square$

Finally, if $(X_i, \mathcal{M}_i)_{i \in I}$ are measurable spaces, the product σ -algebra $\bigotimes_{i \in I} \mathcal{M}_i$ is precisely the smallest σ -algebra on $\prod_i X_i$ for which every coordinate projection

$$\pi_j: \prod_{i \in I} X_i \longrightarrow X_j$$

is measurable. This is the measurable-space analogue of the fact that the product topology is the smallest topology making all coordinate projections continuous.

8. SEPTEMBER 17, 2026

We continue the study of measurable functions and then introduce the building blocks of the Lebesgue integral. Throughout, $(X, \mathcal{M})$ is a measurable space. A real- or complex-valued function is understood to have the Borel σ -algebra on its codomain.

Recall that the Borel σ -algebra on $\overline{\mathbb{R}} = \mathbb{R} \cup \{-\infty, +\infty\}$ is generated by either of the families

$$\{(a, +\infty] : a \in \mathbb{R}\} \quad \text{and} \quad \{[-\infty, a] : a \in \mathbb{R}\}.$$

Thus the half-line criterion from the preceding lecture applies equally well to extended-real-valued functions.

8.1. Operations on measurable functions. Let $(Y_i, \mathcal{N}_i)_{i \in I}$ be measurable spaces, and give

$$Y = \prod_{i \in I} Y_i$$

the product σ -algebra $\mathcal{N} = \bigotimes_{i \in I} \mathcal{N}_i$. If $\pi_i : Y \rightarrow Y_i$ is the i th coordinate projection, then $\mathcal{N}$ is generated by the cylinder sets

$$\pi_i^{-1}(E), \quad i \in I, \quad E \in \mathcal{N}_i.$$

Lemma 4 (Coordinate criterion for product measurability). *A map $f : X \rightarrow Y$ is $(\mathcal{M}, \bigotimes_{i \in I} \mathcal{N}_i)$ -measurable if and only if every coordinate map*

$$\pi_i \circ f : X \longrightarrow Y_i$$

is $(\mathcal{M}, \mathcal{N}_i)$ -measurable.

Proof. If f is measurable, then $\pi_i \circ f$ is measurable because each projection π_i is measurable. Conversely, suppose all the coordinate maps are measurable. For every generating cylinder,

$$f^{-1}(\pi_i^{-1}(E)) = (\pi_i \circ f)^{-1}(E) \in \mathcal{M}.$$

Lemma 3 now shows that f is measurable. □

Corollary 4. *A function $f : X \rightarrow \mathbb{C}$ is measurable if and only if $\operatorname{Re} f$ and $\operatorname{Im} f$ are measurable.*

Proof. Under the identification $\mathbb{C} \cong \mathbb{R}^2$,

$$\mathcal{B}_{\mathbb{C}} = \mathcal{B}_{\mathbb{R}} \otimes \mathcal{B}_{\mathbb{R}}.$$

The claim is therefore the two-coordinate case of Lemma 4. □




FIGURE 3. Measurability into a product can be checked one coordinate at a time.

Proposition 11 (Algebra and lattice operations). *If $f, g: X \rightarrow \mathbb{R}$ are measurable, then so are*

$$f + g, \quad fg, \quad f^+ = \max\{f, 0\}, \quad f^- = \max\{-f, 0\}, \quad |f|,$$

and $\text{sgn}(f)$.

Proof. By Lemma 4, the map

$$F = (f, g): X \longrightarrow \mathbb{R}^2$$

is measurable. Addition and multiplication are continuous maps from $\mathbb{R}^2$ to $\mathbb{R}$, so they are Borel measurable. Hence

$$f + g = \Sigma \circ F, \quad fg = \Pi \circ F,$$

where $\Sigma(s, t) = s + t$ and $\Pi(s, t) = st$, are measurable. The remaining functions follow by composition with the Borel maps

$$t \longmapsto \max\{t, 0\}, \quad t \longmapsto \max\{-t, 0\}, \quad t \longmapsto |t|,$$

and with the Borel measurable sign function. Notice also that $f = f^+ - f^-$ and $|f| = f^+ + f^-$. $\square$

The same argument, together with Corollary 4, shows that sums and products of complex-valued measurable functions are measurable.

Proposition 12 (Countable suprema and limits). *Let $f_n: X \rightarrow \overline{\mathbb{R}}$ be measurable for every $n \in \mathbb{N}$. Then the functions*

$$\sup_n f_n, \quad \inf_n f_n, \quad \limsup_{n \rightarrow \infty} f_n, \quad \liminf_{n \rightarrow \infty} f_n$$

are measurable. In particular, if $f_n \rightarrow f$ pointwise, then f is measurable.

Proof. For $a \in \mathbb{R}$,

$$\left\{ \sup_n f_n > a \right\} = \bigcup_{n=1}^{\infty} \{f_n > a\}, \quad \left\{ \inf_n f_n < a \right\} = \bigcup_{n=1}^{\infty} \{f_n < a\}.$$

The half-line criterion implies that the pointwise supremum and infimum are measurable. Now use the identities

$$\limsup_{n \rightarrow \infty} f_n = \inf_{N \geq 1} \sup_{n \geq N} f_n, \quad \liminf_{n \rightarrow \infty} f_n = \sup_{N \geq 1} \inf_{n \geq N} f_n.$$

If $f_n \rightarrow f$ pointwise, both expressions equal f . $\square$

value at a fixed x




FIGURE 4. The tail supremum decreases and the tail infimum increases. Their limits are the pointwise lim sup and lim inf; the drawing shows the convergent case.

8.2. Simple functions. For $E \in \mathcal{M}$, continue to write $\mathbf{1}_E$ for the characteristic function of E .

Definition 13. A measurable function $\phi: X \rightarrow \mathbb{C}$ is simple if it has finite range. Equivalently, it can be written as a finite linear combination

$$\phi = \sum_{j=1}^N a_j \mathbf{1}_{E_j}, \quad a_j \in \mathbb{C}, \quad E_j \in \mathcal{M}.$$

If $\{a_1, \dots, a_N\}$ is the set of distinct values of ϕ and $E_j = \phi^{-1}(\{a_j\})$, then the sets E_j are pairwise disjoint and cover X . This is the standard representation of ϕ .

Sums, products, absolute values, maxima, and minima of simple functions are again simple. The standard representation makes a simple function look like a finite collection of horizontal levels over a measurable partition of its domain.




FIGURE 5. A simple function $\phi = \sum_{j=1}^4 a_j \mathbf{1}_{E_j}$ in standard form. The sets E_j need not be intervals in a general measurable space.

Simple functions are sufficient to approximate every measurable function.

Theorem 9 (Approximation by simple functions). *(i) If $f: X \rightarrow [0, +\infty]$ is measurable, then there are nonnegative simple functions ϕ_n such that*

$$0 \leq \phi_1 \leq \phi_2 \leq \cdots \leq f, \quad \phi_n(x) \longrightarrow f(x)$$

for every $x \in X$. The convergence is uniform on every set on which f is bounded.

(ii) If $f: X \rightarrow \mathbb{C}$ is measurable, then there are simple functions $\phi_n: X \rightarrow \mathbb{C}$ such that

$$0 \leq |\phi_1| \leq |\phi_2| \leq \cdots \leq |f|, \quad \phi_n(x) \longrightarrow f(x)$$

for every $x \in X$. Again, convergence is uniform on every set on which $|f|$ is bounded.

Proof. For (i), divide the interval $[0, n]$ into dyadic pieces of length 2^{-n} . Set

$$E_{n,k} = f^{-1}((k2^{-n}, (k+1)2^{-n}]), \quad 0 \leq k < n2^n,$$

and $F_n = f^{-1}((n, +\infty])$. Define

$$(6) \quad \phi_n = \sum_{k=0}^{n2^n-1} k2^{-n} \mathbf{1}_{E_{n,k}} + n \mathbf{1}_{F_n}.$$

Each ϕ_n is a nonnegative simple function. The dyadic grids refine one another and the truncation height increases, so $\phi_n \leq \phi_{n+1} \leq f$. Moreover,

$$0 \leq f(x) - \phi_n(x) \leq 2^{-n} \quad \text{whenever } f(x) \leq n.$$

It follows that $\phi_n \uparrow f$ pointwise. If $f \leq M$ on a set E , then for $n \geq M$,

$$\sup_{x \in E} |f(x) - \phi_n(x)| \leq 2^{-n},$$

which proves the asserted uniform convergence.

For (ii), write

$$f = (\operatorname{Re} f)^+ - (\operatorname{Re} f)^- + i(\operatorname{Im} f)^+ - i(\operatorname{Im} f)^-.$$

Apply (i) to these four nonnegative measurable functions to obtain increasing simple approximants u_n, v_n, w_n, z_n , respectively, and set

$$\phi_n = (u_n - v_n) + i(w_n - z_n).$$

At each point, at most one of u_n, v_n is nonzero and at most one of w_n, z_n is nonzero. Consequently $|\phi_n|$ increases with n and is bounded by $|f|$. The convergence statements follow from those for the four real components. $\square$




FIGURE 6. The approximation in (6) rounds f down to a dyadic level and truncates it at height n . Below the truncation height, the vertical error is at most 2^{-n} .

8.3. Nonnegative integration. Now fix a measure space $(X, \mathcal{M}, \mu)$ and write

$$L^+ = \{f: X \rightarrow [0, +\infty] : f \text{ is measurable}\}.$$

Definition 14 (Integral of a nonnegative simple function). *Let $\phi \in L^+$ be simple, with standard representation*

$$\phi = \sum_{j=1}^N a_j \mathbf{1}_{E_j}.$$

Define

$$\int_X \phi \, d\mu = \sum_{j=1}^N a_j \mu(E_j),$$

using the convention $0 \cdot \infty = 0$. For $A \in \mathcal{M}$, set

$$\int_A \phi \, d\mu = \int_X \phi \mathbf{1}_A \, d\mu.$$

The standard representation is unique up to the ordering of its nonempty level sets, so this definition is unambiguous. Geometrically, each term $a_j \mu(E_j)$ is the measure-theoretic analogue of the area of a rectangle.




FIGURE 7. For Lebesgue measure on an interval, the simple integral is literally the sum of the shaded rectangular areas. The definition $\sum_j a_j \mu(E_j)$ makes sense on every measure space.

Definition 15 (Integral of a nonnegative measurable function). *For $f \in L^+$, define*

$$\int_X f \, d\mu = \sup \left\{ \int_X \phi \, d\mu : \phi \text{ is simple and } 0 \leq \phi \leq f \right\}.$$

This value belongs to $[0, +\infty]$. For $A \in \mathcal{M}$, define

$$\int_A f \, d\mu = \int_X f \mathbf{1}_A \, d\mu.$$

Theorem 9 explains why the supremum uses precisely the right class of building blocks: every $f \in L^+$ is the pointwise increasing limit of simple functions below it.

Example 14 (Three measures and their integrals). (i) *If $\mu = \delta_{x_0}$ is the Dirac measure at $x_0 \in X$, then*

$$\int_X f \, d\delta_{x_0} = f(x_0) \quad (f \in L^+).$$

Thus a Dirac integral samples the function at one point.

(ii) *If $\mu = m$ is Lebesgue measure on $\mathbb{R}$, then the integral of a non-negative simple function is the usual sum of rectangular areas, as in Figure 7.*

(iii) Since $\mathbb{Q}$ is null but the fat Cantor set K from Example 13 has measure $1/2$,

$$\int_{\mathbb{R}} \mathbf{1}_{\mathbb{Q}} d\mu = 0, \quad \int_{\mathbb{R}} \mathbf{1}_K d\mu = \frac{1}{2}.$$

Figure 2 shows the construction of K .

We finish with properties that follow directly from the finite sum defining the integral of a simple function.

Proposition 13 (Elementary properties of the simple integral). *Let $\phi, \psi \in L^+$ be simple and let $c \geq 0$.*

- (a) $\int_X c\phi d\mu = c \int_X \phi d\mu$.
- (b) $\int_X (\phi + \psi) d\mu = \int_X \phi d\mu + \int_X \psi d\mu$.
- (c) If $\phi \leq \psi$, then $\int_X \phi d\mu \leq \int_X \psi d\mu$.
- (d) The set function

$$\nu(A) = \int_A \phi d\mu, \quad A \in \mathcal{M},$$

is a measure on $(X, \mathcal{M})$.

Proof. Part (a) is immediate. For (b), write the standard representations as

$$\phi = \sum_{j=1}^N a_j \mathbf{1}_{E_j}, \quad \psi = \sum_{k=1}^M b_k \mathbf{1}_{F_k}.$$

The sets $E_j \cap F_k$ give a common measurable partition, and

$$\phi + \psi = \sum_{j,k} (a_j + b_k) \mathbf{1}_{E_j \cap F_k}.$$

Finite additivity of μ gives

$$\begin{aligned} \int_X (\phi + \psi) d\mu &= \sum_{j,k} (a_j + b_k) \mu(E_j \cap F_k) \\ &= \sum_j a_j \mu(E_j) + \sum_k b_k \mu(F_k) \\ &= \int_X \phi d\mu + \int_X \psi d\mu. \end{aligned}$$

The same common partition proves (c): whenever $E_j \cap F_k \neq \emptyset$, the inequality $\phi \leq \psi$ implies $a_j \leq b_k$, so comparison term by term gives the result.

Finally,

$$\nu(A) = \int_X \phi \mathbf{1}_A d\mu = \sum_{j=1}^N a_j \mu(E_j \cap A).$$

Each map $A \mapsto \mu(E_j \cap A)$ is a measure. Therefore their finite nonnegative linear combination ν is a measure, proving (d). $\square$