

MATH 535: LINEAR ALGEBRA

RUIZHE QIAN

CONTENTS

1. August 24, 2026	3
1.1. Fields	3
2. August 26, 2026	5
2.1. Further properties of fields	5
2.2. Characteristic of a field	5
2.3. Algebraically closed fields	6
2.4. Vectors and vector spaces	7
3. August 28, 2026	7
3.1. Examples of vector spaces	7
3.2. Linear combinations and spanning sets	8
3.3. Linear independence	9
4. August 31, 2026	11
4.1. Bases and coordinates	11
4.2. Linear transformations	13
5. September 2, 2026	15
5.1. Linear transformations determined by a basis	15
5.2. Matrix representations of linear transformations	16
6. September 4, 2026	18
6.1. Matrix–vector multiplication	18
6.2. Finding matrix representations	20
6.3. The vector space of linear maps	21
6.4. Composition of linear maps	22
7. September 9, 2026	22
7.1. Matrix operations and linear maps	22
7.2. Composition and matrix multiplication	23
7.3. Vector subspaces	25

8. September 11, 2026	25
8.1. The subspace criterion	25
8.2. Kernels and images	26
8.3. Sums of subspaces	27
8.4. Direct sums	28
8.5. Product vector spaces	29
9. September 14, 2026	29
9.1. Direct sums and injectivity	29
9.2. Quotient vector spaces	30
9.3. Finite-dimensional vector spaces	31
10. September 16, 2026	32
10.1. Comparing independent and spanning collections	32
10.2. Subspaces of finite-dimensional spaces	33
10.3. Extending a linearly independent collection	34
11. September 18, 2026	35
11.1. Complements of subspaces	35
11.2. Dimension	36
11.3. Consequences of dimension	37
12. September 21, 2026	38
12.1. Spanning collections and dimension	38
12.2. Dimension of a sum of subspaces	38
12.3. Linear maps on finite-dimensional spaces	40
13. September 23, 2026	41
13.1. Injectivity, surjectivity, and dimension	41
13.2. Proof of the fundamental theorem of linear maps	41
13.3. Dimension restrictions on linear maps	43
13.4. A complement to the kernel	43
14. September 25, 2026	44
14.1. Invertible linear maps	44
14.2. Invertibility and bijectivity	46
14.3. Why the dimension hypotheses matter	47
14.4. One-sided inverses in equal dimensions	48

1. AUGUST 24, 2026

1.1. Fields.

Definition 1 (Binary operation). *A binary operation on a set S is a function*

$$F: S \times S \longrightarrow S.$$

Thus, for every $x, y \in S$, the value $F(x, y)$ is again an element of S . We usually write $x \cdot y$ or $x + y$ in place of $F(x, y)$.

Example 1. *Addition on $\mathbb{Z}$ is a binary operation: the map*

$$\mathbb{Z} \times \mathbb{Z} \longrightarrow \mathbb{Z}, \quad (m, n) \longmapsto m + n$$

takes every pair of integers to an integer.

Definition 2. *Let F be a binary operation on S . A subset $T \subseteq S$ is closed under F if*

$$F(x, y) \in T$$

whenever $x, y \in T$.

Example 2. *The positive integers $\mathbb{Z}_{>0}$ are closed under addition. The odd integers are not closed under addition, but they are closed under multiplication.*

Definition 3 (Field). *Let $\mathbb{F}$ be a set equipped with two binary operations, called addition and multiplication. The sum and product of $a, b \in \mathbb{F}$ are denoted by $a + b$ and ab , respectively. Suppose that for every $a, b, c \in \mathbb{F}$,*

- (i) $a + b = b + a$;
- (ii) $(a + b) + c = a + (b + c)$;
- (iii) $ab = ba$;
- (iv) $a(bc) = (ab)c$;
- (v) $a(b + c) = ab + ac$.

Suppose further that

(vi) $\mathbb{F}$ contains an additive identity 0 and a multiplicative identity 1, so that

$$a + 0 = a \quad \text{and} \quad 1a = a$$

for every $a \in \mathbb{F}$;

(vii) $1 \neq 0$;

(viii) every $a \in \mathbb{F}$ has an additive inverse $-a \in \mathbb{F}$ such that $a + (-a) = 0$;

(ix) every nonzero $a \in \mathbb{F}$ has a multiplicative inverse $a^{-1} \in \mathbb{F}$ such that $aa^{-1} = 1$.

Then $\mathbb{F}$ is called a field.

Example 3. The integers $\mathbb{Z}$ do not form a field: for example, 2 has no multiplicative inverse in $\mathbb{Z}$. The sets $\mathbb{Q}$, $\mathbb{R}$, and $\mathbb{C}$, with their usual operations, are fields.

There are also finite fields. For example,

$$\mathbb{F}_2 = \{0, 1\} \quad \text{and} \quad \mathbb{F}_3 = \{0, 1, 2\},$$

where the elements are the possible remainders upon division by 2 and 3, respectively. Addition and multiplication in $\mathbb{F}_3$ are given by the following tables:

$$\begin{array}{c|ccc} + & 0 & 1 & 2 \\ \hline 0 & 0 & 1 & 2 \\ 1 & 1 & 2 & 0 \\ 2 & 2 & 0 & 1 \end{array} \quad \begin{array}{c|ccc} \cdot & 0 & 1 & 2 \\ \hline 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 2 \\ 2 & 0 & 2 & 1 \end{array}.$$

More generally, if p is prime, then

$$\mathbb{F}_p = \{0, 1, \dots, p-1\},$$

with addition and multiplication performed modulo p , is a field.

Lemma 1. If a is an element of a field $\mathbb{F}$, then $a \cdot 0 = 0$.

Proof. Since $0 + 0 = 0$, distributivity gives

$$a \cdot 0 = a(0 + 0) = a \cdot 0 + a \cdot 0.$$

Adding the additive inverse of $a \cdot 0$ to both sides yields $0 = a \cdot 0$. $\square$

2. AUGUST 26, 2026

2.1. Further properties of fields. The preceding lecture showed that $a \cdot 0 = 0$ for every element a of a field. We now obtain the zero-product property.

Proposition 1 (Zero-product property). *Let $a, b \in \mathbb{F}$. If $ab = 0$, then either $a = 0$ or $b = 0$.*

Proof. Suppose that $ab = 0$ and $a \neq 0$. Since $\mathbb{F}$ is a field, a^{-1} exists. Therefore

$$0 = a^{-1}0 = a^{-1}(ab) = (a^{-1}a)b = 1b = b.$$

Thus $a = 0$ or $b = 0$. $\square$

2.2. Characteristic of a field. For a positive integer n and $a \in \mathbb{F}$, write

$$na = \underbrace{a + a + \cdots + a}_{n \text{ terms}}.$$

Definition 4 (Characteristic). *Let $\mathbb{F}$ be a field. If $n1 = 0$ for some positive integer n , then the least such n is called the characteristic of $\mathbb{F}$, denoted $\text{char}(\mathbb{F})$. If $n1 \neq 0$ for every positive integer n , then $\mathbb{F}$ has characteristic zero.*

The fields $\mathbb{Q}$, $\mathbb{R}$, and $\mathbb{C}$ have characteristic zero. For every prime p , the field $\mathbb{F}_p$ has characteristic p . The rational function field $\mathbb{F}_p(t)$ is an infinite field of characteristic p .

Proposition 2. *If $\mathbb{F}$ has positive characteristic r , then r is prime.*

Proof. We have $r \neq 1$, since $1 \neq 0$. Suppose that r is composite, so

$$r = st$$

for integers s, t satisfying $1 < s, t < r$. Distributivity gives

$$(s1)(t1) = (st)1 = r1 = 0.$$

By the zero-product property, either $s1 = 0$ or $t1 = 0$. Both alternatives contradict the minimality of r . Hence r is prime. $\square$

Proposition 3. *If $\text{char}(\mathbb{F}) = p > 0$, then*

$$pa = 0$$

for every $a \in \mathbb{F}$.

Proof. Since $p1 = 0$, associativity and distributivity imply

$$pa = p(1a) = (p1)a = 0a = 0.$$

□

Proposition 4. *Every finite field has positive characteristic.*

Proof. Let $\mathbb{F}$ be finite. The sequence

$$1, \quad 2 \cdot 1, \quad 3 \cdot 1, \quad \dots$$

cannot consist of distinct elements. Hence there are positive integers $m > n$ such that

$$m1 = n1.$$

Adding the additive inverse of $n1$ to both sides gives

$$(m - n)1 = 0.$$

Thus $\mathbb{F}$ has positive characteristic. □

2.3. Algebraically closed fields.

Definition 5 (Algebraically closed field). *A field $\mathbb{F}$ is algebraically closed if every nonconstant polynomial $f \in \mathbb{F}[x]$ has a root in $\mathbb{F}$.*

The field $\mathbb{Q}$ is not algebraically closed, because $x^2 - 2$ has no root in $\mathbb{Q}$. The field $\mathbb{R}$ is not algebraically closed, because $x^2 + 1$ has no real root. By the fundamental theorem of algebra, $\mathbb{C}$ is algebraically closed.

Proposition 5. *No finite field is algebraically closed.*

Proof. Let

$$\mathbb{F} = \{a_1, \dots, a_q\}$$

be finite. Consider the nonconstant polynomial

$$f(x) = 1 + \prod_{i=1}^q (x - a_i).$$

For each $i = 1, \dots, q$,

$$f(a_i) = 1 \neq 0.$$

Thus f has no root in $\mathbb{F}$, so $\mathbb{F}$ is not algebraically closed. $\square$

2.4. Vectors and vector spaces. The spaces $\mathbb{R}^2$, $\mathbb{R}^3$, and, more generally,

$$\mathbb{R}^n = \left\{ \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} : x_1, \dots, x_n \in \mathbb{R} \right\}$$

are the motivating examples of vector spaces.

Definition 6 (Vector space). *Let $\mathbb{F}$ be a field. A vector space over $\mathbb{F}$ is a set V equipped with two operations,*

$$V \times V \longrightarrow V, \quad (u, v) \longmapsto u + v,$$

and

$$\mathbb{F} \times V \longrightarrow V, \quad (\alpha, v) \longmapsto \alpha v,$$

satisfying the following axioms for all $u, v, w \in V$ and $\alpha, \beta \in \mathbb{F}$:

$$(V1) \quad u + (v + w) = (u + v) + w;$$

$$(V2) \quad u + v = v + u;$$

$$(V3) \quad \text{there is an element } 0 \in V \text{ such that } v + 0 = 0 + v = v;$$

$$(V4) \quad \text{for every } v \in V, \text{ there is an element } -v \in V \text{ such that } v + (-v) = (-v) + v = 0;$$

$$(V5) \quad 1v = v;$$

$$(V6) \quad (\alpha\beta)v = \alpha(\beta v);$$

$$(V7) \quad \alpha(u + v) = \alpha u + \alpha v;$$

$$(V8) \quad (\alpha + \beta)v = \alpha v + \beta v.$$

The elements of V are called *vectors*, and the elements of $\mathbb{F}$ are called *scalars*.

3. AUGUST 28, 2026

3.1. Examples of vector spaces.

Example 4 (Coordinate spaces). *Let $\mathbb{F}$ be a field. The set*

$$\mathbb{F}^n = \left\{ \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} : x_1, \dots, x_n \in \mathbb{F} \right\}$$

is a vector space over $\mathbb{F}$ under componentwise addition and scalar multiplication:

$$\begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} + \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix} = \begin{bmatrix} x_1 + y_1 \\ \vdots \\ x_n + y_n \end{bmatrix}, \quad \alpha \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} \alpha x_1 \\ \vdots \\ \alpha x_n \end{bmatrix}.$$

In particular, $\mathbb{R}^n$ is a vector space over $\mathbb{R}$ and $\mathbb{C}^n$ is a vector space over $\mathbb{C}$.

Example 5 (Matrix spaces). *The set $M_{m \times n}(\mathbb{F})$ of $m \times n$ matrices with entries in $\mathbb{F}$ is a vector space over $\mathbb{F}$. Addition and scalar multiplication are defined entrywise:*

$$[a_{ij}] + [b_{ij}] = [a_{ij} + b_{ij}], \quad \alpha[a_{ij}] = [\alpha a_{ij}].$$

Example 6 (Polynomial spaces). *Let*

$$\mathcal{P}_n(\mathbb{F}) = \{a_0 + a_1t + \dots + a_nt^n : a_0, \dots, a_n \in \mathbb{F}\}.$$

This is the vector space over $\mathbb{F}$ of polynomials of degree at most n . Its operations are

$$\begin{aligned} (a_0 + a_1t + \dots + a_nt^n) + (b_0 + b_1t + \dots + b_nt^n) \\ = (a_0 + b_0) + (a_1 + b_1)t + \dots + (a_n + b_n)t^n, \end{aligned}$$

and

$$\alpha(a_0 + a_1t + \dots + a_nt^n) = \alpha a_0 + \alpha a_1t + \dots + \alpha a_nt^n.$$

3.2. Linear combinations and spanning sets.

Definition 7 (Linear combination and span). *Let V be a vector space over $\mathbb{F}$ and let $v_1, \dots, v_p \in V$. A linear combination of these vectors is a vector of the form*

$$\alpha_1 v_1 + \dots + \alpha_p v_p, \quad \alpha_1, \dots, \alpha_p \in \mathbb{F}.$$

The set of all their linear combinations is called their span:

$$\text{span}\{v_1, \dots, v_p\} = \{\alpha_1 v_1 + \dots + \alpha_p v_p : \alpha_1, \dots, \alpha_p \in \mathbb{F}\}.$$

A linear combination is trivial if all of its coefficients are zero.

In $\mathbb{R}^2$, let

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

Then

$$\text{span}\{e_1, e_2\} = \mathbb{R}^2,$$

because

$$\begin{bmatrix} x \\ y \end{bmatrix} = x e_1 + y e_2.$$

If

$$v = \begin{bmatrix} 1 \\ 1 \end{bmatrix},$$

then $\{e_1, e_2, v\}$ also spans $\mathbb{R}^2$, since

$$\begin{bmatrix} x \\ y \end{bmatrix} = x e_1 + y e_2 + 0 v.$$

The additional vector is unnecessary, which motivates the question of redundancy in a spanning set.

Definition 8 (Spanning set). *A collection $\{v_1, \dots, v_p\} \subseteq V$ is a spanning set for V if every $v \in V$ can be written as a linear combination*

$$v = \alpha_1 v_1 + \dots + \alpha_p v_p.$$

Equivalently,

$$\text{span}\{v_1, \dots, v_p\} = V.$$

3.3. Linear independence.

Definition 9 (Linear independence). *A collection $\{v_1, \dots, v_p\} \subseteq V$ is linearly independent if*

$$\alpha_1 v_1 + \dots + \alpha_p v_p = 0$$

implies

$$\alpha_1 = \cdots = \alpha_p = 0.$$

In other words, only the trivial linear combination of $v_1, \dots, v_p$ equals zero. A collection that is not linearly independent is called linearly dependent.

Example 7. Consider the vectors

$$v_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \quad v_2 = \begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix}, \quad v_3 = \begin{bmatrix} 3 \\ 1 \\ 1 \end{bmatrix}$$

in $\mathbb{R}^3$. To test linear independence, solve

$$\alpha v_1 + \beta v_2 + \gamma v_3 = 0.$$

This gives the system

$$\begin{cases} \alpha + 2\beta + 3\gamma = 0, \\ \alpha + \gamma = 0, \\ \beta + \gamma = 0. \end{cases}$$

Taking $\gamma = 1$ gives $\alpha = \beta = -1$, so

$$-v_1 - v_2 + v_3 = 0.$$

Thus the vectors are linearly dependent; equivalently, $v_3 = v_1 + v_2$.

The following result precisely characterizes redundancy in a finite collection of vectors.

Proposition 6. Let $v_1, \dots, v_p \in V$. The collection $\{v_1, \dots, v_p\}$ is linearly dependent if and only if one of its vectors is a linear combination of the others. Equivalently, there is an index k and scalars $\beta_j \in \mathbb{F}$ such that

$$v_k = \sum_{\substack{j=1 \\ j \neq k}}^p \beta_j v_j.$$

Proof. Suppose first that $\{v_1, \dots, v_p\}$ is linearly dependent. Then there are scalars $\alpha_1, \dots, \alpha_p$, not all zero, such that

$$\alpha_1 v_1 + \dots + \alpha_p v_p = 0.$$

Choose k such that $\alpha_k \neq 0$. Since $\mathbb{F}$ is a field, α_k^{-1} exists, and we can solve for v_k :

$$v_k = - \sum_{\substack{j=1 \\ j \neq k}}^p \alpha_k^{-1} \alpha_j v_j.$$

Thus v_k is a linear combination of the other vectors.

Conversely, suppose that for some k ,

$$v_k = \sum_{\substack{j=1 \\ j \neq k}}^p \beta_j v_j.$$

Moving all terms to one side gives

$$v_k - \sum_{\substack{j=1 \\ j \neq k}}^p \beta_j v_j = 0.$$

This is a nontrivial linear combination of $v_1, \dots, v_p$, because the coefficient of v_k is $1 \neq 0$. Hence the collection is linearly dependent. $\square$

4. AUGUST 31, 2026

4.1. Bases and coordinates.

Definition 10 (Basis). *A collection of vectors $\{v_1, \dots, v_n\}$ in a vector space V is a basis for V if it is linearly independent and spans V .*

The two conditions in the definition can be combined into a statement about representing vectors uniquely.

Proposition 7. *A collection $\{v_1, \dots, v_n\}$ is a basis for V if and only if every vector $v \in V$ has a unique representation of the form*

$$v = a_1 v_1 + \dots + a_n v_n, \quad a_1, \dots, a_n \in \mathbb{F}.$$

When $\{v_1, \dots, v_n\}$ is a basis, the scalars $a_1, \dots, a_n$ are called the coordinates of v with respect to this basis.

Proof. Suppose first that $\{v_1, \dots, v_n\}$ is a basis. Since the collection spans V , every $v \in V$ has a representation

$$v = a_1v_1 + \dots + a_nv_n.$$

To prove uniqueness, suppose that

$$v = \tilde{a}_1v_1 + \dots + \tilde{a}_nv_n$$

is another such representation. Subtracting the two equations gives

$$(a_1 - \tilde{a}_1)v_1 + \dots + (a_n - \tilde{a}_n)v_n = 0.$$

Because $\{v_1, \dots, v_n\}$ is linearly independent,

$$a_i - \tilde{a}_i = 0$$

for each i . Thus $a_i = \tilde{a}_i$ for every i , and the representation is unique.

Conversely, suppose every $v \in V$ has a unique representation as a linear combination of $v_1, \dots, v_n$. The existence of such a representation for every v shows that the collection spans V . Now suppose

$$a_1v_1 + \dots + a_nv_n = 0.$$

The zero vector also has the representation

$$0 = 0v_1 + \dots + 0v_n.$$

Uniqueness therefore gives $a_1 = \dots = a_n = 0$. Hence the collection is linearly independent, so it is a basis for V . $\square$

Example 8 (The standard basis of $\mathbb{F}^n$). For $i = 1, \dots, n$, let $e_i \in \mathbb{F}^n$ be the vector whose i th coordinate is 1 and whose other coordinates are 0. Thus

$$e_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \quad \dots \quad e_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix}.$$

Every vector in $\mathbb{F}^n$ has the unique representation

$$\begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = x_1 e_1 + x_2 e_2 + \cdots + x_n e_n.$$

Consequently, $\{e_1, \dots, e_n\}$ is a basis for $\mathbb{F}^n$, called its standard basis.

Example 9 (The standard basis of $\mathcal{P}_n(\mathbb{F})$). Recall that $\mathcal{P}_n(\mathbb{F})$ is the vector space of polynomials over $\mathbb{F}$ of degree at most n . Each polynomial has a unique expression

$$p(t) = a_0 + a_1 t + \cdots + a_n t^n.$$

Hence

$$\{1, t, t^2, \dots, t^n\}$$

is a basis for $\mathcal{P}_n(\mathbb{F})$, called its standard basis.

Proposition 8. Every finite nonempty spanning set for a vector space contains a basis.

Proof. Let $\{v_1, \dots, v_p\}$ span V . If this collection is linearly independent, then it is already a basis. Otherwise, by Proposition 6, some v_k is a linear combination of the other vectors. It follows that removing v_k does not change the span:

$$\text{span}\{v_1, \dots, v_p\} = \text{span}\{v_1, \dots, v_{k-1}, v_{k+1}, \dots, v_p\}.$$

If the remaining collection is linearly independent, it is a basis. If not, repeat the same argument. The original spanning set is finite, so this process must terminate. The final collection still spans V and is linearly independent, and is therefore a basis contained in the original spanning set. $\square$

4.2. Linear transformations. A *transformation*, also called a map or function, from a set X to a set Y is a rule $T: X \rightarrow Y$ that assigns to every $x \in X$ an element $T(x) \in Y$.

Definition 11 (Linear transformation). *Let V and W be vector spaces over the same field $\mathbb{F}$. A transformation $T: V \rightarrow W$ is linear if*

- (i) $T(u + v) = T(u) + T(v)$ for all $u, v \in V$;
- (ii) $T(\alpha u) = \alpha T(u)$ for all $u \in V$ and $\alpha \in \mathbb{F}$.

Equivalently, $T: V \rightarrow W$ is linear if

$$T(\alpha u + \beta v) = \alpha T(u) + \beta T(v)$$

for all $u, v \in V$ and $\alpha, \beta \in \mathbb{F}$.

Example 10 (Reflection). *Define $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by*

$$T \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} x_1 \\ -x_2 \end{bmatrix}.$$

Geometrically, T reflects each point across the first coordinate axis. To check additivity, observe that

$$\begin{aligned} T \left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} \right) &= T \begin{bmatrix} x_1 + y_1 \\ x_2 + y_2 \end{bmatrix} \\ &= \begin{bmatrix} x_1 + y_1 \\ -x_2 - y_2 \end{bmatrix} \\ &= \begin{bmatrix} x_1 \\ -x_2 \end{bmatrix} + \begin{bmatrix} y_1 \\ -y_2 \end{bmatrix} \\ &= T \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + T \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}. \end{aligned}$$

For $\alpha \in \mathbb{R}$,

$$\begin{aligned} T \left(\alpha \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \right) &= T \begin{bmatrix} \alpha x_1 \\ \alpha x_2 \end{bmatrix} \\ &= \begin{bmatrix} \alpha x_1 \\ -\alpha x_2 \end{bmatrix} \\ &= \alpha \begin{bmatrix} x_1 \\ -x_2 \end{bmatrix} = \alpha T \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}. \end{aligned}$$

Thus T is linear.

Example 11 (The differentiation operator). *Let $n \geq 1$, $V = \mathcal{P}_n(\mathbb{F})$, and $W = \mathcal{P}_{n-1}(\mathbb{F})$. Define*

$$T: V \longrightarrow W, \quad T(p) = p'.$$

Since

$$(p + q)' = p' + q' \quad \text{and} \quad (\alpha p)' = \alpha p',$$

the differentiation operator T is linear.

5. SEPTEMBER 2, 2026

5.1. Linear transformations determined by a basis. The following result shows that prescribing the values of a linear transformation on a basis completely determines the transformation.

Lemma 2 (Linear map lemma). *Suppose that $\{v_1, \dots, v_n\}$ is a basis for V , and let $w_1, \dots, w_n$ be arbitrary vectors in W . Then there exists a unique linear transformation $T: V \rightarrow W$ such that*

$$T(v_k) = w_k$$

for each $k = 1, \dots, n$.

Proof. Every $v \in V$ has a unique representation

$$v = \sum_{i=1}^n c_i v_i.$$

Define

$$T(v) = \sum_{i=1}^n c_i w_i.$$

The uniqueness of the coordinates $c_1, \dots, c_n$ shows that this definition is well-defined. In particular, taking $v = v_k$ gives $T(v_k) = w_k$.

To prove that T is linear, write

$$u = \sum_{i=1}^n a_i v_i \quad \text{and} \quad v = \sum_{i=1}^n c_i v_i.$$

Then

$$\begin{aligned}
 T(u + v) &= T\left(\sum_{i=1}^n (a_i + c_i)v_i\right) \\
 &= \sum_{i=1}^n (a_i + c_i)w_i \\
 &= \sum_{i=1}^n a_i w_i + \sum_{i=1}^n c_i w_i \\
 &= T(u) + T(v).
 \end{aligned}$$

Similarly, for every $\alpha \in \mathbb{F}$,

$$\begin{aligned}
 T(\alpha v) &= T\left(\sum_{i=1}^n \alpha c_i v_i\right) \\
 &= \sum_{i=1}^n \alpha c_i w_i = \alpha \sum_{i=1}^n c_i w_i = \alpha T(v).
 \end{aligned}$$

Thus T is linear.

It remains to prove uniqueness. Suppose that $T': V \rightarrow W$ is another linear transformation satisfying $T'(v_k) = w_k$ for every k . If $v = \sum_{i=1}^n c_i v_i$, then

$$T'(v) = \sum_{i=1}^n c_i T'(v_i) = \sum_{i=1}^n c_i w_i = T(v).$$

Hence $T' = T$. □

5.2. Matrix representations of linear transformations. An *ordered basis* is a basis whose vectors are listed in a fixed order. Let

$$\mathcal{B} = (v_1, \dots, v_n) \quad \text{and} \quad \mathcal{C} = (w_1, \dots, w_m)$$

be ordered bases for V and W , respectively. For $v = x_1 v_1 + \dots + x_n v_n$, write

$$[v]_{\mathcal{B}} = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$$

for the coordinate vector of v with respect to $\mathcal{B}$.

$$A = [A_{ij}] = \begin{bmatrix} A_{11} & \cdots & A_{1n} \\ \vdots & & \vdots \\ A_{m1} & \cdots & A_{mn} \end{bmatrix} \in M_{m \times n}(\mathbb{F})$$
$$T(v_j) = \sum_{i=1}^m A_{ij} w_i$$

Proof. Let $T: V \rightarrow W$ be linear. Because $\mathcal{C}$ is a basis for W , each vector $T(v_j)$ has a unique representation

$$T(v_j) = A_{1j}w_1 + \cdots + A_{mj}w_m = \sum_{i=1}^m A_{ij}w_i.$$

$$A = \begin{bmatrix} | & & | \\ [T(v_1)]_{\mathcal{C}} & \cdots & [T(v_n)]_{\mathcal{C}} \\ | & & | \end{bmatrix}.$$
$$\sum_{i=1}^m B_{ij} w_i = T(v_j) = \sum_{i=1}^m A_{ij} w_i.$$

Conversely, let $A = [A_{ij}]$ be an arbitrary $m \times n$ matrix and define

$$\tilde{w}_j = \sum_{i=1}^m A_{ij} w_i \quad (j = 1, \dots, n).$$

By the linear map lemma, there is a unique linear transformation $T: V \rightarrow W$ such that $T(v_j) = \tilde{w}_j$ for every j . Thus A determines a unique linear transformation with the required property. $\square$

If

$$v = x_1 v_1 + \cdots + x_n v_n,$$

then linearity gives

$$\begin{aligned} T(v) &= \sum_{j=1}^n x_j T(v_j) \\ &= \sum_{j=1}^n x_j \sum_{i=1}^m A_{ij} w_i \\ &= \sum_{i=1}^m \left(\sum_{j=1}^n A_{ij} x_j \right) w_i. \end{aligned}$$

Consequently,

$$[T(v)]_{\mathcal{C}} = A[v]_{\mathcal{B}}.$$

Thus multiplication by the matrix A computes the coordinates of the image of a vector from the coordinates of the original vector.

6. SEPTEMBER 4, 2026

6.1. Matrix–vector multiplication. Let V and W be vector spaces over $\mathbb{F}$ with ordered bases

$$\mathcal{B} = (v_1, \dots, v_n) \quad \text{and} \quad \mathcal{C} = (w_1, \dots, w_m),$$

respectively, and let $T: V \rightarrow W$ be linear. Recall that the matrix representing T with respect to these bases is

$$A = \begin{bmatrix} A_{11} & \cdots & A_{1n} \\ \vdots & & \vdots \\ A_{m1} & \cdots & A_{mn} \end{bmatrix},$$

where

$$T(v_j) = A_{1j} w_1 + \cdots + A_{mj} w_m.$$

Thus the j th column of A is the coordinate vector $[T(v_j)]_{\mathcal{C}}$.

Suppose that

$$v = x_1 v_1 + \cdots + x_n v_n, \quad [v]_{\mathcal{B}} = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}.$$

By linearity,

$$\begin{aligned} [T(v)]_{\mathcal{C}} &= x_1 [T(v_1)]_{\mathcal{C}} + \cdots + x_n [T(v_n)]_{\mathcal{C}} \\ &= x_1 \begin{bmatrix} A_{11} \\ A_{21} \\ \vdots \\ A_{m1} \end{bmatrix} + \cdots + x_n \begin{bmatrix} A_{1n} \\ A_{2n} \\ \vdots \\ A_{mn} \end{bmatrix} \\ &= \begin{bmatrix} A_{11}x_1 + \cdots + A_{1n}x_n \\ \vdots \\ A_{m1}x_1 + \cdots + A_{mn}x_n \end{bmatrix}. \end{aligned}$$

This motivates the following definition.

Definition 12 (Matrix–vector product). *Let $A = [A_{ij}] \in M_{m \times n}(\mathbb{F})$, and write its columns as $A_1, \dots, A_n$. For*

$$x = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} \in \mathbb{F}^n,$$

define

$$Ax = x_1 A_1 + \cdots + x_n A_n.$$

Equivalently, the k th coordinate of Ax is

$$(Ax)_k = \sum_{j=1}^n A_{kj} x_j, \quad k = 1, \dots, m.$$

With this notation, the coordinate formula for a linear transformation is

$$[T(v)]_{\mathcal{C}} = A[v]_{\mathcal{B}}.$$

Example 12. *Matrix–vector multiplication can be computed either as a linear combination of the columns or one coordinate at a time. For*

instance,

$$\begin{bmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = 1 \begin{bmatrix} 1 \\ 3 \end{bmatrix} + 2 \begin{bmatrix} 2 \\ 2 \end{bmatrix} + 3 \begin{bmatrix} 3 \\ 1 \end{bmatrix} = \begin{bmatrix} 14 \\ 10 \end{bmatrix},$$

while

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = \begin{bmatrix} 1 \cdot 1 + 2 \cdot 2 + 3 \cdot 3 \\ 4 \cdot 1 + 5 \cdot 2 + 6 \cdot 3 \end{bmatrix} = \begin{bmatrix} 14 \\ 32 \end{bmatrix}.$$

6.2. Finding matrix representations.

Example 13. Define $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$ by

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x + 2y \\ 2x - 5y \\ 7y \end{bmatrix}.$$

With respect to the standard bases, the columns of the representing matrix are

$$T(e_1) = \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix} \quad \text{and} \quad T(e_2) = \begin{bmatrix} 2 \\ -5 \\ 7 \end{bmatrix}.$$

Hence

$$A = \begin{bmatrix} 1 & 2 \\ 2 & -5 \\ 0 & 7 \end{bmatrix}.$$

Example 14 (Differentiation). Let $n \geq 1$, and let

$$T: \mathcal{P}_n(\mathbb{R}) \longrightarrow \mathcal{P}_n(\mathbb{R}), \quad T(p) = p',$$

and use the ordered basis

$$\mathcal{B} = (1, t, t^2, \dots, t^n)$$

for both the domain and codomain. Since

$$T(1) = 0, \quad T(t) = 1, \quad T(t^j) = jt^{j-1},$$

the matrix representing T is

$$A = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 2 & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & 0 & n \\ 0 & \cdots & \cdots & 0 & 0 \end{bmatrix}.$$

6.3. The vector space of linear maps. Let V and W be vector spaces over $\mathbb{F}$. If $T: V \rightarrow W$ is linear and $\alpha \in \mathbb{F}$, define $\alpha T: V \rightarrow W$ pointwise by

$$(\alpha T)(v) = \alpha T(v).$$

This map is again linear. Indeed, for $u, v \in V$,

$$\begin{aligned} (\alpha T)(u + v) &= \alpha T(u + v) \\ &= \alpha(T(u) + T(v)) \\ &= (\alpha T)(u) + (\alpha T)(v), \end{aligned}$$

and, for $\beta \in \mathbb{F}$,

$$(\alpha T)(\beta u) = \alpha T(\beta u) = \alpha \beta T(u) = \beta(\alpha T)(u).$$

Similarly, if $T_1, T_2: V \rightarrow W$ are linear, define their sum by

$$(T_1 + T_2)(v) = T_1(v) + T_2(v).$$

The map $T_1 + T_2$ is linear because

$$\begin{aligned} (T_1 + T_2)(u + v) &= T_1(u + v) + T_2(u + v) \\ &= (T_1 + T_2)(u) + (T_1 + T_2)(v) \end{aligned}$$

and

$$(T_1 + T_2)(\alpha u) = \alpha(T_1 + T_2)(u).$$

Definition 13 (Space of linear maps). *The collection of all linear transformations from V to W is denoted by*

$$\mathcal{L}(V, W) = \{T: V \rightarrow W : T \text{ is linear}\}.$$

Proposition 9. *The set $\mathcal{L}(V, W)$, equipped with pointwise addition and scalar multiplication, is a vector space over $\mathbb{F}$.*

Proof. The preceding calculations show that the two operations are closed in $\mathcal{L}(V, W)$. All vector-space identities follow pointwise from the corresponding identities in W . The additive identity is the zero transformation $0: V \rightarrow W$, defined by $0(v) = 0$, and the additive inverse of T is the map $-T$, defined by $(-T)(v) = -T(v)$. $\square$

6.4. Composition of linear maps. Let V , W , and Z be vector spaces over $\mathbb{F}$. Suppose that

$$T_2: V \longrightarrow W \quad \text{and} \quad T_1: W \longrightarrow Z$$

are linear. Their *composition* is the map

$$T_1 \circ T_2: V \longrightarrow Z, \quad (T_1 \circ T_2)(v) = T_1(T_2(v)).$$

Proposition 10. *The composition $T_1 \circ T_2$ is linear.*

Proof. For $u, v \in V$ and $\alpha \in \mathbb{F}$,

$$\begin{aligned} (T_1 \circ T_2)(u + v) &= T_1(T_2(u) + T_2(v)) \\ &= T_1(T_2(u)) + T_1(T_2(v)), \end{aligned}$$

and

$$(T_1 \circ T_2)(\alpha u) = T_1(\alpha T_2(u)) = \alpha T_1(T_2(u)).$$

$\square$

7. SEPTEMBER 9, 2026

7.1. Matrix operations and linear maps. Let V and W be vector spaces over $\mathbb{F}$ with ordered bases

$$\mathcal{B} = (v_1, \dots, v_n) \quad \text{and} \quad \mathcal{C} = (w_1, \dots, w_m).$$

Suppose that $T, S \in \mathcal{L}(V, W)$ are represented by the matrices $A = [A_{ij}]$ and $B = [B_{ij}]$, respectively. Thus

$$T(v_j) = \sum_{i=1}^m A_{ij} w_i \quad \text{and} \quad S(v_j) = \sum_{i=1}^m B_{ij} w_i.$$

For each basis vector v_j ,

$$\begin{aligned}(T + S)(v_j) &= T(v_j) + S(v_j) \\ &= \sum_{i=1}^m A_{ij}w_i + \sum_{i=1}^m B_{ij}w_i \\ &= \sum_{i=1}^m (A_{ij} + B_{ij})w_i.\end{aligned}$$

Consequently, the matrix representing $T + S$ is obtained by adding the matrices representing T and S entrywise.

Definition 14 (Matrix addition). *If $A, B \in M_{m \times n}(\mathbb{F})$, their sum is the matrix $A + B \in M_{m \times n}(\mathbb{F})$ defined by*

$$(A + B)_{ij} = A_{ij} + B_{ij}.$$

Similarly, if $\alpha \in \mathbb{F}$, then

$$(\alpha T)(v_j) = \alpha T(v_j) = \sum_{i=1}^m \alpha A_{ij}w_i.$$

Thus scalar multiplication of linear maps corresponds to entrywise scalar multiplication of matrices.

Definition 15 (Scalar multiplication of matrices). *If $A \in M_{m \times n}(\mathbb{F})$ and $\alpha \in \mathbb{F}$, then $\alpha A \in M_{m \times n}(\mathbb{F})$ is defined by*

$$(\alpha A)_{ij} = \alpha A_{ij}.$$

7.2. Composition and matrix multiplication. Let Z be a third vector space with ordered basis

$$\mathcal{D} = (z_1, \dots, z_p),$$

and consider linear maps

$$T_2: V \longrightarrow W \quad \text{and} \quad T_1: W \longrightarrow Z.$$

Suppose T_2 is represented by $B = [B_{kj}] \in M_{m \times n}(\mathbb{F})$ and T_1 is represented by $A = [A_{ik}] \in M_{p \times m}(\mathbb{F})$. For $j = 1, \dots, n$,

$$\begin{aligned} (T_1 \circ T_2)(v_j) &= T_1 \left(\sum_{k=1}^m B_{kj} w_k \right) \\ &= \sum_{k=1}^m B_{kj} T_1(w_k) \\ &= \sum_{k=1}^m B_{kj} \sum_{i=1}^p A_{ik} z_i \\ &= \sum_{i=1}^p \left(\sum_{k=1}^m A_{ik} B_{kj} \right) z_i. \end{aligned}$$

The coefficient of z_i in this expression is the (i, j) entry of the matrix representing $T_1 \circ T_2$.

Definition 16 (Matrix multiplication). *If $A \in M_{p \times m}(\mathbb{F})$ and $B \in M_{m \times n}(\mathbb{F})$, their product is the matrix $AB \in M_{p \times n}(\mathbb{F})$ whose entries are*

$$(AB)_{ij} = \sum_{k=1}^m A_{ik} B_{kj}.$$

Thus composition of linear maps corresponds to multiplication of their representing matrices, in the same order:

$$[T_1 \circ T_2] = [T_1][T_2].$$

Whenever all the products and sums involved are defined, matrix multiplication satisfies

$$\begin{aligned} (AB)C &= A(BC), \\ A(B + C) &= AB + AC, \\ (A + B)C &= AC + BC, \\ A(\alpha B) &= \alpha(AB) = (\alpha A)B. \end{aligned}$$

In general, however, $AB \neq BA$; the two products may even have different sizes, or one may be undefined. In particular, $M_{n \times n}(\mathbb{F})$, with

matrix addition, scalar multiplication, and matrix multiplication, is an associative algebra over $\mathbb{F}$.

7.3. Vector subspaces.

Definition 17 (Subspace). *Let V be a vector space over $\mathbb{F}$. A subset $U \subseteq V$ is a subspace of V if U , with the addition and scalar multiplication inherited from V , is itself a vector space over $\mathbb{F}$.*

Checking all the vector-space axioms is unnecessary. The following criterion reduces the verification to three conditions.

Proposition 11 (Subspace criterion). *A subset $U \subseteq V$ is a subspace of V if and only if*

- (i) $0 \in U$;
- (ii) $u + v \in U$ whenever $u, v \in U$;
- (iii) $\alpha u \in U$ whenever $u \in U$ and $\alpha \in \mathbb{F}$.

Equivalently, if U is known to be nonempty, it is enough to check the two closure conditions (ii) and (iii).

Example 15. *The affine line*

$$H = \left\{ \begin{bmatrix} x \\ y \end{bmatrix} \in \mathbb{R}^2 : x = 2 \right\}$$

is not a subspace of $\mathbb{R}^2$, because it does not contain the zero vector.

The subspaces of $\mathbb{R}^2$ are precisely the zero subspace $\{0\}$, the lines through the origin, and $\mathbb{R}^2$ itself.

8. SEPTEMBER 11, 2026

8.1. The subspace criterion.

Proof of Proposition 11. If U is a subspace of V , then U is a vector space under the operations inherited from V . It therefore contains the additive identity and is closed under addition and scalar multiplication.

Conversely, suppose conditions (i)–(iii) hold. The additive identity belongs to U by (i). If $u \in U$, then condition (iii), applied with the

scalar -1 , gives

$$-u = (-1)u \in U.$$

Hence U contains the additive inverse of each of its elements. The remaining vector-space axioms are identities already valid in V , so they remain valid when the vectors are restricted to U . Thus U is a subspace of V .

Finally, if U is nonempty and satisfies the two closure conditions, choose $u \in U$. Closure under scalar multiplication gives $0 = 0u \in U$, so condition (i) follows automatically. $\square$

8.2. Kernels and images.

Definition 18 (Kernel and image). *Let $T \in \mathcal{L}(V, W)$. The kernel and image of T are*

$$\ker T = \{v \in V : T(v) = 0\}$$

and

$$\operatorname{im} T = \{T(v) : v \in V\},$$

respectively. The image is also called the range of T .

Proposition 12. *If $T \in \mathcal{L}(V, W)$, then $\ker T$ is a subspace of V and $\operatorname{im} T$ is a subspace of W .*

Proof. Since $T(0) = 0$, both sets contain the appropriate zero vector. If $u, v \in \ker T$ and $\alpha \in \mathbb{F}$, then

$$T(u + v) = T(u) + T(v) = 0 \quad \text{and} \quad T(\alpha u) = \alpha T(u) = 0.$$

Thus $\ker T$ is a subspace. If $y_1 = T(v_1)$ and $y_2 = T(v_2)$ are in $\operatorname{im} T$, then

$$y_1 + y_2 = T(v_1 + v_2) \quad \text{and} \quad \alpha y_1 = T(\alpha v_1),$$

so the image is also a subspace. $\square$

Example 16. *The continuous real-valued functions on $[0, 1]$ form a subspace*

$$C([0, 1]) \subseteq \{f : [0, 1] \rightarrow \mathbb{R}\}$$

of the vector space of all real-valued functions on $[0, 1]$.

Example 17. Fix $b \in \mathbb{F}$ and let

$$U_b = \left\{ \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} \in \mathbb{F}^4 : x_3 = 5x_4 + b \right\}.$$

Then U_b is a subspace of $\mathbb{F}^4$ if and only if $b = 0$. Indeed, the zero vector belongs to U_b only when $b = 0$, and when $b = 0$ the defining equation $x_3 = 5x_4$ is preserved by addition and scalar multiplication.

8.3. Sums of subspaces.

Definition 19 (Sum of subspaces). If $V_1, \dots, V_m$ are subspaces of V , their sum is

$$V_1 + \dots + V_m = \{v_1 + \dots + v_m : v_i \in V_i \text{ for } i = 1, \dots, m\}.$$

Example 18. Let

$$U = \left\{ \begin{bmatrix} x \\ 0 \\ 0 \end{bmatrix} : x \in \mathbb{F} \right\}, \quad W = \left\{ \begin{bmatrix} 0 \\ y \\ 0 \end{bmatrix} : y \in \mathbb{F} \right\}.$$

Then

$$U + W = \left\{ \begin{bmatrix} x \\ y \\ 0 \end{bmatrix} : x, y \in \mathbb{F} \right\}.$$

Proposition 13. If $V_1, \dots, V_m$ are subspaces of V , then $V_1 + \dots + V_m$ is the smallest subspace of V containing each V_i .

Proof. The sum contains $0 = 0 + \dots + 0$. Adding two elements of the sum, or multiplying one by a scalar, can be done componentwise inside each V_i , so the sum is a subspace. For each i , any $v_i \in V_i$ can be written as

$$v_i = 0 + \dots + 0 + v_i + 0 + \dots + 0,$$

and hence $V_i \subseteq V_1 + \dots + V_m$.

Conversely, let W be any subspace of V containing all the V_i . If $v = v_1 + \dots + v_m$ with $v_i \in V_i$, then every v_i belongs to W . Since W is closed under addition, $v \in W$. Therefore $V_1 + \dots + V_m \subseteq W$. $\square$

8.4. Direct sums.

Definition 20 (Direct sum). *Let $V_1, \dots, V_m$ be subspaces of V . Their sum is a direct sum if every element of $V_1 + \dots + V_m$ has a unique representation*

$$v = v_1 + \dots + v_m, \quad v_i \in V_i.$$

In this case we write

$$V_1 \oplus \dots \oplus V_m.$$

Proposition 14 (Zero-sum criterion). *The sum $V_1 + \dots + V_m$ is direct if and only if the equation*

$$0 = v_1 + \dots + v_m, \quad v_i \in V_i,$$

implies $v_1 = \dots = v_m = 0$.

Proof. If the sum is direct, then $0 = 0 + \dots + 0$ is one representation of the zero vector, so uniqueness forces every other representation to have all components equal to zero.

Conversely, suppose the zero vector has only the trivial representation. If

$$v_1 + \dots + v_m = u_1 + \dots + u_m,$$

where $v_i, u_i \in V_i$, then

$$0 = (v_1 - u_1) + \dots + (v_m - u_m).$$

The hypothesis gives $v_i - u_i = 0$ for each i , so $v_i = u_i$. Thus every representation is unique. $\square$

Proposition 15. *If U and W are subspaces of V , then*

$$U + W = U \oplus W \iff U \cap W = \{0\}.$$

Proof. Suppose first that $U + W$ is a direct sum, and let $v \in U \cap W$. Then

$$0 = v + (-v) = 0 + 0$$

gives two representations with one term in U and one term in W . Uniqueness implies $v = 0$.

Conversely, assume $U \cap W = \{0\}$. If

$$u + w = u' + w', \quad u, u' \in U, \quad w, w' \in W,$$

then

$$u - u' = w' - w \in U \cap W.$$

Hence $u = u'$ and $w = w'$, so the representation is unique. $\square$

For the coordinate-axis subspaces in the preceding example, $U \cap W = \{0\}$, and hence their sum is $U \oplus W$.

8.5. Product vector spaces.

Definition 21 (Product vector space). *Let $V_1, \dots, V_m$ be vector spaces over $\mathbb{F}$. Their Cartesian product is*

$$V_1 \times \cdots \times V_m = \{(v_1, \dots, v_m) : v_i \in V_i\}.$$

Addition and scalar multiplication are defined coordinatewise:

$$\begin{aligned} (v_1, \dots, v_m) + (u_1, \dots, u_m) &= (v_1 + u_1, \dots, v_m + u_m), \\ \alpha(v_1, \dots, v_m) &= (\alpha v_1, \dots, \alpha v_m). \end{aligned}$$

With these operations, $V_1 \times \cdots \times V_m$ is a vector space over $\mathbb{F}$; all vector-space axioms hold coordinatewise.

Example 19. *In $\mathcal{P}_5(\mathbb{R}) \times \mathbb{R}^3$,*

$$\begin{aligned} \left(5 - 6t + 4t^2, \begin{bmatrix} 3 \\ 8 \\ 7 \end{bmatrix} \right) + \left(t + 9t^5, \begin{bmatrix} 2 \\ 2 \\ 2 \end{bmatrix} \right) \\ = \left(5 - 5t + 4t^2 + 9t^5, \begin{bmatrix} 5 \\ 10 \\ 9 \end{bmatrix} \right). \end{aligned}$$

9. SEPTEMBER 14, 2026

9.1. Direct sums and injectivity. Let $V_1, \dots, V_m$ be subspaces of a vector space V . Define the summation map

$$\Sigma: V_1 \times \cdots \times V_m \longrightarrow V_1 + \cdots + V_m$$

by

$$\Sigma(v_1, \dots, v_m) = v_1 + \dots + v_m.$$

This map is linear and surjective.

Proposition 16. *The sum $V_1 + \dots + V_m$ is direct if and only if the summation map Σ is injective.*

Proof. The map Σ is injective precisely when

$$\Sigma(v_1, \dots, v_m) = \Sigma(u_1, \dots, u_m)$$

implies

$$(v_1, \dots, v_m) = (u_1, \dots, u_m).$$

Equivalently, every vector in $V_1 + \dots + V_m$ has only one representation as a sum of vectors from the respective subspaces. This is exactly the definition of a direct sum. $\square$

9.2. Quotient vector spaces. Let U be a subset of V . For $v \in V$, define

$$v + U = \{v + u : u \in U\}.$$

The set $v + U$ is called a *translate* of U . When U is a subspace, a translate is also called a *coset* of U .

Example 20. *Let*

$$U = \left\{ \begin{bmatrix} x \\ 2x \end{bmatrix} : x \in \mathbb{R} \right\} \subseteq \mathbb{R}^2.$$

Then

$$\begin{bmatrix} 2 \\ 1 \end{bmatrix} + U = \left\{ \begin{bmatrix} 2+x \\ 1+2x \end{bmatrix} : x \in \mathbb{R} \right\}.$$

The original subspace is the line $y = 2x$, while this translate is the parallel line $y = 2x - 3$.

Definition 22 (Quotient space). *If U is a subspace of V , the quotient space of V by U is the set of all cosets of U :*

$$V/U = \{v + U : v \in V\}.$$

For the subspace $U \subseteq \mathbb{R}^2$ in the preceding example, $\mathbb{R}^2/U$ is the collection of all lines in $\mathbb{R}^2$ with slope 2.

Proposition 17. *Let U be a subspace of V and let $v, w \in V$. The following are equivalent:*

- (i) $v - w \in U$;
- (ii) $v + U = w + U$;
- (iii) $(v + U) \cap (w + U) \neq \emptyset$.

Proof. Suppose first that $v - w \in U$. If $v + u \in v + U$, then

$$v + u = w + (v - w + u) \in w + U,$$

because $v - w + u \in U$. Thus $v + U \subseteq w + U$. Since $w - v = -(v - w) \in U$, the reverse inclusion follows in the same way, so $v + U = w + U$.

If $v + U = w + U$, then the two cosets plainly have nonempty intersection.

Finally, suppose $(v + U) \cap (w + U) \neq \emptyset$. There are $u_1, u_2 \in U$ such that

$$v + u_1 = w + u_2.$$

Therefore

$$v - w = u_2 - u_1 \in U.$$

□

Thus two cosets of a subspace are either equal or disjoint. We define addition and scalar multiplication on V/U by

$$\begin{aligned} (v + U) + (w + U) &= (v + w) + U, \\ \alpha(v + U) &= \alpha v + U. \end{aligned}$$

Proposition 17 shows that these operations do not depend on the choice of representatives. With these operations, V/U is a vector space over $\mathbb{F}$, whose zero vector is the coset $U = 0 + U$.

9.3. Finite-dimensional vector spaces.

Definition 23 (Finite-dimensional vector space). *A vector space V is finite-dimensional if some finite collection of vectors spans V . It is infinite-dimensional if it is not finite-dimensional.*

Example 21. *The coordinate space $\mathbb{F}^n$ is finite-dimensional because*

$$\mathbb{F}^n = \text{span}\{e_1, \dots, e_n\}.$$

Likewise, the space of polynomials of degree at most m is finite-dimensional because

$$\mathcal{P}_m(\mathbb{F}) = \text{span}\{1, t, t^2, \dots, t^m\}.$$

Example 22. *The vector space $\mathcal{P}(\mathbb{F})$ of all polynomials with coefficients in $\mathbb{F}$ is infinite-dimensional. Indeed, suppose a finite collection of polynomials spanned $\mathcal{P}(\mathbb{F})$, and let m be the largest degree among the nonzero polynomials in the collection. Every polynomial in their span would have degree at most m , so t^{m+1} could not belong to the span, a contradiction.*

10. SEPTEMBER 16, 2026

10.1. Comparing independent and spanning collections.

Proposition 18. *Suppose that $u_1, \dots, u_m$ is a linearly independent collection in V and that $w_1, \dots, w_n$ spans V . Then*

$$m \leq n.$$

Thus, in a finite-dimensional vector space, every linearly independent collection has cardinality at most that of every finite spanning collection.

Proof. We successively insert the vectors $u_1, u_2, \dots$ into the spanning list $w_1, \dots, w_n$, removing one of the w 's after each insertion.

First adjoin u_1 to obtain

$$u_1, w_1, \dots, w_n.$$

This collection is linearly dependent because $w_1, \dots, w_n$ spans V . Since $u_1 \neq 0$, one of the w 's is a linear combination of the other vectors.

Remove that vector. The resulting collection still spans V and has n vectors.

Inductively, suppose that after $k - 1$ steps we have a spanning collection of n vectors consisting of $u_1, \dots, u_{k-1}$ and $n - k + 1$ of the original w 's. Adjoin u_k . The resulting collection is dependent. In a nontrivial dependence relation, some remaining w must have a nonzero coefficient; otherwise the relation would contradict the linear independence of $u_1, \dots, u_k$. We can therefore remove that w without changing the span.

If $m > n$, then after n steps the collection $u_1, \dots, u_n$ would span V . In particular, u_{n+1} would lie in their span, contradicting the linear independence of $u_1, \dots, u_m$. Hence $m \leq n$. $\square$

Example 23. *No collection of four vectors in $\mathbb{R}^3$ is linearly independent, because the three standard basis vectors span $\mathbb{R}^3$. Similarly, no collection of three vectors can span $\mathbb{R}^4$, because the four standard basis vectors are linearly independent in $\mathbb{R}^4$.*

10.2. Subspaces of finite-dimensional spaces.

Proposition 19. *Every subspace of a finite-dimensional vector space is finite-dimensional.*

Proof. Let U be a subspace of a finite-dimensional vector space V . If $U = \{0\}$, then the empty collection spans U , so U is finite-dimensional.

Otherwise choose a nonzero vector $u_1 \in U$. If $U = \text{span}\{u_1\}$, we are done. If not, choose

$$u_2 \in U \setminus \text{span}\{u_1\}.$$

Continue in this fashion: once $u_1, \dots, u_{k-1}$ have been chosen, stop if they span U ; otherwise choose

$$u_k \in U \setminus \text{span}\{u_1, \dots, u_{k-1}\}.$$

At every stage the chosen collection is linearly independent. Since V has a finite spanning collection, Proposition 18 bounds the number of vectors in any linearly independent collection in V . The process

must therefore stop after finitely many steps, at which point the chosen vectors span U . $\square$

Corollary 1. *Every finite-dimensional vector space has a basis.*

Proof. By definition, a finite-dimensional vector space has a finite spanning collection. Removing redundant vectors from that collection produces a basis. $\square$

10.3. Extending a linearly independent collection.

Proposition 20 (Basis extension theorem). *Every finite linearly independent collection in a finite-dimensional vector space can be extended to a basis of the space.*

Proof. Let $u_1, \dots, u_m$ be linearly independent in V , and choose a finite spanning collection $w_1, \dots, w_n$ for V . The combined collection

$$u_1, \dots, u_m, w_1, \dots, w_n$$

spans V . Examine the w 's in order. Whenever a vector w_j lies in the span of the vectors preceding it, remove it. Removing such a vector does not change the span, and none of the u_i is removed. After all the w 's have been considered, the remaining collection spans V and is linearly independent. It is therefore a basis containing $u_1, \dots, u_m$. $\square$

Example 24. *Consider the linearly independent vectors*

$$v_1 = \begin{bmatrix} -2 \\ 3 \\ 4 \end{bmatrix}, \quad v_2 = \begin{bmatrix} -9 \\ 6 \\ 8 \end{bmatrix}$$

in $\mathbb{R}^3$. Adjoin the standard basis vectors e_1, e_2, e_3 . Since

$$e_1 = -\frac{1}{5}(v_2 - 2v_1),$$

the vector e_1 is redundant. On the other hand, e_2 is not in $\text{span}\{v_1, v_2\}$, so we retain it. Finally,

$$e_3 = -\frac{1}{10} \left(v_2 - \frac{9}{2}v_1 + \frac{15}{2}e_2 \right),$$

so e_3 is redundant. Therefore

$$\left\{ \begin{bmatrix} -2 \\ 3 \\ 4 \end{bmatrix}, \begin{bmatrix} -9 \\ 6 \\ 8 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \right\}$$

is a basis of $\mathbb{R}^3$ extending the original collection.

11. SEPTEMBER 18, 2026

11.1. Complements of subspaces.

Proposition 21. *Suppose that V is finite-dimensional and that U is a subspace of V . Then there is a subspace W of V such that*

$$V = U \oplus W.$$

Proof. Because V is finite-dimensional, so is U . Let $u_1, \dots, u_m$ be a basis of U . This collection is linearly independent in V , so the basis extension theorem gives a basis

$$u_1, \dots, u_m, w_1, \dots, w_n$$

of V . Let

$$W = \text{span}\{w_1, \dots, w_n\}.$$

Certainly $U + W \subseteq V$. Conversely, if $v \in V$, then there are scalars $a_1, \dots, a_m, b_1, \dots, b_n$ such that

$$v = \underbrace{a_1u_1 + \dots + a_mu_m}_{\in U} + \underbrace{b_1w_1 + \dots + b_nw_n}_{\in W}.$$

Hence $v \in U + W$, and therefore $V = U + W$.

It remains to show that $U \cap W = \{0\}$. Let $v \in U \cap W$. Then

$$v = a_1u_1 + \dots + a_mu_m = b_1w_1 + \dots + b_nw_n$$

for some scalars a_i and b_j . Thus

$$a_1u_1 + \dots + a_mu_m - b_1w_1 - \dots - b_nw_n = 0.$$

Because the combined collection is a basis of V , all the coefficients are zero. In particular, $v = 0$, so $U \cap W = \{0\}$. Thus $V = U \oplus W$. $\square$

11.2. Dimension.

Proposition 22. *Any two bases of a finite-dimensional vector space have the same cardinality.*

Proof. Let $\mathcal{B}_1$ and $\mathcal{B}_2$ be bases of a finite-dimensional vector space V . Because $\mathcal{B}_1$ is linearly independent and $\mathcal{B}_2$ spans V , Proposition 18 gives

$$|\mathcal{B}_1| \leq |\mathcal{B}_2|.$$

Interchanging the roles of the two bases gives

$$|\mathcal{B}_2| \leq |\mathcal{B}_1|.$$

Therefore $|\mathcal{B}_1| = |\mathcal{B}_2|$. □

Definition 24 (Dimension). *The dimension of a finite-dimensional vector space V is the cardinality of any basis of V . It is denoted by $\dim V$.*

Example 25. *The standard basis $\{e_1, \dots, e_n\}$ of $\mathbb{F}^n$ has n elements, so*

$$\dim \mathbb{F}^n = n.$$

Example 26. *The standard basis $\{1, x, \dots, x^m\}$ of $\mathcal{P}_m(\mathbb{F})$ has $m + 1$ elements, so*

$$\dim \mathcal{P}_m(\mathbb{F}) = m + 1.$$

Example 27. *Let*

$$U = \left\{ \begin{bmatrix} x \\ 0 \\ y \end{bmatrix} : x, y \in \mathbb{F} \right\}.$$

Because

$$U = \text{span} \left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$$

and the two displayed vectors are linearly independent, they form a basis of U . Hence $\dim U = 2$.

Example 28. As a vector space over $\mathbb{R}$, the space $\mathbb{R}^2$ has dimension 2. As a vector space over $\mathbb{C}$, the space $\mathbb{C}$ has dimension 1. Although $\mathbb{R}^2$ and $\mathbb{C}$ are often identified as sets, the choice of scalar field is essential when determining dimension.

11.3. Consequences of dimension.

Proposition 23. If V is finite-dimensional and U is a subspace of V , then

$$\dim U \leq \dim V.$$

Proof. A basis of U is linearly independent in V , while a basis of V spans V . The result follows from Proposition 18. $\square$

Proposition 24. Suppose that V is finite-dimensional. Every linearly independent collection of $\dim V$ vectors in V is a basis of V .

Proof. Write $\dim V = n$, and suppose that $v_1, \dots, v_n$ is linearly independent in V . By the basis extension theorem, this collection can be extended to a basis of V . Every basis of V has exactly n vectors, however, so no vectors need to be adjoined. Thus $v_1, \dots, v_n$ is already a basis of V . $\square$

Corollary 2. Suppose that V is finite-dimensional and that U is a subspace of V . If $\dim U = \dim V$, then $U = V$.

Proof. Let $u_1, \dots, u_n$ be a basis of U . Then $n = \dim U = \dim V$, and $u_1, \dots, u_n$ is linearly independent in V . By Proposition 24, it is a basis of V . Hence every vector in V is a linear combination of vectors in U , so $V \subseteq U$. The reverse inclusion holds because U is a subspace of V . Therefore $U = V$. $\square$

Example 29. Let

$$U = \{p \in \mathcal{P}_3(\mathbb{R}) : p'(5) = 0\}.$$

We find a basis of U . Write

$$p(x) = a_0 + a_1x + a_2x^2 + a_3x^3.$$

Because

$$p'(x) = a_1 + 2a_2x + 3a_3x^2,$$

the condition $p'(5) = 0$ is equivalent to

$$a_1 + 10a_2 + 75a_3 = 0, \quad \text{or equivalently} \quad a_1 = -10a_2 - 75a_3.$$

Therefore every $p \in U$ has the form

$$\begin{aligned} p(x) &= a_0 + (-10a_2 - 75a_3)x + a_2x^2 + a_3x^3 \\ &= a_0 + a_2(x^2 - 10x) + a_3(x^3 - 75x). \end{aligned}$$

Thus the collection

$$\{1, x^2 - 10x, x^3 - 75x\}$$

spans U . It is linearly independent because its members have distinct degrees. Consequently,

$$\boxed{\{1, x^2 - 10x, x^3 - 75x\}}$$

is a basis of U , and $\dim U = 3$.

12. SEPTEMBER 21, 2026

12.1. Spanning collections and dimension.

Proposition 25. *Suppose that V is finite-dimensional. Every collection of $\dim V$ vectors that spans V is a basis of V .*

Proof. Write $\dim V = n$, and suppose that $v_1, \dots, v_n$ spans V . This collection can be reduced to a basis by removing redundant vectors. However, every basis of V has exactly n vectors, so no vectors can be removed. Thus $v_1, \dots, v_n$ is already a basis of V . $\square$

12.2. Dimension of a sum of subspaces.

Proposition 26. *If V_1 and V_2 are subspaces of a finite-dimensional vector space, then*

$$\dim(V_1 + V_2) = \dim V_1 + \dim V_2 - \dim(V_1 \cap V_2).$$

Proof. Let $v_1, \dots, v_m$ be a basis of $V_1 \cap V_2$, so that $\dim(V_1 \cap V_2) = m$. This collection is linearly independent in V_1 , so the basis extension

theorem gives a basis

$$v_1, \dots, v_m, u_1, \dots, u_j$$

of V_1 . Similarly, extend $v_1, \dots, v_m$ to a basis

$$v_1, \dots, v_m, w_1, \dots, w_k$$

of V_2 . Hence

$$\dim V_1 = m + j \quad \text{and} \quad \dim V_2 = m + k.$$

We will show that

$$v_1, \dots, v_m, u_1, \dots, u_j, w_1, \dots, w_k$$

is a basis of $V_1 + V_2$.

All the displayed vectors belong to $V_1 + V_2$, so their span is contained in $V_1 + V_2$. Conversely, every vector in $V_1 + V_2$ is a sum of a vector in V_1 and a vector in V_2 . Expressing each summand in the corresponding basis shows that the combined collection spans $V_1 + V_2$.

To prove linear independence, suppose that

$$(1) \quad a_1 v_1 + \dots + a_m v_m + b_1 u_1 + \dots + b_j u_j + c_1 w_1 + \dots + c_k w_k = 0.$$

Rearranging gives

$$c_1 w_1 + \dots + c_k w_k = -a_1 v_1 - \dots - a_m v_m - b_1 u_1 - \dots - b_j u_j.$$

The right-hand side belongs to V_1 , while the left-hand side belongs to V_2 . Therefore

$$c_1 w_1 + \dots + c_k w_k \in V_1 \cap V_2.$$

Since $v_1, \dots, v_m$ is a basis of the intersection, there are scalars $d_1, \dots, d_m$ such that

$$c_1 w_1 + \dots + c_k w_k = d_1 v_1 + \dots + d_m v_m.$$

Thus

$$c_1 w_1 + \dots + c_k w_k - d_1 v_1 - \dots - d_m v_m = 0.$$

The collection $v_1, \dots, v_m, w_1, \dots, w_k$ is a basis of V_2 and hence linearly independent. Consequently, all the c 's and d 's are zero. Equation (1)

now becomes

$$a_1v_1 + \cdots + a_mv_m + b_1u_1 + \cdots + b_ju_j = 0.$$

Because $v_1, \dots, v_m, u_1, \dots, u_j$ is a basis of V_1 , all the a 's and b 's are also zero. This proves linear independence.

The combined collection is therefore a basis of $V_1 + V_2$, and

$$\begin{aligned} \dim(V_1 + V_2) &= m + j + k \\ &= (m + j) + (m + k) - m \\ &= \dim V_1 + \dim V_2 - \dim(V_1 \cap V_2). \end{aligned}$$

□

12.3. Linear maps on finite-dimensional spaces.

Theorem 2 (Fundamental theorem of linear maps). *Suppose that V is finite-dimensional and that $T \in \mathcal{L}(V, W)$. Then $\text{im } T$ is finite-dimensional and*

$$\dim V = \dim \ker T + \dim \text{im } T.$$

Recall that

$$\ker T = \{v \in V : T(v) = 0\} \quad \text{and} \quad \text{im } T = \{T(v) : v \in V\}.$$

Before proving the theorem, we connect the kernel and image with injectivity and surjectivity, beginning with injectivity.

Proposition 27. *Let $T \in \mathcal{L}(V, W)$. Then T is injective if and only if $\ker T = \{0\}$.*

Proof. First note that every linear map sends zero to zero, since

$$T(0) = T(0 + 0) = T(0) + T(0)$$

implies $T(0) = 0$. This does not require injectivity. In particular, $\{0\} \subseteq \ker T$.

Suppose that T is injective. If $v \in \ker T$, then

$$T(v) = 0 = T(0).$$

Injectivity gives $v = 0$, so $\ker T \subseteq \{0\}$. Therefore $\ker T = \{0\}$.

Conversely, suppose that $\ker T = \{0\}$. Let $v_1, v_2 \in V$ satisfy $T(v_1) = T(v_2)$. By linearity,

$$T(v_1 - v_2) = T(v_1) - T(v_2) = 0.$$

Hence $v_1 - v_2 \in \ker T = \{0\}$, so $v_1 = v_2$. Thus T is injective. $\square$

13. SEPTEMBER 23, 2026

13.1. Injectivity, surjectivity, and dimension. Recall from Proposition 27 that a linear map $T \in \mathcal{L}(V, W)$ is injective if and only if $\ker T = \{0\}$. In particular, if $\ker T = \{0\}$, then

$$T(v) = T(v') \implies T(v - v') = 0 \implies v - v' = 0 \implies v = v'.$$

The corresponding criterion for surjectivity follows directly from the definition of the image:

$$T \text{ is surjective} \iff \operatorname{im} T = W.$$

13.2. Proof of the fundamental theorem of linear maps. We now prove Theorem 2: if V is finite-dimensional and $T \in \mathcal{L}(V, W)$, then $\operatorname{im} T$ is finite-dimensional and

$$\dim V = \dim \ker T + \dim \operatorname{im} T.$$

This identity is also called the *rank-nullity formula*; $\dim \operatorname{im} T$ is the rank and $\dim \ker T$ is the nullity.

Proof of Theorem 2. Since $\ker T$ is a subspace of the finite-dimensional space V , it has a basis $u_1, \dots, u_m$. Extend this collection to a basis

$$u_1, \dots, u_m, v_1, \dots, v_n$$

of V . Thus $\dim \ker T = m$ and $\dim V = m + n$. We claim that

$$T(v_1), \dots, T(v_n)$$

is a basis of $\operatorname{im} T$.

First, every $v \in V$ can be written as

$$v = a_1 u_1 + \dots + a_m u_m + b_1 v_1 + \dots + b_n v_n.$$

Because each u_i belongs to $\ker T$, linearity gives

$$T(v) = b_1T(v_1) + \cdots + b_nT(v_n).$$

Hence $\operatorname{im} T$ is contained in the span of the displayed images. The reverse inclusion holds because each $T(v_j)$ belongs to the subspace $\operatorname{im} T$. Therefore

$$\operatorname{im} T = \operatorname{span}\{T(v_1), \dots, T(v_n)\}.$$

To show linear independence, suppose that

$$c_1T(v_1) + \cdots + c_nT(v_n) = 0.$$

Then $T(c_1v_1 + \cdots + c_nv_n) = 0$, so $c_1v_1 + \cdots + c_nv_n \in \ker T$. There are scalars $d_1, \dots, d_m$ such that

$$c_1v_1 + \cdots + c_nv_n = d_1u_1 + \cdots + d_mu_m.$$

Rearranging gives

$$-d_1u_1 - \cdots - d_mu_m + c_1v_1 + \cdots + c_nv_n = 0.$$

The chosen basis of V is linearly independent, so all the coefficients are zero. In particular, $c_1 = \cdots = c_n = 0$.

Thus the image has a basis of n vectors, so $\dim \operatorname{im} T = n$. It follows that

$$\dim V = m + n = \dim \ker T + \dim \operatorname{im} T.$$

The same argument includes $m = 0$ or $n = 0$, using the empty basis for the zero subspace. $\square$




FIGURE 1. The kernel basis maps to zero; the images of the remaining basis vectors form a basis of the image.

13.3. Dimension restrictions on linear maps.

Corollary 3. *Let V and W be finite-dimensional vector spaces. If $\dim V > \dim W$, then no linear map from V to W is injective.*

Proof. If $T \in \mathcal{L}(V, W)$, then $\operatorname{im} T \subseteq W$, so $\dim \operatorname{im} T \leq \dim W$. Rank-nullity gives

$$\dim \ker T = \dim V - \dim \operatorname{im} T \geq \dim V - \dim W > 0.$$

Hence $\ker T \neq \{0\}$, and T is not injective. $\square$

Example 30. Define $T: \mathbb{R}^4 \rightarrow \mathbb{R}^3$ by

$$T \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} \sqrt{7}x_1 + \pi x_2 + x_4 \\ 97x_1 + 3x_2 + 2x_3 \\ x_2 + 6x_3 + 7x_4 \end{bmatrix}.$$

Since $4 > 3$, this map is not injective, regardless of the particular coefficients. There is no need to solve for its kernel explicitly. The same example works over $\mathbb{C}$.

Proposition 28. *Suppose that V and W are finite-dimensional and $\dim V < \dim W$. Then no linear map from V to W is surjective.*

Proof. For $T \in \mathcal{L}(V, W)$, rank-nullity gives

$$\dim \operatorname{im} T = \dim V - \dim \ker T \leq \dim V < \dim W.$$

Thus $\operatorname{im} T \neq W$, so T is not surjective. $\square$

13.4. A complement to the kernel.

Proposition 29. *Let V be finite-dimensional, and let $T \in \mathcal{L}(V, W)$. There is a subspace $U \subseteq V$ such that*

$$V = \ker T \oplus U \quad \text{and} \quad \operatorname{im} T = \{T(u) : u \in U\}.$$

In particular, $U \cap \ker T = \{0\}$, and the restriction $T|_U: U \rightarrow \operatorname{im} T$ is bijective.

Proof. As in the proof of rank–nullity, extend a basis $u_1, \dots, u_m$ of $\ker T$ to a basis $u_1, \dots, u_m, v_1, \dots, v_n$ of V . Set

$$U = \text{span}\{v_1, \dots, v_n\}.$$

The combined basis spans V , so $V = \ker T + U$. If $w \in U \cap \ker T$, then

$$w = b_1 v_1 + \dots + b_n v_n = a_1 u_1 + \dots + a_m u_m.$$

Linear independence of the combined basis forces every coefficient to vanish. Thus $w = 0$, and the sum is direct.

Every $v \in V$ has the form $v = k + u$, where $k \in \ker T$ and $u \in U$. Therefore $T(v) = T(u)$, proving that $T|_U$ maps onto $\text{im } T$. Its kernel is $U \cap \ker T = \{0\}$, so it is also injective. $\square$




FIGURE 2. For $T: \mathbb{R}^2 \rightarrow \mathbb{R}$, $T(x, y) = x$, take U to be the horizontal axis. The decomposition $v = k + u$ gives $T(v) = T(u) = a$, while $T(k) = 0$.

14. SEPTEMBER 25, 2026

14.1. Invertible linear maps. Write Id_V for the identity map on V , defined by $\text{Id}_V(v) = v$. As with matrix multiplication, we sometimes write ST for the composition $S \circ T$.

Definition 25 (Invertibility). *A linear map $T \in \mathcal{L}(V, W)$ is invertible if there exists a linear map $S \in \mathcal{L}(W, V)$ such that*

$$S \circ T = \text{Id}_V \quad \text{and} \quad T \circ S = \text{Id}_W.$$

Such a map S is called an inverse of T .

Proposition 30 (Uniqueness of the inverse). *An invertible linear map has a unique inverse.*

Proof. Suppose $S_1, S_2 \in \mathcal{L}(W, V)$ are both inverses of T . By associativity of composition,

$$\begin{aligned} S_1 &= S_1 \circ \text{Id}_W \\ &= S_1 \circ (T \circ S_2) \\ &= (S_1 \circ T) \circ S_2 \\ &= \text{Id}_V \circ S_2 = S_2. \end{aligned}$$

□

We denote the unique inverse by $T^{-1} \in \mathcal{L}(W, V)$. Thus

$$T^{-1}T = \text{Id}_V \quad \text{and} \quad TT^{-1} = \text{Id}_W.$$

Example 31. Define $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ by

$$T \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -y \\ x \\ 4z \end{bmatrix}.$$

Its values on the standard basis are

$$T(e_1) = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \quad T(e_2) = \begin{bmatrix} -1 \\ 0 \\ 0 \end{bmatrix}, \quad T(e_3) = \begin{bmatrix} 0 \\ 0 \\ 4 \end{bmatrix}.$$

Hence its matrix with respect to the standard basis is

$$A = \begin{bmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 4 \end{bmatrix}.$$

To recover the input from an output (x, y, z) , reverse the first two coordinates with the appropriate sign and divide the third coordinate by

4. *This gives*

$$T^{-1} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} y \\ -x \\ z/4 \end{bmatrix}, \quad A^{-1} = \begin{bmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & \frac{1}{4} \end{bmatrix}.$$

Indeed, direct multiplication gives $AA^{-1} = A^{-1}A = I_3$. Equivalently, augment A by the identity matrix and perform the row operations $R_1 \leftrightarrow R_2$, $R_2 \leftarrow -R_2$, and $R_3 \leftarrow \frac{1}{4}R_3$:

$$\left[\begin{array}{ccc|ccc} 0 & -1 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 4 & 0 & 0 & 1 \end{array} \right] \longrightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & \frac{1}{4} \end{array} \right].$$




FIGURE 3. In the xy -plane, T rotates by 90° counter-clockwise and T^{-1} rotates clockwise. In the third coordinate, T multiplies by 4 and T^{-1} divides by 4.

14.2. Invertibility and bijectivity.

Proposition 31. *A linear map is invertible if and only if it is bijective.*

Proof. Let $T \in \mathcal{L}(V, W)$. Suppose first that T is invertible. If $T(u) = T(v)$, applying T^{-1} gives

$$u = T^{-1}(T(u)) = T^{-1}(T(v)) = v.$$

Thus T is injective. For every $w \in W$, the vector $v = T^{-1}(w)$ satisfies

$$T(v) = T(T^{-1}(w)) = w,$$

so T is also surjective.

Conversely, suppose that T is bijective. For each $w \in W$, surjectivity provides $v \in V$ with $T(v) = w$, and injectivity makes this vector unique. Define $S: W \rightarrow V$ by assigning to w this unique vector. By construction, $T \circ S = \text{Id}_W$. Also, for $v \in V$,

$$T(S(T(v))) = T(v),$$

so injectivity gives $S(T(v)) = v$. Thus $S \circ T = \text{Id}_V$.

It remains to check that S is linear. For $w_1, w_2 \in W$,

$$\begin{aligned} T(S(w_1) + S(w_2)) &= T(S(w_1)) + T(S(w_2)) \\ &= w_1 + w_2 \\ &= T(S(w_1 + w_2)). \end{aligned}$$

Injectivity of T yields

$$S(w_1 + w_2) = S(w_1) + S(w_2).$$

Similarly, for $\lambda \in \mathbb{F}$ and $w \in W$,

$$T(\lambda S(w)) = \lambda T(S(w)) = \lambda w = T(S(\lambda w)),$$

so $S(\lambda w) = \lambda S(w)$. Therefore $S \in \mathcal{L}(W, V)$ and S is an inverse of T . $\square$

14.3. Why the dimension hypotheses matter. In general, injectivity alone or surjectivity alone does not imply invertibility.

Example 32. On the space $\mathcal{P}(\mathbb{R})$ of all real polynomials, define

$$M: \mathcal{P}(\mathbb{R}) \rightarrow \mathcal{P}(\mathbb{R}), \quad (Mp)(x) = x^2 p(x).$$

This map is linear and injective: if $x^2 p(x)$ is the zero polynomial, then p is the zero polynomial. It is not surjective, because the constant polynomial 1 does not belong to its image. Thus M is not invertible, even though its domain and codomain are the same vector space.

Example 33 (A complementary example). *Differentiation*

$$D: \mathcal{P}(\mathbb{R}) \rightarrow \mathcal{P}(\mathbb{R}), \quad D(p) = p',$$

is surjective, since every real polynomial has a polynomial antiderivative. It is not injective, since every constant polynomial belongs to $\ker D$. Therefore D is not invertible.

Proposition 32. *Suppose that V and W are finite-dimensional with $\dim V = \dim W$, and let $T \in \mathcal{L}(V, W)$. Then the following are equivalent:*

- (i) T is invertible;
- (ii) T is injective;
- (iii) T is surjective.

Proof. By rank–nullity and the equality of dimensions,

$$\begin{aligned}
 T \text{ is injective} &\iff \ker T = \{0\} \\
 &\iff \dim \ker T = 0 \\
 &\iff \dim \operatorname{im} T = \dim V = \dim W \\
 &\iff \operatorname{im} T = W \\
 &\iff T \text{ is surjective.}
 \end{aligned}$$

Here we used the fact that a subspace of a finite-dimensional space with the same dimension is the whole space. Thus either injectivity or surjectivity implies bijectivity, which is equivalent to invertibility by Proposition 31. $\square$

14.4. One-sided inverses in equal dimensions.

Proposition 33. *Suppose that V and W are finite-dimensional with $\dim V = \dim W$. If $T \in \mathcal{L}(V, W)$ and $S \in \mathcal{L}(W, V)$, then*

$$ST = \operatorname{Id}_V \iff TS = \operatorname{Id}_W.$$

In either case, $S = T^{-1}$.

Proof. Suppose $ST = \operatorname{Id}_V$. If $v \in \ker T$, then

$$v = (ST)(v) = S(T(v)) = S(0) = 0.$$

Hence T is injective, and Proposition 32 makes it invertible. Consequently,

$$S = S(TT^{-1}) = (ST)T^{-1} = T^{-1},$$

which implies $TS = \text{Id}_W$.

Conversely, suppose $TS = \text{Id}_W$. For every $w \in W$,

$$w = T(S(w)),$$

so T is surjective and hence invertible by the same proposition. Therefore

$$S = (T^{-1}T)S = T^{-1}(TS) = T^{-1},$$

and $ST = \text{Id}_V$. □