

MATH 535: LINEAR ALGEBRA

RUIZHE QIAN

CONTENTS

1. August 24, 2026	2
1.1. Fields	2
2. August 26, 2026	4
2.1. Further properties of fields	4
2.2. Characteristic of a field	4
2.3. Algebraically closed fields	5
2.4. Vectors and vector spaces	6
3. August 28, 2026	6
3.1. Examples of vector spaces	6
3.2. Linear combinations and spanning sets	7
3.3. Linear independence	8
4. August 31, 2026	10
4.1. Bases and coordinates	10
4.2. Linear transformations	12
5. September 2, 2026	14
5.1. Linear transformations determined by a basis	14
5.2. Matrix representations of linear transformations	15
6. September 4, 2026	17
6.1. Matrix–vector multiplication	17
6.2. Finding matrix representations	19
6.3. The vector space of linear maps	20
6.4. Composition of linear maps	21

1. AUGUST 24, 2026

1.1. Fields.

Definition 1 (Binary operation). *A binary operation on a set S is a function*

$$F: S \times S \longrightarrow S.$$

Thus, for every $x, y \in S$, the value $F(x, y)$ is again an element of S . We usually write $x \cdot y$ or $x + y$ in place of $F(x, y)$.

Example 1. *Addition on $\mathbb{Z}$ is a binary operation: the map*

$$\mathbb{Z} \times \mathbb{Z} \longrightarrow \mathbb{Z}, \quad (m, n) \longmapsto m + n$$

takes every pair of integers to an integer.

Definition 2. *Let F be a binary operation on S . A subset $T \subseteq S$ is closed under F if*

$$F(x, y) \in T$$

whenever $x, y \in T$.

Example 2. *The positive integers $\mathbb{Z}_{>0}$ are closed under addition. The odd integers are not closed under addition, but they are closed under multiplication.*

Definition 3 (Field). *Let $\mathbb{F}$ be a set equipped with two binary operations, called addition and multiplication. The sum and product of $a, b \in \mathbb{F}$ are denoted by $a + b$ and ab , respectively. Suppose that for every $a, b, c \in \mathbb{F}$,*

- (i) $a + b = b + a$;
- (ii) $(a + b) + c = a + (b + c)$;
- (iii) $ab = ba$;
- (iv) $a(bc) = (ab)c$;
- (v) $a(b + c) = ab + ac$.

Suppose further that

(vi) $\mathbb{F}$ contains an additive identity 0 and a multiplicative identity 1, so that

$$a + 0 = a \quad \text{and} \quad 1a = a$$

for every $a \in \mathbb{F}$;

(vii) $1 \neq 0$;

(viii) every $a \in \mathbb{F}$ has an additive inverse $-a \in \mathbb{F}$ such that $a + (-a) = 0$;

(ix) every nonzero $a \in \mathbb{F}$ has a multiplicative inverse $a^{-1} \in \mathbb{F}$ such that $aa^{-1} = 1$.

Then $\mathbb{F}$ is called a field.

Example 3. The integers $\mathbb{Z}$ do not form a field: for example, 2 has no multiplicative inverse in $\mathbb{Z}$. The sets $\mathbb{Q}$, $\mathbb{R}$, and $\mathbb{C}$, with their usual operations, are fields.

There are also finite fields. For example,

$$\mathbb{F}_2 = \{0, 1\} \quad \text{and} \quad \mathbb{F}_3 = \{0, 1, 2\},$$

where the elements are the possible remainders upon division by 2 and 3, respectively. Addition and multiplication in $\mathbb{F}_3$ are given by the following tables:

$$\begin{array}{c|ccc} + & 0 & 1 & 2 \\ \hline 0 & 0 & 1 & 2 \\ 1 & 1 & 2 & 0 \\ 2 & 2 & 0 & 1 \end{array} \quad \begin{array}{c|ccc} \cdot & 0 & 1 & 2 \\ \hline 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 2 \\ 2 & 0 & 2 & 1 \end{array}.$$

More generally, if p is prime, then

$$\mathbb{F}_p = \{0, 1, \dots, p-1\},$$

with addition and multiplication performed modulo p , is a field.

Lemma 1. If a is an element of a field $\mathbb{F}$, then $a \cdot 0 = 0$.

Proof. Since $0 + 0 = 0$, distributivity gives

$$a \cdot 0 = a(0 + 0) = a \cdot 0 + a \cdot 0.$$

Adding the additive inverse of $a \cdot 0$ to both sides yields $0 = a \cdot 0$. $\square$

2. AUGUST 26, 2026

2.1. Further properties of fields. The preceding lecture showed that $a \cdot 0 = 0$ for every element a of a field. We now obtain the zero-product property.

Proposition 1 (Zero-product property). *Let $a, b \in \mathbb{F}$. If $ab = 0$, then either $a = 0$ or $b = 0$.*

Proof. Suppose that $ab = 0$ and $a \neq 0$. Since $\mathbb{F}$ is a field, a^{-1} exists. Therefore

$$0 = a^{-1}0 = a^{-1}(ab) = (a^{-1}a)b = 1b = b.$$

Thus $a = 0$ or $b = 0$. $\square$

2.2. Characteristic of a field. For a positive integer n and $a \in \mathbb{F}$, write

$$na = \underbrace{a + a + \cdots + a}_{n \text{ terms}}.$$

Definition 4 (Characteristic). *Let $\mathbb{F}$ be a field. If $n1 = 0$ for some positive integer n , then the least such n is called the characteristic of $\mathbb{F}$, denoted $\text{char}(\mathbb{F})$. If $n1 \neq 0$ for every positive integer n , then $\mathbb{F}$ has characteristic zero.*

The fields $\mathbb{Q}$, $\mathbb{R}$, and $\mathbb{C}$ have characteristic zero. For every prime p , the field $\mathbb{F}_p$ has characteristic p . The rational function field $\mathbb{F}_p(t)$ is an infinite field of characteristic p .

Proposition 2. *If $\mathbb{F}$ has positive characteristic r , then r is prime.*

Proof. We have $r \neq 1$, since $1 \neq 0$. Suppose that r is composite, so

$$r = st$$

for integers s, t satisfying $1 < s, t < r$. Distributivity gives

$$(s1)(t1) = (st)1 = r1 = 0.$$

By the zero-product property, either $s1 = 0$ or $t1 = 0$. Both alternatives contradict the minimality of r . Hence r is prime. $\square$

Proposition 3. *If $\text{char}(\mathbb{F}) = p > 0$, then*

$$pa = 0$$

for every $a \in \mathbb{F}$.

Proof. Since $p1 = 0$, associativity and distributivity imply

$$pa = p(1a) = (p1)a = 0a = 0.$$

□

Proposition 4. *Every finite field has positive characteristic.*

Proof. Let $\mathbb{F}$ be finite. The sequence

$$1, \quad 2 \cdot 1, \quad 3 \cdot 1, \quad \dots$$

cannot consist of distinct elements. Hence there are positive integers $m > n$ such that

$$m1 = n1.$$

Adding the additive inverse of $n1$ to both sides gives

$$(m - n)1 = 0.$$

Thus $\mathbb{F}$ has positive characteristic. □

2.3. Algebraically closed fields.

Definition 5 (Algebraically closed field). *A field $\mathbb{F}$ is algebraically closed if every nonconstant polynomial $f \in \mathbb{F}[x]$ has a root in $\mathbb{F}$.*

The field $\mathbb{Q}$ is not algebraically closed, because $x^2 - 2$ has no root in $\mathbb{Q}$. The field $\mathbb{R}$ is not algebraically closed, because $x^2 + 1$ has no real root. By the fundamental theorem of algebra, $\mathbb{C}$ is algebraically closed.

Proposition 5. *No finite field is algebraically closed.*

Proof. Let

$$\mathbb{F} = \{a_1, \dots, a_q\}$$

be finite. Consider the nonconstant polynomial

$$f(x) = 1 + \prod_{i=1}^q (x - a_i).$$

For each $i = 1, \dots, q$,

$$f(a_i) = 1 \neq 0.$$

Thus f has no root in $\mathbb{F}$, so $\mathbb{F}$ is not algebraically closed. $\square$

2.4. Vectors and vector spaces. The spaces $\mathbb{R}^2$, $\mathbb{R}^3$, and, more generally,

$$\mathbb{R}^n = \left\{ \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} : x_1, \dots, x_n \in \mathbb{R} \right\}$$

are the motivating examples of vector spaces.

Definition 6 (Vector space). *Let $\mathbb{F}$ be a field. A vector space over $\mathbb{F}$ is a set V equipped with two operations,*

$$V \times V \longrightarrow V, \quad (u, v) \longmapsto u + v,$$

and

$$\mathbb{F} \times V \longrightarrow V, \quad (\alpha, v) \longmapsto \alpha v,$$

satisfying the following axioms for all $u, v, w \in V$ and $\alpha, \beta \in \mathbb{F}$:

- (V1) $u + (v + w) = (u + v) + w$;
- (V2) $u + v = v + u$;
- (V3) there is an element $0 \in V$ such that $v + 0 = 0 + v = v$;
- (V4) for every $v \in V$, there is an element $-v \in V$ such that $v + (-v) = (-v) + v = 0$;
- (V5) $1v = v$;
- (V6) $(\alpha\beta)v = \alpha(\beta v)$;
- (V7) $\alpha(u + v) = \alpha u + \alpha v$;
- (V8) $(\alpha + \beta)v = \alpha v + \beta v$.

The elements of V are called *vectors*, and the elements of $\mathbb{F}$ are called *scalars*.

3. AUGUST 28, 2026

3.1. Examples of vector spaces.

Example 4 (Coordinate spaces). *Let $\mathbb{F}$ be a field. The set*

$$\mathbb{F}^n = \left\{ \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} : x_1, \dots, x_n \in \mathbb{F} \right\}$$

is a vector space over $\mathbb{F}$ under componentwise addition and scalar multiplication:

$$\begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} + \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix} = \begin{bmatrix} x_1 + y_1 \\ \vdots \\ x_n + y_n \end{bmatrix}, \quad \alpha \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} \alpha x_1 \\ \vdots \\ \alpha x_n \end{bmatrix}.$$

In particular, $\mathbb{R}^n$ is a vector space over $\mathbb{R}$ and $\mathbb{C}^n$ is a vector space over $\mathbb{C}$.

Example 5 (Matrix spaces). *The set $M_{m \times n}(\mathbb{F})$ of $m \times n$ matrices with entries in $\mathbb{F}$ is a vector space over $\mathbb{F}$. Addition and scalar multiplication are defined entrywise:*

$$[a_{ij}] + [b_{ij}] = [a_{ij} + b_{ij}], \quad \alpha[a_{ij}] = [\alpha a_{ij}].$$

Example 6 (Polynomial spaces). *Let*

$$\mathcal{P}_n(\mathbb{F}) = \{a_0 + a_1t + \dots + a_nt^n : a_0, \dots, a_n \in \mathbb{F}\}.$$

This is the vector space over $\mathbb{F}$ of polynomials of degree at most n . Its operations are

$$\begin{aligned} (a_0 + a_1t + \dots + a_nt^n) + (b_0 + b_1t + \dots + b_nt^n) \\ = (a_0 + b_0) + (a_1 + b_1)t + \dots + (a_n + b_n)t^n, \end{aligned}$$

and

$$\alpha(a_0 + a_1t + \dots + a_nt^n) = \alpha a_0 + \alpha a_1t + \dots + \alpha a_nt^n.$$

3.2. Linear combinations and spanning sets.

Definition 7 (Linear combination and span). *Let V be a vector space over $\mathbb{F}$ and let $v_1, \dots, v_p \in V$. A linear combination of these vectors is a vector of the form*

$$\alpha_1 v_1 + \dots + \alpha_p v_p, \quad \alpha_1, \dots, \alpha_p \in \mathbb{F}.$$

The set of all their linear combinations is called their span:

$$\text{span}\{v_1, \dots, v_p\} = \{\alpha_1 v_1 + \dots + \alpha_p v_p : \alpha_1, \dots, \alpha_p \in \mathbb{F}\}.$$

A linear combination is trivial if all of its coefficients are zero.

In $\mathbb{R}^2$, let

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

Then

$$\text{span}\{e_1, e_2\} = \mathbb{R}^2,$$

because

$$\begin{bmatrix} x \\ y \end{bmatrix} = x e_1 + y e_2.$$

If

$$v = \begin{bmatrix} 1 \\ 1 \end{bmatrix},$$

then $\{e_1, e_2, v\}$ also spans $\mathbb{R}^2$, since

$$\begin{bmatrix} x \\ y \end{bmatrix} = x e_1 + y e_2 + 0 v.$$

The additional vector is unnecessary, which motivates the question of redundancy in a spanning set.

Definition 8 (Spanning set). *A collection $\{v_1, \dots, v_p\} \subseteq V$ is a spanning set for V if every $v \in V$ can be written as a linear combination*

$$v = \alpha_1 v_1 + \dots + \alpha_p v_p.$$

Equivalently,

$$\text{span}\{v_1, \dots, v_p\} = V.$$

3.3. Linear independence.

Definition 9 (Linear independence). *A collection $\{v_1, \dots, v_p\} \subseteq V$ is linearly independent if*

$$\alpha_1 v_1 + \dots + \alpha_p v_p = 0$$

implies

$$\alpha_1 = \cdots = \alpha_p = 0.$$

In other words, only the trivial linear combination of $v_1, \dots, v_p$ equals zero. A collection that is not linearly independent is called linearly dependent.

Example 7. Consider the vectors

$$v_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \quad v_2 = \begin{bmatrix} 2 \\ 0 \\ 1 \end{bmatrix}, \quad v_3 = \begin{bmatrix} 3 \\ 1 \\ 1 \end{bmatrix}$$

in $\mathbb{R}^3$. To test linear independence, solve

$$\alpha v_1 + \beta v_2 + \gamma v_3 = 0.$$

This gives the system

$$\begin{cases} \alpha + 2\beta + 3\gamma = 0, \\ \alpha + \gamma = 0, \\ \beta + \gamma = 0. \end{cases}$$

Taking $\gamma = 1$ gives $\alpha = \beta = -1$, so

$$-v_1 - v_2 + v_3 = 0.$$

Thus the vectors are linearly dependent; equivalently, $v_3 = v_1 + v_2$.

The following result precisely characterizes redundancy in a finite collection of vectors.

Proposition 6. Let $v_1, \dots, v_p \in V$. The collection $\{v_1, \dots, v_p\}$ is linearly dependent if and only if one of its vectors is a linear combination of the others. Equivalently, there is an index k and scalars $\beta_j \in \mathbb{F}$ such that

$$v_k = \sum_{\substack{j=1 \\ j \neq k}}^p \beta_j v_j.$$

Proof. Suppose first that $\{v_1, \dots, v_p\}$ is linearly dependent. Then there are scalars $\alpha_1, \dots, \alpha_p$, not all zero, such that

$$\alpha_1 v_1 + \dots + \alpha_p v_p = 0.$$

Choose k such that $\alpha_k \neq 0$. Since $\mathbb{F}$ is a field, α_k^{-1} exists, and we can solve for v_k :

$$v_k = - \sum_{\substack{j=1 \\ j \neq k}}^p \alpha_k^{-1} \alpha_j v_j.$$

Thus v_k is a linear combination of the other vectors.

Conversely, suppose that for some k ,

$$v_k = \sum_{\substack{j=1 \\ j \neq k}}^p \beta_j v_j.$$

Moving all terms to one side gives

$$v_k - \sum_{\substack{j=1 \\ j \neq k}}^p \beta_j v_j = 0.$$

This is a nontrivial linear combination of $v_1, \dots, v_p$, because the coefficient of v_k is $1 \neq 0$. Hence the collection is linearly dependent. $\square$

4. AUGUST 31, 2026

4.1. Bases and coordinates.

Definition 10 (Basis). *A collection of vectors $\{v_1, \dots, v_n\}$ in a vector space V is a basis for V if it is linearly independent and spans V .*

The two conditions in the definition can be combined into a statement about representing vectors uniquely.

Proposition 7. *A collection $\{v_1, \dots, v_n\}$ is a basis for V if and only if every vector $v \in V$ has a unique representation of the form*

$$v = a_1 v_1 + \dots + a_n v_n, \quad a_1, \dots, a_n \in \mathbb{F}.$$

When $\{v_1, \dots, v_n\}$ is a basis, the scalars $a_1, \dots, a_n$ are called the coordinates of v with respect to this basis.

Proof. Suppose first that $\{v_1, \dots, v_n\}$ is a basis. Since the collection spans V , every $v \in V$ has a representation

$$v = a_1v_1 + \cdots + a_nv_n.$$

To prove uniqueness, suppose that

$$v = \tilde{a}_1v_1 + \cdots + \tilde{a}_nv_n$$

is another such representation. Subtracting the two equations gives

$$(a_1 - \tilde{a}_1)v_1 + \cdots + (a_n - \tilde{a}_n)v_n = 0.$$

Because $\{v_1, \dots, v_n\}$ is linearly independent,

$$a_i - \tilde{a}_i = 0$$

for each i . Thus $a_i = \tilde{a}_i$ for every i , and the representation is unique.

Conversely, suppose every $v \in V$ has a unique representation as a linear combination of $v_1, \dots, v_n$. The existence of such a representation for every v shows that the collection spans V . Now suppose

$$a_1v_1 + \cdots + a_nv_n = 0.$$

The zero vector also has the representation

$$0 = 0v_1 + \cdots + 0v_n.$$

Uniqueness therefore gives $a_1 = \cdots = a_n = 0$. Hence the collection is linearly independent, so it is a basis for V . $\square$

Example 8 (The standard basis of $\mathbb{F}^n$). For $i = 1, \dots, n$, let $e_i \in \mathbb{F}^n$ be the vector whose i th coordinate is 1 and whose other coordinates are 0. Thus

$$e_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \quad \dots \quad e_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix}.$$

Every vector in $\mathbb{F}^n$ has the unique representation

$$\begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = x_1 e_1 + x_2 e_2 + \cdots + x_n e_n.$$

Consequently, $\{e_1, \dots, e_n\}$ is a basis for $\mathbb{F}^n$, called its standard basis.

Example 9 (The standard basis of $\mathcal{P}_n(\mathbb{F})$). Recall that $\mathcal{P}_n(\mathbb{F})$ is the vector space of polynomials over $\mathbb{F}$ of degree at most n . Each polynomial has a unique expression

$$p(t) = a_0 + a_1 t + \cdots + a_n t^n.$$

Hence

$$\{1, t, t^2, \dots, t^n\}$$

is a basis for $\mathcal{P}_n(\mathbb{F})$, called its standard basis.

Proposition 8. Every finite nonempty spanning set for a vector space contains a basis.

Proof. Let $\{v_1, \dots, v_p\}$ span V . If this collection is linearly independent, then it is already a basis. Otherwise, by Proposition 6, some v_k is a linear combination of the other vectors. It follows that removing v_k does not change the span:

$$\text{span}\{v_1, \dots, v_p\} = \text{span}\{v_1, \dots, v_{k-1}, v_{k+1}, \dots, v_p\}.$$

If the remaining collection is linearly independent, it is a basis. If not, repeat the same argument. The original spanning set is finite, so this process must terminate. The final collection still spans V and is linearly independent, and is therefore a basis contained in the original spanning set. $\square$

4.2. Linear transformations. A *transformation*, also called a map or function, from a set X to a set Y is a rule $T: X \rightarrow Y$ that assigns to every $x \in X$ an element $T(x) \in Y$.

Definition 11 (Linear transformation). *Let V and W be vector spaces over the same field $\mathbb{F}$. A transformation $T: V \rightarrow W$ is linear if*

- (i) $T(u + v) = T(u) + T(v)$ for all $u, v \in V$;
- (ii) $T(\alpha u) = \alpha T(u)$ for all $u \in V$ and $\alpha \in \mathbb{F}$.

Equivalently, $T: V \rightarrow W$ is linear if

$$T(\alpha u + \beta v) = \alpha T(u) + \beta T(v)$$

for all $u, v \in V$ and $\alpha, \beta \in \mathbb{F}$.

Example 10 (Reflection). *Define $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by*

$$T \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} x_1 \\ -x_2 \end{bmatrix}.$$

Geometrically, T reflects each point across the first coordinate axis. To check additivity, observe that

$$\begin{aligned} T \left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} \right) &= T \begin{bmatrix} x_1 + y_1 \\ x_2 + y_2 \end{bmatrix} \\ &= \begin{bmatrix} x_1 + y_1 \\ -x_2 - y_2 \end{bmatrix} \\ &= \begin{bmatrix} x_1 \\ -x_2 \end{bmatrix} + \begin{bmatrix} y_1 \\ -y_2 \end{bmatrix} \\ &= T \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + T \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}. \end{aligned}$$

For $\alpha \in \mathbb{R}$,

$$\begin{aligned} T \left(\alpha \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \right) &= T \begin{bmatrix} \alpha x_1 \\ \alpha x_2 \end{bmatrix} \\ &= \begin{bmatrix} \alpha x_1 \\ -\alpha x_2 \end{bmatrix} \\ &= \alpha \begin{bmatrix} x_1 \\ -x_2 \end{bmatrix} = \alpha T \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}. \end{aligned}$$

Thus T is linear.

Example 11 (The differentiation operator). Let $n \geq 1$, $V = \mathcal{P}_n(\mathbb{F})$, and $W = \mathcal{P}_{n-1}(\mathbb{F})$. Define

$$T: V \longrightarrow W, \quad T(p) = p'.$$

Since

$$(p + q)' = p' + q' \quad \text{and} \quad (\alpha p)' = \alpha p',$$

the differentiation operator T is linear.

5. SEPTEMBER 2, 2026

5.1. Linear transformations determined by a basis. The following result shows that prescribing the values of a linear transformation on a basis completely determines the transformation.

Lemma 2 (Linear map lemma). Suppose that $\{v_1, \dots, v_n\}$ is a basis for V , and let $w_1, \dots, w_n$ be arbitrary vectors in W . Then there exists a unique linear transformation $T: V \rightarrow W$ such that

$$T(v_k) = w_k$$

for each $k = 1, \dots, n$.

Proof. Every $v \in V$ has a unique representation

$$v = \sum_{i=1}^n c_i v_i.$$

Define

$$T(v) = \sum_{i=1}^n c_i w_i.$$

The uniqueness of the coordinates $c_1, \dots, c_n$ shows that this definition is well-defined. In particular, taking $v = v_k$ gives $T(v_k) = w_k$.

To prove that T is linear, write

$$u = \sum_{i=1}^n a_i v_i \quad \text{and} \quad v = \sum_{i=1}^n c_i v_i.$$

Then

$$\begin{aligned}
 T(u + v) &= T\left(\sum_{i=1}^n (a_i + c_i)v_i\right) \\
 &= \sum_{i=1}^n (a_i + c_i)w_i \\
 &= \sum_{i=1}^n a_i w_i + \sum_{i=1}^n c_i w_i \\
 &= T(u) + T(v).
 \end{aligned}$$

Similarly, for every $\alpha \in \mathbb{F}$,

$$\begin{aligned}
 T(\alpha v) &= T\left(\sum_{i=1}^n \alpha c_i v_i\right) \\
 &= \sum_{i=1}^n \alpha c_i w_i = \alpha \sum_{i=1}^n c_i w_i = \alpha T(v).
 \end{aligned}$$

Thus T is linear.

It remains to prove uniqueness. Suppose that $T': V \rightarrow W$ is another linear transformation satisfying $T'(v_k) = w_k$ for every k . If $v = \sum_{i=1}^n c_i v_i$, then

$$T'(v) = \sum_{i=1}^n c_i T'(v_i) = \sum_{i=1}^n c_i w_i = T(v).$$

Hence $T' = T$. □

5.2. Matrix representations of linear transformations. An *ordered basis* is a basis whose vectors are listed in a fixed order. Let

$$\mathcal{B} = (v_1, \dots, v_n) \quad \text{and} \quad \mathcal{C} = (w_1, \dots, w_m)$$

be ordered bases for V and W , respectively. For $v = x_1 v_1 + \dots + x_n v_n$, write

$$[v]_{\mathcal{B}} = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$$

for the coordinate vector of v with respect to $\mathcal{B}$.

$$A = [A_{ij}] = \begin{bmatrix} A_{11} & \cdots & A_{1n} \\ \vdots & & \vdots \\ A_{m1} & \cdots & A_{mn} \end{bmatrix} \in M_{m \times n}(\mathbb{F})$$
$$T(v_j) = \sum_{i=1}^m A_{ij} w_i$$

Proof. Let $T: V \rightarrow W$ be linear. Because $\mathcal{C}$ is a basis for W , each vector $T(v_j)$ has a unique representation

$$T(v_j) = A_{1j}w_1 + \cdots + A_{mj}w_m = \sum_{i=1}^m A_{ij}w_i.$$

$$A = \begin{bmatrix} | & & | \\ [T(v_1)]_{\mathcal{C}} & \cdots & [T(v_n)]_{\mathcal{C}} \\ | & & | \end{bmatrix}.$$
$$\sum_{i=1}^m B_{ij} w_i = T(v_j) = \sum_{i=1}^m A_{ij} w_i.$$

Conversely, let $A = [A_{ij}]$ be an arbitrary $m \times n$ matrix and define

$$\tilde{w}_j = \sum_{i=1}^m A_{ij} w_i \quad (j = 1, \dots, n).$$

By the linear map lemma, there is a unique linear transformation $T: V \rightarrow W$ such that $T(v_j) = \tilde{w}_j$ for every j . Thus A determines a unique linear transformation with the required property. $\square$

If

$$v = x_1 v_1 + \cdots + x_n v_n,$$

then linearity gives

$$\begin{aligned} T(v) &= \sum_{j=1}^n x_j T(v_j) \\ &= \sum_{j=1}^n x_j \sum_{i=1}^m A_{ij} w_i \\ &= \sum_{i=1}^m \left(\sum_{j=1}^n A_{ij} x_j \right) w_i. \end{aligned}$$

Consequently,

$$[T(v)]_{\mathcal{C}} = A[v]_{\mathcal{B}}.$$

Thus multiplication by the matrix A computes the coordinates of the image of a vector from the coordinates of the original vector.

6. SEPTEMBER 4, 2026

6.1. Matrix–vector multiplication. Let V and W be vector spaces over $\mathbb{F}$ with ordered bases

$$\mathcal{B} = (v_1, \dots, v_n) \quad \text{and} \quad \mathcal{C} = (w_1, \dots, w_m),$$

respectively, and let $T: V \rightarrow W$ be linear. Recall that the matrix representing T with respect to these bases is

$$A = \begin{bmatrix} A_{11} & \cdots & A_{1n} \\ \vdots & & \vdots \\ A_{m1} & \cdots & A_{mn} \end{bmatrix},$$

where

$$T(v_j) = A_{1j} w_1 + \cdots + A_{mj} w_m.$$

Thus the j th column of A is the coordinate vector $[T(v_j)]_{\mathcal{C}}$.

Suppose that

$$v = x_1 v_1 + \cdots + x_n v_n, \quad [v]_{\mathcal{B}} = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}.$$

By linearity,

$$\begin{aligned} [T(v)]_{\mathcal{C}} &= x_1 [T(v_1)]_{\mathcal{C}} + \cdots + x_n [T(v_n)]_{\mathcal{C}} \\ &= x_1 \begin{bmatrix} A_{11} \\ A_{21} \\ \vdots \\ A_{m1} \end{bmatrix} + \cdots + x_n \begin{bmatrix} A_{1n} \\ A_{2n} \\ \vdots \\ A_{mn} \end{bmatrix} \\ &= \begin{bmatrix} A_{11}x_1 + \cdots + A_{1n}x_n \\ \vdots \\ A_{m1}x_1 + \cdots + A_{mn}x_n \end{bmatrix}. \end{aligned}$$

This motivates the following definition.

Definition 12 (Matrix–vector product). *Let $A = [A_{ij}] \in M_{m \times n}(\mathbb{F})$, and write its columns as $A_1, \dots, A_n$. For*

$$x = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} \in \mathbb{F}^n,$$

define

$$Ax = x_1 A_1 + \cdots + x_n A_n.$$

Equivalently, the k th coordinate of Ax is

$$(Ax)_k = \sum_{j=1}^n A_{kj} x_j, \quad k = 1, \dots, m.$$

With this notation, the coordinate formula for a linear transformation is

$$[T(v)]_{\mathcal{C}} = A[v]_{\mathcal{B}}.$$

Example 12. *Matrix–vector multiplication can be computed either as a linear combination of the columns or one coordinate at a time. For*

instance,

$$\begin{bmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = 1 \begin{bmatrix} 1 \\ 3 \end{bmatrix} + 2 \begin{bmatrix} 2 \\ 2 \end{bmatrix} + 3 \begin{bmatrix} 3 \\ 1 \end{bmatrix} = \begin{bmatrix} 14 \\ 10 \end{bmatrix},$$

while

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = \begin{bmatrix} 1 \cdot 1 + 2 \cdot 2 + 3 \cdot 3 \\ 4 \cdot 1 + 5 \cdot 2 + 6 \cdot 3 \end{bmatrix} = \begin{bmatrix} 14 \\ 32 \end{bmatrix}.$$

6.2. Finding matrix representations.

Example 13. Define $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$ by

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x + 2y \\ 2x - 5y \\ 7y \end{bmatrix}.$$

With respect to the standard bases, the columns of the representing matrix are

$$T(e_1) = \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix} \quad \text{and} \quad T(e_2) = \begin{bmatrix} 2 \\ -5 \\ 7 \end{bmatrix}.$$

Hence

$$A = \begin{bmatrix} 1 & 2 \\ 2 & -5 \\ 0 & 7 \end{bmatrix}.$$

Example 14 (Differentiation). Let $n \geq 1$, and let

$$T: \mathcal{P}_n(\mathbb{R}) \longrightarrow \mathcal{P}_n(\mathbb{R}), \quad T(p) = p',$$

and use the ordered basis

$$\mathcal{B} = (1, t, t^2, \dots, t^n)$$

for both the domain and codomain. Since

$$T(1) = 0, \quad T(t) = 1, \quad T(t^j) = jt^{j-1},$$

the matrix representing T is

$$A = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 2 & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & 0 & n \\ 0 & \cdots & \cdots & 0 & 0 \end{bmatrix}.$$

6.3. The vector space of linear maps. Let V and W be vector spaces over $\mathbb{F}$. If $T: V \rightarrow W$ is linear and $\alpha \in \mathbb{F}$, define $\alpha T: V \rightarrow W$ pointwise by

$$(\alpha T)(v) = \alpha T(v).$$

This map is again linear. Indeed, for $u, v \in V$,

$$\begin{aligned} (\alpha T)(u + v) &= \alpha T(u + v) \\ &= \alpha(T(u) + T(v)) \\ &= (\alpha T)(u) + (\alpha T)(v), \end{aligned}$$

and, for $\beta \in \mathbb{F}$,

$$(\alpha T)(\beta u) = \alpha T(\beta u) = \alpha \beta T(u) = \beta(\alpha T)(u).$$

Similarly, if $T_1, T_2: V \rightarrow W$ are linear, define their sum by

$$(T_1 + T_2)(v) = T_1(v) + T_2(v).$$

The map $T_1 + T_2$ is linear because

$$\begin{aligned} (T_1 + T_2)(u + v) &= T_1(u + v) + T_2(u + v) \\ &= (T_1 + T_2)(u) + (T_1 + T_2)(v) \end{aligned}$$

and

$$(T_1 + T_2)(\alpha u) = \alpha(T_1 + T_2)(u).$$

Definition 13 (Space of linear maps). *The collection of all linear transformations from V to W is denoted by*

$$\mathcal{L}(V, W) = \{T: V \rightarrow W : T \text{ is linear}\}.$$

Proposition 9. *The set $\mathcal{L}(V, W)$, equipped with pointwise addition and scalar multiplication, is a vector space over $\mathbb{F}$.*

Proof. The preceding calculations show that the two operations are closed in $\mathcal{L}(V, W)$. All vector-space identities follow pointwise from the corresponding identities in W . The additive identity is the zero transformation $0: V \rightarrow W$, defined by $0(v) = 0$, and the additive inverse of T is the map $-T$, defined by $(-T)(v) = -T(v)$. $\square$

6.4. Composition of linear maps. Let V , W , and Z be vector spaces over $\mathbb{F}$. Suppose that

$$T_2: V \longrightarrow W \quad \text{and} \quad T_1: W \longrightarrow Z$$

are linear. Their *composition* is the map

$$T_1 \circ T_2: V \longrightarrow Z, \quad (T_1 \circ T_2)(v) = T_1(T_2(v)).$$

Proposition 10. *The composition $T_1 \circ T_2$ is linear.*

Proof. For $u, v \in V$ and $\alpha \in \mathbb{F}$,

$$\begin{aligned} (T_1 \circ T_2)(u + v) &= T_1(T_2(u) + T_2(v)) \\ &= T_1(T_2(u)) + T_1(T_2(v)), \end{aligned}$$

and

$$(T_1 \circ T_2)(\alpha u) = T_1(\alpha T_2(u)) = \alpha T_1(T_2(u)).$$

$\square$