# A DECADE OF SHORT-DURATION GAMMA-RAY BURST BROAD-BAND AFTERGLOWS: ENERGETICS, CIRCUMBURST DENSITIES, AND JET OPENING ANGLES

W. FONG<sup>1,2</sup>, E. BERGER<sup>3</sup>, R. MARGUTTI<sup>3</sup>, & B. A. ZAUDERER<sup>3</sup>

Draft version September 11, 2015

#### **ABSTRACT**

We present a comprehensive catalog and analysis of broad-band afterglow observations for 103 short-duration gamma-ray bursts (GRBs), comprised of all short GRBs from November 2004 to March 2015 with prompt follow-up observations in the X-ray, optical, near-infrared and/or radio bands. These afterglow observations have uncovered 71 X-ray detections, 30 optical/NIR detections, and 4 radio detections. Employing the standard afterglow synchrotron model, we perform joint probability analyses for a subset of 38 short GRBs with well-sampled light curves to infer the burst isotropic-equivalent energies and circumburst densities. For this subset, we find median isotropic-equivalent  $\gamma$ -ray and kinetic energies of  $E_{\gamma,\rm iso}\approx 2\times 10^{51}$  erg, and  $E_{K,\rm iso}\approx (1-3)\times 10^{51}$  erg, respectively, depending on the values of the model input parameters. We further find that short GRBs occur in low-density environments, with a median density of  $n\approx (3-15)\times 10^{-3}$  cm<sup>-3</sup>, and that  $\approx 80-95\%$  of bursts have densities of  $n\lesssim 1$  cm<sup>-3</sup>. We investigate trends between the circumburst densities and host galaxy properties, and find that events located at large projected offsets of  $\gtrsim 10$  effective radii from their hosts exhibit particularly low densities of  $n\lesssim 10^{-4}$  cm<sup>-3</sup>, consistent with an IGM-like environment. Using late-time afterglow data for 11 events, we find a median jet opening angle of  $\theta_j=16\pm 10^\circ$ . We also calculate a median beaming factor of  $f_b\approx 0.04$ , leading to a beaming-corrected total energy elease of  $E_{\rm true}\approx 1.6\times 10^{50}$  erg. Furthermore, we calculate a beaming-corrected event rate of  $\Re_{\rm true}=270^{+1580}_{-180}$  Gpc<sup>-3</sup> yr<sup>-1</sup>, or  $\approx 8^{+47}_{-5}$  yr<sup>-1</sup> within a 200 Mpc volume, the Advanced LIGO/Virgo typical detection distance for NS-NS binaries.

#### 1. INTRODUCTION

The afterglow emission from gamma-ray bursts (GRBs) provides a unique way to probe their basic properties: the energy scale, circumburst densities, and jet opening angles. Nearly two decades of long-duration GRB afterglow observations established a median beaming-corrected kinetic energy release of  $\approx 10^{51}$  erg (Frail et al. 2001; Berger et al. 2003; Friedman & Bloom 2005; Gehrels et al. 2008; Nysewander et al. 2009; Laskar et al. 2014), circumburst densities of  $\approx 0.1 - 100 \text{ cm}^{-3}$  (Panaitescu & Kumar 2002; Yost et al. 2003), and opening angles of  $\approx 2-20^{\circ}$ (Bloom et al. 2003; Frail et al. 2001; Friedman & Bloom 2005; Kocevski & Butler 2008; Racusin et al. Ryan et al. 2015). In some cases, the radial density profiles of their circumburst environments reflect the wind environments expected for massive stars (Chevalier & Li 2000; Yost et al. 2003).

The afterglows of short-duration GRBs ( $T_{90} \lesssim 2$  sec; Kouveliotou et al. 1993) are uniformly fainter (Berger 2007; Nakar 2007; Gehrels et al. 2008; Nysewander et al. 2009; Berger 2010, 2014), and have thus far been primarily utilized for providing precise burst localization, and therefore robust associations to host galaxies (e.g., Fong et al. 2013). These host studies have revealed that at least some short GRBs originate from older stellar populations, and are distinct in their global properties from the hosts of long GRBs (Berger 2009; Leibler & Berger 2010; Fong et al. 2013; Berger 2014). The local and galactic environments of short GRBs (Fong et al. 2010; Berger 2010; Fong & Berger 2013), together with

the lack of associated supernovae (e.g., Fox et al. 2005; Soderberg et al. 2006; Kocevski et al. 2010; Fong et al. 2014), and the discovery of a near-IR "kilonova" from the short GRB 130603B (Berger et al. 2013; Tanvir et al. 2013), have provided strong observational support for a compact object binary merger progenitor involving two neutron stars or a neutron star and a stellar mass black hole (NS-NS/NS-BH; Eichler et al. 1989; Narayan et al. 1992).

As studies of the host galaxies of short GRBs have progressed at a rapid pace, knowledge of their basic explosion properties has been limited by both the paucity of afterglow detections and the relatively low detection rate of well-localized (~few arcsec uncertainty) short GRBs from the Swift satellite. Furthermore, afterglow studies greatly benefit from observations across several orders of frequency, which serve to provide significantly tighter constraints on the energy scales and circumburst densities than single band observations. Thus far, short GRB afterglow studies have either focused on a small number of events in a single band (e.g., Berger 2007; Nakar 2007; Kann et al. 2008; Nicuesa Guelbenzu et al. 2012a), or on radio through X-ray observations of individual bursts (Berger et al. 2005; Soderberg et al. 2006; Fong et al. 2014). Comparative studies relative to long GRBs have only served to argue for lower energy scales and circumburst densities, but have not provided actual distributions (e.g., Berger 2014). Thus, there have been no attempts to utilize the full set of broad-band afterglow observations of short GRBs.

The relative difficulty of detecting short GRB afterglows is likely a direct reflection of a combination of low explosion energies and low circumburst densities. Predictions for NS-NS/NS-BH mergers span several orders of magnitude in density, depending on the precise distribution of natal kick velocities, merger timescales, and host galaxy type (Perna & Belczynski 2002; Belczynski et al. 2006). Simi-

<sup>&</sup>lt;sup>1</sup> Einstein Fellow

<sup>&</sup>lt;sup>2</sup> Steward Observatory, University of Arizona, 933 North Cherry Avenue, Tucson, AZ 85721

<sup>&</sup>lt;sup>3</sup> Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA

larly, different mechanisms of energy extraction to power a relavistic blast-wave can produce energy scales which differ by three orders of magnitude (e.g., Ruffert & Janka 1999b; Rosswog 2005). Furthermore, the calculation of the true short GRB energy scale and event rate directly depends on the geometry of their jets. Constraints on the energetics, densities, and jet opening angles from short GRB afterglows thus offer a way to study these fundamental questions. These basic properties also serve as critical inputs for the detectability of other electromagnetic counterparts to compact object mergers, and will directly affect follow-up strategies to gravitational wave signals detected with Advanced LIGO/VIRGO (e.g., Metzger & Berger 2012; Nissanke et al. 2013).

Taking advantage of the dedicated searches for short GRB afterglows at all wavelengths, we are now in a position to explore these basic properties for a large population of events for the first time. Here, we present the first comprehensive broad-band catalog of short GRB afterglows, representing a decade of observations since the launch of Swift in 2004, and use this sample to constrain short GRB energies and circumburst densities. In Section 2, we introduce the sample and data reduction methods for X-ray through radio observations. In Section 3, we model the temporal and spectral behavior of the afterglows, and use the observations to infer constraints on the energetics and circumburst densities for each burst. In Section 4, we present the energetics and circumburst densities for 38 events with well-sampled afterglow data sets. In Section 5, we discuss the observational afterglow properties, jet opening angles, and investigate trends between the bursts and their galactic environments. Finally, in the Appendix, we provide X-ray, optical, and radio afterglow catalogs. In an upcoming work, we will address the effect of the basic inferred properties on the detectability of electromagnetic counterparts to gravitational wave sources.

Throughout the paper, all magnitudes are in the AB system and are corrected for Galactic extinction in the direction of each burst (Schlegel et al. 1998; Schlafly & Finkbeiner 2011). Unless otherwise noted, uncertainties correspond to  $1\sigma$  confidence. We employ a standard  $\Lambda$ CDM cosmology with  $\Omega_M=0.27$ ,  $\Omega_\Lambda=0.73$ , and  $H_0=71~{\rm km~s^{-1}~Mpc^{-1}}$ .

### 2. OBSERVATIONS

#### 2.1. Sample

We present afterglow observations for 103 short GRBs discovered by the Swift satellite (Gehrels et al. 2004), Fermi satellite (Atwood et al. 2009; Meegan et al. 2009), Konus-Wind (Aptekar et al. 1995), High Energy Transient Explorer 2 (HETE-2; Ricker et al. 2003), or the Interplanetary Network (IPN; Hurley et al. 2010) between November 2004 and March 2015. We restrict our sample to all bursts with  $T_{90} \lesssim 2$  s and follow-up observations in any of the X-ray, optical, nearinfrared (NIR) or radio bands on timescales of  $\delta t \lesssim$  few days (where  $\delta t$  corresponds to the time after the  $\gamma$ -ray trigger). We also include three events (GRBs 050724A, 090607 and 100213A) which have  $T_{90} \approx 2.5 - 3$  s but which exhibit the spectral hardness and negligible spectral lags typical of short GRBs (Marshall et al. 2009; Grupe et al. 2010). For bursts with optical/NIR follow-up, we only include data from bursts with afterglow detections or meaningful limits of  $\gtrsim 20$  mag at  $\delta t \lesssim 1$  day. Basic information for the sample of 103 events, including durations, redshifts, and the available follow-up in each observing band is presented in Table 1.

Of the 103 bursts in our sample, 96 (93%) have X-ray

**Table 1**Short GRB Basic Information

GRB	T <sub>90</sub> (s)	Z	X-ray	Opt/NIR	Radio
050202 050509B	0.27 0.04	0.225	 Y	N N	N N
050309 <b>Б</b> 050709	0.04	0.223	Y	Y	N
050724A	3.0	0.257	Ÿ	Ÿ	Y
050813	0.6	0.72/1.8	Y	N	N
050906	0.26	•••	N	N	N
050925	0.07	• • •	Ŋ	• • •	N
051105A	0.09		$N^b$		N
051210 051221A	1.3 1.4	> 1.4 0.546	Y Y	N Y	 Y
060121	2.0	< 4.1	Y	<u>Y</u>	
060313	0.7	< 1.7	Ÿ	Ÿ	N
060502B	0.09	0.287	Y	N	
060801	0.5	1.130	Y	N	N
061006	0.4	0.438	Y	Y	• • •
061201	0.8	0.111	Y	Y	
061210 061217	0.2 0.2	0.41 0.827	Y Y	N N	N 
070209	0.2	0.827	N	N	•••
$070406^a$	1.20		N	N	
070429B	0.5	0.902	Y	N	N
$070707^{a}$	1.1	< 3.6	Y	Y	• • •
070714B	2.0	0.923	Y	Y	N
070724A	0.4	0.457	Y	Y	N
070729 070809	0.9 1.3	0.8 0.473	Y Y	N	N 
070809 070810B	0.08	0.473	N N	Y N	
070910B	0.05				N
$071017^a$	0.5		N		
$071112B^a$	0.30	• • •	N	N	N
071227	1.8	0.381	Y	Y	• • •
080121 <sup>a</sup>	0.7	• • •	N	N	• • • •
080123 080426	0.4 1.7		Y Y	N N	
080503	0.3	< 4.2	Y	Y	N
080702A	0.5		Ŷ	N	N
080905A	1.0	0.122	Y	Y	
080919	0.6	• • • •	Y	N	• • •
081024A	1.8	• • •	Y	N	N
$081024B^{a}$	0.4		$N^b$	N	N
081226A	0.4	< 4.1	Y	Y	N
081226B <sup>a</sup>	0.7	···	N	N	N
090305A 090417A	0.4 0.07	< 4.1	Y 	Y 	N
090426	1.3	2.609	Y	Y	
090510	0.3	0.903	Y	Y	N
090515	0.04	0.403	Y	Y	N
090607	$2.3^{c}$	• • • •	Y	N	
090621B	0.14	• • •	Y	N N	N
090715A 090916 <sup>a</sup>	0.5 0.3		 N	N 	N 
090910 091109B	0.3	< 4.4	Y	Y	
$091117^{a}$	0.43		$N^b$	N	N
100117A	0.30	0.915	Y	Y	•••
100206A	0.1	0.407	Y	N	• • •
100213A	$2.4^{d}$	• • •	Y	• • •	• • •
100625A	0.3	0.452	Y	N	N
100628A	0.04	• • • •	N	N	N
100702A	0.16	0.719	Y	N	• • •
101219A 101224A	0.6 0.2	0.718	Y Y	N 	 N
101224A 110112A	0.2	< 5.3	Y	Y	N
$110112A$ $110112B^a$	0.5		N	N	N
	0.0			- 1	

Table 1 — Continued

GRB	T <sub>90</sub> (s)	z	X-ray	Opt/NIR	Radio
110420B <sup>a</sup>	0.08		$N^b$	N	N
111020A	0.4		Y	N	N
111117A	0.5	1.3	Y	N	N
111121A	0.45		Y		N
111222A <sup>a</sup>	0.3		Y		
120229A	0.22			N	N
120305A	0.1		Y	N	N
120521A	0.45		Y	N	N
$120630A^{e}$	0.6		Y		
120804A	0.81	1.3	Y	Y	N
$120817B^a$	0.19		N	N	
121226A	1.0		Y	N	N
130313A	0.26		Y	N	N
130515A	0.29		Y	N	
130603B	0.18	0.356	Y	Y	Y
130626A	0.16		N		
130716A	0.8		Y	N	N
130822A	0.04		Y	N	N
130912A	0.28	< 4.1	Y	Y	N
131004A	1.54	0.717	Y	Y	N
$131125A^{f}$	0.5			N	
$131126A^{f}$	0.3			N	
131224A <sup>a</sup>	0.8		N	•••	N
140129B	1.36	< 1.5	Y	Y	
$140320A^{e}$	0.45		Ÿ	N	
$140402A^{a}$	0.03		N	N	
140414A <sup>a</sup>	0.7		N	N	
140516A	0.19		Y	N	N
140606A <sup>a</sup>	0.34		N	N	
$140619B^{a}$	0.5		N	N	N
140622A	0.13	0.959	Y	N	N
140903A	0.30	0.351	Y	Y	Y
140930B	0.30	< 4.1	Y	Y	N
$141202A^{a}$	1.3		N		
141205A <sup>a</sup>	1.1		N	N	
141212A	0.30	0.596	Y	N	N
150101A	0.06		Y	•••	N
150101B	0.018	0.134	Ÿ	Y	N
150120A	1.20	0.460	Ÿ	N	N
$150301A^{a}$	0.48		Ÿ	•••	

**Note.** — Short GRBs with X-ray, optical/NIR or radio follow-up observations. "Y" = detection, "N" = non-detection, and  $\cdots$  means there is no follow-up in that band.

follow-up observations, 87 (84%) have optical/NIR observations, and 60 (58%) have radio observations. These observations have uncovered 71 X-ray afterglows, 30 optical/NIR afterglows, and 4 radio afterglows, leading to detection fractions of 74%, 34%, and 7%, respectively (Table 1). The observations for these bursts are catalogued in the Appendix (Tables A1-A3).

Thirty-one bursts in this sample have redshifts deter-

mined from their host galaxies (Table 1), while two events have spectroscopic redshifts from absorption in their afterglows, GRB 090426A: (Antonelli et al. 2009; Levesque et al. 2010) and GRB 130603B (Cucchiara et al. 2013; de Ugarte Postigo et al. 2014). Furthermore, we place upper limits on the redshift from the detection of the optical afterglow, and therefore the lack of suppression blueward of the Lyman limit ( $\lambda_0 = 912 \, \text{Å}$ ) or Lyman- $\alpha$  line ( $\lambda_0 = 1216 \, \text{Å}$ ), for 11 bursts (Table 1). We also place a lower limit of z > 1.4 for GRB 051210 from the lack of detection of emission or absorption features in the spectrum of the host galaxy (Berger et al. 2007). In addition to the broad-band afterglow observations that have been published thus far, we present new optical/NIR observations for 11 bursts (Table A2), and new radio observations for 25 events (Table A3).

### 2.1.1. Observing Constraints

Of the 25 bursts with X-ray observations and no detected X-ray afterglow, 18 events had a delayed Swift/XRT response due to an observing constraint or burst discovery from another satellite (indicated in Table 1); thus, the non-detection of X-ray afterglows for these events are due to factors unrelated to the bursts themselves. Likewise, 12 bursts with optical observations and no detected optical afterglow have an observing constraint (e.g., delayed precise localization from Swift, crowded field, high Galactic extinction sightline, position contaminated by a bright star), making the detection of an optical afterglow particularly challenging (Table A2). Only one burst (GRB 150101B) with radio observations is severely contaminated by a neighboring bright source, thus preventing a deep radio limit at the afterglow position. Taking these observing constraints into account, we find that the vast majority (91%) of bursts that have X-ray follow-up and no observing constraints result in an X-ray afterglow detection. Similarly, the fraction of detected optical afterglows increases to 40% after correcting for observing constraints. The radio detection fraction remains at  $\approx 7\%$ .

#### 2.2. *X-rays*

We gather all available X-ray afterglow observations from the Swift light curve repository<sup>4</sup> (Evans et al. 2007a, 2009), the GRB Coordinates Network (GCN) Circulars, and the literature (Table A1). The data were taken with the X-ray Telescope (XRT) on-board Swift, the Chandra X-ray Observatory, and the X-ray Multi-Mirror Mission (XMM-Newton). Ten bursts have Chandra observations, while three bursts have XMM-Newton observations (Table A1). We use unabsorbed fluxes and uncertainties in the 0.3 – 10 keV energy band when they are available; otherwise, we use the count-rate light curves in the same energy range and convert to fluxes (described later in this section). For bursts with multiple upper limits, we only include those which help to constrain the temporal behavior of the X-ray light curve. Of the 96 short GRBs with X-ray observations, four events have no reported measurements or upper limits<sup>5</sup>. Therefore, the resulting late-time X-ray afterglow catalog is comprised of 92 events (Table A1).

When applicable, we convert the count rate light curves to unabsorbed fluxes using the count-rate-to-unabsorbed-flux conversion factors provided by the *Swift* light curve repository. These factors are derived from the automatic spectral

<sup>&</sup>lt;sup>a</sup> Observing constraint or delayed *Swift*/XRT observations.

<sup>&</sup>lt;sup>b</sup> There are XRT observations but no X-ray flux upper limit is reported

<sup>&</sup>lt;sup>c</sup> This burst has  $T_{90} = 2.3 \pm 0.1$  s but is spectrally hard (Marshall et al. 2009).


<sup>&</sup>lt;sup>d</sup> This burst has  $T_{90} = 2.4 \pm 0.4$  s, but the spectral lag of  $5 \pm 15$  ms indicates this is a short-hard burst (Grupe et al. 2010).

<sup>&</sup>lt;sup>e</sup> Delayed reporting of burst position preventing immediate ground-based follow-up.

f IPN-localized burst with no Swift follow-up.

<sup>4</sup> http://www.swift.ac.uk/xrt\_curves

<sup>&</sup>lt;sup>5</sup> These events are GRBs 051105A, 081024B, 091117, and 110420B.



**Figure 1.** Broad-band afterglow light curves of all short GRBs with follow-up observations between November 2004 and March 2015. In all panels, circles denote detections, triangles indicate  $3\sigma$  upper limits, and solid lines connect observations for the same burst. *Top left:* 0.3−10 keV X-ray afterglow light curves for 92 short GRBs with X-ray observations corresponding to the events in Table A1. Included are  $3\sigma$  upper limits for 29 bursts with no detected X-ray emission at  $\delta t \gtrsim 1000$  sec (blue triangles). The data for  $\delta t \gtrsim 1000$  sec (grey vertical band) are used in our subsequent afterglow analysis. *Top right:* Optical afterglow observations for 87 short GRBs corresponding to the events in Table A2. The 30 bursts with detected optical afterglows are shown (orange circles), along with  $3\sigma$  upper limits for bursts with an X-ray afterglow detection (filled triangles), and bursts with no X-ray afterglow detection (open triangles). *Bottom left:* Radio afterglow data for 60 short GRBs corresponding to the events in Table A3. The light curves for four short GRBs with radio afterglow detections are shown: GRBs 050724A (black), 051221A (red), 130603B (light red), and 140903A (maroon). Upper limits for the remaining 56 events with no detected radio afterglows are shown (3 $\sigma$ ; grey triangles), including 25 bursts for which radio observations have not been published in the literature thus far.

fitting routine (Evans et al. 2009). This routine fits the X-ray spectrum for each burst to an absorbed power law model characterized by photon index,  $\Gamma$ , and the intrinsic neutral hydrogen absorption column,  $N_{\rm H,int}$ , in excess of the Galactic column density in the direction of the burst (Kalberla et al. 2005; Wakker et al. 2011; Willingale et al. 2013). We use spectral parameters extracted in the Photon Counting (PC) mode when available; otherwise, we use parameters from the Windowed Timing (WT) mode. In ten cases, the value of  $N_{\rm H,int}$  is highly uncertain, but consistent with zero. Therefore, utilizing the median value of  $N_{\rm H,int}$  may result in an over-estimate of the true unabsorbed flux. Instead of using the given conversion factors for these bursts, we calculate the unabsorbed fluxes using WebPIMMS<sup>6</sup>, setting  $N_{\rm H,int} = 0$ . For 16 events, no countrate-to-unabsorbed-flux conversion factor is available, so we

employ a fiducial value of  $1 \times 10^{-11}$  erg cm<sup>-2</sup> s<sup>-1</sup> ct<sup>-1</sup> set by the median value for all of the events in our sample. Applying these conversion factors to each of the count-rate light curves from *Swift/XRT*, *Chandra* and *XMM-Newton*, we obtain the unabsorbed fluxes,  $1\sigma$  uncertainties, and  $3\sigma$  upper limits for each burst (Table A1).

To enable comparison of the X-ray light curves to the optical and radio data, we convert the X-ray fluxes to flux densities,  $F_{\nu,X}$ , at a fiducial energy of 1 keV ( $F_{\nu,X} \propto \nu^{\beta x}$  where  $\beta_X \equiv 1 - \Gamma$ ). When no spectral information is available, we use a fiducial spectral index of  $\beta_{X,\text{med}} = -1.0$ , set by the median value of the events in our sample. The flux densities,  $1\sigma$  uncertainties, and  $3\sigma$  upper limits are listed in Table A1, and the resulting light curves are shown in Figure 1.

Finally, we present *Chandra* observations for three bursts that have not been published in the literature thus far: GRBs 120804A (PI: Burrows), 140930B (PI: Fong) and

<sup>6</sup> https://heasarc.gsfc.nasa.gov/Tools/w3pimms.html

150101B (PIs: Troja, Levan). We retrieve the pre-processed Level 2 data from the *Chandra* archive. We use the CIAO data reduction package to extract a count-rate within a 2.5"-radius source aperture centered on the X-ray afterglow position, and utilize source-free regions on the same chip to estimate the background. For GRB 120804A, we obtain spectral parameters from earlier *Chandra* epochs of the same burst (Berger et al. 2013). For GRBs 140930B and 150101B, we use CIAO/specextract to extract a spectrum and obtain the spectral parameters. We then apply these parameters to the count rate to convert to flux density as described above (Table A1).

#### 2.3. *Optical/NIR*

For each burst we gather all available optical and NIR afterglow observations from the literature and GCN Circulars (see Table A2 for references). When there are multiple upper limits for a given burst, we include only the deepest limits at  $\delta t \lesssim 1$  day. We convert all magnitudes to the AB system using instrument-specific conversion factors when available, or the standard conversions following Blanton & Roweis (2007). We correct all observations for Galactic extinction in the direction of each burst (Schlegel et al. 1998; Schlafly & Finkbeiner 2011), and convert AB magnitudes to flux densities,  $F_{\nu, \text{opt}}$ . A log of observations for 87 events with optical/NIR follow-up is provided in Table A2, and the light curves and upper limits are shown in Figure 1. Bursts with no detected optical afterglow are further classified by the detection of an X-ray afterglow (Figure 1).

We also present optical/NIR observations of 11 short GRBs that have not been published in the literature thus far: GRBs 070724A, 100628A, 110420B, 120229A, 130716A, 140402A, 140606A, 140619B, 140930B, 150101B, and 150120A. These observations were enabled by target-ofopportunity programs on the twin 6.5-m Magellan telescopes (PI: Berger), the twin 8-m Gemini telescopes (PIs: Berger, Cucchiara, Fox, Tanvir), and the 6.5-m MMT (PIs: Berger, Fong). We use standard tasks in the IRAF/ccdred package to process the Magellan and MMT data, and the IRAF/gemini package to process the Gemini data. For each of these bursts, we obtained the first epoch of observations at  $\delta t \approx 1-20$  hr and at least one additional set of observations at  $\delta t \gtrsim 24$  hr to provide a template (Table A2). To assess any fading between the two epochs, we perform digital image subtraction for each burst and filter using the ISIS software (Alard 2000). With the exception of GRBs 070724A, 140930B, and 150101B, we find no significant emission in any of the subtracted images. We therefore employ aperture photometry using standard tasks in IRAF to place  $3\sigma$  upper limits on the optical/NIR afterglow brightness. To measure the afterglow brightness for GRBs 070724A and 140903B, we employ aperture photometry on the detected point source in the subtracted image. For GRB 150101B, the afterglow position in the subtracted image is contaminated by residual emission from its bright host galaxy; thus, we use PSF photometry to measure the afterglow brightness. Details of the photometry will be outlined in an upcoming paper (Fong et al., in prep.) The observational details, afterglow brightness and  $3\sigma$  upper limits for these bursts are presented in Figure 1 and listed in Table A2.

#### 2.4. Radio

We gather all available radio afterglow data taken with the Karl G. Jansky Very Large Array (VLA), Westerbork Synthesis Radio Telescope (WSRT), Australia Telescope Compact Array (ATCA), and Combined Array for Research in Millimeter-wave Astronomy (CARMA). The resulting radio afterglow catalog is comprised of 60 short GRBs (Table A3). The large majority of events, 53 (88%), were observed with the VLA (Table A3), 28 of which were observed with the upgraded VLA, which has a ten-fold improvement in sensitivity (Perley et al. 2011). We present new observations enabled by target-of-opportunity programs with the upgraded VLA (PI: Berger) and CARMA (PI: Zauderer) for 25 bursts (Table A3). In 11 cases, we obtained multiple sets of observations to probe the radio emission on timescales spanning  $\delta t \approx 1-10$  days.

For data calibration and analysis of VLA observations, we follow standard procedures in the Astronomical Image Processing System (AIPS; Greisen 2003). For CARMA observations, we use the Multichannel Image Reconstruction, Image Analysis and Display (MIRIAD) software package (Sault et al. 1995). For the majority of cases, we do not find any uncatalogued radio sources in or around the X-ray or  $\gamma$ -ray positions. To calculate the  $3\sigma$  upper limits on the radio afterglow brightness, we measured the RMS noise in the map using a large source-free central region utilizing AIPS/IMSTAT for VLA data and MIRIAD/IMSTAT for CARMA data. The radio afterglow detections and upper limits for 60 short GRBs with radio observations are listed in Table A3 and displayed in Figure 1. Flux densities reported here supercede those reported in GCN Circulars.

Four bursts have detected radio afterglows, and all discoveries were made with the VLA (GRB 050724A: Berger et al. 2005; Panaitescu 2006; GRB 051221A: Soderberg et al. 2006; GRB 130603B: Fong et al. 2014; GRB 140903A: this work). In particular, we present the discovery of the radio afterglow of GRB 140903A, which is detected at three frequencies: 1.4, 6.0 and 9.8 GHz (Figure 1 and Table A3), and will be further discussed in an upcoming work.

#### 2.5. Afterglow Brightness and TOO Response Times

Thanks to Swift, X-ray and optical follow-up typically commences within  $\approx 60$  sec after the GRB is detected. While Swift/XRT is responsible for nearly all of the X-ray afterglow detections, only three short GRBs have detected optical afterglows with Swift/UVOT (Table A2). Thus, the detection of short GRB afterglows, or placement of meaningful upper limits, in the optical and radio bands relies on ground-based Target-of-Opportunity programs. For the 30 short GRBs with detected optical afterglows, the median optical afterglow brightness is  $\approx 23.0$  mag at  $\delta t \approx 7.0$  hr. The median  $3\sigma$  limit placed on bursts with an X-ray afterglow and no detected optical afterglow is  $\geq 23.8$  mag with a median response time of  $\delta t \approx 7.4$  hr, while the limit placed on bursts with no detected X-ray or optical afterglow is less constraining and more delayed,  $\gtrsim 22.7$  mag at  $\delta t \approx 12.2$  hr (Figure 1). This is likely due to the more limited number of facilities with instruments that can cover the comparatively larger  $\gamma$ -ray positional error circles. However, since we only include bursts with optical limits of  $\gtrsim 20$  mag, we are excluding a fraction of bursts with very shallow follow-up; thus the limits here for bursts with only  $\gamma$ -ray localizations are an optimistic representation of the entire population. In the radio band, the median  $3\sigma$  upper limit for all observations is  $\lesssim 74.1 \,\mu\text{Jy}$ , and the median response time to the first observation is  $\delta t \approx 24.7$  hr. In more recent cases, we set unprecedented  $3\sigma$  limits of  $15-20\,\mu\text{Jy}$ using the upgraded VLA on timescales of  $\delta t \approx 1-10$  days.

#### 3. BROAD-BAND AFTERGLOW ANALYSIS

We utilize the broad-band afterglow observations to constrain the explosion properties and circumburst environment of each burst. We adopt the standard synchrotron model for a relativistic blastwave in a constant density medium (Sari et al. 1998; Granot & Sari 2002), as expected for a non-massive star progenitor. This model provides a mapping from the broadband afterglow flux densities to the burst physical parameters: the isotropic-equivalent kinetic energy ( $E_{\rm K,iso}$ ), circumburst density (n), fractions of post-shock energy in radiating electrons ( $\epsilon_e$ ) and magnetic fields ( $\epsilon_B$ ), and the electron power-law distribution index (p), with  $N(\gamma) \propto \gamma^{-p}$  for  $\gamma \gtrsim \gamma_{\rm min}$ , where  $\gamma_{\rm min}$  is the minimum Lorentz factor of the electron distribution.

The synchrotron spectrum is characterized by a flux normalization and three break frequencies: the self-absorption frequency  $(\nu_a)$ , the peak frequency  $(\nu_m)$ , and the cooling frequency  $(\nu_c)$ . Constraints on the physical parameters require knowledge of where the synchrotron break frequencies are located with respect to the observing bands. In most cases, there is not enough information to constrain the locations of  $\nu_a$  and  $\nu_m$  with respect to the observing bands, so we make assumptions about their locations (detailed in the next sections). However, there are several cases in which we can use the available data to determine the location of  $\nu_c$  with respect to the X-ray and optical bands. To determine the location of  $\nu_c$ , we first determine the temporal and spectral power-law indices ( $\alpha$  and  $\beta$ , respectively, where  $F_{\nu} \propto t^{\alpha} \nu^{\beta}$ ) from the X-ray and optical light curves and spectra. We then compare these indices to the standard relations given by the synchrotron model to determine whether  $\nu_c$  is located above or below the X-ray band. This also allows us to calculate the value of p, and governs how the fluxes map to the burst physical properties (Granot & Sari 2002).

## 3.1. Temporal and Spectral Behavior

3.1.1. *X-rays* 

To investigate the temporal behavior of the X-ray afterglows, we utilize  $\chi^2$ -minimization to fit a single power law model to each light curve in the form  $F_{\nu,X} \propto t^{\alpha_X}$ , with temporal index  $\alpha_X$  as the single free parameter and the best-fit flux normalization  $C_0$  given by

$$C_{0} = \frac{\sum_{i=1}^{N} \frac{F_{\text{model},i} \times F_{X,i}}{\sigma_{X,i}^{2}}}{\sum_{i=1}^{N} \frac{F_{\text{model},i}^{2}}{\sigma_{X,i}^{2}}},$$
(1)

where  $F_{\text{model},i}$  are the un-normalized model fluxes,  $F_{X,i}$  and  $\sigma_{X,i}$  are the observed fluxes and uncertainties, respectively, and N is the number of data points. Since early-time Xray afterglow light curves are often subject to steep decays, plateaus, or flares which may contaminate the afterglow emission (Nousek et al. 2006; Zhang et al. 2006; Margutti et al. 2011, 2013), we only utilize X-ray data at  $\delta t \gtrsim 1000$  s, when bursts have typically settled into the power-law afterglow phase. For the X-ray light curves, we initially include all of the available data at  $\delta t \gtrsim 1000$  s in the fit. In a few cases, there are light curve features beyond  $\delta t \approx 1000 \text{ s}$  which significantly affect the fit: flares (GRBs 050724A and 111121A), plateaus (GRB 051221A), or steepenings (GRBs 051221A and 111020A). For these bursts, we exclude the time intervals that contain such features in the fits. In 10 cases, there only exists a single detection and an upper limit at  $\delta t \gtrsim 1000$  s,

Table 2
Short GRB Spectral and Temporal Power-law Indices

GRB	$\alpha_X$	$\beta_X$	$lpha_{ m opt}$	$eta_{ m opt}$
050509B	$-1.10 \pm 0.25$	$-0.88 \pm 0.34$		
050709	$-1.23\pm0.10$	$-1.24 \pm 0.35$	$-1.42\pm0.08$	• • •
050724A	$-0.93 \pm 0.08$	$-0.81 \pm 0.15$	$-1.74 \pm 0.11^a$	$-0.82 \pm 0.03^a$
050813	< -0.004	$-1.3^{+2.1}_{-1.3}$	• • •	
051210	• • •	$-2.1 \pm 0.5$	• • •	
051221A	$-1.08 \pm 0.12$	$-1.0 \pm 0.2$	$-0.97\pm0.06$	• • •
060121	$-1.23 \pm 0.20$	$-1.07\pm0.16$	$-0.60\pm0.24$	
060313	$-1.47 \pm 0.39$	$-0.96 \pm 0.09$	$-0.70 \pm 0.19$	$-1.35 \pm 0.19$
060502B	• • •	$-2.07^{+1.50}_{-0.54}$	• • •	
060801	• • •	$-0.68\pm0.12$	• • •	• • •
061006	$-0.85 \pm 0.30$	$-0.90 \pm 0.25$	$-0.43 \pm 0.08$	
061201	$-1.96 \pm 1.18$	$-0.66 \pm 0.12$	• • •	
061210	$-1.71 \pm 1.15$	$-1.86^{+1.26}_{-0.61}$		
070429B	< -1.12	$-2.0 \pm 0.63$	• • •	• • •
070707	< -0.65	• • •	$-2.55\pm0.22$	• • •
070714B	$-1.96 \pm 0.69$	$-1.07 \pm 0.19$	$-0.81 \pm 0.11$	
070724A	$-0.95 \pm 0.33$	$-0.60 \pm 0.25$	< -0.15	$-0.58\pm0.02$
070729		$-0.5^{+0.44}_{-0.25}$		
070809	$-1.09 \pm 1.10$	$-0.37^{+0.13}_{-0.07}$	$-0.73 \pm 0.33$	
071227	$-0.97 \pm 0.27$	$-0.90 \pm 0.31$	< -0.24	
080123	$-0.77 \pm 0.19$	$-1.6 \pm 0.4$		
080426	$-1.54 \pm 0.33$	$-1.03 \pm 0.16$		
080503		$-1.53^{+0.26}_{-0.12}$		
080702A	< -0.38	$-1.00 \pm 0.43$		
080905A		$-0.53 \pm 0.18$	$-0.39 \pm 1.36$	
080919	$-1.23 \pm 0.69$	$-1.9^{+0.63}_{-0.38}$		
081024A		$-0.70 \pm 0.38$		
081226A		$-2.27^{+1.24}_{-0.32}$	$-0.95 \pm 0.30$	$-1.67 \pm 0.67^b$
090305A			$-0.74 \pm 0.09$	$-0.71 \pm 0.27$
090426A	$-1.15 \pm 0.16$	$-1.04 \pm 0.09$	$-0.58 \pm 0.16$	$-0.94 \pm 0.06^b$
090510	$-1.79 \pm 0.63$	$-0.75 \pm 0.08$	$-2.37 \pm 0.29$	$-0.85 \pm 0.05$
090515		$-1.53^{+0.78}_{-0.28}$	< -0.12	
090607	< -0.72	$-1.2 \pm 0.4$		
090621B	$-1.48 \pm 0.54$	$-2.7^{+0.68}_{-1.0}$		
091109B	$-0.83 \pm 0.28$	$-1.1 \pm 0.3$	$-0.49 \pm 0.45$	
100117A		$-1.6 \pm 0.3$	$-1.60 \pm 0.33$	
100206A		$-2.05^{+1.07}_{-0.53}$		
100213		$-2.0^{+1.2}_{-1.0}$		
100625A	< 0.37	$-1.5 \pm 0.2$		
100702A		$-1.7 \pm 0.3$		
101219A	$-1.37 \pm 0.13$	$-0.8 \pm 0.1$		
101224A		$-2.4^{+1.8}_{-0.6}$		
110112A	$-1.10 \pm 0.05$	$-1.2 \pm 0.2$	< -0.32	
111020A	$-0.78 \pm 0.05$	$-1.04 \pm 0.16$		
111117A	$-1.21\pm0.05$	$-1.0 \pm 0.2$		
111121A	$-1.55 \pm 0.30$	$-0.90 \pm 0.12$		
111222A	< -0.30	$-0.7^{+1.4}_{-0.4}$		
120305A		$-2.2^{+0.6}_{-0.4}$		
120521A		$-0.8 \pm 0.25$		
120630A		$-0.8 \pm 0.3$		
120804A	$-1.02 \pm 0.10$	$-1.1 \pm 0.1$		
121226A	$-1.12 \pm 0.28$	$-1.5 \pm 0.25$		
130313A		$-1.6^{+2.1}_{-2.5}$		
130515A		$-0.7 \pm 0.31$		
130603B	$-1.88 \pm 0.15$	$-1.2 \pm 0.1$	$-1.26 \pm 0.05$	$-2.0 \pm 0.1^{b}$
130716A	< -0.52	$-1.1^{+0.56}_{-0.38}$		
		0.38		

Table 2 — Continued

GRB	$\alpha_X$	$eta_X$	$lpha_{ m opt}$	$eta_{ m opt}$
130822A 130912A 131004A 140129B 140320A 140516A 140622A 140903A 140930B	$\begin{array}{c} \dots \\ -1.33 \pm 0.18 \\ -1.1 \pm 0.61 \\ -1.6 \pm 0.26 \\ < -0.23 \\ -0.42 \pm 0.14 \\ < -0.16 \\ -1.05 \pm 0.20 \\ -1.27 \pm 0.15 \end{array}$	$\begin{array}{c} -0.89^{+1.1}_{-0.34} \\ -0.57 \pm 0.13 \\ -0.8 \pm 0.3 \\ -0.97 \pm 0.11 \\ -3.8^{+2.6}_{-0.88} \\ -0.70 \pm 0.44 \\ -0.55^{+0.42}_{-0.18} \\ -0.60 \pm 0.10 \\ -0.73 \pm 0.28 \end{array}$	$ \begin{array}{c}\\ < -0.42\\ -1.94 \pm 0.25\\ -1.54^c\\\\\\\\\\\\\\\\ .$	$$ $-1.52 \pm 0.13^{b}$ $$ $$ $$ $$ $$
141212A 150101A 150101B 150120A 150301A	<-0.12 -1.07 ± 0.15 	$-1.7^{+2.75}_{-0.81}$ $-0.40 \pm 0.56$ $-0.67 \pm 0.17$ $-1.0 \pm 0.31$ $-0.7 \pm 0.13$	-1.01 ± 0.62 	

**Note.** — Error bars correspond to  $1\sigma$  confidence. When applicable,  $\alpha_X$  and  $\alpha_{\rm opt}$  represent pre-jet break values. Values of  $\beta_{\rm opt}$  are observed values and are uncorrected for intrinsic rest-frame extinction,  $A_V$ .

so we can only extract an upper limit for  $\alpha_X$ . The resulting best-fit values for  $\alpha_X$ , along with  $1\sigma$  uncertainties, are listed in Table 2. Also listed are the X-ray spectral indices,  $\beta_X$ , from the relation  $\beta_X \equiv 1-\Gamma$ . The X-ray afterglows have weighted mean values and  $1\sigma$  uncertainties of  $\langle \alpha_X \rangle = -1.07^{+0.31}_{-0.37}$  and  $\langle \beta_X \rangle = -0.89^{+0.38}_{-0.78}$ , and median values of  $\alpha_{X,\text{med}} \approx -1.16$  and  $\beta_{X,\text{med}} \approx -1.06$ .

#### 3.1.2. Optical

We determine the temporal index of the optical observations  $(\alpha_{\rm opt}, \text{ where } F_{\nu, {\rm opt}} \propto t^{\alpha_{\rm opt}})$  in the same manner as described in Section 3.1.1, using the filter with the most well-sampled light curve for each burst. If there are multiple filters for which we can determine  $\alpha_{opt}$ , we independently fit  $\alpha_{opt}$  for each filter and report the weighted mean (Table 2). A few short GRBs have well-measured optical spectral indices from contemporaneous multi-band data, but do not have a well-sampled light curve in a single filter, preventing a measurement of  $\alpha_{\rm opt}$ . For these events, we use the measured value of  $\beta_{\text{opt}}$  (see below) to extrapolate all of the available afterglow data to a single filter, and then determine the temporal decay index from these observations. In this manner, we are able to measure the optical temporal decay index for 19 short GRBs, and place upper limits in five cases for bursts with only a single detection and an upper limit. The best-fit optical temporal indices and  $1\sigma$ uncertainties are listed in Table 2. The weighted mean for the 19 short GRBs with measured values is  $\langle \alpha_{\rm opt} \rangle = -1.07^{+0.31}_{-0.61}$ ,

and the median is  $\alpha_{\rm opt,med} \approx -0.99$ .

If there are contemporaneous observations in multiple filters, we use these to determine the observed spectral slope,  $\beta_{\rm opt}~(F_{\nu,\rm opt} \propto \nu^{\beta_{\rm opt}}).$  For a few bursts, there are multi-band observations taken at different times, as well as a measurement of  $\alpha_{\rm opt}$  from a well-sampled light curve in a single filter; in such cases, we use  $\alpha_{\rm opt}$  to interpolate the data from multiple filters to a common time. To determine  $\beta_{\rm opt}$ , we then use  $\chi^2$ -minimization to fit the optical/NIR photometry to a power law model. The values of  $\beta_{\rm opt}$  are listed in Table 2. The weighted mean for all short GRBs with determined spectral indices after incorporating rest-frame extinction (Section 3.1.3) is  $\langle\beta_{\rm opt}\rangle=-0.71^{+0.25}_{-0.51}$ , and the median is  $\beta_{\rm opt,med}\approx-0.88$ .

#### 3.1.3. Rest-Frame Extinction

In four cases where there are contemporaneous observations in multiple optical/NIR filters, a single power law model provides a poor fit to the broad-band photometry ( $\chi^2_{\nu} \gtrsim 3$ ). For these bursts, we include the line-of-sight rest-frame extinction from the host galaxy ( $A_V^{\rm host}$ ) as a second free parameter in the fit. We constrain the extinction using the Milky Way extinction curve (Cardelli et al. 1989), and employ the burst redshifts (Table 1) to obtain the rest-frame extinction. For bursts with no determined spectroscopic redshift, we assume z=0.5 set by the median of the short GRB population (Berger 2014). We find non-zero extinction values for GRBs 060121, 070724A, 081226A, and 130603B (Table 3). The resulting values for the optical spectral indices, uncertainties, and  $A_V^{\rm host}$  are listed in Table 3.

For bursts where we do not have enough information from the optical/NIR bands alone to constrain the spectral behavior of the afterglow, we initially assume that there is no rest-frame extinction,  $A_V^{\text{host}} = 0$ . However, in eight cases, the difference in slope between the X-ray and optical bands is shallower than expected (e.g.,  $|\beta_{OX}| \lesssim |\beta_X| - 0.5$ , where  $\beta_{OX}$  is the spectral index between the X-ray and optical bands), suggesting that either there is intrinsic extinction, or that the X-rays do not originate from the forward shock. Assuming the former explanation, we include the minimum amount of extinction required until the X-ray and optical solutions agree to within the  $1\sigma$  uncertainties. In two cases, GRBs 080919 and 101219A, there is only an upper limit on the optical afterglow brightness; thus we can only determine a lower limit on the restframe extinction. We note that GRB 080919 has the highest value of rest-frame extinction, with  $A_V^{\text{host}} \gtrsim 6$  mag. However, this burst has a sightline close to the Galactic plane and therefore has a highly uncertain Galactic extinction, which likely affects the inferred value for  $A_V^{\text{host}}$ . For the 12 events with rest-frame extinction, we find values of  $A_V^{\text{host}} \approx 0.3 - 1.5 \text{ mag.}$ These  $A_V^{\text{host}}$  values are also listed in Table 3.

# 3.2. Determination of Electron Power Law Index and the Location of the Cooling Frequency

<sup>&</sup>lt;sup>a</sup> These values are computed over the same time interval as the X-ray flare that is super-imposed on the underlying afterglow power-law decay. No optical detections exist for the underlying afterglow.

<sup>&</sup>lt;sup>b</sup> This value corresponds to the spectral behavior after the jet break.

<sup>&</sup>lt;sup>c</sup> No uncertainties are given for the optical light curve.

FONG ET AL.

Table 3
Inferred Properties

	li							
GRB	$A_V^{ m host}$	$\nu_c < \nu_X$ ?	p	$\epsilon_B$	$E_{\gamma, \text{iso}, 52}$	$\langle E_{ m K,iso,52} \rangle$	$\eta_{\gamma}$	< n >
	(mag)				(10 <sup>52</sup> erg)	(10 <sup>52</sup> erg)		(cm <sup>-3</sup> )
050709	0	Y	$2.31 \pm 0.13$	0.1	0.09	$2.6^{+0.8}_{-0.5} \times 10^{-3}$	0.97	$1.0^{+0.5}_{-0.4}$
	0	Y	$2.31 \pm 0.13$	0.01	0.09	$6.2^{+0.4}_{-0.6} \times 10^{-3}$	0.93	$1.6^{+0.2}_{-0.2}$
050724A	0	N	$2.29 \pm 0.10$	$10^{-4}$	0.24	$0.18^{+0.08}_{-0.05}$	0.58	$0.89^{+0.58}_{-0.49}$
051221A	0	Y	$2.24\pm0.07$	0.1	1.3	$0.16^{+0.01}_{-0.01}$	0.89	$0.03^{+0.006}_{-0.005}$
	0	Y	$2.24\pm0.07$	0.01	1.3	$0.27^{+0.03}_{-0.03}$	0.83	$0.14^{+0.05}_{-0.04}$
060121	1.6	Y	$2.24\pm0.20$	0.1	4.5	$0.20^{+0.05}_{-0.04}$	0.96	$5.4^{+5.6}_{-2.2} \times 10^{-3}$
	1.6	Y	$2.24\pm0.20$	0.01	4.5	$0.23^{+0.05}_{-0.04}$	0.95	$0.16^{+0.17}_{-0.06}$
$060313^{ab}$	0	Y	$2.03 \pm 0.20$	0.1	2.9	$0.45^{+0.05}_{-0.05}$	0.87	$3.3^{+1.0}_{-0.5} \times 10^{-3}$
061006	0	N	$2.39 \pm 0.31$	0.1	1.1	$0.64^{+0.85}_{-0.37}$	0.63	$2.2^{+16}_{-1.9}\times10^{-5}$
	0	N	$2.39 \pm 0.31$	0.01	1.1	$1.1^{+2.1}_{-0.7}$	0.50	$1.2^{+29}_{-1.1}\times10^{-4}$
061201	0	N	$2.35 \pm 0.24$	0.1	0.05	$0.05^{+0.10}_{-0.03}$	0.47	$5.0^{+66}_{-4.6}\times10^{-5}$
	0	N	$2.35 \pm 0.24$	0.01	0.05	$0.1^{+0.4}_{-0.09}$	0.29	$2.7^{+120}_{-2.6}\times10^{-4}$
$070714\mathrm{B}^{bc}$	0.5	Y	$2.30\pm0.35$	0.1	1.7	$0.1^{+0.02}_{-0.02}$	0.94	$0.056^{+0.024}_{-0.011}$
070724A	1.5	N	$2.24\pm0.33$	0.1	0.03	$0.35^{+0.49}_{-0.20}$	0.07	$1.9^{+12}_{-1.6}\times10^{-5}$
	2.0	N	$2.24\pm0.33$	0.01	0.03	$1.1^{+3.0}_{-0.8}$	0.02	$9.3^{+210}_{-9.2}\times10^{-5}$
070809	0	N	$2.12\pm1.47^d$	0.1	0.09	$0.5^{+0.7}_{-0.3}$	0.14	$2.2^{+15}_{-1.9}\times10^{-5}$
	0	N	$2.12\pm1.47^d$	0.01	0.09	$1.1^{+2.7}_{-0.8}$	0.07	$1.2^{+30}_{-1.1}\times10^{-4}$
071227	0	Y	$\boldsymbol{1.92 \pm 0.31}$	0.1	0.14	$8.4^{+2.5}_{-1.9}\times10^{-3}$	0.94	$1.9^{+2.4}_{-1.1}$
	0	Y	$1.92\pm0.31$	0.01	0.14	$8.9^{+2.6}_{-1.8}\times10^{-3}$	0.94	$60^{+75}_{-33}$
080426	0	Y	$2.29 \pm 0.26$	0.1	0.35	$0.05^{+0.01}_{-0.01}$	0.87	$0.04^{+0.04}_{-0.02}$
	0	Y	$2.29 \pm 0.26$	0.01	0.35	$0.06^{+0.01}_{-0.01}$	0.85	$1.2^{+1.2}_{-0.5}$
080905A	0	N	$2.06\pm0.36$	0.1	0.02	$0.04^{+0.12}_{-0.03}$	0.34	$1.3^{+33}_{-1.2} \times 10^{-4}$
	0	N	$2.06\pm0.36$	0.01	0.02	$0.08^{+0.44}_{-0.07}$	0.21	$7.1^{+610}_{-7.1}\times10^{-4}$
080919	$\gtrsim 6$	Y	$2.97 \pm 0.68$	0.1	0.07	$0.019^{+0.005}_{-0.004}$	0.78	$0.20^{+0.19}_{-0.07}$
	$\gtrsim 6$	Y	$2.97 \pm 0.68$	0.01	0.07	$0.029^{+0.007}_{-0.006}$	0.70	$5.1^{+4.5}_{-1.7}$
081024A	0	N	$2.40\pm0.76$	0.1	0.11	$0.11^{+0.22}_{-0.07}$	0.51	$8.1^{+150}_{-7.7} \times 10^{-5}$
	0	N	$2.40\pm0.76$	0.01	0.11	$0.25^{+0.92}_{-0.20}$	0.31	$4.3^{260}_{-4.2} \times 10^{-4}$
081226A	1.0	N	$2.27 \pm 0.39$	0.1	0.09	$0.20^{+0.36}_{-0.13}$	0.32	$3.2^{+29}_{-2.9} \times 10^{-5}$
	1.0	N	$2.27 \pm 0.39$	0.01	0.09	$0.43^{+1.5}_{-0.33}$	0.18	$1.7^{+54}_{-1.6} \times 10^{-4}$
090426A	0	Y	$2.13 \pm 0.14$	0.1	2.0	$1.4^{+0.4}_{-0.3}$	0.59	$0.04^{+0.04}_{-0.02}$
	0	Y	$2.13 \pm 0.14$	0.01	2.0	$1.5^{+0.4}_{-0.3}$	0.57	$1.2^{+1.4}_{-0.6}$
090510	0	N	$2.65 \pm 0.08$	0.1	0.77	$0.77^{+0.57}_{-0.33}$	0.50	$1.2^{+5.5}_{-1.0} \times 10^{-5}$
	0	N	$2.65 \pm 0.08$	0.01	0.77	$1.9^{+3.0}_{-1.2}$	0.29	$6.4^{+100}_{-6.0} \times 10^{-5}$
090607	0	Y	$2.40\pm0.76$	0.1	0.10	$2.1^{+0.4}_{-0.3}\times10^{-3}$	0.98	$0.84^{+0.52}_{-0.22}$
	0	Y	$2.40\pm0.76$	0.01	0.10	$2.6^{+0.5}_{-0.4}\times10^{-3}$	0.98	$24^{+15}_{-6.3}$
$090621B^c$	2.5	Y	$2.64 \pm 0.72$	0.1	0.07	$0.023^{+0.007}_{-0.006}$	0.75	$0.05^{+0.06}_{-0.02}$
	2.5	Y	$2.64 \pm 0.72$	0.01	0.07	$0.031^{+0.009}_{-0.007}$	0.68	$1.0^{+0.52}_{-0.27}$

 Table 3
 — Continued

GRB	A <sub>V</sub> <sup>host</sup> (mag)	$ u_c < \nu_X$ ?	p	$\epsilon_B$	$E_{\gamma,\mathrm{iso},52}$ (10 <sup>52</sup> erg)	$< E_{K,iso,52} >$ (10 <sup>52</sup> erg)	$\eta_{\gamma}$	< n >  (cm <sup>-3</sup> )
091109B	0	N	$2.40 \pm 0.32$	0.1	0.18	$0.25^{+0.18}_{-0.16}$	0.42	$2.8^{+24}_{-2.5} \times 10^{-5}$
	0	N	$2.40 \pm 0.32$	0.01	0.18	$0.8^{+2.2}_{-0.6}$	0.18	$9.4^{200}_{-9.0} \times 10^{-5}$
100117A	0	Y	$2.36 \pm 0.30$	0.1	0.22	$0.019^{+0.003}_{-0.003}$	0.92	$0.04^{+0.03}_{-0.01}$
	0	Y	$2.36 \pm 0.30$	0.01	0.22	$0.023^{+0.004}_{-0.004}$	0.90	$1.2^{+0.9}_{-0.3}$
101219A	$\gtrsim 2.5$	N	$2.73 \pm 0.13$	0.1	0.74	$0.3^{+0.5}_{-0.2}$	0.68	$4.6^{+59}_{-4.3} \times 10^{-5}$
	$\gtrsim 2.5$	N	$2.73 \pm 0.13$	0.01	0.74	$0.87^{+2.3}_{-0.64}$	0.46	$2.4^{+97}_{-2.3} \times 10^{-4}$
$110112A^{b}$	0	Y	$2.49 \pm 0.07$	0.1	0.03	$0.064^{+0.008}_{-0.005}$	0.31	$2.4^{+0.4}_{-0.4} \times 10^{-2}$
111020A <sup>abc</sup>	0.5	Y	$2.08 \pm 0.32$	0.1	0.17	$0.48^{+0.09}_{-0.08}$	0.26	$4.5^{+6.0}_{-3.8} \times 10^{-3}$
111117A	0	Y	$2.27 \pm 0.07$	0.1	0.55	$0.06^{+0.01}_{-0.01}$	0.90	$8.3^{+5.9}_{-2.3} \times 10^{-3}$
	0	Y	$2.27 \pm 0.07$	0.01	0.55	$0.07^{+0.02}_{-0.01}$	0.89	$0.25^{+0.16}_{-0.07}$
111121A	N/A	N	$2.87 \pm 0.21$	0.1	2.1	$3.1^{+1.9}_{-1.2}$	0.40	$8.2^{+27}_{-6.3} \times 10^{-6}$
	N/A	N	$2.87 \pm 0.21$	0.01	2.1	$8.3^{+11.1}_{-4.8}$	0.20	$4.2^{+48}_{-3.8} \times 10^{-5}$
120804A <sup>c</sup>	2.5	Y	$2.08 \pm 0.11$	0.1	3.4	$1.1^{+0.3}_{-0.2}$	0.76	$3.2^{+3.1}_{-1.5} \times 10^{-3}$
	2.5	Y	$2.08 \pm 0.11$	0.01	3.4	$2.3^{+0.2}_{-0.2}$	0.60	$0.014^{+0.002}_{-0.001}$
$121226 \mathrm{A}^{ab}$	1	Y	$2.50 \pm 0.37$	0.1	0.37	$0.6^{+0.08}_{-0.09}$	0.37	$4.0^{+1.0}_{-0.6} \times 10^{-3}$
130603B	1.2	Y	$2.70\pm0.06$	0.1	0.37	$0.11^{+0.02}_{-0.01}$	0.77	$0.09^{+0.04}_{-0.03}$
	0.3	Y	$2.70 \pm 0.06$	0.01	0.37	$0.15^{+0.02}_{-0.02}$	0.72	$0.31^{+0.08}_{-0.04}$
$130912A^{c}$	1.3	N	$2.49 \pm 0.17$	0.1	0.16	$0.50^{+0.37}_{-0.20}$	0.25	$1.7^{+9.9}_{-1.5}\times10^{-5}$
	1.3	N	$2.49 \pm 0.17$	0.01	0.16	$1.4^{+0.2}_{-0.8}$	0.11	$5.2^{+7.6}_{-4.9}\times10^{-5}$
131004A	0	N	$2.57 \pm 0.48$	0.1	0.45	$1.2^{+0.56}_{-0.36}$	0.28	$1.2^{+5.7}_{-1.0}\times10^{-5}$
	0	N	$2.57 \pm 0.48$	0.01	0.45	$2.8^{+3.4}_{-1.5}$	0.45	$6.5^{+11}_{-6.2} \times 10^{-4}$
140129B	0	N	$3.00 \pm 0.19$	0.1	0.07	$0.98^{+1.14}_{-0.53}$	0.06	$3.1^{+29.4}_{-2.8} \times 10^{-5}$
	0	N	$3.00\pm0.19$	0.01	0.07	$3.8^{+12.0}_{-2.9}$	0.02	$1.6^{+47}_{-1.5}\times10^{-4}$
140516A	0	N	$2.40\pm0.88$	0.1	0.02	$0.02^{+0.04}_{-0.01}$	0.54	$1.0^{+24}_{-0.99}\times10^{-4}$
	0	N	$2.40\pm0.88$	0.01	0.02	$0.04^{+0.18}_{-0.03}$	0.34	$5.5^{+41}_{-5.4} \times 10^{-4}$
140622A	0	N	$2.10\pm0.60$	0.1	0.07	$0.12^{+0.26}_{-0.08}$	0.36	$5.8^{+8.8}_{-5.4} \times 10^{-5}$
	0	N	$2.10\pm0.60$	0.01	0.07	$0.25^{+0.97}_{-0.20}$	0.21	$3.2^{+16.2}_{-3.2} \times 10^{-4}$
140903A	0	N	$2.27 \pm 0.16$	$10^{-3}$	0.08	$2.9^{+0.92}_{-0.74}$	0.03	$3.4^{+2.9}_{-1.6} \times 10^{-3}$
140930B	0	N	$2.67 \pm 0.19$	0.1	0.40	$0.28^{+0.42}_{-0.17}$	0.58	$4.6^{+58}_{-4.3}\times10^{-5}$
	0	N	$2.67 \pm 0.19$	0.01	0.40	$1.8^{+1.7}_{-0.9}$	0.18	$1.8^{11}_{-1.5}\times10^{-5}$
$150101B^c$	0.5	N	$2.40\pm0.17$	0.1	$4.0 \times 10^{-3}$	$0.61^{+0.45}_{-0.25}$	$6.6 \times 10^{-3}$	$8.0^{+24}_{-6.1}\times10^{-6}$
	0.5	N	$2.40 \pm 0.17$	0.1	$4.0 \times 10^{-3}$	$3.0^{+1.7}_{-1.1}$	$1.3 \times 10^{-3}$	$5.3^{+11}_{-3.6} \times 10^{-6}$

**Note.** — Quoted uncertainties are  $1\sigma$ . All solutions presented here are for a fixed  $\epsilon_e = 0.1$  and assume a lower density bound of  $n_{\min} = 10^{-6} \text{ cm}^{-3}$ .


To determine the electron power law index p, we use  $\alpha_X$  and  $\beta_X$  to constrain the location of the cooling frequency,  $\nu_c$ , with respect to the X-ray band. We use the relations given by Granot & Sari (2002) which relate  $\alpha_X$  and  $\beta_X$  to the value of p for the scenarios  $\nu_m < \nu_X < \nu_c$  and  $\nu_c < \nu_X$  by

<sup>&</sup>lt;sup>a</sup> We assume a redshift of z = 1 for this burst.

<sup>&</sup>lt;sup>b</sup> No valid solution is found for  $\epsilon_B = 0.01$ .

 $<sup>^{</sup>c}$  Value of  $A_{V}^{\mathrm{host}}$  is determined from a comparison of the optical and X-ray bands, and not directly from the optical/NIR SED.

 $<sup>^</sup>d$  Determined from  $\alpha_X$  alone.



**Figure 2.** Histogram of electron power law index, p, for 38 short GRBs as inferred from the X-ray temporal and spectral indices. The weighted mean and  $1\sigma$  uncertainties for the population (blue arrow) is  $\langle p \rangle = 2.43^{+0.36}_{-0.28}$ . These values correspond to those listed in Table 3.

$$p = \begin{cases} 1 - 2\beta_X \\ 1 - \frac{4}{3}\alpha_X \end{cases} \qquad \nu_m < \nu_X < \nu_c \tag{2}$$

$$p = \left\{ \begin{array}{c} -2\beta_X \\ \frac{2 - 4\alpha_X}{3} \end{array} \right\} \qquad \nu_c < \nu_X. \tag{3}$$

For a given burst, we calculate the values of p and  $1\sigma$  uncertainties using Equations 2 and 3 and standard propagation of errors. We select the valid scenario under the condition that the values of p independently determined from  $\alpha_X$  and  $\beta_X$  for a given scenario agree within the  $1\sigma$  uncertainties. Following this condition, we can constrain the location of  $\nu_c$  with respect to the X-ray band for 38 bursts (Table 3). Using Equations 2 and 3, we calculate the weighted mean for the value of p in the valid scenario; the resulting values and uncertainties are listed in Table 3 and displayed in Figure 2. We note that in the case of GRB 071227, the condition is satisfied for  $\nu_c < \nu_X$ , but has a median of  $p = 1.92 \pm 0.31$ , which yields a divergent total integrated energy. Thus for this burst, we employ p = 2.05 in our subsequent analysis (Table 3).

Under the reasonable assumption that the optical band lies between the peak frequency,  $\nu_m$ , and the cooling frequency,  $\nu_c$  (i.e.,  $\nu_m < \nu_{\rm opt} < \nu_c$ ), we use the available values for  $\alpha_{\rm opt}$  and  $\beta_{\rm opt}$  and Equation 2 to independently determine the value of p. We then include this in our weighted average of p for each burst (Figure 2). The weighted mean for the sample of 38 bursts is  $\langle p \rangle = 2.43^{+0.36}_{-0.28} \, (1\sigma; \text{Figure 2})$ . For the remaining bursts in the afterglow catalog, there is

For the remaining bursts in the afterglow catalog, there is not enough information to determine the location of the cooling frequency with respect to the X-ray band, and therefore the correct value of p. We thus concentrate on the subset of 38 bursts with determined values of p for our subsequent analysis. We find  $\nu_c < \nu_X$  for 18 cases, while  $\nu_c > \nu_X$  for 20 cases.

# 3.3. The Isotropic-Equivalent Kinetic Energies and Densities for Individual Bursts

In the standard synchrotron model from Granot & Sari (2002), the dependencies on the isotropic-equivalent kinetic energy, circumburst density, and the microphysical parameters are as follows for a given flux density,  $F_{\nu,i}$  and observing band,  $\nu_i$ :

$$F_{\nu,i} \propto \begin{cases} n_0^{1/2} E_{\text{K,iso,52}}^{5/6} \epsilon_{e,-1}^{-2/3} \epsilon_{B,-1}^{1/3} & \nu_a < \nu_i < \nu_m \\ n_0^{1/2} E_{\text{K,iso,52}}^{\frac{3+p}{4}} \epsilon_{e,-1}^{p-1} \epsilon_{B,-1}^{\frac{1+p}{4}} & \nu_m < \nu_i < \nu_c \\ E_{\text{K,iso,52}}^{\frac{2+p}{4}} \epsilon_{e,-1}^{p-1} \epsilon_{B,-1}^{\frac{p-2}{4}} & \nu_i > \nu_c \end{cases}$$
(4)

where  $n_0$  is in units of cm<sup>-3</sup>,  $E_{K,iso,52}$  is in units of  $10^{52}$  erg, and  $\epsilon_{e,-1}$  and  $\epsilon_{B,-1}$  are in units of 0.1. In addition to these four parameters,  $F_{\nu,i}$  is also dependent on the redshift, luminosity distance,  $\nu_i$ , time after the burst,  $\delta t$ , and the value of p; the exact dependencies are provided in Granot & Sari (2002). We note that for  $\nu_i > \nu_c$ , the flux density is independent of circumburst density. For bursts with no spectroscopic redshift, we assume z=0.5. In all cases, we cannot independently constrain  $\epsilon_e$  and  $\epsilon_B$  since this requires knowledge of the locations of the three break frequencies ( $\nu_a$ ,  $\nu_m$ , and  $\nu_c$ ), which generally necessitates well-sampled light curves and spectra in multiple bands. Thus, in order to determine ranges for  $E_{K,iso}$  and n, we fix the values of the microphysical parameters. We first consider the fiducial case that  $\epsilon_e = 0.1$  and  $\epsilon_B = 0.1$ .

For each burst, we determine the constraints on  $E_{\rm K,iso}$  and n by computing individual probability distributions for each observation. We then assign the probabilities to a grid of values, and use a joint probability analysis to calculate the distributions in each parameter. For the grid, the ranges of the density and isotropic-equivalent kinetic energy are  $n_i = 10^{-6} - 10^3$  cm<sup>-3</sup> and  $E_i = 10^{46} - 10^{54}$  erg, with 1000 logarithmically, uniformly-spaced steps in each parameter. We choose the lower bound of the density range,  $n_{\rm min} = 10^{-6}$  cm<sup>-3</sup> to match the typical density of the intergalactic medium (IGM).

#### 3.3.1. Detections

To calculate the individual probability distributions for afterglow detections, we apply Equation 4 to the observations using the relevant regime for each observing band,  $\nu_i$ . For the X-ray band, we use the location of  $\nu_c$  as determined in Section 3.1.1 to determine which branch of Equation 4 to use. Since the value of p is primarily determined from the X-ray band, the  $E_{K,iso}$  – n relation remains unchanged when using different X-ray observations that follow the same temporal decline; thus, we only use one X-ray observation per burst. For the optical band, we make the reasonable assumption that  $\nu_m < \nu_{\rm opt} < \nu_c$  and utilize the second branch of Equation 4. In some cases, individual optical/NIR observations lead to  $E_{\rm K.iso} - n$  relations that do not overlap within their  $1\sigma$  uncertainties. In these cases, we use the weighted mean and standard deviation of these relations (i.e., systematic uncertainty) as the optical/NIR solution. For the radio band, we assume that  $\nu_a < \nu_{\rm radio} < \nu_m$ , (first branch of Equation 4) to calculate the  $E_{K,iso}$  – n relations. Since the radio band lies on a different segment of the synchrotron spectrum than the other bands, it contributes a  $E_{K,iso}$  – n relation with a different slope, and thus enables tighter constraints on the physical parameters (Fig-

After calculating the unique probability distribution from each of the radio, optical/NIR and X-ray bands, we normalize the area under each of the distributions to unity. We assume that the uncertainties in the flux densities are Gaussian, and




Figure 3. Isotropic-equivalent kinetic energy versus circumburst density for four short GRBs with radio afterglow detections. In each panel, the X-rays (light blue), optical (orange) and radio (red) provide independent constraints on the parameter space. Measurements are shown as solid regions, where the width of the region corresponds to the  $1\sigma$  uncertainty. Upper (lower) limits set by the cooling frequency are denoted by blue dotted (dash-dotted) lines. The regions of parameter space ruled out by the observations are denoted (grey hatched regions). The median solution and  $1\sigma$  uncertainty is indicated by the black cross in each panel, corresponding to the values listed in Table 3. For each burst, the joint probability distributions in n (bottom panel) and  $E_{K,iso}$  (right panel) are shown. Red lines correspond to the median, and dotted lines are the  $1\sigma$  uncertainty about the median. The green line corresponds to  $E_{\gamma,iso}$ . The only optical observations available for GRB 050724A are during a flare; thus we do not include the optical data in our analysis. The addition of a detection in the radio band is crucial in constraining the best-fit solution, and in three cases constrains  $\epsilon_B$ .

thus each band contributes a unique, log-normal distribution. These distributions are shown for the four bursts with radio afterglow detections (Figure 3), and 34 bursts with radio afterglow non-detections: 16 bursts with  $\nu_c < \nu_X$  (Figure 4) and 18 bursts with  $\nu_c > \nu_X$  (Figure 5).

### 3.3.2. Upper Limits

For flux density upper limits, we use Equation 4 to determine the  $E_{\rm K,iso}-n$  relation at the  $3\sigma$  upper limit. We then assign zero probability to the  $E_{\rm K,iso}-n$  parameter space above the relationship, and assign a constant probability to the allowed parameter space below the relationship, normalized to unity. The upper limits are denoted by dashed lines in Figures 3–5, where the colors of the lines correspond to the relevant observing band (X-rays: blue, optical: orange, radio: red). Regions of parameter space that have been ruled out by the upper limits are marked as hatched regions.

### 3.3.3. Cooling Frequency Constraint

We utilize the relative location of the cooling frequency as a final constraint, since it depends on a combination of energy and density, by

$$\nu_c \propto n_0^{-1} E_{\text{K,iso},52}^{-1/2} \epsilon_{B,-1}^{-3/2},$$
 (5)

additional dependencies on  $\delta t$  and redshift (Granot & Sari 2002). For the cases in which the X-ray band is located above the cooling frequency ( $\nu_c < \nu_X$ ), we employ a maximum value at the lower edge of the X-ray band,  $\nu_{c,\text{max}} = 7.3 \times 10^{16} \, \text{Hz}$  (0.3 keV), to obtain a lower limit on the combination of energy and density. The corresponding probability distribution has zero value in the  $E_{K,iso}$  – n parameter space below the relation, and a constant value above the relation, where the area in the allowed parameter space is normalized to unity. The lower limits are shown as as blue dot-dashed lines for GRBs 051221A and 130603B in Figure 3 and for all bursts in Figure 4. In cases where the X-ray band is below the cooling frequency ( $\nu_m < \nu_X < \nu_c$ ), we set the cooling break to a minimum value,  $\nu_{c, \rm min} = 2.4 \times 10^{18} \ \rm Hz$  (10 keV) at the upper edge of the X-ray band, and determine the  $E_{K,iso}$  – n relation for each burst using Equation 5. This constraint sets an upper limit on the combination of energy




Figure 4. Isotropic-equivalent kinetic energy versus circumburst density for 16 short GRBs with solutions for  $\nu_c < \nu_X$  assuming fiducial values for the microphysical parameters of  $\epsilon_e = \epsilon_B = 0.1$ . In each panel, the X-rays (light blue), optical (orange) and radio (red) provide independent constraints on the parameter space. In particular, the X-ray band provides an estimate of  $E_{K,iso}$ . Measurements are shown as solid regions, where the width of the region corresponds to the  $1\sigma$  uncertainty, while upper limits are denoted as dashed lines. Setting the cooling frequency to a maximum value of  $\nu_{c,max} = 7.2 \times 10^{16}$  Hz (0.3 keV) provides an additional constraint (dark blue dot-dashed line). The regions of parameter space ruled out by the observations are denoted (grey hatched regions). The median solution and  $1\sigma$  uncertainty is indicated by the black cross in each panel, corresponding to the values listed in Table 3. For each burst, the joint probability distributions in n (bottom panel) and  $E_{K,iso}$  (right panel) are shown. Red lines correspond to the median, and dotted lines are the  $1\sigma$  uncertainty about the median. The green line corresponds to  $E_{\gamma,iso}$ .

and density. We form the probability distribution in the same manner as for afterglow upper limits. The limits for each burst set by the cooling frequency are shown as blue dashed lines for GRBs 050724A and 140903A in Figure 3 and all bursts in Figure 5.

### 3.4. Joint Probability Distributions

Since each of the observing bands, as well as the location of the cooling frequency, contribute an independent probability distribution, we calculate the joint probability from a product of these distributions for each burst. To obtain 1-dimensional probability distributions in  $E_{\rm K,iso}$  and n, we integrate over each of the parameters. Finally, we normalize the area under each 1-dimensional distributions, P(n) and  $P(E_{\rm K,iso})$ , for 34 bursts are shown in Figures 4–5 for the fiducial microphysical parameters,  $\epsilon_e = \epsilon_B = 0.1$ . The median values and  $1\sigma$  uncertainties in isotropic-equivalent kinetic energy and circumburst density are also shown in these figures and listed in Table 3.

Figure 3 shows the probability distributions for the four events with radio afterglow detections. In three of the four

cases, we can use the available data to place additional constraints on  $\epsilon_B$ , fixing  $\epsilon_e=0.1$ . For GRBs 050724A and 140903A, the afterglow data require that  $\epsilon_B\lesssim 10^{-4}$  and  $\lesssim 10^{-3}$ , respectively. For larger values of  $\epsilon_B$ , the constraint from the cooling frequency becomes more stringent and conflicts with the solutions obtained from the afterglow observations on the grid of allowed values. For GRB 051221A, we find a significantly better fit at  $\epsilon_B=0.01$ . In all cases, the addition of the radio band enables tighter constraints on both the energy and the density.

### 4. DENSITY AND ENERGY SCALE FOR SHORT GRBS

To quantify the distributions of circumburst densities and isotropic-equivalent kinetic energies for the entire sample, we calculate the combined probability distributions from the sum of the 1-dimensional probability distributions, P(n) and  $P(E_{K,iso})$ . We sum the individual distributions for all bursts with valid solutions at  $\epsilon_B = 0.1$ , as well as GRB 050724A ( $\epsilon_B = 10^{-4}$ ) and GRB 140903A ( $\epsilon_B = 10^{-3}$ ), to create cumulative probability distributions for both density and kinetic energy, shown in Figure 6.




Figure 5. Isotropic-equivalent kinetic energy versus circumburst density for 18 short GRBs with solutions for  $\nu_c > \nu_X$  assuming fiducial values for the microphysical parameters of  $\epsilon_e = \epsilon_B = 0.1$ . In each panel, the X-rays (light blue), optical (orange) and radio (red) provide independent constraints on the parameter space. Measurements are shown as solid regions, where the width of the region corresponds to the  $1\sigma$  uncertainty, while upper limits are denoted as dashed lines. Setting the cooling frequency to a *minimum* value of  $\nu_{c,min} = 2.4 \times 10^{18}$  Hz (10 keV) provides an additional constraint (dark blue dashed line). The regions of parameter space ruled out by the observations are denoted (grey hatched regions). The median solution and  $1\sigma$  uncertainty is indicated by the black cross in each panel, corresponding to the values listed in Table 3. For each burst, the joint probability distributions in n, with an imposed lower bound of  $n_{min} = 10^{-6}$  cm<sup>-3</sup> (bottom panel), and  $E_{K,iso}$  (right panel) are shown. Red lines correspond to the median, and dotted lines are the  $1\sigma$  uncertainty about the median. The green line corresponds to  $E_{\gamma,iso}$ .

The median values for the population of 38 bursts are  $\langle n \rangle \approx 2.9 \times 10^{-3} \ {\rm cm^{-3}}$  and  $\langle E_{\rm K,iso} \rangle \approx 1.9 \times 10^{51} \ {\rm erg}$  (Figure 6 and Table 4). The density which corresponds to 90% of the cumulative distribution  $(n_{90})$  is  $\approx 0.4 \ {\rm cm^{-3}}$ , and the fraction of probability that lies at densities of  $\lesssim 1 \ {\rm cm^{-3}}$  ( $f_{n<1 \ {\rm cm^{-3}}}$ ) is 0.95 (Figure 6 and Table 4). We also calculate these statistics for the sub-sample of bursts with  $\nu_c < \nu_X$ , corresponding to the events in Figure 4, that utilize all three branches of Equation 4 and therefore have relatively well-constrained energies and densities (compared to the broader probability distributions for events with  $\nu_c > \nu_X$ ; see Section 4.2.3). We find that this distribution has a median of  $\langle n \rangle \approx 3.7 \times 10^{-2} \ {\rm cm^{-3}}$ ,

 $n_{90} \approx 1.0~{\rm cm^{-3}}$ , and that  $f_{n<1~{\rm cm^{-3}}} \approx 0.90$ . Furthermore, the median value for kinetic energy decreases by a factor of 2, to  $E_{\rm K,iso} \approx 9.6 \times 10^{50}$  erg. The cumulative distributions in density and energy are shown in Figure 6 and the population statistics are listed in Table 4.

#### 4.1. Isotropic-equivalent $\gamma$ -ray energy and Efficiency

We compare the inferred isotropic-equivalent kinetic energy to the  $\gamma$ -ray energy by computing the isotropic-equivalent  $\gamma$ -ray energy,  $E_{\gamma,\rm iso}$ , to represent the energy range  $\approx 1-10^4$  keV (to match the widest energy ranges for current GRB detection satellites),




Figure 6. Combined and cumulative probability distributions in n and  $E_{K,iso}$  assuming  $\epsilon_e = \epsilon_B = 0.1$  (top) and  $\epsilon_e = 0.1$ ,  $\epsilon_B = 0.01$  (bottom). All scenarios also include GRBs 050724A and 140903A, with  $\epsilon_B = 10^{-4}$  and  $10^{-3}$ , respectively. Each panel shows three populations: all bursts with an imposed lower bound of  $n_{\min} = 10^{-6}$  cm<sup>-3</sup> (black dot-dashed), and the sub-sample of events with  $\nu_c < \nu_X$  (light grey dashed). Color-coded arrows from the bottom denote the median for each distribution (and in some cases, staggered for the sake of clarity), and lines denote 90% upper limits. For  $\epsilon_B = 0.1$  ( $\epsilon_B = 0.01$ ), the cumulative distributions indicate that for  $n \gtrsim 3 \times 10^{-3}$  cm<sup>-3</sup> ( $n \gtrsim 0.3$  cm<sup>-3</sup>), the distributions are virtually independent of the choice of  $n_{\min}$  provided that  $n_{\min} \gtrsim 10^{-4}$  cm<sup>-3</sup>. This allows us to place robust 90% upper limits of  $n \lesssim 0.5 - 1.0$  cm<sup>-3</sup> ( $n \lesssim 3 - 23$  cm<sup>-3</sup>) for our sample of bursts. In addition,  $\approx 95\%$  ( $\approx 80\%$ ) of the total probability for all events lies below densities of 1 cm<sup>-3</sup>. The range of median isotropic-equivalent kinetic energies for all scenarios is  $\approx (2-20) \times 10^{51}$  erg (c.f., Table 4).

$$E_{\gamma,\text{iso}} = k_{\text{bol}} \times \frac{4\pi d_L^2}{1+z} f_{\gamma} \text{ erg}$$
 (6)

where  $k_{\rm bol}$  is the bolometric correction factor to convert the fluence to an energy range of  $\approx 1-10^4$  keV,  $d_L$  is the luminosity distance in cm, and  $f_{\gamma}$  is the fluence in units of erg cm<sup>-2</sup>. For cases in which the fluence is calculated over the 15-150 keV *Swift* energy range, we use  $k_{\rm bol}=5$ . If a burst is detected by other  $\gamma$ -ray satellites which cover a wider energy range of  $\approx 10-1000$  keV (e.g., *Fermi, Konus-Wind, Suzaku*), we utilize the measured fluences from these satellites and  $k_{\rm bol}=1$  to calculate the  $\gamma$ -ray energy. Individual values of  $E_{\gamma,\rm iso}$  are shown in Figures 3–5 and listed in Table 3, while the distribution for 38 bursts is shown in Figure 7). We find that the range in isotropic-equivalent  $\gamma$ -ray energy is  $\approx (0.04-45)\times 10^{51}$  erg, with a median value of  $\langle E_{\gamma,\rm iso}\rangle \approx 1.8\times 10^{51}$  erg, similar to the ranges and median

values of the isotropic-equivalent kinetic energy (Figure 7) We also calculate the  $\gamma$ -ray efficiency,

$$\eta_{\gamma} = \frac{E_{\gamma, \text{iso}}}{E_{\gamma, \text{iso}} + E_{K, \text{iso}}},\tag{7}$$

as well as the  $1\sigma$  uncertainties in  $\eta_{\gamma}$  for each burst, following standard propagation of errors from the  $1\sigma$  uncertainties in  $E_{\rm K,iso}$ . The resulting values of  $\eta_{\gamma}$  are listed in Table 3 and the cumulative distributions, after incorporating the  $1\sigma$  uncertainties, are shown in Figure 7. We find a wide range in  $\eta_{\gamma}$ ,  $\approx 10^{-3}-1$ , and note that the lower bound is set by the single outlier, GRB 150101B (Table 3). Excluding this burst, the lower bound is  $\approx 0.03$  (GRB 140622A).

The distribution for  $\epsilon_B = 0.1$  has a median of  $\langle \eta_{\gamma} \rangle = 0.56^{+0.36}_{-0.37}$  (1 $\sigma$  uncertainties). The isotropic-equivalent kinetic energy scale for the  $\epsilon_B = 0.01$  case is comparatively high (c.f., Figure 6), and thus the median value for the  $\gamma$ -ray efficiency is




Figure 7. Left: Histograms of isotropic-equivalent  $\gamma$ -ray energy (red), and inferred isotropic-equivalent kinetic energies assuming  $\epsilon_B = 0.1$  (dark grey, thick bars) and  $\epsilon_B = 0.01$  (light grey, thin bars) for 38 short GRBs from broad-band afterglow modeling. Median values for each population are denoted by the color-coded arrows from the top. Values for the isotropic-equivalent energies correspond to those listed in Table 3. Right: Cumulative distributions of  $\gamma$ -ray efficiency ( $\eta_{\gamma}$ ) assuming  $\epsilon_B = 0.1$  (dark grey) and  $\epsilon_B = 0.01$  (light grey), taking into account the  $1\sigma$  uncertainties in  $\eta_{\gamma}$  for each burst. Median values are listed in the figure and denoted by the color-coded arrows from the bottom. Arrows are staggered for clarity.

relatively low,  $\langle \eta_{\gamma} \rangle = 0.40^{+0.49}_{-0.35}$  (1 $\sigma$  uncertainties; Figure 7).

#### 4.2. Alternative Cases

We have thus far made assumptions about the values of the microphysical parameters ( $\epsilon_e = \epsilon_B = 0.1$ ), the redshifts (z = 0.5 unless otherwise determined), the minimum allowed density ( $n_{\rm min} = 10^{-6}~{\rm cm}^{-3}$ ), and the cooling frequency (0.3 keV for  $\nu_c < \nu_X$ ; 10 keV for  $\nu_c > \nu_X$ ). To explore the impact of these assumptions on our resulting distributions for the kinetic energy and circumburst density, we consider alternative values for these parameters.

#### 4.2.1. The Value of $\epsilon_B$

In some cases, a valid solution using the fiducial input of  $\epsilon_B=0.1$  (with fixed  $\epsilon_e=0.1$ ) cannot be found. For instance, three of the four bursts with radio afterglows require that  $\epsilon_B\lesssim 0.1$  (Figure 3 and Table 3). At a fixed value of  $\epsilon_e=0.1$ ,  $\epsilon_B$  is constrained to  $\lesssim 10^{-4}$  for GRB 050724A,  $\lesssim 10^{-2}$  for GRB 051221A, and  $\lesssim 5\times 10^{-3}$  for GRB 140903A (Figure 3). For the remaining bursts, we do not have enough information to rule out the  $\epsilon_B=0.1$  scenario. Thus, we consider two additional cases for all bursts:  $\epsilon_B=0.01$  and  $\epsilon_B=10^{-4}$  (with fixed  $\epsilon_e=0.1$ ). Of the 38 bursts in our sample, 33 have valid solutions for  $\epsilon_B=0.01$ ; the median and uncertainties in circumburst density and kinetic energy for each burst are listed in Table 3. To create cumulative probability distributions, we repeat the same exercise as described in Section 4 for the 33 events with valid solutions, displayed in Figure 6.

Using  $\epsilon_B = 10^{-4}$  as the fiducial value, the constraint from the cooling frequency conflicts with the solutions allowed by the afterglow observations in 16 cases, indicating that data do not allow such a low value of  $\epsilon_B$  for these bursts. The population statistics for the 22 bursts with valid solutions are listed in Table 4.

#### 4.2.2. Redshift

For bursts with no determined spectroscopic redshift, we have assumed z = 0.5, set by the median of the short GRB

population with known redshifts (Berger 2014). However, in three cases, GRBs 060313, 111020A, and 121226A, we do not find a valid joint solution at z = 0.5. We find that there are valid solutions at redshifts of  $z \gtrsim 1$ , and thus assume z = 1 for these bursts.

#### 4.2.3. The Value of $n_{\min}$

In the 20 cases in which the cooling frequency lies above the X-ray band ( $\nu_m < \nu_X < \nu_c$ ), the X-ray and optical/NIR bands occupy the same spectral regime (branch 2 of Equation 4) and the resulting  $E_{K,iso} - n$  relations have the same slope (Figure 3 and 5). Thus, the lower bound on the density is set by our minimum grid value of  $n_{\min} = 10^{-6} \text{ cm}^{-3}$ ; the density is otherwise unconstrained at the low end and results in broad probability distributions in both circumburst density and energy. To understand the impact of our choice of  $n_{\min}$  on the resulting distributions, we repeat the individual probability analysis, employing a more stringent lower bound of  $n_{\text{min}} = 10^{-4} \text{ cm}^{-3}$ , at the low end of gas densities for the interstellar medium (ISM; Korpi et al. 1999; Murali 2000; Gent et al. 2013). Since kinetic energy and density are inversely related, the upper bound on  $E_{K,iso}$  for each burst is naturally set by our choice of  $n_{\min}$ . We consider this alternative value of  $n_{\min}$  for  $\epsilon_B = 0.1$  and  $\epsilon_B = 0.01$ .

#### 4.2.4. The Value of $\nu_c$

In all cases, we have assumed that the cooling frequency is on the edge of the X-ray band (0.3 keV for  $\nu_c < \nu_X$ ; 10 keV for  $\nu_c > \nu_X$ ). To test whether this assumption has any impact on our results, we repeat the individual and joint probability analysis assuming that the cooling frequency is at the logarithmic mean of the 0.3–10 keV *Swift* X-ray band,  $\nu_{c,\text{mid}} = 4.1 \times 10^{17}$  Hz (1.7 keV). We consider this alternative value of the cooling frequency for  $\epsilon_B = 0.1$  and  $\epsilon_B = 0.01$ .

#### 4.2.5. Trends

Taking these alternative values into account, we repeat the individual and joint probability analysis for each burst for nine

Table 4
Circumburst Density and Kinetic Energy Population Statistics

Scenario	No. of Events	$\langle n \rangle$ (cm <sup>-3</sup> )	$n_{90}^a$ (cm <sup>-3</sup> )	$f_{n<1\mathrm{cm}^{-3}}^{b}$	$\langle E_{ m K,iso}  angle$ (erg)
	$\epsilon_B = 0.1$				
All bursts, $n_{\min} = 10^{-6} \text{ cm}^{-3}$	38	$2.9 \times 10^{-3}$	0.40	0.95	$1.9 \times 10^{51}$
All bursts, $n_{\min} = 10^{-4} \text{ cm}^{-3}$	37	$3.7 \times 10^{-3}$	0.49	0.95	$1.2 \times 10^{51}$
All bursts, $n_{\min} = 10^{-6} \text{ cm}^{-3}$ , $\nu_c = 1.7 \text{ keV}$	38	$2.2 \times 10^{-3}$	0.20	0.95	$1.6 \times 10^{51}$
Bursts with $\nu_c < \nu_X$	19	$3.7 \times 10^{-2}$	1.01	0.90	$9.6 \times 10^{50}$
	$\epsilon_B = 0.01$				
All bursts, $n_{\text{min}} = 10^{-6} \text{ cm}^{-3}$	33	$5.2 \times 10^{-3}$	2.7	0.79	$2.9 \times 10^{51}$
All bursts, $n_{\min} = 10^{-4} \text{ cm}^{-3}$	32	$1.4 \times 10^{-2}$	3.1	0.78	$1.7 \times 10^{51}$
All bursts, $n_{\min} = 10^{-6} \text{ cm}^{-3}$ , $\nu_c = 1.7 \text{ keV}$	34	$1.6 \times 10^{-2}$	2.1	0.83	$2.4 \times 10^{51}$
Bursts with $\nu_c < \nu_X$	14	0.96	23.3	0.51	$6.4 \times 10^{50}$
	$\epsilon_B = 1 \times 10^{-4}$				
All bursts, $n_{\text{min}} = 10^{-6} \text{ cm}^{-3}$	22	$3.0 \times 10^{-2}$	771	0.68	$1.8 \times 10^{52}$

**Note.** — All scenarios include GRBs 050724A and 140903A, with  $\epsilon_B = 10^{-4}$  and  $10^{-3}$ , respectively.

different sets of input parameters in total. The population medians, as well as values for  $n_{90}$  and  $f_{n<1\,\mathrm{cm}^{-3}}$  are listed in Table 4. In addition, cumulative distributions for kinetic energy and circumburst density for varying values of  $\epsilon_B$  and  $n_{\min}$  are shown in Figure 6.

Overall, we find that a change in  $\epsilon_B$  results in an increase in the circumburst density. For instance, assuming  $n_{\min}$  =  $10^{-6}$  cm<sup>-3</sup> for all bursts, decreasing  $\epsilon_B$  by a factor of 10 to  $\epsilon_B = 0.01$  results in an increase in the median density by a factor of  $\approx 1.8$ . This trend becomes more drastic for other scenarios: when assuming  $n_{\text{min}} = 10^{-4} \text{ cm}^{-3}$ , the median density in the scenarios when assuming  $n_{\text{min}} = 10^{-4} \text{ cm}^{-3}$ , the median density in the scenarios. sity increases by a factor of  $\approx 3.8$ , and when considering only the bursts with  $\nu_c < \nu_X$ , the factor of increase is  $\approx 26$ . When comparing the values of  $n_{90}$  and  $f_{n<1 \text{ cm}^{-3}}$ , we also find overall trends commensurate with an increase in circumburst density. In particular, the factor of ten decrease in  $\epsilon_B$  causes  $f_{n<1\,\mathrm{cm}^{-3}}$ to decrease by a larger factor, from  $\approx 0.95$  to  $\approx 0.8$  for all bursts. When considering only bursts with  $\nu_c < \nu_X$ ,  $f_{n<1\,\mathrm{cm}^{-3}}$ decreases from  $\approx 0.90$  to  $\approx 0.51$ . The effect of  $\epsilon_B$  on the median kinetic energy is less pronounced, with increases by factors of  $\approx 1.5$  in all cases except when considering only bursts with  $\nu_c < \nu_X$ ; in this case, the median undergoes a slight decrease by a factor of  $\approx 1.5$  (Table 4). Overall, we find that a decrease from  $\epsilon_B = 0.1$  to 0.01 results in higher median densities by factors of  $\approx 2-25$  depending on the considered scenario, and a uniform increase in the density cumulative distributions (Figure 6). When considering the more extreme input of  $\epsilon_B = 10^{-4}$  for all bursts, the median density and kinetic energy both increase by factors of  $\approx 10$ , compared to the  $\epsilon_B = 0.1$ case (Table 4).

We next investigate the effects of the values of  $n_{\rm min}$  and  $\nu_c$  on the parameter distributions. We find that at constant  $\epsilon_B$ , the upper  $\approx 30-50\%$  of the density cumulative distributions are virtually independent of our choice of  $n_{\rm min}$ , provided that

 $n_{\rm min} \lesssim 10^{-4}~{\rm cm}^{-3}$  (Figure 6). Importantly, this also allows us to place robust 90% upper limits on the density that are largely unaffected by our choice of  $n_{\rm min}$ :  $n_{90} \approx 0.4-0.5~{\rm cm}^{-3}$  for  $\epsilon_B=0.1$  and  $n_{90}\approx 2.7-3.1~{\rm cm}^{-3}$  for  $\epsilon_B=0.01$  (Table 4). We find that the choice of cooling frequency within the X-ray band has a minor effect on the circumburst density, either a  $\approx 1.3$  factor of decrease ( $\epsilon_B=0.1$ ) or a factor of  $\approx 3$  increase ( $\epsilon_B=0.01$ ), while the median kinetic energy only decreases by a factor of  $\approx 1.2$  in both cases (Table 4).

When considering all cases with less extreme values of  $\epsilon_B$  for all bursts, the median density range is  $(3-15)\times 10^{-3}$  cm<sup>-3</sup>, with 90% upper limits of  $n_{90}\approx 0.4-3$  cm<sup>-3</sup>. Furthermore,  $\approx 80-95\%$  of the probability is below  $\approx 1$  cm<sup>-3</sup> regardless of the input parameters considered (Table 4). The median isotropic-equivalent kinetic energy ranges from  $\approx (1.2-2.9)\times 10^{51}$  erg. Including more extreme scenarios like the subset of bursts with  $\nu_c < \nu_X$  and  $\epsilon_B = 10^{-4}$ , the full median density range is  $\approx (3-1000)\times 10^{-3}$  cm<sup>-3</sup> and the median isotropic-equivalent kinetic energy range is  $\approx (0.6-20)\times 10^{52}$  erg (Table 4).

### 5. DISCUSSION AND IMPLICATIONS

#### 5.1. Afterglow Properties

The X-ray, optical, and radio afterglow catalogs (Tables A1-A3) allow us to analyze the observational afterglow properties of short GRBs as a population. By fitting the afterglow light curves, we find a weighted mean prejet break decline rate of  $\langle \alpha_X \rangle \approx -1.07$  at  $\delta t \gtrsim 1000$  s for bursts with measured temporal indices, similar to the prejet break declines measured from long GRB light curves (Nysewander et al. 2009; Racusin et al. 2009; Kann et al. 2010; Zaninoni et al. 2013), and slightly shallower than the value of  $\alpha_X \approx -1.2$  found for 11 short GRBs in an earlier study (Nysewander et al. 2009). We measure the optical decline rates and find the same weighted mean decline rate of  $\langle \alpha_{\rm opt} \rangle \approx -1.07$  from 19 well-sampled bursts with measured indices.

From spectral fitting of the optical, near-IR, and X-ray data, we find 12 short GRBs which require rest-frame extinction, with measured values of  $A_V^{\text{host}} \approx 0.3 - 1.5$  mag, and two

<sup>&</sup>lt;sup>a</sup> This is the circumburst density which corresponds to 90% of the cumulative distribution.

<sup>&</sup>lt;sup>b</sup> This is the fraction of the circumburst density cumulative distribution below a value of 1 cm<sup>-3</sup>.

bursts with lower limits of  $A_V^{\text{host}} \gtrsim 2.5 - 6$  mag (Table 3). We note that GRB 080919 has the highest value of rest-frame extinction; however, the sightline to this burst is close to the Galactic plane and therefore has a highly uncertain Galactic extinction which likely affects the measurement of  $A_V^{\text{host}}$ . Prior to this study, evidence for  $\gtrsim 0.5$  mag of extinction has only been reported in three short GRBs: GRB 070724A (Berger et al. 2009; Kocevski et al. 2010), GRB 111020A (Fong et al. 2012b), and GRB 120804A (Berger et al. 2013). Afterglow modeling in this work results in the same conclusions for those three events, and includes nine additional events with rest-frame extinction. We note that most of the events with non-zero extinction and robust host associations are in star-forming host galaxies; the single exception is GRB 150101B which has an inferred value of  $A_V^{\text{host}} \approx$ 0.5 mag and is located on the outskirts of an early-type galaxy (Fong et al., in prep). In comparison,  $\approx 15-20\%$  of Swift long GRBs have optically sub-luminous afterglows that have been attributed to dust extinction. For long GRBs, inferred values of  $A_V^{
m host} pprox 0.5$  mag are common, with a substantial fraction of events with  $A_V^{\text{host}} \gtrsim 1 - 2.5$  mag (Cenko et al. 2009; Perley et al. 2009a, 2013).

While rest-frame extinction can be explained by dust in star-forming regions in the local environments of long GRBs, substantial extinction in short GRBs cannot be easily explained in the context of the double compact object merger progenitor. It is possible that these events are "impostors" which in actuality have massive star progenitors; in that case, we might expect them to be distinct in their  $\gamma$ -ray properties with longer durations and softer  $\gamma$ -ray spectra. However, there does not appear to be a correlation between short GRBs with extinction and their durations as they span the full range, with  $T_{90} \approx 0.02 - 2$  sec, and only three events with non-zero extinction have  $T_{90} \gtrsim 1$  sec (GRBs 060121, 070714B, and 121226A). Furthermore, a study conducted by Bromberg et al. (2013) assigned six of these events probabilities of having non-collapsar progenitors based on their  $\gamma$ -ray properties. According to this study, 4/6 events have  $\gtrsim 60\%$ probabilities that they do not originate from collapsars. Thus, it is unlikely that the majority of these events are in fact "impostors". Instead, this suggests that some short GRBs may originate in a star-forming regions, or have progenitor systems that can produce dust.

#### 5.2. Opening Angles

Most well-sampled short GRBs exhibit a single afterglow decline rate in the X-ray (at  $\delta t \gtrsim 1000~{\rm sec}$ ) and optical bands. However, there are four short GRBs (GRBs 051221A, 090426A, 111020A, and 130603B) which have temporal steepenings on timescales of  $\delta t \approx 2-5~{\rm days}$ , attributed to jet breaks (Table 5; Soderberg et al. 2006; Nicuesa Guelbenzu et al. 2011; Fong et al. 2012b, 2014). Jet breaks are achromatic features, and in principle can be detected in the X-ray through radio bands. The measurement of jet break time, in conjunction with the energy, density, and redshift, can be converted to a jet opening angle,  $\theta_j$ , using (Sari et al. 1999; Frail et al. 2001)

$$\theta_j = 9.5 t_{j,d}^{3/8} (1+z)^{-3/8} E_{K,iso,52}^{-1/8} n_0^{1/8} \text{ deg},$$
 (8)

where  $t_{j,d}$  is in days. The opening angle measurements for the four short GRBs with jet break detections are listed in Table 5. Using these four short GRB opening angle measure-

ments alone, taking into account the published range of angles for individual bursts, the median is  $\langle \theta_j \rangle = 6 \pm 1$  deg (Figure 8).

**Table 5**Short GRB Opening Angles

GRB	Band <sup>a</sup>	$\theta_j$ (deg)	$\delta t_{ m last}^b$ (days)	Reference
050709	О	≥ 15°	16.2	1
050724A	X	$\gtrsim 15^{\circ}$ $\gtrsim 25^{\circ}$	22.0	2
051221A	X	6-7°	26.6	3
090426A	O	$5-7^{\circ}$	2.7	4
101219A	X	$\gtrsim 4^{\circ}$ $3-8^{\circ}$	3.9	5, This work
111020A	X	$3-8^{\circ}$	10.2	6
111117A	X	$\gtrsim 3-10^{\circ}$	3.0	7, 8
120804A	X	$\gtrsim 13^{\circ}$	45.9	9, This work
130603B	OR	$4 - 8^{\circ}$	6.5	10
140903A	X	$\gtrsim 6^{\circ}$ $\gtrsim 9^{\circ}$	3.0	11, This work
140930B	X	$\gtrsim 9^{\circ}$	23.1	This work

**Note.** — Bursts with opening angle measurements are in bold.

**References:** (1) Fox et al. 2005; (2) Grupe et al. 2006; (3) Soderberg et al. 2006; (4) Nicuesa Guelbenzu et al. 2011; (5) Fong et al. 2013; (6) Fong et al. 2012b; (7) Margutti et al. 2012; (8) Sakamoto et al. 2013; (9) Berger et al. 2013; (10) Fong et al. 2014; (11) Evans et al. 2009


However, the majority of short GRBs do not have detected jet breaks and instead exhibit a single power-law decline as long as they are detected. In these cases, the time of the last observation ( $\delta t_{\rm last}$ ) can be used to place lower limits on the opening angles. The inclusion of these bursts is essential in understanding the true opening angle distribution. In most cases, Swift/XRT observes short GRBs until they fade below the detection threshhold at  $\delta t \lesssim 1$  day, and enables relatively shallow lower limits of  $\theta_j \gtrsim 2-5^\circ$  (Coward et al. 2012; Fong et al. 2012b). The inclusion of such limits will not have a significant effect on the opening angle distribution, as they virtually span the entire range of allowable angles.

Therefore, in order to have a more complete understanding of the opening angle distribution, we collect all existing published lower limits of  $\theta_j \gtrsim 5^\circ$ , and calculate lower limits for GRBs 101219A, 120804A, 140903A, and 140903B using the observations and physical parameters presented in this work. The inferred lower limits, the band in which the jet break was detected, and the value of  $\delta t_{\rm last}$  used to compute the limits, are listed in Table 5. These seven events demonstrate that multiwavelength afterglow observations to  $\delta t_{\rm last} \approx 3-25$  days enable more meaningful lower limits on the opening angles of  $\gtrsim 5-25^\circ$  (Table 5).

To calculate the opening angle distribution, we give each of the 11 events equal weighting, where measurements are given Gaussian probability distributions to represent their allowed range of angles (Table 5), while lower limits are given probability that is evenly distributed between the lower limit and the maximum possible opening angle,  $\theta_{j,max} = 90^{\circ}$ . The resulting cumulative distribution for 11 short GRBs, including measurements and lower limits, is shown in Figure 8. Assuming an upper bound of  $\theta_{j,max} = 90^{\circ}$ , the short GRB

<sup>&</sup>lt;sup>a</sup> This indicates the band in which the jet break was detected or the lower limit was placed. X=X-ray, O=optical, R=radio. A range of angles is due to uncertainty in the redshift, kinetic energy or circumburst density.

<sup>&</sup>lt;sup>b</sup> This is the time elapsed between burst detection and the last observation.



**Figure 8.** Cumulative histograms of 11 short GRBs (dark red dash-dotted) and 265 long GRBs (dark blue) with opening angle measurements or limits, assuming a maximum opening angle of  $\theta_{\rm j,max} = 90^{\circ}$ . Also shown is the distribution for 11 short GRBs assuming a more realistic maximum opening angle of  $\theta_{\rm j,max} = 30^{\circ}$  (red). The distributions for the subset of four short GRBs (light red dotted) and 248 long GRBs (light blue) with opening angle measurements are also shown. Medians are denoted by color-coded arrows, and listed with their  $1\sigma$  uncertainties in units of degrees.

population median is  $\langle \theta_j \rangle = 33^{+38}_{-27}$  deg  $(1\sigma)$ . Motivated by simulations of post-merger black hole accretion predict jets with  $\theta_j \sim 5-30^\circ$  (Ruffert & Janka 1999b; Aloy et al. 2005; Rosswog 2005; Rezzolla et al. 2011), we also calculate the cumulative distribution employing a more realistic maximum value of  $\theta_{\rm j,max} = 30^\circ$ , and find a median of  $\langle \theta_j \rangle = 16 \pm 10$  deg  $(1\sigma)$ .

To compare these distributions to those for long GRBs, we collect opening angle measurements for 265 long GRBs, including 17 events with limits (Frail et al. 2001; Berger et al. 2003; Bloom et al. 2003; Ghirlanda et al. 2004; Friedman & Bloom 2005; Racusin et al. 2009; Cenko et al. 2010, 2011; Filgas et al. 2011; Goldstein et al. 2011; Ryan et al. 2015) and calculate the cumulative distributions in the same manner (Figure 8). We find a median value for the 248 long GRBs with measurements of  $\langle \theta_j \rangle = 13^{+5}_{-9}$  deg. Including the 17 events with limits  $(\theta_{j,max} = 90^\circ)$ , the median becomes  $14^{+10}_{-10}$  deg.

The opening angle distribution of short GRBs impacts the true energy scale, as the true energy is lower than the isotropic-equivalent value by the beaming factor,  $f_b$ , where  $f_b \equiv 1 - \cos(\theta_j)$  and therefore  $E_{\rm true} = f_b E_{\rm iso}$ . To calculate the cumulative beaming factor distribution, we use the individual opening angle probability distributions for each burst to convert to individual distributions in beaming factor. We then sum the individual distributions in a cumulative sense and calculate the median and  $1\sigma$  uncertainties about the median. Including all short GRBs with opening angle measurements and limits and assuming the more realistic scenario of  $\theta_{\rm j,max} = 30^\circ$ , the median beaming factor is  $f_b = 0.04^{+0.07}_{-0.03}$ . The beaming correction is less substantial if we assume  $\theta_{\rm j,max} = 90^\circ$ ,  $f_b = 0.17^{+0.52}_{-0.16}$ , and is much more substantial if we only include short GRBs with measurements,  $f_b = 0.005 \pm 0.002$ .

We find median isotropic-equivalent  $\gamma$ -ray and kinetic energy scales of  $E_{\gamma,\rm iso}\approx 2\times 10^{51}$  erg and  $E_{\rm K,iso}\approx (1-3)\times 10^{51}$  erg. Applying the beaming correction for the most re-

alistic scenario gives median beaming-corrected  $\gamma$ -ray and kinetic energy scales of  $\langle E_{\gamma} \rangle = 0.8^{+1.4}_{-0.6} \times 10^{50}$  erg and  $\langle E_K \rangle = 0.8^{+2.5}_{-0.7} \times 10^{50}$  erg, resulting in a total beaming-corrected energy release of  $\langle E_{\rm tot} \rangle = 1.6^{+3.9}_{-1.3} \times 10^{50}$  erg. The inferred energy scales can be used to constrain the mechanism of energy extraction to power the relativistic jet: the thermal energy release from  $\nu\bar{\nu}$  annihilation in a baryonic outflow (Jaroszynski 1993; Mochkovitch et al. 1993) and magnetohydrodynamic (MHD) processes in the black hole's accretion remnant (e.g. Blandford & Znajek 1977; Rosswog et al. 2003). The general consensus is that  $\nu\bar{\nu}$  annihilation can only produce beamingcorrected total energy releases of  $10^{48} - 10^{49}$  erg, while MHD processes can more easily produce energy releases in excess of 10<sup>49</sup> erg (Ruffert & Janka 1999b,a; Popham et al. 1999; Rosswog 2005; Birkl et al. 2007; Lee & Ramirez-Ruiz 2007). Thus, if the majority of short GRBs have wider opening angles than the four short GRBs with measurements, and thus have a smaller overall correction to the isotropic-equivalent energy scale, it will be necessary to invoke MHD processes to explain the observed energy releases.

The opening angles also impact the event rate, as the true event rate is elevated compared to the observed rate by a factor of  $f_b^{-1}$ , so  $\Re_{\text{true}} = f_b^{-1} \Re_{\text{obs}}$ . The current estimated observed short GRB volumetric rate is  $\Re_{\text{obs}} \approx 10$  Gpc<sup>-3</sup> yr<sup>-1</sup> (Nakar et al. 2006). Using  $f_b^{-1} = 27_{-18}^{+158}$ , which corresponds to all short GRBs with opening angle measurements and limits  $(\theta_{\rm j,max}=30^\circ)$ , we find a true event rate of  $\Re_{\rm true}\approx 270^{+1580}_{-180}~{\rm Gpc}^{-3}~{\rm yr}^{-1}$ . The observed all-sky event rate of  $\approx 0.3~{\rm yr}^{-1}$  within 200 Mpc (Guetta & Piran 2005) then becomes  $8^{+47}_{-5}~{\rm yr}^{-1}$ . We note that this rate is conservative compared to previously reported rates based on short GRB observations (Coward et al. 2012; Fong et al. 2012b, 2014), as it properly incorporates opening angle lower limits, with the only assumption of an upper bound on the opening angle of 30°. This range is fully consistent with the expected detection rates of neutron star mergers within a volume of 200 Mpc by Advanced LIGO/VIRGO of  $\approx 0.2 - 200 \text{ yr}^{-1}$ (LIGO Scientific Collaboration et al. 2013). Since there are a limited number of short GRBs with meaningful information on the opening angles, any additional measurements will greatly help to elucidate the true opening angle distribution, and therefore true energy scale and event rate.

#### 5.3. Connection to Galactic Environments

We connect the afterglow properties of short GRBs to their larger-scale, galactic environments. In particular, we investigate trends between the circumburst densities, which probe the explosion environment on sub-parsec-scales, to the predicted distributions for NS-NS mergers, host galaxy type, and locations within the host galaxies.

We find a wide range of inferred circumburst densities for the 38 bursts that we have studied in detail, and can compare the properties of the sub-parsec-scale environment with the global host galaxy environment. Separating the bursts by host galaxy type according to Fong et al. (2013) and additional data collected since, we find that bursts in both star-forming and elliptical host galaxies span a wide range of densities,  $10^{-6}$ -5 cm<sup>-3</sup> (Figure 9). We compare the inferred densities to predictions for NS-NS mergers from population synthesis for varying Galactic potentials, which have input distributions for merger timescales and kick velocities (Belczynski et al. 2006). Considering only the four




Figure 9. Top: Projected physical offset,  $\delta R$ , versus circumburst density for 22 short GRBs with inferred circumburst densities and sub-arcsecond localization allowing for precise offset measurements. The sample is separated by host galaxy type: star-forming hosts (blue asterisks) and elliptical hosts with no signs of star formation (red squares) according to Fong et al. (2013). For bursts with no spectroscopic redshift, we assume z=0.5 to calculate the physical offset, except in two cases, GRBs 060313 and 111020A, where we have assumed z=1 as suggested by the afterglow modeling. Middle: Host-normalized offset,  $\delta R/r_e$ , versus circumburst density for 16 bursts with measured host galaxy sizes. Physical and host-normalized offsets are from Fong et al. (2010), Margutti et al. (2012), Berger et al. (2013), Fong & Berger (2013), Sakamoto et al. (2013), and this work. Error bars correspond to  $1\sigma$  confidence. Bottom: Histogram of inferred median circumburst densities for 22 short GRBs with physical offsets.

bursts with elliptical hosts, we find particularly good agreement with the distributions for the large elliptical galaxy model ( $M_* = 10^{11} M_{\odot}$ ,  $M_{\rm halo} = 10^{12} M_{\odot}$ ) which has probability peaks at  $10^{-5}$  cm<sup>-3</sup> and 1 cm<sup>-3</sup>, and poor agreement with the small elliptical galaxy model ( $M_* = 10^8 M_{\odot}$ ), which is dominated by very low densities of  $\lesssim 10^{-6} \text{ cm}^{-3}$  (Belczynski et al. 2006). This conclusion is commensurate with the known stellar masses of short GRB elliptical hosts, which have a median of  $M_* \approx 10^{11.0}~M_{\odot}$  (Leibler & Berger 2010; Berger 2014). There are no predicted density distributions for star-forming galaxies; however, we would expect short GRBs which originate in elliptical galaxies to have lower inferred densities due to the lower average ISM densities in elliptical galaxies (e.g., Fukazawa et al. 2006). Based on the small number of events, we do not find any significant difference between the inferred circumburst densities of short GRBs in star-forming versus elliptical hosts.

We next investigate the relationship between the inferred circumburst densities and burst offsets from their host galaxies. If short GRBs trace the large-scale distribution of the ISM, we expect the inferred circumburst densities to decrease as a function of offset. In this vein, we gather all available projected physical offsets,  $\delta R$ , between the afterglow location and host galaxy center, derived from ground-based (Margutti et al.

2012; Berger et al. 2013; Sakamoto et al. 2013) and Hubble Space Telescope (HST) observations (Fong et al. 2010; Fong & Berger 2013). For bursts with no spectroscopic redshift, we calculate the physical offsets at z = 0.5 to be consistent with this work. However, for GRBs 060313 and 111020A, afterglow modeling implies that z > 0.5 (c.f. Section 3.3; Fong et al. 2012b; Roming et al. 2006) so we keep the original fiducial value of z = 1 for these bursts for complete uniformity. The distribution of circumburst densities with respect to projected physical offset for 22 bursts is shown in Figure 9. We find that four bursts with  $\delta R \gtrsim 10$  kpc have very low densities of  $\lesssim 10^{-3}$  cm<sup>-3</sup>, while bursts with  $\delta R \lesssim 10$  kpc have a wider range of densities, spanning  $\approx$  6 orders of magnitude (Figure 9). Overall, we find that for  $\delta R \approx 1-15$  kpc, there is no obvious trend between circumburst density and projected physical offset. We also find no obvious trends when considering only bursts with relatively well-measured densities

To analyze the relationship with offsets in a more uniform manner, we utilize offsets that have been normalized by the sizes of their host galaxies ( $\delta R/r_e$  where  $r_e$  is the galaxy halflight radius). The sample of bursts with host-normalized offsets is smaller since precise galaxy size measurements require the resolution of *HST*; thus the sample comprises 16 events (Fong et al. 2010; Fong & Berger 2013). The circumburst densities as a function of projected host-normalized offset is provided in Figure 9. Our analysis suggests that for  $\lesssim 5 r_e$ , the inferred densities are largely independent of host-normalized offset. We discuss a couple of possible contributing factors. First, since we can only measure projected offsets, we are not sensitive to the distance component along our line of sight, which could contribute a significant amount to the absolute distance. This may explain the case of GRB 061006, which has a small projected offset  $\approx 0.4r_e$  but has a very low density of  $\approx 2 \times 10^{-5}$  cm<sup>-3</sup> (Table 3 and Figure 9). Second, the afterglow only probes the sub-parsec circumburst environment and to a certain extent will be more sensitive to small-scale fluctuations in the ISM rather than the average ISM density on kiloparsec scales.

However, bursts that appear to have no coincident host galaxy to deep optical/NIR limits of  $\gtrsim 26$  mag and are located  $\approx 30-75$  kpc from the nearest most probable host galaxy ("host-less" bursts; Berger 2010; Fong & Berger 2013; Tunnicliffe et al. 2013) are expected to have low inferred densities. Indeed, the three bursts located at offsets of  $\gtrsim 10r_e$  have low densities of  $\lesssim 10^{-4}$  cm<sup>-3</sup>, as expected if these bursts occur in the IGM or outer halos of their hosts.

#### 6. CONCLUSIONS

We present the most comprehensive catalog of short GRB afterglows to date, representing a decade of observations since the launch of *Swift* in 2004. This catalog is comprised of 103 short GRBs with prompt X-ray, optical/NIR and radio follow-up, enabled by broad-band Target-of-Opportunity programs. Applying the synchrotron afterglow model to the observations, we also place constraints on the isotropic-equivalent kinetic energies and circumburst densities for a subset of 38 events with well-sampled data sets. While a handful of short GRB afterglows have been studied in detail on an individual basis, our work presents the energy and density scales for a large population of events for the first time. We come to the following key conclusions:

• The afterglow observations presented in this work in-

clude 71 X-ray detections, 30 optical/NIR detections, and 4 radio detections. The detection fractions are 91%, 40% and 7%, respectively, after accounting for observing constraints. We present new optical/NIR observations for 11 events, and new radio observations for 25 bursts.

- Applying a synchrotron model to the broad-band afterglows of 38 bursts, we calculate the inferred circumburst density. Considering a range of scenarios with varying values for the microphysical parameters, cooling frequency, and minimum circumburst density, the median circumburst density is  $(3-15) \times 10^{-3}$  cm<sup>-3</sup>, with 90% upper limits of  $n_{90} \approx 0.4-3$  cm<sup>-3</sup>. Furthermore,  $\approx 80-95\%$  of the probability is below  $\approx 1$  cm<sup>-3</sup>. This indicates that overall short GRBs explode in low-density environments.
- Depending on the set of assumptions in our analysis, we infer a median isotropic-equivalent kinetic energy of  $\approx (1-3) \times 10^{51}$  erg (considering all scenarios), and an isotropic-equivalent  $\gamma$ -ray energy scale of  $\approx 2 \times 10^{51}$  erg. We find a median  $\gamma$ -ray efficiency of  $\approx 0.40-0.56$ .
- We find no obvious trends between circumburst density and host galaxy offset for projected offsets of  $\lesssim 10$  kpc (or  $\lesssim 5~r_e$ ), and no trend between density and host galaxy type, indicating that the circumburst density is not strongly dependent on the average ISM density. However, three bursts in our sample with offsets of  $\gtrsim 10$  kpc have low densities of  $\lesssim 10^{-4}$  cm<sup>-3</sup>, as expected if these bursts explode in the IGM.
- Using 11 short GRBs with opening angle measurements and lower limits, and assuming a maximum value on the opening angle of 30°, we calculate a median jet opening angle of  $16\pm10$  deg and a median beaming factor of  $0.04^{+0.07}_{-0.03}$ . This results in a beaming-corrected total energy release of  $\approx 1.6^{+3.9}_{-1.3} \times 10^{50}$  erg ( $1\sigma$  range), which is broadly consistent with the two primary proposed mechanisms of energy extraction,  $\nu\bar{\nu}$  annihilation and MHD processes. The beaming-corrected volumetric rate is  $\approx 270^{+1580}_{-180}$  Gpc<sup>-3</sup> yr<sup>-1</sup> with an all-sky event rate within a volume of 200 Mpc of  $8^{+47}_{-5}$  yr<sup>-1</sup>. This range is fully consistent with the expected detection rates of gravitational wave signals from neutron star mergers by Advanced LIGO/VIRGO within the same volume.

Our study highlights the importance of broad-band observations in constraining the basic properties of short GRBs. For bursts with detected radio afterglows, we can start to constrain the microphysical parameters, which has thus far only been possible for long GRBs. While our study provides the isotropic-equivalent  $\gamma$ -ray and kinetic energy scales, the true energy release depends on the degree of jet collimation for short GRBs. Current knowledge of the collimation of short GRBs relies on only a handful of events with measured opening angles from their light curves, primarily due to the faintness of short GRB afterglows which prevent temporal monitoring on timescales longer than 1-2 days. Therefore, it is imperative to use the most sensitive ground- and space-based resources to uncover additional collimated events or

place meaningful lower limits on the opening angles. It is especially important to undertake these studies while *Swift* is in operation, since this satellite has the unique capability of providing multi-wavelength light curves within minutes after the bursts.

The past decade of short GRB observations has enabled significant progress in understanding the basic properties of short GRBs, namely their energetics, circumburst densities, and opening angles. Furthermore, in addition to informing the behavior of on-axis afterglows, the circumburst density and energy are key parameters which feed in to predictions for electromagnetic counterparts to compact object mergers, such as off-axis afterglows (Granot et al. 2002; van Eerten et al. 2010) and long-lived radio flares from mildly relativistic ejecta (Nakar & Piran 2011). Advanced LIGO/VIRGO will detect NS-NS mergers within a horizon distance of  $\approx 200$  Mpc (LIGO Scientific Collaboration et al. 2013), making these alternative electromagnetic signatures promising for joint detection with gravitational waves. In a subsequent work, we will assess the detectability of such counterparts by using the distributions of circumburst densities and energies of on-axis short GRBs as inputs, which will help to inform search strategies in the upcoming revolutionary era of gravitational wave discovery.

We acknowledge Matthew Bayliss, Kathy Cooksey, Francesco Di Mille, Steven Elhert, Michael Florian, Tolga Guver, Traci Johnson, Dan Kelson, Michael McDonald, Andy Monson, David Osip, Benjamin Rackham, Nathan Sanders, Anil Seth, Meghin Spencer, Brian Stalder, Tony Stark, Rik Williams, and Amanda Zangari for their assistance in Magellan Target-of-Opportunity observations. Partial support for this work was provided by NASA through Einstein Postdoctoral Fellowship grant number PF4-150121 and Chandra Award Number G04-15055X issued by the Chandra X-ray Observatory Center, which is operated by the Smithsonian Astrophysical Observatory for NASA under contract NAS8-03060. BAZ acknowledges support from NSF Grant AST-1302954, EB acknowledges partial support from NSF Grant AST-1107973, and RM acknowledges support from the James Arthur Fellowship at NYU. Additional support was provided by several NASA/Swift grants. This work made use of data supplied by the UK Swift Science Data Centre at the University of Leicester. The scientific results reported in this article are in part based on observations made by the Chandra X-ray Observatory and data obtained from the Chandra Data Archive. This paper includes data gathered with the 6.5 meter Magellan Telescopes located at Las Campanas Observatory, Chile. Observations reported here were obtained at the MMT Observatory, a joint facility of the University of Arizona and the Smithsonian Institution. Based on observations obtained at the Gemini Observatory acquired through the Gemini Science Archive and processed using the Gemini IRAF package which is operated by the Association of Universities for Research in Astronomy, Inc., under a cooperative agreement with the NSF on behalf of the Gemini partnership: the National Science Foundation (United States), the National Research Council (Canada), CONICYT (Chile), the Australian Research Council (Australia), Ministério da Ciência, Tecnologia e Inovação (Brazil) and Ministerio de Ciencia, Tecnología e Innovación Productiva (Argentina). The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc.

Alard, C. 2000, A&AS, 144, 363

#### REFERENCES

Aloy, M. A., Janka, H.-T., & Müller, E. 2005, A&A, 436, 273 Antonelli, L. A., et al. 2009, A&A, 507, L45 Aptekar, R. L., et al. 1995, Space Sci. Rev., 71, 265 Atwood, W. B., et al. 2009, ApJ, 697, 1071 Beardmore, A. P., Page, K. L., Evans, P. A., Pagani, C., Kennea, J., & Burrows, D. N. 2009, GRB Coordinates Network, 8937, 1 Beardmore, A. P., Page, K. L., Gehrels, N., Greiner, J., Kennea, J., Nousek, J., Osborne, J. P., & Tagliaferri, G. 2005, GRB Coordinates Network, 4043, 1 Belczynski, K., Perna, R., Bulik, T., Kalogera, V., Ivanova, N., & Lamb, D. Q. 2006, ApJ, 648, 1110 Berger, E. 2007, ApJ, 670, 1254 Berger, E. 2009, ApJ, 690, 231 Berger, E. 2010, ApJ, 722, 1946 Berger, E. 2014, ARA&A, 52, 43 Berger, E., Cenko, S. B., Fox, D. B., & Cucchiara, A. 2009, ApJ, 704, 877 Berger, E., & Challis, P. 2007, GRB Coordinates Network, 7085, 1 Berger, E., & Fong, W. 2009a, GRB Coordinates Network, 9373, 1 Berger, E., & Fong, W. 2009b, GRB Coordinates Network, 10182, 1 Berger, E., Fong, W., & Chornock, R. 2013, ApJ, 774, L23 Berger, E., & Fong, W.-F. 2009c, GRB Coordinates Network, 9555, 1 Berger, E., & Fong, W.-F. 2009d, GRB Coordinates Network, 9683, 1 Berger, E., et al. 2007, ApJ, 664, 1000 Berger, E., Kulkarni, S. R., & Frail, D. A. 2003, ApJ, 590, 379 Berger, E., & Mulchaey, J. 2009, GRB Coordinates Network, 10174, 1 Berger, E., et al. 2005, Nature, 438, 988 Berger, E., et al. 2013, ApJ, 765, 121 Birkl, R., Aloy, M. A., Janka, H.-T., & Müller, E. 2007, A&A, 463, 51 Blandford, R. D., & Znajek, R. L. 1977, MNRAS, 179, 433 Blanton, M. R., & Roweis, S. 2007, AJ, 133, 734 Bloom, J. S., Frail, D. A., & Kulkarni, S. R. 2003, ApJ, 594, 674 Bloom, J. S., et al. 2006, ApJ, 638, 354 Bromberg, O., Nakar, E., Piran, T., & Sari, R. 2013, ApJ, 764, 179 Butler, N., et al. 2013, GRB Coordinates Network, 15226, 1 Butler, N., et al. 2014, GRB Coordinates Network, 16297, 1 Cameron, P. B., & Frail, D. A. 2005a, GRB Coordinates Network, 3815, 1 Cameron, P. B., & Frail, D. A. 2005b, GRB Coordinates Network, 3933, 1 Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, ApJ, 345, 245 Castro-Tirado, A. J., Bond, I., Kilmartin, P., de Ugarte Postigo, A., Gorosabel, J., Jelinek, M., & Yock, P. 2005, GRB Coordinates Network, 3018.1 Cenko, S. B., Capone, J., Toy, V., Cucchiara, A., Troja, E., Kutyrev, A., Veilleux, S., & Gezari, S. 2014, GRB Coordinates Network, 16785, 1 Cenko, S. B., & Cucchiara, A. 2013, GRB Coordinates Network, 14670, 1 Cenko, S. B., et al. 2011, ApJ, 732, 29 Cenko, S. B., et al. 2010, ApJ, 711, 641 Cenko, S. B., & Kasliwal, M. M. 2008, GRB Coordinates Network, 8417, 1 Cenko, S. B., et al. 2009, ApJ, 693, 1484 Cenko, S. B., Perley, D. A., Cucchiara, A., Fong, W., & Levan, A. J. 2013a, GRB Coordinates Network, 15121, 1 Cenko, S. B., Sudilovsky, V., Tanga, M., & Greiner, J. 2013b, GRB Coordinates Network, 15222, 1 Chandra, P., Cenko, S. B., Frail, D. A., & Harrison, F. 2008, GRB Coordinates Network, 8451, 1 Chandra, P., & Frail, D. A. 2006, GRB Coordinates Network, 5910, 1 Chandra, P., & Frail, D. A. 2007a, GRB Coordinates Network, 6667, 1 Chandra, P., & Frail, D. A. 2007b, GRB Coordinates Network, 6367, 1 Chandra, P., & Frail, D. A. 2007c, GRB Coordinates Network, 6831, 1 Chandra, P., & Frail, D. A. 2007d, GRB Coordinates Network, 6685, 1 Chandra, P., & Frail, D. A. 2007e, GRB Coordinates Network, 6742, 1 Chandra, P., & Frail, D. A. 2008a, GRB Coordinates Network, 8419, 1 Chandra, P., & Frail, D. A. 2008b, GRB Coordinates Network, 7934, 1 Chandra, P., & Frail, D. A. 2009, GRB Coordinates Network, 9160, 1 Chevalier, R. A., & Li, Z.-Y. 2000, ApJ, 536, 195 Covino, S., et al. 2006, A&A, 447, L5 Covino, S., Piranomonte, S., D'Avanzo, P., A., A. L., de Luise, F., Pedani, M., & Pinilla Alfonso, N. 2007, GRB Coordinates Network, 6635, 1

Coward, D. M., et al. 2012, MNRAS, 425, 2668 Cucchiara, A., et al. 2013, ApJ, 777, 94

Network, 16113, 1

D'Avanzo, P., Bernardini, M. G., & Campana, S. 2014, GRB Coordinates

D'Avanzo, P., Covino, S., Melandri, A., Padilla Torres, C. P., & Andreuzzi, G. 2013, GRB Coordinates Network, 14307, 1 D'Avanzo, P., D'Elia, V., Lorenzi, V., Mainella, G., Boschin, W., Garcia de Gurtubai Escudero, A., & Levan, A. J. 2015, GRB Coordinates Network, D'Avanzo, P., Fugazza, D., di Fabrizio, L., & Mainella, G. 2014a, GRB Coordinates Network, 16112, 1 D'Avanzo, P., et al. 2009, A&A, 498, 711 D'Avanzo, P., Melandri, A., Bernardini, M. G., Cecconi, M., & Mainella, G. 2014b, GRB Coordinates Network, 16286, 1 de Ugarte Postigo, A., et al. 2006, ApJ, 648, L83 de Ugarte Postigo, A., Kann, D. A., Castro-Tirado, A. J., Aguirre, A., & Jordi, C. 2008, GRB Coordinates Network, 7644, 1 de Ugarte Postigo, A., et al. 2013, ArXiv e-prints de Ugarte Postigo, A., et al. 2014, A&A, 563, A62 Eichler, D., Livio, M., Piran, T., & Schramm, D. N. 1989, Nature, 340, 126 Evans, P. A., et al. 2009, MNRAS, 397, 1177 Evans, P. A., et al. 2007a, A&A, 469, 379 Evans, P. A., & Hoversten, E. A. 2008, GRB Coordinates Network, 8740, 1 Evans, P. A., et al. 2007b, GRB Coordinates Network, 6942, 1 Ferrero, P., et al. 2007, AJ, 134, 2118 Filgas, R., et al. 2011, A&A, 526, A113 Fong, W., & Berger, E. 2013, ApJ, 776, 18 Fong, W., et al. 2013, ApJ, 769, 56 Fong, W., et al. 2011, ApJ, 730, 26 Fong, W., Berger, E., & Fox, D. B. 2010, ApJ, 708, 9 Fong, W., Berger, E., Kelson, D., & Williams, R. 2012a, GRB Coordinates Network, 13700, 1 Fong, W., et al. 2012b, ApJ, 756, 189 Fong, W., et al. 2014, ApJ, 780, 118 Fox, D. B., et al. 2005, Nature, 437, 845 Frail, D. A., & Cameron, P. B. 2005, GRB Coordinates Network, 4199, 1 Frail, D. A., & Chandra, P. 2008, GRB Coordinates Network, 7684, 1 Frail, D. A., & Chandra, P. 2009, GRB Coordinates Network, 9354, 1 Frail, D. A., et al. 2001, ApJ, 562, L55 Frail, D. A., & Soderberg, A. M. 2005, GRB Coordinates Network, 3007, 1 Friedman, A. S., & Bloom, J. S. 2005, ApJ, 627, 1 Fukazawa, Y., Botoya-Nonesa, J. G., Pu, J., Ohto, A., & Kawano, N. 2006, ApJ, 636, 698 Galeev, A., et al. 2009, GRB Coordinates Network, 9549, 1 Gehrels, N., et al. 2008, ApJ, 689, 1161 Gehrels, N., et al. 2004, ApJ, 611, 1005 Gehrels, N., et al. 2005, Nature, 437, 851 Gent, F. A., Shukurov, A., Fletcher, A., Sarson, G. R., & Mantere, M. J. 2013, MNRAS, 432, 1396 Ghirlanda, G., Ghisellini, G., & Lazzati, D. 2004, ApJ, 616, 331 Goldstein, A., et al. 2011, ArXiv e-prints Gompertz, B. P., Page, K. L., & de Pasquale, M. 2013, GRB Coordinates Network, 15610, 1 Gorosabel, J., Sanchez-Ramirez, R., Hellmich, S., & Mottola, S. 2014a, GRB Coordinates Network, 17167, 1 Gorosabel, J., et al. 2014b, GRB Coordinates Network, 16290, 1 Graham, J., Nicuesa Guelbenzu, A., Bolmer, J., & Greiner, J. 2014, GRB Coordinates Network, 16872, 1 Graham, J. F., et al. 2009, ApJ, 698, 1620 Granot, J., Panaitescu, A., Kumar, P., & Woosley, S. E. 2002, ApJ, 570, L61 Granot, J., & Sari, R. 2002, ApJ, 568, 820 Greco, G., et al. 2008, GRB Coordinates Network, 7977, 1 Greisen, E. W. 2003, Information Handling in Astronomy - Historical Vistas, 285, 109 Grupe, D., et al. 2010, GCN Report, 268, 1 Grupe, D., Burrows, D. N., Patel, S. K., Kouveliotou, C., Zhang, B., Mészáros, P., Wijers, R. A. M., & Gehrels, N. 2006, ApJ, 653, 462 Guetta, D., & Piran, T. 2005, A&A, 435, 421 Guidorzi, C., Melandri, A., O'Brien, P., & Tanvir, N. 2009, GRB Coordinates Network, 9492, 1 Haislip, J., et al. 2009, GRB Coordinates Network, 9921, 1 Hancock, P., Murphy, T., Gaensler, B., Bell, M., Burlon, D., & de Ugarte Postigo, A. 2012a, GRB Coordinates Network, 13338, 1 Hancock, P., Murphy, T., Gaensler, B., Bell, M., Burlon, D., & de Ugarte Postigo, A. 2012b, GRB Coordinates Network, 13611, 1 Hjorth, J., et al. 2005, Nature, 437, 859 Hurley, K., et al. 2010, in American Institute of Physics Conference Series, Vol. 1279, American Institute of Physics Conference Series, ed. N. Kawai & S. Nagataki, 330 Jaroszynski, M. 1993, AcA, 43, 183 Johnson, S., et al. 2007, GRB Coordinates Network, 6218, 1

Kalberla, P. M. W., Burton, W. B., Hartmann, D., Arnal, E. M., Bajaja, E., Morras, R., & Pöppel, W. G. L. 2005, A&A, 440, 775

Kann, D. A., et al. 2010, ApJ, 720, 1513

Kann, D. A., et al. 2008, ArXiv e-prints

Kocevski, D., Bloom, J. S., Thoene, C. C., & Prochaska, J. 2007, GRB Coordinates Network, 6771, 1

Kocevski, D., & Butler, N. 2008, ApJ, 680, 531

Kocevski, D., et al. 2010, MNRAS, 404, 963

Kopac, D., Smith, R. J., Mundell, C. G., Tanvir, N. R., & Gomboc, A. 2013, GRB Coordinates Network, 15306, 1

Korpi, M. J., Brandenburg, A., Shukurov, A., Tuominen, I., & Nordlund, Å. 1999, ApJ, 514, L99

Kouveliotou, C., Meegan, C. A., Fishman, G. J., Bhat, N. P., Briggs, M. S.,Koshut, T. M., Paciesas, W. S., & Pendleton, G. N. 1993, ApJ, 413, L101Laskar, T., et al. 2014, ApJ, 781, 1

Lee, W. H., & Ramirez-Ruiz, E. 2007, New Journal of Physics, 9, 17

Leibler, C. N., & Berger, E. 2010, ApJ, 725, 1202

Levan, A. J., et al. 2006, ApJ, 648, L9

Levan, A. J., et al. 2008, MNRAS, 384, 541

Levesque, E. M., et al. 2010, MNRAS, 401, 963

LIGO Scientific Collaboration, et al. 2013, ArXiv e-prints

Littlejohns, O., et al. 2013, GRB Coordinates Network, 15312, 1

Littlejohns, O., et al. 2014, GRB Coordinates Network, 16026, 1

Littlejohns, O. M., de Pasquale, M., & Morris, D. C. 2011, GRB Coordinates Network, 11567, 1

Malesani, D., et al. 2007a, A&A, 473, 77

Malesani, D., D'Avanzo, P., di Fabrizio, L., & Tessicini, G. 2013, GRB Coordinates Network, 15305, 1

Malesani, D., Uthas, H., Thoene, C. C., Fynbo, J. P. U., Hjorth, J., & Andersen, M. I. 2007b, GRB Coordinates Network, 6254, 1

Margutti, R., et al. 2012, ApJ, 756, 63

Margutti, R., et al. 2011, MNRAS, 417, 2144

Margutti, R., et al. 2013, MNRAS, 428, 729

Marshall, F. E., Curran, P. A., Page, K. L., Barthelmy, S. D., Burrows, D. N., Roming, P., & Gehrels, N. 2009, GCN Report, 220, 1

Maselli, A., & D'Avanzo, P. 2014, GRB Coordinates Network, 16424, 1 McLeod, B., & Williams, G. 2009, GRB Coordinates Network, 9370, 1 Meegan, C., et al. 2009, ApJ, 702, 791

Melandri, A., Burgdorf, M., Smith, R. J., Steele, I. A., Bersier, D. F., Mundell, C. G., Gomboc, A., & Guidorzi, C. 2008, GRB Coordinates Network, 8402, 1

Metzger, B. D., & Berger, E. 2012, ApJ, 746, 48

Minowa, Y., Pyo, T.-S., Kim, J.-H., & Kawai, N. 2014, GRB Coordinates Network, 16296, 1

Mochkovitch, R., Hernanz, M., Isern, J., & Martin, X. 1993, Nature, 361, 236

Moin, A., Tingay, S., Phillips, C., Taylor, G., Wieringa, M., & Martin, R. 2009a, GRB Coordinates Network, 8952, 1

Moin, A., Tingay, S., Phillips, C., Taylor, G., Wieringa, M., & Martin, R. 2009b, GRB Coordinates Network, 8953, 1

Murali, C. 2000, ApJ, 529, L81

Nakar, E. 2007, Phys. Rep., 442, 166

Nakar, E., Gal-Yam, A., & Fox, D. B. 2006, ApJ, 650, 281

Nakar, E., & Piran, T. 2011, Nature, 478, 82

Narayan, R., Paczynski, B., & Piran, T. 1992, ApJ, 395, L83

Nayana, A. J., & Chandra, P. 2014, GRB Coordinates Network, 16815, 1

Nicuesa Guelbenzu, A., et al. 2012a, A&A, 548, A101

Nicuesa Guelbenzu, A., et al. 2012b, A&A, 538, L7

Nicuesa Guelbenzu, A., et al. 2011, A&A, 531, L6

Nissanke, S., Kasliwal, M., & Georgieva, A. 2013, ApJ, 767, 124

Nousek, J. A., et al. 2006, ApJ, 642, 389

Nysewander, M., Fruchter, A., & Graham, J. 2007, GRB Coordinates Network, 6604, 1

Nysewander, M., Fruchter, A. S., & Pe'er, A. 2009, ApJ, 701, 824

Pagani, C. 2012, GRB Coordinates Network, 13680, 1

Pagani, C. 2014, GRB Coordinates Network, 16075, 1

Pagani, C., & Evans, P. A. 2014, GRB Coordinates Network, 17134, 1

Pagani, C., La Parola, V., & Burrows, D. N. 2005, GRB Coordinates Network, 3934, 1

Page, K. L., & de Pasquale, M. 2013, GRB Coordinates Network, 15004, 1 Panaitescu, A. 2006, MNRAS, 367, L42

Panaitescu, A., & Kumar, P. 2002, ApJ, 571, 779

Perley, D. A., Bloom, J. S., Modjaz, M., Poznanski, D., & Thoene, C. C. 2007, GRB Coordinates Network, 7140, 1

Perley, D. A., et al. 2009a, AJ, 138, 1690

Perley, D. A., & Jencson, J. 2014, GRB Coordinates Network, 16867, 1

Perley, D. A., et al. 2013, ApJ, 778, 128

Perley, D. A., et al. 2009b, ApJ, 696, 1871

Perley, D. A., Modjaz, M., Morgan, A. N., Cenko, S. B., Bloom, J. S., Butler, N. R., Filippenko, A. V., & Miller, A. A. 2012, ApJ, 758, 122

Perley, D. A., Thoene, C. C., & Bloom, J. S. 2007, GRB Coordinates Network, 6774, 1

Perley, R. A., Chandler, C. J., Butler, B. J., & Wrobel, J. M. 2011, ApJ, 739, L1

Perna, R., & Belczynski, K. 2002, ApJ, 570, 252

Perri, M., Stratta, G., Fenimore, E., Schady, P., Barthelmy, S. D., Burrows, D. N., Roming, P., & Gehrels, N. 2007, GCN Report, 103, 1

Piranomonte, S., et al. 2008, A&A, 491, 183

Poleshchuk, V. A., et al. 2014, GRB Coordinates Network, 15776, 1

Popham, R., Woosley, S. E., & Fryer, C. 1999, ApJ, 518, 356

Price, P. A., Berger, E., Fox, D. B., Cenko, S. B., & Rau, A. 2006, GRB Coordinates Network, 5077, 1

Racusin, J. L., et al. 2009, ApJ, 698, 43

Rezzolla, L., Giacomazzo, B., Baiotti, L., Granot, J., Kouveliotou, C., & Aloy, M. A. 2011, ApJ, 732, L6

Ricker, G. R., et al. 2003, in American Institute of Physics Conference Series, Vol. 662, Gamma-Ray Burst and Afterglow Astronomy 2001: A Workshop Celebrating the First Year of the HETE Mission, ed. G. R. Ricker & R. K. Vanderspek, 3

Roming, P. W. A., et al. 2006, ApJ, 651, 985

Rossi, A., Nardini, M., Klose, S., & Greiner, J. 2012, GRB Coordinates Network, 13335, 1

Rosswog, S. 2005, Nuovo Cimento C Geophysics Space Physics C, 28, 607 Rosswog, S., Ramirez-Ruiz, E., & Davies, M. B. 2003, MNRAS, 345, 1077 Rowlinson, A., et al. 2010a, MNRAS, 409, 531

Rowlinson, A., et al. 2010b, MNRAS, 408, 383

Ruffert, M., & Janka, H. 1999a, Progress of Theoretical Physics Supplement, 136, 287

Ruffert, M., & Janka, H.-T. 1999b, A&A, 344, 573

Ryan, G., van Eerten, H., MacFadyen, A., & Zhang, B.-B. 2015, ApJ, 799, 3 Sakamoto, T., et al. 2013, ApJ, 766, 41

Sari, R., Piran, T., & Halpern, J. P. 1999, ApJ, 519, L17

Sari, R., Piran, T., & Narayan, R. 1998, ApJ, 497, L17

Sault, R. J., Teuben, P. J., & Wright, M. C. H. 1995, in Astronomical Society of the Pacific Conference Series, Vol. 77, Astronomical Data Analysis Software and Systems IV, ed. R. A. Shaw, H. E. Payne, & J. J. E. Hayes, 422

Schlafly, E. F., & Finkbeiner, D. P. 2011, ApJ, 737, 103

Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, ApJ, 500, 525

Singer, L. P., Kasliwal, M. M., & Cenko, S. B. 2013a, GRB Coordinates Network, 15552, 1

Singer, L. P., Kasliwal, M. M., & Cenko, S. B. 2013b, GRB Coordinates Network, 15572, 1

Soderberg, A. M., et al. 2006, ApJ, 650, 261

Soderberg, A. M., & Frail, D. A. 2006, GRB Coordinates Network, 4884, 1 Soderberg, A. M., Frail, D. A., & Chandra, P. 2006, GRB Coordinates Network, 5388, 1

Starling, R., Markwardt, C., Marshall, F. E., & Sakamoto, T. 2007, GRB Coordinates Network, 6754, 1

Starling, R. L. C., & Page, K. L. 2014, GRB Coordinates Network, 17141, 1 Stratta, G., et al. 2007a, A&A, 474, 827

Stratta, G., Perri, M., Conciatore, M. L., Burrows, D. N., & Sato, G. 2007b, GRB Coordinates Network, 6119, 1

Stroh, M. C., Kennea, J. A., Gehrels, N., & Malesani, D. 2014, GRB Coordinates Network, 16357, 1

Tanga, M., Delvaux, C., & Greiner, J. 2014, GRB Coordinates Network, 16435, 1

Tanga, M., Klose, S., Sudilovsky, V., Filgas, R., & Greiner, J. 2013, GRB Coordinates Network, 15214, 1

Tanvir, N. R., Levan, A. J., & Fraser, M. 2014, GRB Coordinates Network, 16861. 1

Tanvir, N. R., Levan, A. J., Fruchter, A. S., Hjorth, J., Hounsell, R. A., Wiersema, K., & Tunnicliffe, R. L. 2013, Nature, 500, 547

Tanvir, N. R., Wiersema, K., & Levan, A. J. 2013, GRB Coordinates Network, 15224, 1

Troja, E., Cummings, J. R., Palmer, D. M., Cucchiara, A., Barthelmy, S. D.,Burrows, D. N., Roming, P., & Gehrels, N. 2008, GCN Report, 118, 1Troja, E., et al. 2009, GCN Report, 248, 1

Troja, E., Page, K., & McBreen, S. 2007, GRB Coordinates Network, 6255,

Tunnicliffe, R. L., et al. 2013, MNRAS

Ukwatta, T. N., et al. 2008, GCN Report, 111, 1

Updike, A. C., Bryngelson, G., & Milne, P. A. 2009, GRB Coordinates Network, 9361, 1

van der Horst, A. J. 2005, GRB Coordinates Network, 4039, 1

van der Horst, A. J. 2006, GRB Coordinates Network, 5408, 1

van der Horst, A. J., Levan, A. J., Wiersema, K., Tanvir, N. R., & Hjorth, J. 2015, GRB Coordinates Network, 17309, 1

van der Horst, A. J., Wiersema, K., & Wijers, R. A. M. J. 2005, GRB Coordinates Network, 3405, 1

van Eerten, H., Zhang, W., & MacFadyen, A. 2010, ApJ, 722, 235 Virgili, F. J., Guidorzi, C., Melandri, A., & Mundell, C. G. 2012, GRB Coordinates Network, 13006, 1

Volnova, A., Inasaridze, R., Kvaratskhelia, O., Ayvazian, V., Krugly, Y., Molotov, I., & Pozanenko, A. 2014, GRB Coordinates Network, 16371, 1 Wakker, B. P., Lockman, F. J., & Brown, J. M. 2011, ApJ, 728, 159

Wieringa, M. H., Chandra, P., & Frail, D. A. 2007, GRB Coordinates Network, 7095, 1 Willingale, R., Starling, R. L. C., Beardmore, A. P., Tanvir, N. R., & O'Brien, P. T. 2013, MNRAS, 431, 394

Xin, L.-P., et al. 2011, MNRAS, 410, 27

Xu, D., Malesani, D., Hjorth, J., Jakobsson, P., Kruehler, T., Santiago, E., Groeneboom, N., & Hafreager, A. 2013, GRB Coordinates Network, 15304

Xu, D., Niu, H.-B., Esamdin, A., & Ma, L. 2014, GRB Coordinates Network, 17142, 1

Xu, D., Smirnova, O., Purismo, T., Fynbo, J. P. U., & Jakobsson, P. 2012, GRB Coordinates Network, 14110, 1

Yost, S. A., Harrison, F. A., Sari, R., & Frail, D. A. 2003, ApJ, 597, 459
 Zaninoni, E., Bernardini, M. G., Margutti, R., Oates, S., & Chincarini, G. 2013, A&A, 557, A12

Zhang, B., Fan, Y. Z., Dyks, J., Kobayashi, S., Mészáros, P., Burrows, D. N., Nousek, J. A., & Gehrels, N. 2006, ApJ, 642, 354

# APPENDIX BROAD-BAND AFTERGLOW CATALOGS

Table A1
Short GRB Late-time X-ray afterglow catalog

				-time A-ray aftergrow cata	
GRB	$\frac{\delta t}{(\mathrm{s})}$	Exposure Time (s)	$F_X \ (\mu \mathbf{J} \mathbf{y})$	$\sigma_X \ (\mu  ext{Jy})$	References
050509B	831	381	0.07	0.02	Gehrels et al. (2005)
	$6.2 \times 10^{3}$	$2.0 \times 10^{3}$	< 0.02		
	$^{+}2.1 \times 10^{5}$	$5.0 \times 10^{4}$	$< 8.0 \times 10^{-3}$		
050709	$1.4 \times 10^{5}$	$1.5 \times 10^{4}$	$1.9 \times 10^{-3}$	$1.4 \times 10^{-4}$	Fox et al. (2005)
	$2.1 \times 10^{5}$	$5.0 \times 10^{3}$	$< 1.5 \times 10^{-3}$		
	$^{+}2.2 \times 10^{5}$	$3.8 \times 10^{4}$	$1.1 \times 10^{-3}$	$2.0 \times 10^{-5}$	
	$2.8 \times 10^{5}$	$6.4 \times 10^{3}$	$< 7.3 \times 10^{-4}$		
	$3.7 \times 10^{5}$	$2.1 \times 10^{3}$	$< 5.9 \times 10^{-3}$		
	$^{+}1.4 \times 10^{6}$	$6.1 \times 10^{3}$	$1.3 \times 10^{-3}$	$6.0 \times 10^{-5}$	
	$^{+}1.4 \times 10^{6}$	$1.2 \times 10^{4}$	$1.1 \times 10^{-4}$	$2.0 \times 10^{-5}$	
050724A	$1.0 \times 10^{3}$	514	0.43	0.086	Grupe et al. (2006)
	$1.4 \times 10^{3}$	323	0.68	0.14	
	$1.9 \times 10^{3}$	493	0.44	0.090	
	$6.6 \times 10^{3}$	$2.0 \times 10^{3}$	0.091	0.022	
	$1.3 \times 10^4$	$3.4 \times 10^{3}$	0.050	0.013	
	$2.5 \times 10^4$	$1.9 \times 10^{3}$	0.10	0.023	
	$3.5 \times 10^{4}$	805	0.27	0.056	
	$3.7 \times 10^4$	$1.2 \times 10^{3}$	0.19	0.039	
	$4.1 \times 10^{4}$	830	0.26	0.053	
	$4.2 \times 10^{4}$	652	0.32	0.068	
	$4.2 \times 10^{4}$	772	0.27	0.057	
	$4.7 \times 10^{4}$	$1.2 \times 10^{3}$	0.18	0.038	
	$4.8 \times 10^{4}$	880	0.24	0.050	
	$4.8 \times 10^{4}$	790	0.27	0.056	
	$5.2 \times 10^4$	587	0.37	0.075	
	$5.3 \times 10^4$	800	0.27	0.055	
	$5.8 \times 10^4$	759	0.29	0.058	
	$5.9 \times 10^4$	554	0.39	0.080	
	$6.3 \times 10^4$	972	0.22	0.046	
	$6.6 \times 10^4$	$1.0 \times 10^{3}$	0.20	0.044	
	$7.7 \times 10^4$	$1.6 \times 10^{3}$	0.11	0.027	
	$8.8 \times 10^{4}$	$2.0 \times 10^{3}$	0.10	0.023	
	$9.9 \times 10^{4}$	$1.2 \times 10^{3}$	0.090	0.028	
	$1.4 \times 10^{5}$	$4.4 \times 10^{3}$	0.025	$8.4 \times 10^{-3}$	
	$1.5 \times 10^{5}$	$4.0 \times 10^{3}$	0.027	$9.5 \times 10^{-3}$	
	$1.8 \times 10^{5}$	$7.0 \times 10^{3}$	0.012	$5.5 \times 10^{-3}$	
	$2.2 \times 10^{5}$	$2.1 \times 10^{3}$	0.014	$8.4 \times 10^{-3}$	
	$5.6 \times 10^{5}$	$4.7 \times 10^4$	$1.4 \times 10^{-3}$	$7.8 \times 10^{-4}$	
	$^{+}2.1 \times 10^{5}$	$8.3 \times 10^{3}$	$6.0 \times 10^{-3}$	$1.1 \times 10^{-3}$	
	$^{+}2.1 \times 10^{5}$	$9.6 \times 10^{3}$	$5.1 \times 10^{-3}$	$9.7 \times 10^{-4}$	
	$^{+}2.3 \times 10^{5}$	$1.2 \times 10^4$	$3.9 \times 10^{-3}$	$7.5 \times 10^{-4}$	
	$^{+}2.4 \times 10^{5}$	$1.3 \times 10^4$	$3.8 \times 10^{-3}$	$7.2 \times 10^{-4}$	
	$^{+}2.5 \times 10^{5}$	$6.5 \times 10^3$	$5.4 \times 10^{-3}$	$1.2 \times 10^{-3}$	
050012	$^{+}1.9 \times 10^{6}$	$4.3 \times 10^4$	$3.3 \times 10^{-4}$	$1.2 \times 10^{-4}$	F (2007 2006)
050813	$4.7 \times 10^{3}$	$2.7 \times 10^{5}$	$3.4 \times 10^{-3}$	$1.0 \times 10^{-3}$	Evans et al. (2007a, 2009)

Table A1 — Continued

GRB	$\frac{\delta t}{(s)}$	Exposure Time (s)	$F_X \ (\mu J y)$	$\sigma_X \ (\mu  ext{Jy})$	References
	$6.5 \times 10^{5}$	$3.4 \times 10^{5}$	$< 2.7 \times 10^{-3}$		
050906	$3.0 \times 10^3$	$5.2 \times 10^3$	< 0.018		Pagani et al. (2005)
050925 051210	$1.4 \times 10^{3}$ 894	$1.4 \times 10^3$ $840$	$<6.6 \times 10^{-3}$ $0.40$	0.15	Beardmore et al. (2005) This work
051210 051221A	$1.4 \times 10^3$	512	0.86	0.10	Soderberg et al. (2006); Belczynski et al. (2006)
03122111	$1.9 \times 10^{3}$	512	0.59	0.085	Boderoeig et di. (2000), Belezyliski et di. (2000)
	$2.4 \times 10^{3}$	512	0.43	0.082	
	$5.3 \times 10^{3}$	$3.5 \times 10^{3}$	0.23	0.033	
	$8.8 \times 10^{3}$	$3.5 \times 10^{3}$	0.24	0.035	
	$1.2 \times 10^4$	$3.5 \times 10^{3}$	0.25	0.026	
	$1.6 \times 10^{4}$	$3.5 \times 10^{3}$	0.20	0.076	
	$1.9 \times 10^4$	$3.5 \times 10^{3}$	0.16	0.021	
	$2.3 \times 10^4$	$3.5 \times 10^3$	0.11	0.023	
	$2.6 \times 10^4$ $3.0 \times 10^4$	$3.5 \times 10^3$ $3.5 \times 10^3$	0.11	0.026	
	$3.0 \times 10^{4}$ $3.7 \times 10^{4}$	$3.5 \times 10^{-3}$ $3.5 \times 10^{3}$	0.088 0.13	0.015 0.020	
	$4.5 \times 10^4$	$1.0 \times 10^4$	0.079	0.020	
	$5.5 \times 10^4$	$1.0 \times 10^4$ $1.0 \times 10^4$	0.066	0.011	
	$1.0 \times 10^{5}$	$4.1 \times 10^4$	0.017	$2.7 \times 10^{-3}$	
	$1.4 \times 10^{5}$	$4.1 \times 10^4$	0.018	$2.9 \times 10^{-3}$	
	$2.0 \times 10^{5}$	$5.7 \times 10^{4}$	0.013	$1.9 \times 10^{-3}$	
	$2.5 \times 10^{5}$	$5.7 \times 10^{4}$	0.014	$2.0 \times 10^{-3}$	
	$3.1 \times 10^{5}$	$5.7 \times 10^{4}$	$9.0 \times 10^{-3}$	$1.7 \times 10^{-3}$	
	$3.8 \times 10^{5}$	$8.6 \times 10^{4}$	$5.2 \times 10^{-3}$	$1.0 \times 10^{-3}$	
	$4.7 \times 10^{5}$	$8.6 \times 10^{4}$	$5.8 \times 10^{-3}$	$1.1 \times 10^{-3}$	
	$5.5 \times 10^{5}$	$8.6 \times 10^{4}$	$4.4 \times 10^{-3}$	$9.4 \times 10^{-4}$	
	$6.4 \times 10^5$	$8.3 \times 10^4$	$3.6 \times 10^{-3}$	$9.4 \times 10^{-4}$	
	$7.3 \times 10^5$	$8.2 \times 10^4$	$5.4 \times 10^{-3}$	$1.0 \times 10^{-3}$	
	$9.8 \times 10^5$ $^{+}1.3 \times 10^5$	$4.1 \times 10^5$ $5.4 \times 10^3$	$2.3 \times 10^{-3}$ $0.022$	$4.1 \times 10^{-4} \\ 2.2 \times 10^{-3}$	
	$^{+}1.4 \times 10^{5}$	$5.5 \times 10^{3}$	0.022	$2.2 \times 10^{-3}$ $2.1 \times 10^{-3}$	
	$^{+1.4}\times10^{5}$	$6.4 \times 10^3$	0.018	$1.9 \times 10^{-3}$	
	$^{+}1.5 \times 10^{5}$	$6.6 \times 10^3$	0.018	$1.8 \times 10^{-3}$	
	$^{+}1.6 \times 10^{5}$	$6.3 \times 10^{3}$	0.017	$1.8 \times 10^{-3}$	
	$^{+}3.9 \times 10^{5}$	$9.5 \times 10^{3}$	$6.1 \times 10^{-3}$	$8.8 \times 10^{-4}$	
	$^{+}4.0 \times 10^{5}$	$1.0 \times 10^{4}$	$5.5 \times 10^{-3}$	$8.0 \times 10^{-4}$	
	$^{+}4.1 \times 10^{5}$	$1.0 \times 10^{4}$	$5.3 \times 10^{-3}$	$7.9 \times 10^{-4}$	
	$^{+}1.3 \times 10^{6}$	$1.8 \times 10^{4}$	$5.5 \times 10^{-4}$	$1.9 \times 10^{-4}$	
	$^{+}1.7 \times 10^{6}$	$2.5 \times 10^4$	$3.8 \times 10^{-4}$	$1.4 \times 10^{-4}$	
	$^{+}2.3 \times 10^{6}$	$4.8 \times 10^4$	$1.9 \times 10^{-4}$	$7.2 \times 10^{-5}$	
060121	$1.1 \times 10^{4}$	153	0.61	0.14	Evans et al. (2007a, 2009)
	$1.1 \times 10^4$	221	0.42	0.095	
	$1.1 \times 10^4$	206	0.45	0.10	
	$1.1 \times 10^4$ $1.1 \times 10^4$	198 208	0.47 0.34	0.11 0.087	
	$1.1 \times 10^{4}$ $1.2 \times 10^{4}$	208 178	0.52	0.087	
	$1.2 \times 10^{4}$ $1.2 \times 10^{4}$	183	0.52	0.12	
	$1.2 \times 10^4$ $1.2 \times 10^4$	155	0.60	0.14	
	$1.2 \times 10^4$	193	0.36	0.094	
	$1.2 \times 10^{4}$	338	0.37	0.072	
	$1.6 \times 10^{4}$	281	0.27	0.069	
	$1.6 \times 10^{4}$	308	0.24	0.063	
	$1.6 \times 10^{4}$	196	0.39	0.10	
	$1.7 \times 10^4$	258	0.29	0.076	
	$1.7 \times 10^4$	246	0.31	0.080	
	$1.7 \times 10^4$	236	0.32	0.083	
	$1.8 \times 10^4$ $1.8 \times 10^4$	221	0.34	0.089	
	$1.8 \times 10^{4}$ $1.8 \times 10^{4}$	201 617	0.38 0.19	0.10 0.039	
	$1.8 \times 10^{-1}$ $2.2 \times 10^{4}$	361	0.19	0.039	
	$7.0 \times 10^4$	$1.3 \times 10^3$	0.23	0.036	
	$7.5 \times 10^4$	$2.5 \times 10^3$	0.030	$7.9 \times 10^{-3}$	
	$8.1 \times 10^4$	$2.3 \times 10^{3}$ $2.3 \times 10^{3}$	0.053	0.011	
	$8.9 \times 10^4$	$7.6 \times 10^{3}$	0.043	$8.0 \times 10^{-3}$	
	$1.6 \times 10^{5}$	$1.2 \times 10^{4}$	0.021	$4.9 \times 10^{-3}$	

Table A1 - Continued

GRB	δ <i>t</i> (s)	Exposure Time (s)	$F_X$ ( $\mu$ Jy)	$\sigma_X \ (\mu  ext{Jy})$	References
	$1.7 \times 10^{5}$	$1.9 \times 10^{3}$	0.038	0.010	
	$1.8 \times 10^{5}$	$7.3 \times 10^{3}$	0.033	$7.2 \times 10^{-3}$	
	$2.0 \times 10^{5}$	$3.6 \times 10^4$	0.012	$3.2 \times 10^{-3}$	
	$2.5 \times 10^5$	$4.9 \times 10^4$	0.013	$3.4 \times 10^{-3}$	
	$3.1 \times 10^5$ $4.2 \times 10^5$	$6.5 \times 10^4$ $1.5 \times 10^5$	$7.7 \times 10^{-3}$ $4.1 \times 10^{-3}$	$2.1 \times 10^{-3} \\ 1.2 \times 10^{-3}$	
	$6.3 \times 10^{5}$	$2.8 \times 10^{5}$	$3.5 \times 10^{-3}$	$9.7 \times 10^{-4}$	
	$1.0 \times 10^6$	$8.3 \times 10^4$	$1.8 \times 10^{-3}$	$8.5 \times 10^{-4}$	
060313	$4.2 \times 10^{3}$	161	1.3	0.52	This work
	$4.3 \times 10^{3}$	92	1.4	0.52	
	$4.4 \times 10^{3}$	97	1.7	0.62	
	$4.5 \times 10^{3}$	128	1.2	0.43	
	$4.7 \times 10^{3}$	113	1.2	0.45	
	$4.8 \times 10^3$	113	1.1	0.39	
	$4.9 \times 10^3$ $5.0 \times 10^3$	128 65	1.1 2.4	0.43 0.91	
	$5.0 \times 10^{3}$ $5.1 \times 10^{3}$	133	0.91	0.33	
	$5.1 \times 10^{3}$ $5.2 \times 10^{3}$	146	0.83	0.30	
	$5.4 \times 10^{3}$	176	0.64	0.23	
	$5.5 \times 10^{3}$	163	0.75	0.27	
	$5.7 \times 10^{3}$	95	1.4	0.54	
	$5.8 \times 10^{3}$	211	0.61	0.23	
	$6.0 \times 10^{3}$	125	1.2	0.46	
	$6.1 \times 10^3$	153	0.85	0.31	
	$6.3 \times 10^3$ $6.5 \times 10^3$	125 286	1.0	0.38 0.30	
	$0.3 \times 10^{4}$ $1.0 \times 10^{4}$	301	0.96 0.38	0.30	
	$1.0 \times 10^{4}$ $1.0 \times 10^{4}$	309	0.38	0.14	
	$1.1 \times 10^4$	228	0.66	0.25	
	$1.1 \times 10^{4}$	243	0.47	0.17	
	$1.1 \times 10^{4}$	266	0.44	0.16	
	$1.1 \times 10^{4}$	384	0.346	0.13	
	$1.2 \times 10^4$	344	0.36	0.13	
	$1.2 \times 10^4$	459	0.37	0.12	
	$1.6 \times 10^4$ $1.6 \times 10^4$	602 412	0.19	0.069	
	$1.6 \times 10^4$ $1.7 \times 10^4$	414	0.29 0.27	0.11 0.10	
	$1.7 \times 10^4$ $1.8 \times 10^4$	$1.1 \times 10^{3}$	0.25	0.078	
	$2.2 \times 10^4$	934	0.11	0.043	
	$2.3 \times 10^{4}$	545	0.20	0.075	
	$2.3 \times 10^{4}$	$1.1 \times 10^{3}$	0.13	0.047	
	$2.8 \times 10^{4}$	781	0.15	0.057	
	$2.9 \times 10^{4}$	$1.8 \times 10^{3}$	0.089	0.031	
	$3.4 \times 10^4$	$2.5 \times 10^3$	0.061	0.022	
	$4.2 \times 10^4$ $5.2 \times 10^4$	$6.3 \times 10^3$ $1.2 \times 10^4$	0.058	0.024	
	$3.2 \times 10^4$ $8.7 \times 10^4$	$5.9 \times 10^4$	0.036 0.013	$0.014$ $5.4 \times 10^{-3}$	
	$2.7 \times 10^{5}$	$3.9 \times 10^{5}$ $3.1 \times 10^{5}$	$3.6 \times 10^{-3}$	$1.2 \times 10^{-3}$	
060502B	$3.8 \times 10^4$	$6.6 \times 10^4$	$<1.0 \times 10^{-3}$	***	Evans et al. (2007a, 2009)
060801	918	954	0.20	0.077	This work
061006	$3.6 \times 10^{3}$	670	0.12	0.067	This work
	$4.4 \times 10^3$	997	0.11	0.060	
	$9.4 \times 10^3$	$4.9 \times 10^3$	0.046	0.026	
	$1.4 \times 10^4$ $6.6 \times 10^4$	$4.7 \times 10^3$ $2.4 \times 10^5$	$0.054$ $6.0 \times 10^{-3}$	$0.030$ $3.3 \times 10^{-3}$	
061201	$5.7 \times 10^{3}$	2.4 × 10 <sup>5</sup> 238	$0.0 \times 10^{-5}$	0.31	This work
001201	$5.7 \times 10^{3}$ $5.9 \times 10^{3}$	188	0.56	0.31	IIIIS WOIK
	$6.1 \times 10^3$	238	0.45	0.26	
	$6.4 \times 10^{3}$	264	0.66	0.34	
	$1.2 \times 10^{4}$	$5.9 \times 10^{3}$	0.11	0.072	
	$2.1 \times 10^{4}$	$1.2 \times 10^{4}$	0.055	0.036	
	$6.7 \times 10^4$	$1.3 \times 10^{5}$	$4.8 \times 10^{-3}$	$2.9 \times 10^{-3}$	
061210	$2.3 \times 10^5$	$5.3 \times 10^4$	0.048	0.013	Evans et al. (2007a, 2009)
	$3.1 \times 10^5$ $4.2 \times 10^5$	$9.4 \times 10^4$ $1.3 \times 10^5$	0.026 0.018	$8.0 \times 10^{-3}$ $6.0 \times 10^{-3}$	
	$4.2 \times 10^{5}$ $7.0 \times 10^{5}$	$4.9 \times 10^{5}$	< 0.018	0.0 × 10 °	
	/.U \ 1U	T.2 A 10	$< 2.5 \times 10^{-3}$		

Table A1 - Continued

GRB	$\frac{\delta t}{(\mathrm{s})}$	Exposure Time (s)	$F_X$ $(\mu Jy)$	$\sigma_X (\mu  ext{Jy})$	References
070209 <sup>b</sup>	$4.4 \times 10^{5}$	$1.9 \times 10^{4}$	$< 8.3 \times 10^{-4}$		C4444 -1 (2007h)
$070209^{b}$	$9.7 \times 10^4$	$1.9 \times 10^{4}$ $1.3 \times 10^{4}$	$< 8.3 \times 10$ $< 1.8 \times 10^{-3}$		Stratta et al. (2007b) Troja et al. (2007)
	$9.7 \times 10^{3}$ $1.6 \times 10^{3}$	$1.3 \times 10^{-1}$ $2.0 \times 10^{3}$		0.000	3 , , ,
070429B	$1.6 \times 10^{4}$ $1.6 \times 10^{4}$	$3.6 \times 10^4$	0.32 < 0.024	0.080	Evans et al. (2007a, 2009)
$070707^{a}$	$4.1 \times 10^4$	$2.2 \times 10^4$	0.024	0.032	Evans et al. (2007a, 2009)
0/0/0/	$4.1 \times 10^{5}$ $4.1 \times 10^{5}$	$1.9 \times 10^{5}$	< 0.032	0.032	Evans et al. (2007a, 2009)
070714B	$8.5 \times 10^{2}$	116	2.0	0.64	This work
070714B	$2.8 \times 10^3$	183	0.40	0.18	THIS WOLK
	$3.1 \times 10^3$	311	0.40	0.16	
	$3.1 \times 10^{3}$ $3.6 \times 10^{3}$	675	0.19	0.09	
	$5.9 \times 10^3$	417	0.15	0.075	
	$6.5 \times 10^3$	795	0.13	0.058	
	$9.2 \times 10^{3}$	$1.4 \times 10^{3}$	0.054	0.027	
	$2.9 \times 10^4$	$5.8 \times 10^4$	$3.0 \times 10^{-3}$	$1.7 \times 10^{-3}$	
070724A	$4.6 \times 10^{3}$	$4.8 \times 10^{3}$	0.055	0.032	This work
07072-71	$1.1 \times 10^4$	$7.6 \times 10^{3}$	0.026	0.014	This work
	$1.8 \times 10^4$	$5.2 \times 10^3$	0.024	0.013	
	$2.6 \times 10^4$	$1.2 \times 10^4$	0.014	$7.5 \times 10^{-3}$	
	$5.3 \times 10^4$	$5.5 \times 10^4$	0.014	$5.4 \times 10^{-3}$	
	$1.8 \times 10^{5}$	$3.0 \times 10^{5}$	$1.8 \times 10^{-3}$	$5.9 \times 10^{-4}$	
070729	$2.0 \times 10^4$	$6.6 \times 10^4$	$<3.8 \times 10^{-3}$		Evans et al. (2007a, 2009)
070809	$1.1 \times 10^{3}$	271	0.27	0.13	This work
0,000	$1.6 \times 10^{3}$	660	0.20	0.080	Time work
	$5.5 \times 10^{3}$	321	0.23	0.11	
	$5.9 \times 10^{3}$	507	0.14	0.068	
	$6.3 \times 10^{3}$	366	0.23	0.11	
	$6.8 \times 10^{3}$	555	0.13	0.062	
	$7.4 \times 10^{3}$	673	0.12	0.058	
	$1.1 \times 10^{4}$	673	0.10	0.047	
	$1.2 \times 10^{4}$	803	0.093	0.044	
	$1.3 \times 10^{4}$	989	0.085	0.039	
	$1.8 \times 10^{4}$	$2.6 \times 10^{3}$	0.038	0.017	
	$2.4 \times 10^{4}$	$2.0 \times 10^{3}$	0.050	0.021	
$070810B^{b}$	983	$7.2 \times 10^{3}$	$< 6.6 \times 10^{-4}$		Starling et al. (2007)
$071017^{b}$	$2.0 \times 10^{5}$	$2.1 \times 10^{3}$	$< 5.9 \times 10^{-3}$		Evans et al. (2007b)
$071112B^{a}$	$3.7 \times 10^{3}$	$2.2 \times 10^{3}$	< 0.023		Perri et al. (2007)
071227	$6.2 \times 10^{3}$	$9.5 \times 10^{3}$	0.022	$9.8 \times 10^{-3}$	This work
	$5.7 \times 10^4$	$2.5 \times 10^{5}$	$2.6 \times 10^{-3}$	$8.6 \times 10^{-4}$	
080121 <sup>a</sup>	$2.0 \times 10^{5}$	$2.2 \times 10^4$	$<4.0 \times 10^{-3}$		Troja et al. (2008)
081023	$1.4 \times 10^{3}$	$1.7 \times 10^{3}$	0.072	0.024	This work
	$1.8 \times 10^{4}$	$5.3 \times 10^{4}$	$9.5 \times 10^{-3}$	$3.2 \times 10^{-3}$	
080426	$1.1 \times 10^{3}$	100	2.13	0.48	Evans et al. (2007a, 2009)
080426	$1.2 \times 10^{3}$	100 148	1.4	0.32	Evans et al. (2007a, 2009)
080426	$1.2 \times 10^3$ $1.3 \times 10^3$	100 148 95	1.4 2.2	0.32 0.50	Evans et al. (2007a, 2009)
080426	$1.2 \times 10^{3}$ $1.3 \times 10^{3}$ $1.4 \times 10^{3}$	100 148 95 108	1.4 2.2 2.1	0.32 0.50 0.45	Evans et al. (2007a, 2009)
080426	$1.2 \times 10^{3}$ $1.3 \times 10^{3}$ $1.4 \times 10^{3}$ $1.5 \times 10^{3}$	100 148 95 108 125	1.4 2.2 2.1 1.7	0.32 0.50 0.45 0.38	Evans et al. (2007a, 2009)
080426	$1.2 \times 10^{3}$ $1.3 \times 10^{3}$ $1.4 \times 10^{3}$ $1.5 \times 10^{3}$ $1.7 \times 10^{3}$	100 148 95 108 125 211	1.4 2.2 2.1 1.7 1.01	0.32 0.50 0.45 0.38 0.23	Evans et al. (2007a, 2009)
080426	$1.2 \times 10^{3}$ $1.3 \times 10^{3}$ $1.4 \times 10^{3}$ $1.5 \times 10^{3}$ $1.7 \times 10^{3}$ $1.9 \times 10^{3}$	100 148 95 108 125 211 191	1.4 2.2 2.1 1.7 1.01 1.44	0.32 0.50 0.45 0.38 0.23 0.29	Evans et al. (2007a, 2009)
080426	$1.2 \times 10^{3}$ $1.3 \times 10^{3}$ $1.4 \times 10^{3}$ $1.5 \times 10^{3}$ $1.7 \times 10^{3}$ $1.9 \times 10^{3}$ $6.2 \times 10^{3}$	100 148 95 108 125 211 191 258	1.4 2.2 2.1 1.7 1.01 1.44 0.65	0.32 0.50 0.45 0.38 0.23 0.29 0.17	Evans et al. (2007a, 2009)
080426	$1.2 \times 10^{3}$ $1.3 \times 10^{3}$ $1.4 \times 10^{3}$ $1.5 \times 10^{3}$ $1.7 \times 10^{3}$ $1.9 \times 10^{3}$ $6.2 \times 10^{3}$ $6.6 \times 10^{3}$	100 148 95 108 125 211 191 258 687	1.4 2.2 2.1 1.7 1.01 1.44 0.65 0.24	0.32 0.50 0.45 0.38 0.23 0.29 0.17 0.063	Evans et al. (2007a, 2009)
080426	$1.2 \times 10^{3}$ $1.3 \times 10^{3}$ $1.4 \times 10^{3}$ $1.5 \times 10^{3}$ $1.7 \times 10^{3}$ $1.9 \times 10^{3}$ $6.2 \times 10^{3}$ $6.6 \times 10^{3}$ $7.5 \times 10^{3}$	100 148 95 108 125 211 191 258 687 978	1.4 2.2 2.1 1.7 1.01 1.44 0.65 0.24 0.19	0.32 0.50 0.45 0.38 0.23 0.29 0.17 0.063 0.047	Evans et al. (2007a, 2009)
080426	$1.2 \times 10^{3}$ $1.3 \times 10^{3}$ $1.4 \times 10^{3}$ $1.5 \times 10^{3}$ $1.7 \times 10^{3}$ $1.9 \times 10^{3}$ $6.2 \times 10^{3}$ $6.6 \times 10^{3}$ $7.5 \times 10^{3}$ $1.3 \times 10^{4}$	$   \begin{array}{c}     100 \\     148 \\     95 \\     108 \\     125 \\     211 \\     191 \\     258 \\     687 \\     978 \\     2.1 \times 10^{3}   \end{array} $	1.4 2.2 2.1 1.7 1.01 1.44 0.65 0.24 0.19 0.11	0.32 0.50 0.45 0.38 0.23 0.29 0.17 0.063 0.047 0.025	Evans et al. (2007a, 2009)
080426	$1.2 \times 10^{3}$ $1.3 \times 10^{3}$ $1.4 \times 10^{3}$ $1.5 \times 10^{3}$ $1.7 \times 10^{3}$ $1.9 \times 10^{3}$ $6.2 \times 10^{3}$ $6.6 \times 10^{3}$ $7.5 \times 10^{3}$ $1.3 \times 10^{4}$ $2.1 \times 10^{4}$	$   \begin{array}{c}     100 \\     148 \\     95 \\     108 \\     125 \\     211 \\     191 \\     258 \\     687 \\     978 \\     2.1 \times 10^3 \\     8.1 \times 10^3   \end{array} $	1.4 2.2 2.1 1.7 1.01 1.44 0.65 0.24 0.19 0.11	0.32 0.50 0.45 0.38 0.23 0.29 0.17 0.063 0.047 0.025 0.012	Evans et al. (2007a, 2009)
080426	$1.2 \times 10^{3}$ $1.3 \times 10^{3}$ $1.4 \times 10^{3}$ $1.5 \times 10^{3}$ $1.7 \times 10^{3}$ $1.9 \times 10^{3}$ $6.2 \times 10^{3}$ $6.6 \times 10^{3}$ $7.5 \times 10^{3}$ $1.3 \times 10^{4}$ $2.1 \times 10^{4}$ $3.8 \times 10^{4}$	$   \begin{array}{c}     100 \\     148 \\     95 \\     108 \\     125 \\     211 \\     191 \\     258 \\     687 \\     978 \\     2.1 \times 10^3 \\     8.1 \times 10^3 \\     2.0 \times 10^4   \end{array} $	1.4 2.2 2.1 1.7 1.01 1.44 0.65 0.24 0.19 0.11 0.056 0.021	$0.32$ $0.50$ $0.45$ $0.38$ $0.23$ $0.29$ $0.17$ $0.063$ $0.047$ $0.025$ $0.012$ $5.7 \times 10^{-3}$	Evans et al. (2007a, 2009)
	$\begin{array}{c} 1.2\times10^{3}\\ 1.3\times10^{3}\\ 1.4\times10^{3}\\ 1.5\times10^{3}\\ 1.7\times10^{3}\\ 1.9\times10^{3}\\ 6.2\times10^{3}\\ 6.6\times10^{3}\\ 7.5\times10^{3}\\ 1.3\times10^{4}\\ 2.1\times10^{4}\\ 3.8\times10^{4}\\ 7.0\times10^{4}\\ \end{array}$	$   \begin{array}{c}     100 \\     148 \\     95 \\     108 \\     125 \\     211 \\     191 \\     258 \\     687 \\     978 \\     2.1 \times 10^{3} \\     8.1 \times 10^{3} \\     2.0 \times 10^{4} \\     4.2 \times 10^{4}   \end{array} $	$\begin{array}{c} 1.4 \\ 2.2 \\ 2.1 \\ 1.7 \\ 1.01 \\ 1.44 \\ 0.65 \\ 0.24 \\ 0.19 \\ 0.11 \\ 0.056 \\ 0.021 \\ 4.9 \times 10^{-3} \end{array}$	$\begin{array}{c} 0.32 \\ 0.50 \\ 0.45 \\ 0.38 \\ 0.23 \\ 0.29 \\ 0.17 \\ 0.063 \\ 0.047 \\ 0.025 \\ 0.012 \\ 5.7 \times 10^{-3} \\ 2.8 \times 10^{-3} \end{array}$	
	$\begin{array}{c} 1.2 \times 10^{3} \\ 1.3 \times 10^{3} \\ 1.4 \times 10^{3} \\ 1.5 \times 10^{3} \\ 1.7 \times 10^{3} \\ 1.9 \times 10^{3} \\ 6.2 \times 10^{3} \\ 6.6 \times 10^{3} \\ 7.5 \times 10^{3} \\ 1.3 \times 10^{4} \\ 2.1 \times 10^{4} \\ 3.8 \times 10^{4} \\ 7.0 \times 10^{4} \\ 920 \end{array}$	$   \begin{array}{c}     100 \\     148 \\     95 \\     108 \\     125 \\     211 \\     191 \\     258 \\     687 \\     978 \\     2.1 \times 10^{3} \\     8.1 \times 10^{3} \\     2.0 \times 10^{4} \\     4.2 \times 10^{4} \\     1.1 \times 10^{3}   \end{array} $	$\begin{array}{c} 1.4 \\ 2.2 \\ 2.1 \\ 1.7 \\ 1.01 \\ 1.44 \\ 0.65 \\ 0.24 \\ 0.19 \\ 0.11 \\ 0.056 \\ 0.021 \\ 4.9 \times 10^{-3} \\ 0.45 \end{array}$	$\begin{array}{c} 0.32 \\ 0.50 \\ 0.45 \\ 0.38 \\ 0.23 \\ 0.29 \\ 0.17 \\ 0.063 \\ 0.047 \\ 0.025 \\ 0.012 \\ 5.7 \times 10^{-3} \\ 2.8 \times 10^{-3} \\ 0.18 \end{array}$	This work
080426 080503	$1.2 \times 10^{3}$ $1.3 \times 10^{3}$ $1.4 \times 10^{3}$ $1.5 \times 10^{3}$ $1.7 \times 10^{3}$ $1.9 \times 10^{3}$ $6.2 \times 10^{3}$ $6.5 \times 10^{3}$ $7.5 \times 10^{3}$ $1.3 \times 10^{4}$ $2.1 \times 10^{4}$ $3.8 \times 10^{4}$ $7.0 \times 10^{4}$ $920$ $+d^{2}3.9 \times 10^{5}$	$   \begin{array}{c}     100 \\     148 \\     95 \\     108 \\     125 \\     211 \\     191 \\     258 \\     687 \\     978 \\     2.1 \times 10^3 \\     8.1 \times 10^3 \\     2.0 \times 10^4 \\     4.2 \times 10^4 \\     1.1 \times 10^3 \\     3.2 \times 10^4 \\   \end{array} $	$\begin{array}{c} 1.4 \\ 2.2 \\ 2.1 \\ 1.7 \\ 1.01 \\ 1.44 \\ 0.65 \\ 0.24 \\ 0.19 \\ 0.11 \\ 0.056 \\ 0.021 \\ 4.9 \times 10^{-3} \\ 0.45 \\ 2.1 \times 10^{-3} \end{array}$	$\begin{array}{c} 0.32 \\ 0.50 \\ 0.45 \\ 0.38 \\ 0.23 \\ 0.29 \\ 0.17 \\ 0.063 \\ 0.047 \\ 0.025 \\ 0.012 \\ 5.7 \times 10^{-3} \\ 2.8 \times 10^{-3} \end{array}$	This work Perley et al. (2009b)
080503	$\begin{array}{c} 1.2\times10^{3}\\ 1.3\times10^{3}\\ 1.4\times10^{3}\\ 1.5\times10^{3}\\ 1.7\times10^{3}\\ 1.9\times10^{3}\\ 6.2\times10^{3}\\ 6.6\times10^{3}\\ 7.5\times10^{3}\\ 1.3\times10^{4}\\ 2.1\times10^{4}\\ 3.8\times10^{4}\\ 7.0\times10^{4}\\ 920\\ ^{+d}3.9\times10^{5}\\ ^{+1.9\times10^{6}} \end{array}$	$\begin{array}{c} 100 \\ 148 \\ 95 \\ 108 \\ 125 \\ 211 \\ 191 \\ 258 \\ 687 \\ 978 \\ 2.1 \times 10^3 \\ 8.1 \times 10^3 \\ 2.0 \times 10^4 \\ 4.2 \times 10^4 \\ 1.1 \times 10^3 \\ 3.2 \times 10^4 \\ 3.3 \times 10^4 \\ \end{array}$	$\begin{array}{c} 1.4 \\ 2.2 \\ 2.1 \\ 1.7 \\ 1.01 \\ 1.44 \\ 0.65 \\ 0.24 \\ 0.19 \\ 0.11 \\ 0.056 \\ 0.021 \\ 4.9 \times 10^{-3} \\ 0.45 \\ 2.1 \times 10^{-3} \\ < 8.0 \times 10^{-5} \\ \end{array}$	$\begin{array}{c} 0.32 \\ 0.50 \\ 0.45 \\ 0.38 \\ 0.23 \\ 0.29 \\ 0.17 \\ 0.063 \\ 0.047 \\ 0.025 \\ 0.012 \\ 5.7 \times 10^{-3} \\ 2.8 \times 10^{-3} \\ 0.18 \\ 9.6 \times 10^{-4} \end{array}$	This work Perley et al. (2009b) Perley et al. (2009b)
	$\begin{array}{c} 1.2\times10^{3}\\ 1.3\times10^{3}\\ 1.4\times10^{3}\\ 1.5\times10^{3}\\ 1.7\times10^{3}\\ 1.9\times10^{3}\\ 6.2\times10^{3}\\ 6.6\times10^{3}\\ 7.5\times10^{3}\\ 1.3\times10^{4}\\ 2.1\times10^{4}\\ 3.8\times10^{4}\\ 7.0\times10^{4}\\ 920\\ ^{+d}3.9\times10^{5}\\ ^{+1.9\times10^{6}}\\ 3.9\times10^{3}\\ \end{array}$	$\begin{array}{c} 100 \\ 148 \\ 95 \\ 108 \\ 125 \\ 211 \\ 191 \\ 258 \\ 687 \\ 978 \\ 2.1 \times 10^3 \\ 8.1 \times 10^3 \\ 2.0 \times 10^4 \\ 4.2 \times 10^4 \\ 1.1 \times 10^3 \\ 3.2 \times 10^4 \\ 3.3 \times 10^4 \\ 2.2 \times 10^4 \end{array}$	$\begin{array}{c} 1.4 \\ 2.2 \\ 2.1 \\ 1.7 \\ 1.01 \\ 1.44 \\ 0.65 \\ 0.24 \\ 0.19 \\ 0.11 \\ 0.056 \\ 0.021 \\ 4.9 \times 10^{-3} \\ 0.45 \\ 2.1 \times 10^{-3} \\ < 8.0 \times 10^{-5} \\ 0.041 \\ \end{array}$	$\begin{array}{c} 0.32 \\ 0.50 \\ 0.45 \\ 0.38 \\ 0.23 \\ 0.29 \\ 0.17 \\ 0.063 \\ 0.047 \\ 0.025 \\ 0.012 \\ 5.7 \times 10^{-3} \\ 2.8 \times 10^{-3} \\ 0.18 \end{array}$	This work Perley et al. (2009b)
080503 080702A	$1.2 \times 10^{3}$ $1.3 \times 10^{3}$ $1.4 \times 10^{3}$ $1.5 \times 10^{3}$ $1.7 \times 10^{3}$ $1.9 \times 10^{3}$ $6.2 \times 10^{3}$ $6.6 \times 10^{3}$ $7.5 \times 10^{3}$ $1.3 \times 10^{4}$ $2.1 \times 10^{4}$ $920$ $1.3 \times 10^{5}$ $1.9 \times 10^{6}$ $3.9 \times 10^{3}$ $7.5 \times 10^{4}$	$\begin{array}{c} 100 \\ 148 \\ 95 \\ 108 \\ 125 \\ 211 \\ 191 \\ 258 \\ 687 \\ 978 \\ 2.1 \times 10^3 \\ 8.1 \times 10^3 \\ 2.0 \times 10^4 \\ 4.2 \times 10^4 \\ 1.1 \times 10^3 \\ 3.2 \times 10^4 \\ 3.3 \times 10^4 \\ 2.2 \times 10^4 \\ 7.6 \times 10^4 \\ \end{array}$	$\begin{array}{c} 1.4 \\ 2.2 \\ 2.1 \\ 1.7 \\ 1.01 \\ 1.44 \\ 0.65 \\ 0.24 \\ 0.19 \\ 0.11 \\ 0.056 \\ 0.021 \\ 4.9 \times 10^{-3} \\ 0.45 \\ 2.1 \times 10^{-3} \\ < 8.0 \times 10^{-5} \\ 0.041 \\ < 0.013 \\ \end{array}$	$0.32 \\ 0.50 \\ 0.45 \\ 0.38 \\ 0.23 \\ 0.29 \\ 0.17 \\ 0.063 \\ 0.047 \\ 0.025 \\ 0.012 \\ 5.7 \times 10^{-3} \\ 2.8 \times 10^{-3} \\ 0.18 \\ 9.6 \times 10^{-4} \\ 8.9 \times 10^{-3}$	This work Perley et al. (2009b) Perley et al. (2009b) Evans et al. (2007a, 2009)
080503 080702A 080905A	$1.2 \times 10^{3}$ $1.3 \times 10^{3}$ $1.4 \times 10^{3}$ $1.5 \times 10^{3}$ $1.7 \times 10^{3}$ $1.9 \times 10^{3}$ $6.2 \times 10^{3}$ $6.6 \times 10^{3}$ $7.5 \times 10^{3}$ $1.3 \times 10^{4}$ $2.1 \times 10^{4}$ $3.8 \times 10^{4}$ $920$ $+^{4} 3.9 \times 10^{5}$ $+1.9 \times 10^{6}$ $3.9 \times 10^{3}$ $7.5 \times 10^{4}$ $758$	$\begin{array}{c} 100 \\ 148 \\ 95 \\ 108 \\ 125 \\ 211 \\ 191 \\ 258 \\ 687 \\ 978 \\ 2.1 \times 10^3 \\ 8.1 \times 10^3 \\ 2.0 \times 10^4 \\ 4.2 \times 10^4 \\ 4.2 \times 10^4 \\ 3.3 \times 10^4 \\ 2.2 \times 10^4 \\ 7.6 \times 10^4 \\ 593 \\ \end{array}$	$\begin{array}{c} 1.4 \\ 2.2 \\ 2.1 \\ 1.7 \\ 1.01 \\ 1.44 \\ 0.65 \\ 0.24 \\ 0.19 \\ 0.11 \\ 0.056 \\ 0.021 \\ 4.9 \times 10^{-3} \\ 0.45 \\ 2.1 \times 10^{-3} \\ < 8.0 \times 10^{-5} \\ 0.041 \\ < 0.013 \\ 0.74 \\ \end{array}$	0.32 0.50 0.45 0.38 0.23 0.29 0.17 0.063 0.047 0.025 0.012 $5.7 \times 10^{-3}$ $2.8 \times 10^{-3}$ 0.18 $9.6 \times 10^{-4}$ $8.9 \times 10^{-3}$ 0.31	This work Perley et al. (2009b) Perley et al. (2009b) Evans et al. (2007a, 2009) This work
080503	$1.2 \times 10^{3}$ $1.3 \times 10^{3}$ $1.4 \times 10^{3}$ $1.5 \times 10^{3}$ $1.7 \times 10^{3}$ $1.9 \times 10^{3}$ $6.2 \times 10^{3}$ $6.6 \times 10^{3}$ $7.5 \times 10^{3}$ $1.3 \times 10^{4}$ $2.1 \times 10^{4}$ $920$ $1.3 \times 10^{5}$ $1.9 \times 10^{6}$ $3.9 \times 10^{3}$ $7.5 \times 10^{4}$	$\begin{array}{c} 100 \\ 148 \\ 95 \\ 108 \\ 125 \\ 211 \\ 191 \\ 258 \\ 687 \\ 978 \\ 2.1 \times 10^3 \\ 8.1 \times 10^3 \\ 2.0 \times 10^4 \\ 4.2 \times 10^4 \\ 1.1 \times 10^3 \\ 3.2 \times 10^4 \\ 3.3 \times 10^4 \\ 2.2 \times 10^4 \\ 7.6 \times 10^4 \\ \end{array}$	$\begin{array}{c} 1.4 \\ 2.2 \\ 2.1 \\ 1.7 \\ 1.01 \\ 1.44 \\ 0.65 \\ 0.24 \\ 0.19 \\ 0.11 \\ 0.056 \\ 0.021 \\ 4.9 \times 10^{-3} \\ 0.45 \\ 2.1 \times 10^{-3} \\ < 8.0 \times 10^{-5} \\ 0.041 \\ < 0.013 \\ \end{array}$	$0.32 \\ 0.50 \\ 0.45 \\ 0.38 \\ 0.23 \\ 0.29 \\ 0.17 \\ 0.063 \\ 0.047 \\ 0.025 \\ 0.012 \\ 5.7 \times 10^{-3} \\ 2.8 \times 10^{-3} \\ 0.18 \\ 9.6 \times 10^{-4} \\ 8.9 \times 10^{-3}$	This work Perley et al. (2009b) Perley et al. (2009b) Evans et al. (2007a, 2009)

Table A1 - Continued

GRB	δ <i>t</i> (s)	Exposure Time (s)	$F_X$ ( $\mu$ Jy)	$\sigma_X \ (\mu Jy)$	References
	1.5 ×10 <sup>4</sup>	$3.7 \times 10^4$	<3.2 ×10 <sup>-3</sup>		
081226A 081226B <sup>a</sup>	$1.6 \times 10^4$ $1.3 \times 10^4$	$4.3 \times 10^4$ $8.0 \times 10^3$	$0.026 < 6.6 \times 10^{-3}$	0.014	Evans et al. (2007a, 2009) Evans & Hoversten (2008)
090305A <sup>b</sup>	$4.8 \times 10^{3}$	1.3 ×10 <sup>4</sup>	$<2.0 \times 10^{-3}$		Beardmore et al. (2009)
090426	$1.0 \times 10^{3}$	95	1.2	0.26	Evans et al. (2007a, 2009)
	$1.1 \times 10^{3}$	108	1.0	0.23	
	$1.2 \times 10^3$ $1.4 \times 10^3$	95 145	1.2	0.26	
	$1.4 \times 10^{3}$ $1.5 \times 10^{3}$	145 92	0.77 1.2	0.17 0.27	
	$1.6 \times 10^3$	100	1.1	0.25	
	$1.7 \times 10^{3}$	178	0.62	0.14	
	$1.9 \times 10^{3}$	243	0.45	0.10	
	$2.2 \times 10^{3}$	221	0.68	0.13	
	$5.6 \times 10^3$ $6.0 \times 10^3$	346 351	0.27 0.27	0.072 0.071	
	$6.0 \times 10^{3}$ $6.4 \times 10^{3}$	481	0.27	0.071	
	$6.9 \times 10^3$	479	0.20	0.051	
	$7.6 \times 10^{3}$	938	0.18	0.035	
	$1.8 \times 10^{4}$	$1.2 \times 10^4$	0.060	0.013	
000510	$2.3 \times 10^5$	$7.5 \times 10^{5}$	$1.5 \times 10^{-3}$	$6.5 \times 10^{-4}$	
090510	$1.0 \times 10^3$ $2.8 \times 10^3$	31 89	2.7 0.61	1.0 0.26	This work
	$2.8 \times 10^{3}$ $2.9 \times 10^{3}$	89 141	0.47	0.26	
	$3.1 \times 10^3$	174	0.37	0.17	
	$3.3 \times 10^{3}$	338	0.18	0.081	
	$3.6 \times 10^{3}$	251	0.22	0.10	
	$3.9 \times 10^3$	321	0.18	0.084	
	$6.2 \times 10^3$ $6.8 \times 10^3$	770 547	0.051 0.073	0.027 0.039	
	$1.3 \times 10^4$	$9.2 \times 10^3$	0.073	0.039	
	$4.1 \times 10^4$	$7.6 \times 10^4$	$4.0 \times 10^{-3}$	$1.3 \times 10^{-3}$	
090515	$8.1 \times 10^{2}$	$2.2 \times 10^{3}$	0.24	0.093	This work
090607	$9.3 \times 10^{2}$	$2.0 \times 10^{3}$	0.060	0.017	Evans et al. (2007a, 2009)
090621B	$2.0 \times 10^4$ $7.2 \times 10^3$	$2.4 \times 10^4$ $8.3 \times 10^3$	$< 6.6 \times 10^{-3}$ $0.04$	0.01	Evans et al. (2007a, 2009)
090021 <b>D</b>	$2.5 \times 10^4$	$2.5 \times 10^4$	$6.0 \times 10^{-3}$	$3.4 \times 10^{-3}$	Evalis et al. (2007a, 2009)
$090916^{b}$	$6.5 \times 10^4$	$1.2 \times 10^4$	$< 9.4 \times 10^{-4}$		Troja et al. (2009)
091109B	$5.7 \times 10^{3}$	$2.4 \times 10^{3}$	0.097	0.019	Evans et al. (2007a, 2009)
	$1.6 \times 10^4$	$1.3 \times 10^4$	0.030	0.023	
	$5.8 \times 10^4$ $2.0 \times 10^5$	$5.8 \times 10^4$ $1.2 \times 10^5$	0.011 0.011	$7.9 \times 10^{-3}$ $9.6 \times 10^{-3}$	
	2.0 × 10	1.2 × 10	0.011	7.0 × 10	
100117A	$3.1 \times 10^4$	$3.6 \times 10^{5}$	$2.2 \times 10^{-3}$	$7.2 \times 10^{-4}$	This work
100206A	$2.4 \times 10^4$	$9.5 \times 10^4$	$<3.3 \times 10^{-3}$		Evans et al. (2007a, 2009)
100213 100625A	$2.0 \times 10^4$ $1.6 \times 10^4$	$3.1 \times 10^4$ $1.3 \times 10^4$	$< 0.033$ $4.4 \times 10^{-3}$	$1.8 \times 10^{-3}$	Evans et al. (2007a, 2009) Fong et al. (2013)
10002JA	$9.0 \times 10^4$	$4.6 \times 10^4$	$< 8.2 \times 10^{-3}$	1.0 ^ 10	Fong et al. (2013)
100628A <sup>a</sup>	$^{+}3.9 \times 10^{5}$	$2.0 \times 10^{4}$	$< 5.9 \times 10^4$		This work
100702A	$1.4 \times 10^4$	$4.3 \times 10^{4}$	< 0.011		Evans et al. (2007a, 2009)
101219A	$7.7 \times 10^2$	317	0.58	0.20	Fong et al. (2013)
	$8.6 \times 10^3$ $^{+}3.4 \times 10^5$	$1.4 \times 10^4$ $2.0 \times 10^5$	$0.038 < 1.9 \times 10^{-4}$	0.014	
101224A	$1.9 \times 10^3$	$4.0 \times 10^4$	0.015	$5.0 \times 10^{-3}$	Evans et al. (2007a, 2009)
110112A	$4.6 \times 10^{3}$	1.0 ×10 <sup>3</sup>	0.083	0.028	Fong et al. (2013)
	$5.4 \times 10^{3}$	605	0.14	0.047	
	$6.2 \times 10^{3}$	941	0.14	0.043	
	$1.1 \times 10^4$	$2.5 \times 10^3$	0.064	0.016	
	$1.9 \times 10^4$	$6.6 \times 10^3$	0.025	$8.8 \times 10^{-3}$	
	$2.8 \times 10^4$ $4.5 \times 10^4$	$1.3 \times 10^4$ $2.3 \times 10^4$	$0.012$ $7.3 \times 10^{-3}$	$4.3 \times 10^{-3}$ $3.1 \times 10^{-3}$	
	$1.4 \times 10^5$	$2.3 \times 10^{5}$ $2.7 \times 10^{5}$	$2.0 \times 10^{-3}$	$1.0 \times 10^{-3}$	
$110112B^b$	$6.1 \times 10^{4}$	$1.2 \times 10^4$	$<2.4 \times 10^{-3}$		Littlejohns et al. (2011)
111020A	$1.1 \times 10^{3}$		1.0	0.20	Fong et al. (2012b)

Table A1 - Continued

Table A1 — Continuea								
GRB	δ <i>t</i> (s)	Exposure Time (s)	$F_X \ (\mu J y)$	$\sigma_{X} \ (\mu  m Jy)$	References			
	$5.8 \times 10^{3}$	$2.5 \times 10^3$	0.19	0.039				
	$1.2 \times 10^{4}$	$2.5 \times 10^{3}$	0.13	0.032				
	$1.9 \times 10^4$	$6.4 \times 10^3$	0.11	0.027				
	$2.6 \times 10^4$	$6.3 \times 10^3$	0.14	0.037				
	$3.2 \times 10^4$ $4.2 \times 10^4$	$5.9 \times 10^3$ $1.6 \times 10^4$	0.12 0.10	0.031 0.028				
	$4.2 \times 10^{5}$ $1.0 \times 10^{5}$	$1.6 \times 10^{-5}$ $1.5 \times 10^{5}$	0.10	$6.4 \times 10^{-3}$				
	$2.9 \times 10^{5}$	$2.1 \times 10^{5}$	0.019	$4.9 \times 10^{-3}$				
	$^{++}6.9 \times 10^4$	$1.4 \times 10^4$	0.013	$2.4 \times 10^{-3}$				
	$^{+}2.6 \times 10^{5}$	$2.0 \times 10^{4}$	$7.0 \times 10^{-3}$	$1.0 \times 10^{-3}$				
	$5.1 \times 10^{5}$	$2.1 \times 10^{5}$	< 0.019					
	$^{+}8.8 \times 10^{5}$	$2.0 \times 10^{4}$	$< 1.1 \times 10^{-3}$					
111117A	$1.0 \times 10^{3}$	178	0.55	0.20	Margutti et al. (2012)			
	$1.2 \times 10^{3}$	211	0.45	0.15				
	$5.0 \times 10^{3}$	881	0.11	0.035				
	$5.5 \times 10^4$	$1.9 \times 10^{5}$	$4.0 \times 10^{-3}$	$1.4 \times 10^{-3}$				
111101 4	$^{+}2.6 \times 10^{5}$	$4.0 \times 10^4$	$4.2 \times 10^{-4}$	$1.6 \times 10^{-4}$	F (1/2007 2000)			
111121A	$4.2 \times 10^3$ $4.3 \times 10^3$	85 133	2.17	0.47 0.30	Evans et al. (2007a, 2009)			
	$4.3 \times 10^{3}$ $4.5 \times 10^{3}$	143	1.3 1.4	0.30				
	$4.3 \times 10^{3}$ $4.7 \times 10^{3}$	228	1.4	0.26				
	$4.9 \times 10^{3}$	113	1.6	0.37				
	$5.1 \times 10^{3}$	196	1.1	0.22				
	$5.3 \times 10^{3}$	246	0.93	0.21				
	$5.6 \times 10^{3}$	326	1.0	0.20				
	$6.0 \times 10^{3}$	133	1.3	0.30				
	$6.2 \times 10^{3}$	143	1.2	0.28				
	$6.3 \times 10^{3}$	160	1.1	0.25				
	$6.5 \times 10^3$	158	1.1	0.25				
	$6.7 \times 10^3$	206	1.1	0.21				
	$1.0 \times 10^4$ $1.0 \times 10^4$	155	1.2 1.0	0.26 0.23				
	$1.0 \times 10^{4}$ $1.1 \times 10^{4}$	183 384	0.34	0.23				
	$1.1 \times 10^4$ $1.1 \times 10^4$	188	0.94	0.090				
	$1.1 \times 10^4$ $1.1 \times 10^4$	181	0.74	0.19				
	$1.1 \times 10^{4}$	243	0.54	0.14				
	$1.1 \times 10^{4}$	223	0.60	0.16				
	$1.2 \times 10^{4}$	268	0.50	0.13				
	$1.2 \times 10^{4}$	657	0.37	0.072				
	$1.6 \times 10^{4}$	985	0.24	0.063				
	$1.8 \times 10^{4}$	$1.6 \times 10^{3}$	0.17	0.043				
	$2.2 \times 10^4$	667	0.22	0.057				
	$2.3 \times 10^4$	$1.8 \times 10^3$	0.12	0.026				
	$4.2 \times 10^4$	$3.7 \times 10^4$	0.045	$8.3 \times 10^{-3}$				
111222A	$1.4 \times 10^5$ $9.7 \times 10^4$	$6.5 \times 10^4$ $7.3 \times 10^3$	$< 0.020 \\ 0.047$	$9.5 \times 10^{-3}$	Evans et al. (2007a, 2009)			
1112221	$7.8 \times 10^{5}$	$5.2 \times 10^4$	< 0.027	<i>9.5</i> ∧ 10	Evalis et al. (2007a, 2009)			
	$4.0 \times 10^6$	$1.7 \times 10^5$	< 0.014					
120305A	2.3 ×10 <sup>4</sup>	9.4 ×10 <sup>4</sup>	0.016	$8.0 \times 10^{-3}$	Evans et al. (2007a, 2009)			
120521A	$6.8 \times 10^{3}$	$7.3 \times 10^{3}$	< 0.020	Evans et al. (2007a, 2009)	, , , , , , , , , , , , , , , , , , , ,			
120630A	$6.6 \times 10^{2}$	$1.2 \times 10^{3}$	0.11	0.024	Evans et al. (2007a, 2009)			
120804A	$4.3 \times 10^{3}$	203	1.1	0.23	Berger et al. (2013)			
	$4.5 \times 10^{3}$	211	0.82	0.20				
	$4.7 \times 10^{3}$	178	0.85	0.22				
	$4.9 \times 10^3$	181	1.1	0.26				
	$5.2 \times 10^3$	281	0.54	0.14				
	$5.4 \times 10^3$ $5.7 \times 10^3$	263	0.58	0.15				
	$5.7 \times 10^{3}$ $6.0 \times 10^{3}$	301 349	0.50 0.43	0.13				
	$6.0 \times 10^{3}$ $6.4 \times 10^{3}$	349 494	0.43	0.12 0.090				
	$0.4 \times 10^{4}$ $1.1 \times 10^{4}$	$2.6 \times 10^{3}$	0.34	0.056				
	$1.7 \times 10^4$	$2.5 \times 10^{3}$	0.20	0.051				
	$2.3 \times 10^4$	$2.6 \times 10^{3}$	0.11	0.021				
	$3.0 \times 10^{4}$	$5.7 \times 10^{3}$	0.096	0.026				
	$3.4 \times 10^{4}$	$2.6 \times 10^{3}$	0.10	0.021				

Table A1 - Continued

GRB	δ <i>t</i> (s)	Exposure Time (s)	<i>F</i> <sub>X</sub> (μJy)	$\sigma_X \ (\mu  ext{Jy})$	References
	$4.3 \times 10^{4}$	$8.2 \times 10^{3}$	0.049	0.013	
	$6.2 \times 10^4$ $1.4 \times 10^5$	$2.3 \times 10^4$ $9.3 \times 10^4$	0.076 0.048	0.016 0.014	
	$3.1 \times 10^5$	$2.8 \times 10^{5}$	$6.8 \times 10^{-3}$	$2.2 \times 10^{-3}$	
	$5.0 \times 10^{5}$	$2.3 \times 10^{4}$	< 0.0456		
	$+8.1 \times 10^{5}$	$2.0 \times 10^{4}$	$3.5 \times 10^{-3}$	$6.2 \times 10^{-4}$	
	$^{++}1.6 \times 10^6$	$2.5 \times 10^4$	$3.0 \times 10^{-3}$	$6.2 \times 10^{-4}$	This are also
$120817B^{b}$	$^{+}4.0 \times 10^{6}$ $6.8 \times 10^{4}$	$5.9 \times 10^4$ $2.8 \times 10^3$	$6.6 \times 10^{-4}$ < $5.9 \times 10^{-3}$	$1.3 \times 10^{-3}$	This work Pagani (2012)
121226A	$4.3 \times 10^{3}$	589	0.58	0.15	Evans et al. (2007a, 2009)
	$4.9 \times 10^{3}$	542	0.62	0.16	
	$5.5 \times 10^3$	679	0.49	0.13	
	$7.9 \times 10^3$ $6.2 \times 10^4$	$4.7 \times 10^3$ $8.3 \times 10^4$	0.36 0.030	$0.079$ $8.6 \times 10^{-3}$	
	$1.2 \times 10^5$	$6.2 \times 10^3$	0.034	0.027	
130313A <sup>a</sup>	2.1 ×10 <sup>3</sup>	5.3 ×10 <sup>4</sup>	< 0.013		Evans et al. (2007a, 2009)
130515A	$1.2 \times 10^4$	$3.0 \times 10^4$	< 0.014	1.2	Evans et al. (2007a, 2009)
130603B	$4.0 \times 10^3$ $4.0 \times 10^3$	33 41	3.8 3.3	1.2 1.0	Fong et al. (2014)
	$4.0 \times 10^{3}$ $4.1 \times 10^{3}$	75	3.3 1.7	0.54	
	$4.1 \times 10^{3}$	44	2.9	0.95	
	$4.2 \times 10^{3}$	44	3.0	0.91	
	$4.2 \times 10^3$ $4.3 \times 10^3$	36 75	3.5	1.2	
	$4.3 \times 10^{3}$ $4.3 \times 10^{3}$	75 44	1.8 3.0	0.53 1.0	
	$4.4 \times 10^{3}$	54	2.5	0.71	
	$4.4 \times 10^{3}$	41	3.2	1.1	
	$4.5 \times 10^3$ $4.5 \times 10^3$	65 31	2.0	0.63 1.3	
	$4.6 \times 10^3$	44	4.1 3.0	0.99	
	$4.6 \times 10^{3}$	52	2.4	0.80	
	$4.7 \times 10^3$	65	2.0	0.68	
	$4.7 \times 10^3$ $4.8 \times 10^3$	52 98	2.6 1.6	0.74 0.46	
	$4.8 \times 10^{3}$ $4.9 \times 10^{3}$	57	2.2	0.72	
	$4.9 \times 10^{3}$	44	3.0	0.98	
	$5.0 \times 10^{3}$	59	2.3	0.77	
	$5.1 \times 10^3$ $5.1 \times 10^3$	83 59	1.5 2.5	0.43 0.71	
	$5.1 \times 10^{3}$ $5.2 \times 10^{3}$	93	1.4	0.43	
	$5.3 \times 10^{3}$	96	1.4	0.45	
	$5.4 \times 10^{3}$	88	1.6	0.49	
	$5.5 \times 10^3$ $5.5 \times 10^3$	72 41	1.8	0.59 0.98	
	$5.6 \times 10^3$	67	3.0 1.9	0.59	
	$5.7 \times 10^{3}$	85	1.5	0.44	
	$5.8 \times 10^3$	78	1.7	0.50	
	$5.8 \times 10^3$ $5.9 \times 10^3$	65 75	2.0 1.6	0.59 0.54	
	$6.0 \times 10^3$	67	1.9	0.56	
	$6.0 \times 10^{3}$	91	1.4	0.42	
	$6.2 \times 10^3$	114	1.1	0.34	
	$6.2 \times 10^3$ $6.3 \times 10^3$	65 59	2.0 2.1	0.59 0.59	
	$6.4 \times 10^3$	91	1.4	0.42	
	$6.5 \times 10^{3}$	122	1.5	0.40	
	$9.8 \times 10^3$	91	1.3	0.36	
	$9.9 \times 10^3$ $1.0 \times 10^4$	143 166	0.83 0.71	0.23 0.19	
	$1.0 \times 10^{4}$ $1.1 \times 10^{4}$	$1.5 \times 10^3$	0.71	0.19	
	$1.2 \times 10^{4}$	166	0.65	0.19	
	$1.2 \times 10^4$	226	0.49	0.13	
	$1.2 \times 10^4$ $1.6 \times 10^4$	205 338	0.72 0.29	0.17 0.076	
	$1.6 \times 10^4$	252	0.38	0.11	

Table A1 — Continued

GRB	$\frac{\delta t}{(s)}$	Exposure Time (s)	$F_X \ (\mu J y)$	$\sigma_X \ (\mu  ext{Jy})$	References
	$1.7 \times 10^{4}$	$2.0 \times 10^{3}$	0.21	0.056	
	$2.1 \times 10^{4}$	450	0.19	0.048	
	$2.2 \times 10^{4}$	463	0.19	0.047	
	$2.8 \times 10^{4}$	$1.1 \times 10^{3}$	0.077	0.029	
	$3.6 \times 10^{4}$	$5.9 \times 10^{3}$	0.038	0.011	
	$4.2 \times 10^4$	$5.5 \times 10^{3}$	0.054	0.014	
	$4.8 \times 10^{4}$	$6.7 \times 10^{3}$	0.021	$5.6 \times 10^{-3}$	
	$1.5 \times 10^{5}$	$2.1 \times 10^{5}$	$4.0 \times 10^{-3}$	$8.8 \times 10^{-4}$	
	$^{++}2.3 \times 10^5$	$1.9 \times 10^4$	$2.4 \times 10^{-3}$	$3.2 \times 10^{-4}$	
h	$^{++}5.6 \times 10^{5}$	$3.0 \times 10^4$	$8.2 \times 10^{-4}$	$5.0 \times 10^{-4}$	
130626A <sup>b</sup>	$8.7 \times 10^{5}$	$7.2 \times 10^5$	$<1.8 \times 10^{-3}$	10 10-3	Page & de Pasquale (2013)
130716A	$1.2 \times 10^4$	$2.5 \times 10^4$	0.014	$4.0 \times 10^{-3}$	Evans et al. (2007a, 2009)
1200221	$4.0 \times 10^4$	$1.3 \times 10^4$	$< 7.2 \times 10^{-3}$		F 1 (2007 2000)
130822A	$3.0 \times 10^4$	$1.9 \times 10^4$	$<4.6 \times 10^{-3}$	1.2	Evans et al. (2007a, 2009)
130912A	$9.7 \times 10^2$	95 103	5.6	1.3	Evans et al. (2007a, 2009)
	$1.1 \times 10^3$	103	2.7	0.60	
	$1.2 \times 10^3$	100	1.3	0.30	
	$1.3 \times 10^3$	105	1.3	0.29	
	$1.4 \times 10^3$ $1.5 \times 10^3$	95 180	1.4 1.6	0.31	
	$1.5 \times 10^3$ $5.2 \times 10^3$	180 780	0.26	0.26	
	$5.2 \times 10^3$ $6.4 \times 10^3$	$1.8 \times 10^{3}$	0.26	0.068 0.04	
	$1.2 \times 10^4$	$2.5 \times 10^3$	0.066	0.04	
	$1.2 \times 10^{4}$ $1.8 \times 10^{4}$	$2.5 \times 10^{3}$ $2.5 \times 10^{3}$	0.057	0.015	
	$2.8 \times 10^4$	$2.3 \times 10^{4}$ $1.4 \times 10^{4}$	0.037	$4.9 \times 10^{-3}$	
	$1.1 \times 10^5$	$5.8 \times 10^4$	$< 3.9 \times 10^{-3}$	4.9 × 10	
131004A	$1.1 \times 10^{2}$ $1.8 \times 10^{2}$	306	13.2	3.1	Evans et al. (2007a, 2009)
131004A	$4.4 \times 10^2$	115	34.8	8.1	Evalis et al. (2007a, 2009)
	$5.5 \times 10^2$	115	33.9	8.1	
	$6.9 \times 10^2$	158	24.8	5.9	
	$8.8 \times 10^2$	228	31.9	5.5	
	$1.1 \times 10^4$	$1.5 \times 10^3$	0.23	0.049	
	$2.4 \times 10^4$	$1.8 \times 10^4$	0.055	0.015	
	$5.9 \times 10^4$	$4.7 \times 10^4$	0.050	0.013	
	$1.1 \times 10^5$	$4.1 \times 10^4$	< 0.017	0.014	
131224A <sup>b</sup>	$1.3 \times 10^4$	$4.0 \times 10^{3}$	$<4.1 \times 10^{-3}$		Gompertz et al. (2013)
1.40120D	0.0 × 102	(5	2.01	0.60	F
140129B	$9.9 \times 10^2$ $1.1 \times 10^3$	65 45	3.01	0.68	Evans et al. (2007a, 2009)
		45	4.62	1.01	
	$1.1 \times 10^3$ $4.7 \times 10^3$	60	3.24	0.73	
		624	0.27	0.052	
1 40220 4	$6.7 \times 10^3$	366	0.20	0.064	F (1/2007 2000)
140320A	$1.1 \times 10^3$	$1.2 \times 10^4$	0.13	0.032	Evans et al. (2007a, 2009)
140402 A h	$2.7 \times 10^4$	$1.2 \times 10^4$	< 0.062		P: (2014)
140402A <sup>b</sup> 140414A <sup>b</sup>	$4.2 \times 10^4$ $4.1 \times 10^4$	$4.5 \times 10^3$ $1.7 \times 10^3$	$<4.2 \times 10^{-3}$		Pagani (2014)
		1 / X IU"	< 0.027	2	D'Avanzo et al. (2014)
			0.012	20 \( 10-3	Evens et al. (2007a, 2000)
140516A	$1.5 \times 10^{3}$	$2.5 \times 10^{4}$	0.013	$2.9 \times 10^{-3}$	Evans et al. (2007a, 2009)
	$1.5 \times 10^3$ $1.9 \times 10^5$	$2.5 \times 10^4$ $3.1 \times 10^5$	$1.7 \times 10^{-3}$	$2.9 \times 10^{-3} \\ 9.7 \times 10^{-4}$	, , ,
140606A <sup>b</sup>	$1.5 \times 10^{3}$ $1.9 \times 10^{5}$ $4.6 \times 10^{3}$	$2.5 \times 10^4$ $3.1 \times 10^5$ $2.4 \times 10^3$	$1.7 \times 10^{-3} < 8.7 \times 10^{-3}$		Stroh et al. (2014)
140606A <sup>b</sup> 140619B <sup>b</sup>	$1.5 \times 10^{3}$ $1.9 \times 10^{5}$ $4.6 \times 10^{3}$ $5.9 \times 10^{4}$	$2.5 \times 10^4$ $3.1 \times 10^5$ $2.4 \times 10^3$ $3.9 \times 10^3$	$1.7 \times 10^{-3}$ $< 8.7 \times 10^{-3}$ $< 3.4 \times 10^{-3}$	$9.7 \times 10^{-4}$	Stroh et al. (2014) Maselli & D'Avanzo (2014)
140606A <sup>b</sup> 140619B <sup>b</sup>	$1.5 \times 10^{3}$ $1.9 \times 10^{5}$ $4.6 \times 10^{3}$ $5.9 \times 10^{4}$ $1.3 \times 10^{2}$	$2.5 \times 10^{4}$ $3.1 \times 10^{5}$ $2.4 \times 10^{3}$ $3.9 \times 10^{3}$ $50$	$1.7 \times 10^{-3} < 8.7 \times 10^{-3} < 3.4 \times 10^{-3}  1.86$	$9.7 \times 10^{-4}$ $0.40$	Stroh et al. (2014)
140606A <sup>b</sup> 140619B <sup>b</sup>	$1.5 \times 10^{3}$ $1.9 \times 10^{5}$ $4.6 \times 10^{3}$ $5.9 \times 10^{4}$ $1.3 \times 10^{2}$ $2.3 \times 10^{3}$	$2.5 \times 10^{4}$ $3.1 \times 10^{5}$ $2.4 \times 10^{3}$ $3.9 \times 10^{3}$ $50$ $3.5 \times 10^{4}$	$1.7 \times 10^{-3}$ $< 8.7 \times 10^{-3}$ $< 3.4 \times 10^{-3}$ $1.86$ $0.015$	$9.7 \times 10^{-4}$	Stroh et al. (2014) Maselli & D'Avanzo (2014)
40606A <sup>b</sup> 40619B <sup>b</sup> 40622A	$1.5 \times 10^{3}$ $1.9 \times 10^{5}$ $4.6 \times 10^{3}$ $5.9 \times 10^{4}$ $1.3 \times 10^{2}$ $2.3 \times 10^{3}$ $8.0 \times 10^{4}$	$2.5 \times 10^{4}$ $3.1 \times 10^{5}$ $2.4 \times 10^{3}$ $3.9 \times 10^{3}$ $50$ $3.5 \times 10^{4}$ $2.3 \times 10^{4}$	$1.7 \times 10^{-3}$ $< 8.7 \times 10^{-3}$ $< 3.4 \times 10^{-3}$ $1.86$ $0.015$ $< 8.4 \times 10^{-3}$	$9.7 \times 10^{-4}$ $0.40$ $2.9 \times 10^{-3}$	Stroh et al. (2014) Maselli & D'Avanzo (2014) Evans et al. (2007a, 2009)
140606A <sup>b</sup> 140619B <sup>b</sup> 140622A	$1.5 \times 10^{3}$ $1.9 \times 10^{5}$ $4.6 \times 10^{3}$ $5.9 \times 10^{4}$ $1.3 \times 10^{2}$ $2.3 \times 10^{3}$ $8.0 \times 10^{4}$ $1.0 \times 10^{3}$	$2.5 \times 10^{4}$ $3.1 \times 10^{5}$ $2.4 \times 10^{3}$ $3.9 \times 10^{3}$ $50$ $3.5 \times 10^{4}$ $2.3 \times 10^{4}$ $221$	$\begin{array}{l} 1.7\times10^{-3}\\ < 8.7\times10^{-3}\\ < 3.4\times10^{-3}\\ 1.86\\ 0.015\\ < 8.4\times10^{-3}\\ 1.02 \end{array}$	$9.7 \times 10^{-4}$ $0.40$ $2.9 \times 10^{-3}$ $0.23$	Stroh et al. (2014) Maselli & D'Avanzo (2014)
140606A <sup>b</sup> 140619B <sup>b</sup> 140622A	$1.5 \times 10^{3}$ $1.9 \times 10^{5}$ $4.6 \times 10^{3}$ $5.9 \times 10^{4}$ $1.3 \times 10^{2}$ $2.3 \times 10^{3}$ $8.0 \times 10^{4}$ $1.0 \times 10^{3}$ $1.2 \times 10^{3}$	$2.5 \times 10^{4}$ $3.1 \times 10^{5}$ $2.4 \times 10^{3}$ $3.9 \times 10^{3}$ $50$ $3.5 \times 10^{4}$ $2.3 \times 10^{4}$ $221$ $176$	$\begin{array}{l} 1.7\times10^{-3}\\ < 8.7\times10^{-3}\\ < 3.4\times10^{-3}\\ 1.86\\ 0.015\\ < 8.4\times10^{-3}\\ 1.02\\ 1.3 \end{array}$	$9.7 \times 10^{-4}$ $0.40$ $2.9 \times 10^{-3}$ $0.23$ $0.30$	Stroh et al. (2014) Maselli & D'Avanzo (2014) Evans et al. (2007a, 2009)
140606A <sup>b</sup> 140619B <sup>b</sup> 140622A	$1.5 \times 10^{3}$ $1.9 \times 10^{5}$ $4.6 \times 10^{3}$ $5.9 \times 10^{4}$ $1.3 \times 10^{2}$ $2.3 \times 10^{3}$ $8.0 \times 10^{4}$ $1.0 \times 10^{3}$	$2.5 \times 10^{4}$ $3.1 \times 10^{5}$ $2.4 \times 10^{3}$ $3.9 \times 10^{3}$ $50$ $3.5 \times 10^{4}$ $2.3 \times 10^{4}$ $221$ $176$ $303$	$\begin{array}{l} 1.7\times10^{-3}\\ < 8.7\times10^{-3}\\ < 3.4\times10^{-3}\\ 1.86\\ 0.015\\ < 8.4\times10^{-3}\\ 1.02\\ 1.3\\ 0.74 \end{array}$	$9.7 \times 10^{-4}$ $0.40$ $2.9 \times 10^{-3}$ $0.23$ $0.30$ $0.17$	Stroh et al. (2014) Maselli & D'Avanzo (2014) Evans et al. (2007a, 2009)
140606A <sup>b</sup> 140619B <sup>b</sup> 140622A	$1.5 \times 10^{3}$ $1.9 \times 10^{5}$ $4.6 \times 10^{3}$ $5.9 \times 10^{4}$ $1.3 \times 10^{2}$ $2.3 \times 10^{3}$ $8.0 \times 10^{4}$ $1.0 \times 10^{3}$ $1.2 \times 10^{3}$ $1.4 \times 10^{3}$	$2.5 \times 10^{4}$ $3.1 \times 10^{5}$ $2.4 \times 10^{3}$ $3.9 \times 10^{3}$ $50$ $3.5 \times 10^{4}$ $2.3 \times 10^{4}$ $221$ $176$ $303$ $223$	$\begin{array}{l} 1.7\times10^{-3}\\ < 8.7\times10^{-3}\\ < 3.4\times10^{-3}\\ 1.86\\ 0.015\\ < 8.4\times10^{-3}\\ 1.02\\ 1.3\\ 0.74\\ 0.93 \end{array}$	$9.7 \times 10^{-4}$ $0.40$ $2.9 \times 10^{-3}$ $0.23$ $0.30$ $0.17$ $0.16$	Stroh et al. (2014) Maselli & D'Avanzo (2014) Evans et al. (2007a, 2009)
140606A <sup>b</sup> 140619B <sup>b</sup> 140622A	$1.5 \times 10^{3}$ $1.9 \times 10^{5}$ $4.6 \times 10^{3}$ $5.9 \times 10^{4}$ $1.3 \times 10^{2}$ $2.3 \times 10^{3}$ $8.0 \times 10^{4}$ $1.0 \times 10^{3}$ $1.2 \times 10^{3}$ $1.4 \times 10^{3}$ $5.0 \times 10^{3}$ $5.4 \times 10^{3}$	$2.5 \times 10^{4}$ $3.1 \times 10^{5}$ $2.4 \times 10^{3}$ $3.9 \times 10^{3}$ $50$ $3.5 \times 10^{4}$ $2.3 \times 10^{4}$ $221$ $176$ $303$ $223$ $158$	$\begin{array}{c} 1.7\times10^{-3}\\ < 8.7\times10^{-3}\\ < 3.4\times10^{-3}\\ 1.86\\ 0.015\\ < 8.4\times10^{-3}\\ 1.02\\ 1.3\\ 0.74\\ 0.93\\ 0.78 \end{array}$	$9.7 \times 10^{-4}$ $0.40$ $2.9 \times 10^{-3}$ $0.23$ $0.30$ $0.17$ $0.16$ $0.18$	Stroh et al. (2014) Maselli & D'Avanzo (2014) Evans et al. (2007a, 2009)
140606A <sup>b</sup> 140619B <sup>b</sup> 140622A	$1.5 \times 10^{3}$ $1.9 \times 10^{5}$ $4.6 \times 10^{3}$ $5.9 \times 10^{4}$ $1.3 \times 10^{2}$ $2.3 \times 10^{3}$ $8.0 \times 10^{4}$ $1.0 \times 10^{3}$ $1.2 \times 10^{3}$ $1.4 \times 10^{3}$ $5.0 \times 10^{3}$ $5.4 \times 10^{3}$ $5.6 \times 10^{3}$	$2.5 \times 10^4$ $3.1 \times 10^5$ $2.4 \times 10^3$ $3.9 \times 10^3$ $50$ $3.5 \times 10^4$ $2.3 \times 10^4$ $221$ $176$ $303$ $223$ $158$ $243$	$\begin{array}{c} 1.7\times10^{-3}\\ < 8.7\times10^{-3}\\ < 3.4\times10^{-3}\\ 1.86\\ 0.015\\ < 8.4\times10^{-3}\\ 1.02\\ 1.3\\ 0.74\\ 0.93\\ 0.78\\ 0.95 \end{array}$	$9.7 \times 10^{-4}$ $0.40$ $2.9 \times 10^{-3}$ $0.23$ $0.30$ $0.17$ $0.16$ $0.18$ $0.16$	Stroh et al. (2014) Maselli & D'Avanzo (2014) Evans et al. (2007a, 2009)
140606A <sup>b</sup> 140619B <sup>b</sup> 140622A	$1.5 \times 10^{3}$ $1.9 \times 10^{5}$ $4.6 \times 10^{3}$ $5.9 \times 10^{4}$ $1.3 \times 10^{2}$ $2.3 \times 10^{3}$ $8.0 \times 10^{4}$ $1.0 \times 10^{3}$ $1.2 \times 10^{3}$ $1.4 \times 10^{3}$ $5.0 \times 10^{3}$ $5.4 \times 10^{3}$ $5.6 \times 10^{3}$ $6.2 \times 10^{3}$	$2.5 \times 10^4$ $3.1 \times 10^5$ $2.4 \times 10^3$ $3.9 \times 10^3$ $50$ $3.5 \times 10^4$ $2.3 \times 10^4$ $221$ $176$ $303$ $223$ $158$ $243$ $130$	$\begin{array}{c} 1.7\times10^{-3}\\ < 8.7\times10^{-3}\\ < 3.4\times10^{-3}\\ 1.86\\ 0.015\\ < 8.4\times10^{-3}\\ 1.02\\ 1.3\\ 0.74\\ 0.93\\ 0.78\\ 0.95\\ 0.95\\ \end{array}$	$9.7 \times 10^{-4}$ $0.40$ $2.9 \times 10^{-3}$ $0.23$ $0.30$ $0.17$ $0.16$ $0.18$ $0.16$ $0.21$	Stroh et al. (2014) Maselli & D'Avanzo (2014) Evans et al. (2007a, 2009)
140606A <sup>b</sup> 140619B <sup>b</sup> 140622A	$1.5 \times 10^{3}$ $1.9 \times 10^{5}$ $4.6 \times 10^{3}$ $5.9 \times 10^{4}$ $1.3 \times 10^{2}$ $2.3 \times 10^{3}$ $8.0 \times 10^{4}$ $1.0 \times 10^{3}$ $1.2 \times 10^{3}$ $1.4 \times 10^{3}$ $5.0 \times 10^{3}$ $5.4 \times 10^{3}$ $6.2 \times 10^{3}$ $6.3 \times 10^{3}$	$2.5 \times 10^4$ $3.1 \times 10^5$ $2.4 \times 10^3$ $3.9 \times 10^3$ $50$ $3.5 \times 10^4$ $2.3 \times 10^4$ $221$ $176$ $303$ $223$ $158$ $243$ $130$ $123$	$\begin{array}{c} 1.7\times10^{-3}\\ < 8.7\times10^{-3}\\ < 3.4\times10^{-3}\\ 1.86\\ 0.015\\ < 8.4\times10^{-3}\\ 1.02\\ 1.3\\ 0.74\\ 0.93\\ 0.78\\ 0.95\\ 0.95\\ 0.86 \end{array}$	$9.7 \times 10^{-4}$ $0.40$ $2.9 \times 10^{-3}$ $0.23$ $0.30$ $0.17$ $0.16$ $0.18$ $0.16$ $0.21$ $0.19$	Stroh et al. (2014) Maselli & D'Avanzo (2014) Evans et al. (2007a, 2009)
140606A <sup>b</sup> 140619B <sup>b</sup> 140622A	$1.5 \times 10^{3}$ $1.9 \times 10^{5}$ $4.6 \times 10^{3}$ $5.9 \times 10^{4}$ $1.3 \times 10^{2}$ $2.3 \times 10^{3}$ $8.0 \times 10^{4}$ $1.0 \times 10^{3}$ $1.2 \times 10^{3}$ $5.0 \times 10^{3}$ $5.4 \times 10^{3}$ $6.2 \times 10^{3}$ $6.3 \times 10^{3}$ $6.5 \times 10^{3}$	$2.5 \times 10^4$ $3.1 \times 10^5$ $2.4 \times 10^3$ $3.9 \times 10^3$ $50$ $3.5 \times 10^4$ $2.3 \times 10^4$ $221$ $176$ $303$ $223$ $158$ $243$ $130$ $123$ $183$	$\begin{array}{l} 1.7\times10^{-3}\\ < 8.7\times10^{-3}\\ < 3.4\times10^{-3}\\ 1.86\\ 0.015\\ < 8.4\times10^{-3}\\ 1.02\\ 1.3\\ 0.74\\ 0.93\\ 0.78\\ 0.95\\ 0.95\\ 0.86\\ 1.05 \end{array}$	$9.7 \times 10^{-4}$ $0.40$ $2.9 \times 10^{-3}$ $0.23$ $0.30$ $0.17$ $0.16$ $0.18$ $0.16$ $0.21$ $0.19$ $0.21$	Stroh et al. (2014) Maselli & D'Avanzo (2014) Evans et al. (2007a, 2009)
140606A <sup>b</sup> 140619B <sup>b</sup> 140622A	$1.5 \times 10^{3}$ $1.9 \times 10^{5}$ $4.6 \times 10^{3}$ $5.9 \times 10^{4}$ $1.3 \times 10^{2}$ $2.3 \times 10^{3}$ $8.0 \times 10^{4}$ $1.0 \times 10^{3}$ $1.2 \times 10^{3}$ $1.4 \times 10^{3}$ $5.0 \times 10^{3}$ $5.4 \times 10^{3}$ $6.2 \times 10^{3}$ $6.3 \times 10^{3}$	$2.5 \times 10^4$ $3.1 \times 10^5$ $2.4 \times 10^3$ $3.9 \times 10^3$ $50$ $3.5 \times 10^4$ $2.3 \times 10^4$ $221$ $176$ $303$ $223$ $158$ $243$ $130$ $123$ $183$ $288$	$\begin{array}{c} 1.7\times10^{-3}\\ < 8.7\times10^{-3}\\ < 3.4\times10^{-3}\\ 1.86\\ 0.015\\ < 8.4\times10^{-3}\\ 1.02\\ 1.3\\ 0.74\\ 0.93\\ 0.78\\ 0.95\\ 0.95\\ 0.86\\ 1.05\\ 0.93\\ \end{array}$	$9.7 \times 10^{-4}$ $0.40$ $2.9 \times 10^{-3}$ $0.23$ $0.30$ $0.17$ $0.16$ $0.18$ $0.16$ $0.21$ $0.19$ $0.21$ $0.16$	Stroh et al. (2014) Maselli & D'Avanzo (2014) Evans et al. (2007a, 2009)
140516A 140606A <sup>b</sup> 140619B <sup>b</sup> 140622A 140903A	$1.5 \times 10^{3}$ $1.9 \times 10^{5}$ $4.6 \times 10^{3}$ $5.9 \times 10^{4}$ $1.3 \times 10^{2}$ $2.3 \times 10^{3}$ $8.0 \times 10^{4}$ $1.0 \times 10^{3}$ $1.2 \times 10^{3}$ $5.0 \times 10^{3}$ $5.4 \times 10^{3}$ $6.2 \times 10^{3}$ $6.3 \times 10^{3}$ $6.5 \times 10^{3}$ $6.8 \times 10^{3}$	$2.5 \times 10^4$ $3.1 \times 10^5$ $2.4 \times 10^3$ $3.9 \times 10^3$ $50$ $3.5 \times 10^4$ $2.3 \times 10^4$ $221$ $176$ $303$ $223$ $158$ $243$ $130$ $123$ $183$	$\begin{array}{l} 1.7\times10^{-3}\\ < 8.7\times10^{-3}\\ < 3.4\times10^{-3}\\ 1.86\\ 0.015\\ < 8.4\times10^{-3}\\ 1.02\\ 1.3\\ 0.74\\ 0.93\\ 0.78\\ 0.95\\ 0.95\\ 0.86\\ 1.05 \end{array}$	$9.7 \times 10^{-4}$ $0.40$ $2.9 \times 10^{-3}$ $0.23$ $0.30$ $0.17$ $0.16$ $0.18$ $0.16$ $0.21$ $0.19$ $0.21$	Stroh et al. (2014) Maselli & D'Avanzo (2014) Evans et al. (2007a, 2009)

Table A1 — Continued

GRB	δ <i>t</i> (s)	Exposure Time (s)	$F_X$ $(\mu Jy)$	$\sigma_X \ (\mu  ext{Jy})$	References
	$1.1 \times 10^4$	178	0.54	0.12	
	$1.1 \times 10^4$	173	0.59	0.13	
	$1.1 \times 10^4$	216	0.55	0.11	
	$1.3 \times 10^{4}$	333	0.49	0.085	
	$1.6 \times 10^{4}$	223	0.42	0.098	
	$1.7 \times 10^4$	228	0.32	0.085	
	$1.7 \times 10^{4}$	203	0.370	0.096	
	$1.7 \times 10^{4}$	133	0.75	0.17	
	$1.7 \times 10^{4}$	333	0.38	0.076	
	$2.2 \times 10^{4}$	517	0.25	0.065	
	$2.3 \times 10^{4}$	354	0.36	0.093	
	$2.3 \times 10^{4}$	461	0.27	0.070	
	$2.8 \times 10^{4}$	354	0.23	0.057	
	$2.8 \times 10^{4}$	253	0.30	0.077	
	$2.9 \times 10^{4}$	366	0.20	0.054	
	$3.1 \times 10^{4}$	$5.1 \times 10^{3}$	0.18	0.033	
	$4.2 \times 10^{4}$	$6.0 \times 10^{3}$	0.13	0.034	
	$4.6 \times 10^{4}$	479	0.21	0.052	
	$5.4 \times 10^{4}$	$6.5 \times 10^{3}$	0.080	0.018	
	$6.8 \times 10^{4}$	$1.2 \times 10^{4}$	0.076	0.0142	
	$1.2 \times 10^{5}$	$8.6 \times 10^{4}$	0.029	$6.5 \times 10^{-3}$	
	$2.6 \times 10^{5}$	$5.8 \times 10^{4}$	< 0.011		
140930B	$1.1 \times 10^{3}$	158	0.56	0.13	Evans et al. (2007a, 2009)
	$1.2 \times 10^{3}$	115	0.77	0.17	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,
	$1.3 \times 10^{3}$	133	0.66	0.15	
	$1.5 \times 10^{3}$	196	0.45	0.10	
	$1.7 \times 10^{3}$	191	0.35	0.090	
	$1.9 \times 10^{3}$	221	0.40	0.090	
	$2.1 \times 10^{3}$	216	0.41	0.093	
	$2.3 \times 10^{3}$	236	0.28	0.074	
	$2.6 \times 10^{3}$	306	0.29	0.066	
	$6.3 \times 10^3$	559	0.13	0.033	
	$9.7 \times 10^{3}$	$7.7 \times 10^{3}$	0.050	$8.7 \times 10^{-3}$	
	$8.0 \times 10^4$	$1.2 \times 10^{5}$	$4.5 \times 10^{-3}$	$1.3 \times 10^{-3}$	
	$1.7 \times 10^{5}$	925	< 0.025	1.5 × 10	
	$^{+}3.2 \times 10^{5}$	$2.3 \times 10^4$	$3.3 \times 10^{-4}$	$1.6 \times 10^{-4}$	This work
	$^{+}2.0 \times 10^{6}$	$3.4 \times 10^4$	$< 2.7 \times 10^{-4}$	1.0 / 10	This work This work
141202A <sup>b</sup>	$1.2 \times 10^{5}$	$5.1 \times 10^4$	$< 6.1 \times 10^{-3}$		Pagani & Evans (2014)
$141202A$ $141205A^b$	$2.9 \times 10^4$	$5.0 \times 10^3$	$< 3.3 \times 10^{-3}$		Starling & Page (2014)
141203A 141212A	$2.9 \times 10^{4}$ $2.1 \times 10^{4}$	$2.9 \times 10^4$	$9.1 \times 10^{-3}$	$3.5 \times 10^{-3}$	Evans et al. (2007a, 2009)
141212A	2.1 × 10	2.9 X 10	9.1 × 10	3.3 × 10	Evans et al. (2007a, 2009)
150101A	$1.3 \times 10^{3}$	$2.3 \times 10^{4}$	0.016	$3.7 \times 10^{-3}$	Evans et al. (2007a, 2009)
	$6.0 \times 10^{4}$	$1.8 \times 10^{4}$	< 0.010		, , ,
150101B	$^{+}6.9 \times 10^{5}$	$1.5 \times 10^{4}$	0.011	$1.1 \times 10^{-3}$	Fong et al., in prep.
	$^{+}3.4 \times 10^{6}$	$1.5 \times 10^{4}$	$1.9 \times 10^{-3}$	$4.7 \times 10^{-4}$	Fong et al., in prep.
150120A	$8.3 \times 10^{3}$	$1.2 \times 10^{4}$	< 0.031		Evans et al. (2007a, 2009)
150301A	$1.4 \times 10^{4}$	$4.0 \times 10^{4}$	< 0.051		Evans et al. (2007a, 2009)

**Note**. — Upper limits correspond to  $3\sigma$ . Unless otherwise stated, all data are taken with *Swift/XRT* and X-ray flux densities are at 1 keV. <sup>a</sup> We employed a fiducial spectral index of  $\beta_X = -1$ . <sup>b</sup> We employed a fiducial count rate-to-unabsorbed flux conversion factor of  $10^{-11}$  erg cm<sup>-2</sup> s<sup>-1</sup> and spectral index  $\beta_X = -1$ .

we employed a fiductar count rate-to-unassorbed flux conversion factor of 10 stell eigent = s stand spectral fidux p<sub>X</sub> = -1.
 Reported flux is in the energy range of 0.7-10 keV.
 d Late-time re-brightening in GRB 080503 light curve is observed in both optical and X-ray bands and is unlikely the afterglow (Perley et al. 2009b).
 + Chandra observation.
 + XMM-Newton observation.

Table A2
Short GRB Optical/near-IR afterglow catalog

GRB	δt (hr)	Telescope	Instrument	Filter	Exposure Time (s)	$F_{\nu}$ $(\mu Jy)$	σ (μ <b>J</b> y)	References
050202	12.6	Mount John	MOA	R	900	< 42.89	(103)	Castro-Tirado et al. (2005)
050509B	2.1	WIYN	OPTIC	r	600	< 0.75		Bloom et al. (2006)
050709	34.0	Danish tel.	DFOSC	R	7200	2.34	0.12	Hjorth et al. (2005)
030707	59.1	VLT	FORS2	V	3600	0.64	0.07	Covino et al. (2006)
	59.3	VLT	FORS2	R	3000	0.90	0.05	Covino et al. (2000)
	60.0	Danish tel.	DFOSC	R	10200	1.17	0.26	Hjorth et al. (2005)
	104.7	VLT	FORS1	V	3600	< 0.36	0.20	Covino et al. (2006)
	134.4	HST	ACS	F814W	6360	0.34	0.006	Fox et al. (2005)
	235.2	HST	ACS	F814W	6360	0.17	0.008	100000000000000000000000000000000000000
	832.8	HST	ACS	F814W	6360	< 0.02	0.000	
050724A	11.4	VLT	FORS1	V	480	16.0	0.45	Malesani et al. (2007a)
	11.6	Magellan/Baade	PANIC	K	1320	45.6	1.4	Berger et al. (2005)
	11.8	VLT	FORS1	R	540	17.8	0.7	Malesani et al. (2007a)
	11.8	VLT	FORS1	I	540	18.37	0.51	,
	12.0	Swope 40-in		I	1800	19.1	0.2	Berger et al. (2005)
	14.2	Swope 40-in		I	1800	25.2	0.9	. , ,
	34.8	VLТ	FORS1	I	540	2.81	0.33	Malesani et al. (2007a)
	34.9	Magellan/Baade	PANIC	K	1320	< 5.4		Berger et al. (2005)
	35.0	VLT	FORS1	R	540	3.35	0.29	Malesani et al. (2007a)
	36.7	Swope 40-in		I	2700	< 9.1		Berger et al. (2005)
	83.1	VLT	FORS1	Ī	540	0.33	0.12	Malesani et al. (2007a)
050813	13.2	CAHA 2.2-m	CAFOS	Ī	6000	< 1.82		Ferrero et al. (2007)
	14.1	CAHA 2.2-m	CAFOS	$\overline{R}$	4140	< 1.43		
050906	21.4	VLT	FORS2	R	1800	< 0.15		Levan et al. (2008)
051210	19.4	Magellan/Clay	LDSS3	r	1200	< 1.6		Berger et al. (2007)
051221A	3.1	Gemini-N	GMOS	r		16.8	1.3	Soderberg et al. (2006)
	3.4	Gemini-N	GMOS	r		15.6	1.2	8 (,
	26.9	Gemini-N	GMOS	r		2.5	0.19	
	27.2	Gemini-N	GMOS	i		2.3	0.54	
	51.0	Gemini-N	GMOS	i		0.83	0.32	
	51.3	Gemini-N	GMOS	Z		1.1	0.47	
	51.6	Gemini-N	GMOS	r		0.93	0.11	
	75.7	Gemini-N	GMOS	r		0.94	0.28	
	123.6	Gemini-N	GMOS	r		0.50	0.11	
	123.0	Ochini-IV	OMOS	,		0.50		
	147.8	Gemini-N	GMOS	r		0.32	0.11	
060121	147.8	Gemini-N		r	120	0.32		de Ugarte Postigo et al. (2006)
060121	0.37	Gemini-N OSN		r I	120 120	19.3	4.4	de Ugarte Postigo et al. (2006)
060121	0.37 0.70	Gemini-N OSN OSN	GMOS	r I I	120 120	0.32 19.3 10.0	4.4 2.8	
060121	0.37 0.70 2.0	Gemini-N OSN OSN NOT	GMOS ALFOSC	r I I R		19.3 10.0 3.3	4.4 2.8 0.70	de Ugarte Postigo et al. (2006) Levan et al. (2006)
060121	0.37 0.70 2.0 2.6	OSN OSN NOT NOT	GMOS	I I R R	120	19.3 10.0 3.3 6.1	4.4 2.8 0.70 1.9	Levan et al. (2006)
060121	0.37 0.70 2.0 2.6 2.7	OSN OSN NOT NOT OSN	GMOS ALFOSC	I I R R I	120 300	19.3 10.0 3.3 6.1 10.8	4.4 2.8 0.70 1.9 2.5	
060121	0.37 0.70 2.0 2.6 2.7 2.8	Gemini-N OSN OSN NOT NOT OSN OSN	GMOS ALFOSC	r I I R R I R	300 600	19.3 10.0 3.3 6.1 10.8 3.7	4.4 2.8 0.70 1.9 2.5 0.97	Levan et al. (2006)
060121	0.37 0.70 2.0 2.6 2.7 2.8 3.9	Gemini-N  OSN OSN NOT NOT OSN OSN CSN	GMOS ALFOSC	r II IR R II R R R IR	300 600 1200	19.3 10.0 3.3 6.1 10.8 3.7 1.1	4.4 2.8 0.70 1.9 2.5 0.97 0.33	Levan et al. (2006)
060121	0.37 0.70 2.0 2.6 2.7 2.8 3.9 4.4	OSN OSN NOT NOT OSN OSN CAHA 2.2-m CAHA 2.2-m	GMOS ALFOSC	r I I R R I R R	300 600 1200 1800	19.3 10.0 3.3 6.1 10.8 3.7 1.1 4.6	4.4 2.8 0.70 1.9 2.5 0.97 0.33 0.83	Levan et al. (2006)
060121	0.37 0.70 2.0 2.6 2.7 2.8 3.9 4.4 5.4	OSN OSN NOT NOT OSN OSN CAHA 2.2-m CAHA 2.2-m	ALFOSC ALFOSC	r I I R R I R R R	300 600 1200	19.3 10.0 3.3 6.1 10.8 3.7 1.1 4.6 1.2	4.4 2.8 0.70 1.9 2.5 0.97 0.33 0.83 0.49	Levan et al. (2006) de Ugarte Postigo et al. (2006
060121	0.37 0.70 2.0 2.6 2.7 2.8 3.9 4.4 5.4 5.6	OSN OSN NOT NOT OSN OSN CAHA 2.2-m CAHA 2.2-m Bok	GMOS  ALFOSC ALFOSC  90Prime	r I I R R I R R R R	300 600 1200 1800	19.3 10.0 3.3 6.1 10.8 3.7 1.1 4.6 1.2	4.4 2.8 0.70 1.9 2.5 0.97 0.33 0.83	Levan et al. (2006)
060121	0.37 0.70 2.0 2.6 2.7 2.8 3.9 4.4 5.6 6.5	OSN OSN NOT NOT OSN OSN CAHA 2.2-m CAHA 2.2-m Bok Bok	GMOS  ALFOSC ALFOSC  90Prime 90Prime	I I I R R I R R R R R R B	300 600 1200 1800	19.3 10.0 3.3 6.1 10.8 3.7 1.1 4.6 1.2 1.1 < 0.96	4.4 2.8 0.70 1.9 2.5 0.97 0.33 0.83 0.49 0.22	Levan et al. (2006) de Ugarte Postigo et al. (2006
060121	0.37 0.70 2.0 2.6 2.7 2.8 3.9 4.4 5.4 5.6	OSN OSN NOT NOT OSN OSN CAHA 2.2-m CAHA 2.2-m Bok	GMOS  ALFOSC ALFOSC  90Prime	r I I R R I R R R R	300 600 1200 1800	19.3 10.0 3.3 6.1 10.8 3.7 1.1 4.6 1.2	4.4 2.8 0.70 1.9 2.5 0.97 0.33 0.83 0.49	Levan et al. (2006) de Ugarte Postigo et al. (2006
060121	0.37 0.70 2.0 2.6 2.7 2.8 3.9 4.4 5.4 5.6 6.5 7.3 7.3	OSN OSN NOT NOT OSN OSN CAHA 2.2-m CAHA 2.2-m CAHA 2.2-m Bok Bok Bok CAHA 2.2-m	GMOS  ALFOSC ALFOSC  90Prime 90Prime	I I R R I R R R R R R	300 600 1200 1800 1200	19.3 10.0 3.3 6.1 10.8 3.7 1.1 4.6 1.2 1.1 < 0.96 1.2	4.4 2.8 0.70 1.9 2.5 0.97 0.33 0.83 0.49 0.22 0.18 0.37	Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)
060121	0.37 0.70 2.0 2.6 2.7 2.8 3.9 4.4 5.4 5.6 6.5 7.3 7.4	OSN OSN NOT NOT OSN OSN CAHA 2.2-m CAHA 2.2-m CAHA 2.2-m Bok Bok Bok CAHA 2.2-m WHT	GMOS  ALFOSC ALFOSC  90Prime 90Prime	I I R R I R R R R R R R	300 600 1200 1800 1200	19.3 10.0 3.3 6.1 10.8 3.7 1.1 4.6 1.2 1.1 < 0.96 1.2 1.4	4.4 2.8 0.70 1.9 2.5 0.97 0.33 0.83 0.49 0.22	Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)
060121	0.37 0.70 2.0 2.6 2.7 2.8 3.9 4.4 5.4 5.6 6.5 7.3 7.3 7.4 7.7	OSN OSN NOT NOT OSN OSN CAHA 2.2-m CAHA 2.2-m CAHA 2.2-m Bok Bok Bok CAHA 2.2-m WHT OSN	ALFOSC ALFOSC 90Prime 90Prime 90Prime	I I R R I R R R R R R R	300 600 1200 1800 1200	19.3 10.0 3.3 6.1 10.8 3.7 1.1 4.6 1.2 1.1 < 0.96 1.2 1.4 17.1 < 2.5	4.4 2.8 0.70 1.9 2.5 0.97 0.33 0.83 0.49 0.22 0.18 0.37 1.4	Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)
060121	0.37 0.70 2.0 2.6 2.7 2.8 3.9 4.4 5.4 5.6 6.5 7.3 7.3 7.4 7.7	OSN OSN NOT NOT OSN OSN CSN CSN CAHA 2.2-m CAHA 2.2-m Bok Bok Bok CAHA 2.2-m WHT OSN Bok	GMOS  ALFOSC ALFOSC  90Prime 90Prime	I I R R I R R R R R R R I R	300 600 1200 1800 1200	19.3 10.0 3.3 6.1 10.8 3.7 1.1 4.6 1.2 1.1 < 0.96 1.2 1.4 17.1 < 2.5 1.6	4.4 2.8 0.70 1.9 2.5 0.97 0.33 0.83 0.49 0.22 0.18 0.37 1.4	Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)
060121	0.37 0.70 2.0 2.6 2.7 2.8 3.9 4.4 5.4 5.6 6.5 7.3 7.3 7.4 7.7 11.3 13.1	OSN OSN NOT NOT OSN OSN CAHA 2.2-m CAHA 2.2-m Bok Bok Bok CAHA 2.2-m WHT OSN Bok WIYN	ALFOSC ALFOSC 90Prime 90Prime 90Prime	I I R R I R R R R R R R R R R R R R R R	300 600 1200 1800 1200	19.3 10.0 3.3 6.1 10.8 3.7 1.1 4.6 1.2 1.1 < 0.96 1.2 1.4 17.1 < 2.5 1.6	4.4 2.8 0.70 1.9 2.5 0.97 0.33 0.83 0.49 0.22 0.18 0.37 1.4	Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)
060121	0.37 0.70 2.0 2.6 2.7 2.8 3.9 4.4 5.6 6.5 7.3 7.3 7.4 7.7 11.3 13.1 30.0	Gemini-N  OSN OSN NOT NOT OSN OSN CAHA 2.2-m CAHA 2.2-m CAHA 2.2-m Bok Bok Bok CAHA 2.2-m WHT OSN Bok WIYN OSN	ALFOSC ALFOSC 90Prime 90Prime 90Prime	I I R R I R R R R R R R R R R R R R R R	300 600 1200 1800 1200 1800 750 1500	19.3 10.0 3.3 6.1 10.8 3.7 1.1 4.6 1.2 1.1 < 0.96 1.2 1.4 17.1 < 2.5 1.6 1.2 < 0.8	4.4 2.8 0.70 1.9 2.5 0.97 0.33 0.83 0.49 0.22 0.18 0.37 1.4	Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)
060121	0.37 0.70 2.0 2.6 2.7 2.8 3.9 4.4 5.6 6.5 7.3 7.4 7.7 11.3 13.1 30.0 31.3	Gemini-N  OSN OSN NOT NOT OSN OSN CAHA 2.2-m CAHA 2.2-m CAHA 2.2-m Bok Bok CAHA 2.2-m WHT OSN Bok WIYN OSN WHT	ALFOSC ALFOSC 90Prime 90Prime 90Prime	I I R R R R R R R R R R R R K I R K K K K	120 300 600 1200 1800 1200 1800 750 1500 10800 1000	19.3 10.0 3.3 6.1 10.8 3.7 1.1 4.6 1.2 1.1 < 0.96 1.2 1.4 17.1 < 2.5 1.6 1.2 < 0.8 6.3	4.4 2.8 0.70 1.9 2.5 0.97 0.33 0.83 0.49 0.22 0.18 0.37 1.4 0.41 0.24	Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)
060121	0.37 0.70 2.0 2.6 2.7 2.8 3.9 4.4 5.4 5.6 6.5 7.3 7.4 7.7 11.3 13.1 30.0 31.3 33.5	Gemini-N  OSN OSN NOT NOT OSN OSN CAHA 2.2-m CAHA 2.2-m CAHA 2.2-m Bok Bok Bok CAHA 2.2-m WHT OSN Bok WIYN OSN WHT APO/ARC	ALFOSC ALFOSC 90Prime 90Prime 90Prime	I I R R R R R R R R R R R K K K K K K K	300 600 1200 1800 1200 1800 750 1500	19.3 10.0 3.3 6.1 10.8 3.7 1.1 4.6 1.2 1.1 < 0.96 1.2 1.4 17.1 < 2.5 1.6 1.2 < 0.8 6.3 7.5	4.4 2.8 0.70 1.9 2.5 0.97 0.33 0.83 0.49 0.22 0.18 0.37 1.4 0.41 0.24	Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)
060121	0.37 0.70 2.0 2.6 2.7 2.8 3.9 4.4 5.4 5.6 6.5 7.3 7.4 7.7 11.3 13.1 30.0 31.3 33.5 33.6	OSN OSN NOT NOT OSN OSN CAHA 2.2-m CAHA 2.2-m Bok Bok Bok Sok Bok WIYN OSN Bok WIYN OSN WHT APO/ARC WIYN	ALFOSC ALFOSC 90Prime 90Prime 90Prime	I I R R R R R R R R R R R R R R R R R R	300 600 1200 1800 1200 1800 750 1500 10800 1000 3600	19.3 10.0 3.3 6.1 10.8 3.7 1.1 4.6 1.2 1.1 < 0.96 1.2 1.4 17.1 < 2.5 1.6 1.2 < 0.8 6.3 7.5 0.41	4.4 2.8 0.70 1.9 2.5 0.97 0.33 0.83 0.49 0.22 0.18 0.37 1.4 0.41 0.24 1.6 0.65 0.06	Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  de Ugarte Postigo et al. (2006)
060121	0.37 0.70 2.0 2.6 2.7 2.8 3.9 4.4 5.4 5.6 6.5 7.3 7.3 7.4 7.7 11.3 13.1 30.0 31.3 33.5 33.6 51.5	OSN OSN OSN NOT NOT OSN OSN CAHA 2.2-m CAHA 2.2-m CAHA 2.2-m Bok Bok Bok CAHA 2.2-m WHT OSN Bok WIYN OSN WHT APO/ARC WIYN CAHA 2.2-m	GMOS  ALFOSC ALFOSC  90Prime 90Prime 90Prime NIC-FPS	I I R R R R R R R R K I R R R R R R R R	300 600 1200 1800 1200 1800 750 1500 10800 1000 3600 5400	19.3 10.0 3.3 6.1 10.8 3.7 1.1 4.6 1.2 1.1 < 0.96 1.2 1.4 17.1 < 2.5 1.6 1.2 < 0.8 6.3 7.5 0.41	4.4 2.8 0.70 1.9 2.5 0.97 0.33 0.83 0.49 0.22 0.18 0.37 1.4 0.41 0.24	Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  de Ugarte Postigo et al. (2006)
	0.37 0.70 2.0 2.6 2.7 2.8 3.9 4.4 5.6 6.5 7.3 7.4 7.7 11.3 13.1 30.0 31.3 33.5 51.5	OSN OSN OSN NOT NOT OSN OSN OSN CAHA 2.2-m CAHA 2.2-m CAHA 2.2-m Bok Bok Bok CAHA 2.2-m WHT OSN Bok WIYN OSN WHT APO/ARC WIYN CAHA 2.2-m	GMOS  ALFOSC ALFOSC  90Prime 90Prime 90Prime NIC-FPS	I I R R R R R R R R K I R R K K K K K K	120 300 600 1200 1800 1200 1800 750 1500 10800 1000 3600 5400 3600	19.3 10.0 3.3 6.1 10.8 3.7 1.1 4.6 1.2 1.1 < 0.96 1.2 1.4 17.1 < 2.5 1.6 1.2 < 0.8 6.3 7.5 0.41 0.55 < 2.13	4.4 2.8 0.70 1.9 2.5 0.97 0.33 0.83 0.49 0.22 0.18 0.37 1.4 0.24 1.6 0.65 0.06 0.14	Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  de Ugarte Postigo et al. (2006)
	0.37 0.70 2.0 2.6 2.7 2.8 3.9 4.4 5.6 6.5 7.3 7.4 7.7 11.3 13.1 30.0 31.3 33.5 33.6 51.5 127.8 0.045	Gemini-N  OSN OSN NOT NOT OSN OSN CAHA 2.2-m CAHA 2.2-m CAHA 2.2-m Bok Bok Bok CAHA 2.2-m WHT OSN Bok WIYN OSN WHT APO/ARC WIYN CAHA 2.2-m APO/ARC Swift	GMOS  ALFOSC ALFOSC  90Prime 90Prime 90Prime  NIC-FPS  NIC-FPS  UVOT	I I R R R R R R R R R R R R R R R R R R	120 300 600 1200 1800 1200 1800 750 1500 10800 1000 3600 5400 3600 200	19.3 10.0 3.3 6.1 10.8 3.7 1.1 4.6 1.2 1.1 < 0.96 1.2 1.4 17.1 < 2.5 1.6 1.2 < 0.8 6.3 7.5 0.41 0.55 < 2.13 55.0	4.4 2.8 0.70 1.9 2.5 0.97 0.33 0.83 0.49 0.22 0.18 0.37 1.4 0.41 0.24 1.6 0.65 0.06 0.14	Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  de Ugarte Postigo et al. (2006)
	0.37 0.70 2.0 2.6 2.7 2.8 3.9 4.4 5.4 5.6 6.5 7.3 7.4 7.7 11.3 13.1 30.0 31.3 33.5 33.6 51.5 127.8 0.045 0.12	OSN OSN OSN NOT NOT OSN OSN CAHA 2.2-m CAHA 2.2-m CAHA 2.2-m Bok Bok Bok CAHA 2.2-m WHT OSN Bok WIYN OSN WHT APO/ARC WIYN CAHA 2.2-m APO/ARC Swift Swift	GMOS  ALFOSC ALFOSC  90Prime 90Prime 90Prime  NIC-FPS  NIC-FPS UVOT UVOT	I I R R R R R R R R R R R K I R R R V U	120 300 600 1200 1800 1200 1800 750 1500 10800 1000 3600 5400 3600 200 50	19.3 10.0 3.3 6.1 10.8 3.7 1.1 4.6 1.2 1.1 < 0.96 1.2 1.4 17.1 < 2.5 1.6 1.2 < 0.8 6.3 7.5 0.41 0.55 < 2.13 55.0 22.7	4.4 2.8 0.70 1.9 2.5 0.97 0.33 0.83 0.49 0.22 0.18 0.37 1.4 0.41 0.24 1.6 0.65 0.06 0.14	Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  de Ugarte Postigo et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)
	0.37 0.70 2.0 2.6 2.7 2.8 3.9 4.4 5.4 5.6 6.5 7.3 7.4 7.7 11.3 13.1 30.0 31.3 33.5 33.6 51.5 127.8 0.045 0.012	OSN OSN OSN NOT NOT OSN OSN CAHA 2.2-m CAHA 2.2-m CAHA 2.2-m Bok Bok Bok CAHA 2.2-m WHT OSN Bok WIYN OSN WHT APO/ARC WIYN CAHA 2.2-m APO/ARC Swift Swift	90Prime 90Prime 90Prime NIC-FPS UVOT UVOT UVOT	I I R R R R R R R R R R R R K I R R C R R W U white	120 300 600 1200 1800 1200 1800 750 1500 10800 1000 3600 5400 3600 200 50 50	19.3 10.0 3.3 6.1 10.8 3.7 1.1 4.6 1.2 1.1 < 0.96 1.2 1.4 17.1 < 2.5 1.6 1.2 < 0.8 6.3 7.5 0.41 0.55 < 2.13 55.0 22.7 30.5	4.4 2.8 0.70 1.9 2.5 0.97 0.33 0.83 0.49 0.22 0.18 0.37 1.4 0.41 0.24 1.6 0.65 0.06 0.14 18.8 13.0 11.6	Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  de Ugarte Postigo et al. (2006)  de Ugarte Postigo et al. (2006)  Roming et al. (2006)
	0.37 0.70 2.0 2.6 2.7 2.8 3.9 4.4 5.6 6.5 7.3 7.3 7.4 7.7 11.3 13.1 30.0 31.3 33.5 33.6 51.5 127.8 0.045 0.12 0.22 1.2	Gemini-N  OSN OSN NOT NOT OSN OSN OSN CAHA 2.2-m CAHA 2.2-m CAHA 2.2-m Bok Bok Bok CAHA 2.2-m WHT OSN Bok WIYN OSN WHT APO/ARC WIYN CAHA 2.2-m APO/ARC Swift Swift Swift Gemini-S	90Prime 90Prime 90Prime NIC-FPS UVOT UVOT GMOS	I I R R R R R R R R R R K I R R K V U white r	120 300 600 1200 1800 1200 1800 750 1500 10800 1000 3600 5400 3600 200 50 50 1800	0.32  19.3 10.0 3.3 6.1 10.8 3.7 1.1 4.6 1.2 1.1 < 0.96 1.2 1.4 17.1 < 2.5 1.6 1.2 < 0.8 6.3 7.5 0.41 0.55 < 2.13 55.0 22.7 30.5 36.6	4.4 2.8 0.70 1.9 2.5 0.97 0.33 0.83 0.49 0.22 0.18 0.37 1.4 0.41 0.24 1.6 0.65 0.06 0.14 18.8 13.0 11.6 6.6	Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Berger et al. (2007)
	0.37 0.70 2.0 2.6 2.7 2.8 3.9 4.4 5.6 6.5 7.3 7.3 7.4 7.7 11.3 13.1 30.0 31.3 33.5 33.6 51.5 127.8 0.045 0.12 0.22 1.2	OSN OSN OSN NOT NOT OSN OSN CAHA 2.2-m CAHA 2.2-m CAHA 2.2-m Bok Bok Bok CAHA 2.2-m WHT OSN Bok WIYN OSN WHT APO/ARC WIYN CAHA 2.2-m APO/ARC Swift Swift Swift Gemini-S Swift	90Prime 90Prime 90Prime 90Prime VIC-FPS UVOT UVOT UVOT GMOS UVOT	I I R R R I R R R R R R R R K I R R R K K V White B	120 300 600 1200 1800 1200 1800 750 1500 10800 1000 3600 5400 3600 200 50 50 1800 886	19.3 10.0 3.3 6.1 10.8 3.7 1.1 4.6 1.2 1.1 < 0.96 1.2 1.4 17.1 < 2.5 1.6 1.2 < 0.8 6.3 7.5 0.41 0.55 < 2.13 55.0 22.7 36.6 18.7	4.4 2.8 0.70 1.9 2.5 0.97 0.33 0.83 0.49 0.22 0.18 0.37 1.4 0.24 1.6 0.65 0.06 0.14	Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Roming et al. (2006)
	0.37 0.70 2.0 2.6 2.7 2.8 3.9 4.4 5.6 6.5 7.3 7.4 7.7 11.3 13.1 30.0 31.3 33.5 33.6 51.5 127.8 0.045 0.12 0.22 1.2	OSN OSN OSN NOT NOT OSN OSN CAHA 2.2-m CAHA 2.2-m CAHA 2.2-m Bok Bok Bok CAHA 2.2-m WHT OSN Bok WIYN OSN WHT APO/ARC WIYN CAHA 2.2-m APO/ARC Swift Swift Swift Swift Swift	90Prime 90Prime 90Prime 90Prime VIC-FPS UVOT UVOT GMOS UVOT UVOT UVOT	I I R R R I R R R R R R R R K I R R K V U white B UVW2	120 300 600 1200 1800 1200 1800 750 1500 10800 1000 3600 5400 3600 200 50 50 1800 886 900	19.3 10.0 3.3 6.1 10.8 3.7 1.1 4.6 1.2 1.1 < 0.96 1.2 1.4 17.1 < 2.5 1.6 1.2 < 0.8 6.3 7.5 0.41 0.55 < 2.13 55.0 22.7 30.5 36.6 18.7 6.4	4.4 2.8 0.70 1.9 2.5 0.97 0.33 0.83 0.49 0.22 0.18 0.37 1.4 0.41 0.24 1.6 0.65 0.06 0.14 18.8 13.0 11.6 6.6 5.3 1.9	Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Berger et al. (2007)
	0.37 0.70 2.0 2.6 2.7 2.8 3.9 4.4 5.4 5.6 6.5 7.3 7.4 7.7 11.3 13.1 30.0 31.3 33.5 33.6 51.5 127.8 0.045 0.12 0.22 1.2 1.3 1.5	OSN OSN OSN NOT NOT OSN OSN CAHA 2.2-m CAHA 2.2-m CAHA 2.2-m Bok Bok CAHA 2.2-m WHT OSN Bok WIYN OSN WHT APO/ARC WIYN CAHA 2.2-m APO/ARC Swift Swift Swift Swift Swift Swift Swift	90Prime 90Prime 90Prime 90Prime VIC-FPS UVOT UVOT UVOT UVOT UVOT UVOT UVOT UVOT	I I R R R R R R R R R R R R R K I R R K V U white r B UVW2	120  300 600 1200 1800 1200  1800 750 1500  10800 1000 3600  5400 3600 200 50 50 1800 886 900 684	19.3 10.0 3.3 6.1 10.8 3.7 1.1 4.6 1.2 1.1 < 0.96 1.2 1.4 17.1 < 2.5 1.6 1.2 < 0.8 6.3 7.5 0.41 0.55 < 2.13 55.0 22.7 30.5 36.6 18.7 6.4 33.7	4.4 2.8 0.70 1.9 2.5 0.97 0.33 0.83 0.49 0.22 0.18 0.37 1.4 0.41 0.24 1.6 0.65 0.06 0.14 18.8 13.0 11.6 6.6 5.3 1.9 9.5	Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Berger et al. (2007)
060121 060313	0.37 0.70 2.0 2.6 2.7 2.8 3.9 4.4 5.4 5.6 6.5 7.3 7.4 7.7 11.3 13.1 30.0 31.3 33.5 33.6 51.5 127.8 0.045 0.012 0.22 1.2 1.3 1.5 1.7 2.9	OSN OSN OSN NOT NOT OSN OSN CAHA 2.2-m CAHA 2.2-m CAHA 2.2-m Bok Bok Bok CAHA 2.2-m WHT OSN Bok WIYN OSN WHT APO/ARC WIYN CAHA 2.2-m APO/ARC Swift Swift Swift Swift Swift Swift Swift Swift Swift	90Prime 90Prime 90Prime 90Prime VIC-FPS UVOT UVOT GMOS UVOT UVOT UVOT UVOT UVOT UVOT UVOT UVOT	I I R R R R R R R R R R R R R R R R R R	300 600 1200 1800 1200 1800 1200 1800 1500 10800 1000 3600 200 50 50 1800 886 900 684 900	19.3 10.0 3.3 6.1 10.8 3.7 1.1 4.6 1.2 1.1 < 0.96 1.2 1.4 17.1 < 2.5 1.6 1.2 < 0.8 6.3 7.5 0.41 0.55 < 2.13 55.0 22.7 30.5 36.6 18.7 6.4 33.7	4.4 2.8 0.70 1.9 2.5 0.97 0.33 0.83 0.49 0.22 0.18 0.37 1.4 0.41 0.24 1.6 0.65 0.06 0.14 18.8 13.0 11.6 6.6 5.3 1.9 9.5 2.3	Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Berger et al. (2007)
	0.37 0.70 2.0 2.6 2.7 2.8 3.9 4.4 5.6 6.5 7.3 7.3 7.4 7.7 11.3 13.1 30.0 31.3 33.5 51.5 127.8 0.045 0.12 0.22 1.2 1.3 1.5 1.7 2.9 3.1	OSN OSN OSN NOT NOT OSN OSN OSN CAHA 2.2-m CAHA 2.2-m CAHA 2.2-m Bok Bok Bok CAHA 2.2-m WHT OSN Bok WIYN OSN WHT APO/ARC WIYN CAHA 2.2-m APO/ARC Swift	GMOS  ALFOSC ALFOSC  90Prime 90Prime 90Prime  NIC-FPS  NIC-FPS UVOT UVOT UVOT UVOT UVOT UVOT UVOT UVOT	I I R R R I R R R R R R R R R K I R R R K K V U white r B UVW2 V UVM2 UVW1	300 600 1200 1800 1200 1800 750 1500 10800 1000 3600 200 50 50 1800 886 900 684 900 900	19.3 10.0 3.3 6.1 10.8 3.7 1.1 4.6 1.2 1.1 < 0.96 1.2 1.4 17.1 < 2.5 1.6 1.2 < 0.8 6.3 7.5 0.41 0.55 < 2.13 55.0 22.7 30.5 36.6 18.7 6.4 33.7 4.7 12.4	4.4 2.8 0.70 1.9 2.5 0.97 0.33 0.83 0.49 0.22 0.18 0.37 1.4 0.41 0.24 1.6 0.65 0.06 0.14 18.8 13.0 11.6 6.6 5.3 1.9 9.5 2.3 2.4	Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Berger et al. (2007)
	0.37 0.70 2.0 2.6 2.7 2.8 3.9 4.4 5.6 6.5 7.3 7.3 7.4 7.7 11.3 13.1 30.0 31.3 33.6 51.5 127.8 0.045 0.12 0.22 1.2 1.3 1.5 1.7 2.9 3.1 3.3	Gemini-N  OSN OSN NOT NOT OSN OSN CAHA 2.2-m CAHA 2.2-m CAHA 2.2-m Bok Bok Bok CAHA 2.2-m WHT OSN Bok WIYN OSN WHT APO/ARC WIYN CAHA 2.2-m APO/ARC Swift	90Prime 90Prime 90Prime 90Prime VIC-FPS VVOT UVOT UVOT UVOT UVOT UVOT UVOT UVOT	I I R R R I R R R R R R R R R R K I R R R K K V U white r B UVW2 UVW1 UVW1 UVW1	300 600 1200 1800 1200 1800 750 1500 10800 1000 3600 200 50 50 1800 886 900 684 900 900 708	19.3 10.0 3.3 6.1 10.8 3.7 1.1 4.6 1.2 1.1 < 0.96 1.2 1.4 17.1 < 2.5 1.6 1.2 < 0.8 6.3 7.5 0.41 0.55 < 2.13 55.0 22.7 30.6 18.7 6.4 33.7 4.7	4.4 2.8 0.70 1.9 2.5 0.97 0.33 0.83 0.49 0.22 0.18 0.37 1.4 0.41 0.24 1.6 0.65 0.06 0.14 18.8 13.0 11.6 6.6 5.3 1.9 9.5 2.3 2.4 1.8	Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Berger et al. (2007)
	0.37 0.70 2.0 2.6 2.7 2.8 3.9 4.4 5.6 6.5 7.3 7.4 7.7 11.3 13.1 30.0 31.3 33.5 33.6 51.5 127.8 0.045 0.12 0.22 1.2 1.3 1.5 1.7 2.9 3.1 3.3 4.5	OSN OSN OSN NOT NOT OSN OSN CAHA 2.2-m CAHA 2.2-m CAHA 2.2-m Bok Bok Bok CAHA 2.2-m WHT OSN Bok WIYN OSN WHT APO/ARC WIYN CAHA 2.2-m APO/ARC Swift	90Prime 90Prime 90Prime 90Prime VIC-FPS VOT UVOT UVOT UVOT UVOT UVOT UVOT UVOT	I I R R R I R R R R R R R R R K I R R R K K V U white R B UVW2 V UVW1 U B	120 300 600 1200 1800 1200 1800 1200  1800 750 1500  10800 1000 3600  5400 3600 200 50 50 1800 886 900 684 900 900 708 886	19.3 10.0 3.3 6.1 10.8 3.7 1.1 4.6 1.2 1.1 < 0.96 1.2 1.4 17.1 < 2.5 1.6 1.2 < 0.8 6.3 7.5 0.41 0.55 < 2.13 55.0 22.7 30.5 36.6 18.7 6.4 33.7 4.7 12.4 7.4 12.2	4.4 2.8 0.70 1.9 2.5 0.97 0.33 0.83 0.49 0.22 0.18 0.37 1.4 0.41 0.24 1.6 0.65 0.06 0.14 18.8 13.0 11.6 6.6 5.3 1.9 9.5 2.3 2.4 1.8 5.8	Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  de Ugarte Postigo et al. (2006)  de Ugarte Postigo et al. (2006)  Roming et al. (2006)  Berger et al. (2007)
	0.37 0.70 2.0 2.6 2.7 2.8 3.9 4.4 5.6 6.5 7.3 7.3 7.4 7.7 11.3 13.1 30.0 31.3 33.6 51.5 127.8 0.045 0.12 0.22 1.2 1.3 1.5 1.7 2.9 3.1 3.3	Gemini-N  OSN OSN NOT NOT OSN OSN CAHA 2.2-m CAHA 2.2-m CAHA 2.2-m Bok Bok Bok CAHA 2.2-m WHT OSN Bok WIYN OSN WHT APO/ARC WIYN CAHA 2.2-m APO/ARC Swift	90Prime 90Prime 90Prime 90Prime VIC-FPS VVOT UVOT UVOT UVOT UVOT UVOT UVOT UVOT	I I R R R I R R R R R R R R R R K I R R R K K V U white r B UVW2 UVW1 UVW1 UVW1	300 600 1200 1800 1200 1800 750 1500 10800 1000 3600 200 50 50 1800 886 900 684 900 900 708	19.3 10.0 3.3 6.1 10.8 3.7 1.1 4.6 1.2 1.1 < 0.96 1.2 1.4 17.1 < 2.5 1.6 1.2 < 0.8 6.3 7.5 0.41 0.55 < 2.13 55.0 22.7 30.6 18.7 6.4 33.7 4.7	4.4 2.8 0.70 1.9 2.5 0.97 0.33 0.83 0.49 0.22 0.18 0.37 1.4 0.41 0.24 1.6 0.65 0.06 0.14 18.8 13.0 11.6 6.6 5.3 1.9 9.5 2.3 2.4 1.8	Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  Levan et al. (2006)  de Ugarte Postigo et al. (2006)  de Ugarte Postigo et al. (2006)  de Ugarte Postigo et al. (2006)  Roming et al. (2006)  Berger et al. (2007)

Table A2 — Continued

GRB	$\delta t$	Telescope	Instrument	Filter	Exposure Time	$F_{\nu}$	σ	References
	(hr)				(s)	(μJy)	(µJy)	
	9.6	Swift	UVOT	UVW1	900	3.0	1.4	
	9.8	Swift	UVOT	U	658	3.4	1.9	
	24.2	Gemini-S	GMOS	r	900	3.7	0.75	Berger et al. (2007)
	48.5	Gemini-S	GMOS	r	1500	1.3	0.18	
0.00502D	240	Gemini-S	GMOS	r	1800	< 0.48		D: 1 (2005)
060502B	16.8	Gemini-N	GMOS	R	1500	< 0.62		Price et al. (2006)
060801	16.0	Hale	LFC	r	1500	< 0.83	0.20	Berger et al. (2007)
061006	14.9	VLT	FORS1	I	1800	4.3	0.20	D'Avanzo et al. (2009)
	45.9	VLT	FORS1	I	1800	2.5	0.17	
	60.0	VLT	FORS1	I	1260	2.6	0.22	
0.61201	49.7	VLT	FORS1	R	645	1.5	0.17	g 1 (2007 )
061201	8.6	VLT	FORS2	I	3640	2.9	0.22	Stratta et al. (2007a)
	9.0 33.1	VLT VLT	FORS2	R I	780 5200	1.9 < 0.91	0.23	
	81.4	VLT VLT	FORS2 FORS2	I	2400	< 0.63		
061210	2.1	Gemini-N	GMOS	r	5400	< 1.44		Berger et al. (2007)
061210	2.1	Magellan/Clay	LDSS3		300	< 2.0		Beiger et al. (2007)
001217	2.0	Magenan/Clay	LDSSS	r	300	< 2.0		
070209	0.29	PROMPT		V	120	< 81.66		Johnson et al. (2007)
070406	24.5	NOT	ALFOSC	R	1800	< 1.26		Malesani et al. (2007b)
070429B	4.8	Gemini-S	GMOS	R	900	< 0.50		Perley et al. (2007)
	27.1	Blanco	ISPI	$\overline{J}$	1080	< 4.0		Nysewander et al. (2007)
070707	11.0	VLT	FORS1	R	1200	2.1	0.039	Piranomonte et al. (2008)
	33.8	VLT	FORS1	R	1200	1.0	0.047	` ,
	37.4	VLT	FORS1	R	180	0.82	0.10	
	59.3	VLT	FORS1	R	1200	0.26	0.020	
	61.0	VLT	ISAAC	J	1800	< 0.60		
	83.2	VLT	FORS1	R	3600	0.078	0.014	
	108.0	VLT	FORS1	R	5100	0.066	0.015	
070714B	0.21	Liverpool		r	60	37.2	6.7	Graham et al. (2009)
	0.27	Liverpool		r	120	43.1	4.6	
	0.30	Liverpool		i	120	50.7	8.6	
	0.40	Liverpool		r	120	37.9	9.8	
	0.44	Liverpool		r	120	38.3	9.0	
	23.6	WHT		R	2400	1.0	0.22	
	24.0	TNG	NICS	J		< 9.1		Covino et al. (2007)
070724A	2.3	Gemini-N	GMOS	g $i$	360	< 1.5		Berger et al. (2009)
	2.3	Gemini-N	GMOS		360	1.1	0.1	
	2.8	Gemini-N	NIRI	Ks	900	9.3	1.5	
	3.1	Gemini-N	NIRI	J	900	3.4	0.3	Berger et al. (2009); this work
	3.4	Gemini-N	NIRI	H	450	7.8	0.44	Berger et al. (2009); this work
	3.7	Gemini-N	NIRI	Ks	900	8.9	1.5	Berger et al. (2009)
070729	8.4	ESO/MPG	GROND	r		< 0.48		Nicuesa Guelbenzu et al. (2012a)
	8.4	ESO/MPG	GROND	J		< 3.02		
070809	11.2	Keck I	LRIS	R	640	0.82	0.17	Perley et al. (2007)
	35.1	Keck I	LRIS	R	222	0.36	0.11	
	11.21	Keck I	LRIS	g	880	0.25	0.051	
700105	35.14	Keck I	LRIS	g	<b>620</b>	< 0.19		V 1' / 1 /2007
070810B	23.4	Keck I	LRIS	R	630	< 0.21		Kocevski et al. (2007)
071112B	6.3	Magellan/Clay	LDSS3	r	1200	< 2.0		Berger & Challis (2007)
071007	9.6	ESO/MPG	GROND	J		< 8.32		Nicuesa Guelbenzu et al. (2012a)
071227	4.2	ESO/MPG	GROND	J	240	< 36.3	0.12	D'A ( 1 (2000)
	7.0 80.9	VLT VLT	FORS2 FORS2	R R	240 540	1.6 0.86	0.12 0.15	D'Avanzo et al. (2009)
080121	55.2	Swift	UVOT	white	2015	< 4.57		Troja et al. (2008)
080123	0.04	Swift	UVOT	white	100	< 22.91		Ukwatta et al. (2008)
080426 <sup>a</sup>	7.5	CAHA 2.2-m	INOT	I white	3300	< 7.06		de Ugarte Postigo et al. (2008)
080503	0.037	Swift Comini N	UVOT	white	98 180	< 14.2		Perley et al. (2009b)
	1.0	Gemini-N	GMOS	r	900	$< 0.20 \\ 0.089$	0.018	
	1.2	Gemini-N Keck I	GMOS	$\stackrel{g}{B}$			0.018	
	1.2	Keck I Keck I	LRIS		300	< 0.21 < 0.21		
	1.3		LRIS	R	630			
	1.5 1.8	Gemini-N Gemini-N	GMOS GMOS	r i	800 800	< 0.081 < 0.078		
	2.2				800 800			
		Gemini-N	GMOS	Z		< 0.16		
	2.4	Gemini-N	GMOS	g	360	< 0.65	0.027	
	<sup>b</sup> 26.0	Gemini-N	GMOS	r	1800	0.27	0.037	
	<sup>b</sup> 47.4	Gemini-N	GMOS	r	1620	0.23	0.038	
	<sup>b</sup> 50.2	Gemini-N	GMOS	g	720	0.12	0.024	
		Gemini-N	GMOS	r	2700	0.19	0.046	
	<sup>b</sup> 74.0	Geillilli-N	GIVIOD	,	2700			
	<sup>b</sup> 74.0 <sup>b</sup> 97.1	Gemini-N Gemini-N Gemini-N	GMOS NIRI	$r K_s$	2880	0.13 < 0.70	0.025	

 Table A2
 — Continued

GRB	δt (hr)	Telescope	Instrument	Filter	Exposure Time (s)	$F_{ u} (\mu \mathbf{J} \mathbf{y})$	$\sigma$ ( $\mu$ Jy)	References
0007024	<sup>b</sup> 128.6	HST	WFPC2	F606W	0.067	0.011		G (2000)
080702A 080905A	12.06 8.5	Loiano NOT	ALFOSC	R R	1800 1800	$< 44.92 \\ 0.72$	0.39	Greco et al. (2008) Rowlinson et al. (2010b)
00070371	14.3	VLT	FORS2	R	2400	0.59	0.20	Rowinison et al. (2010b)
	17.5	ESO/MPG	GROND	r	660	< 3.7		Nicuesa Guelbenzu et al. (2012a)
	17.5 36.0	ESO/MPG VLT	GROND FORS2	J R	660 2400	< 27.5 < 0.30		Rowlinson et al. (2010b)
$080919^{a}$	0.19	ESO/MPG	GROND	J		< 73.3		Nicuesa Guelbenzu et al. (2012a)
081024A	1.9	Faulkes North	LEG	R	600	< 93.93		Melandri et al. (2008)
081024B 081226A	30 0.37	P200 ESO/MPG	LFC GROND	r	R	< 1.36 1.9	0.42	Cenko & Kasliwal (2008) Nicuesa Guelbenzu et al. (2012a)
00122011	1.1	ESO/MPG	GROND	g		0.38	0.12	Triedesa Guersenza et al. (2012a)
	1.1	ESO/MPG	GROND	r		0.63	0.16	
	1.1 1.1	ESO/MPG ESO/MPG	GROND GROND	i z		0.81 1.4	0.31 0.35	
	1.1	ESO/MPG	GROND	$\tilde{J}$		< 7.7	0.55	
081226B	2.7	Swift	UVOT	В	1414	< 14.45		Evans & Hoversten (2008)
090305	0.47	ESO/MPG	GROND	r		3.0	0.084	Nicuesa Guelbenzu et al. (2012a)
	0.56	ESO/MPG	GROND	r		3.1	0.27	
	0.56 0.60	ESO/MPG ESO/MPG	GROND GROND	i r		3.3 2.9	0.59 0.082	
	0.61	Gemini-S	GMOS	r	900	1.8	0.034	Tunnicliffe et al. (2013)
	0.71	ESO/MPG	GROND	g		2.1	0.40	Nicuesa Guelbenzu et al. (2012a)
	0.71 0.79	ESO/MPG ESO/MPG	GROND GROND	r g		2.8 1.9	0.35 0.09	
	0.92	ESO/MPG	GROND	g		2.1	0.31	
	0.92	ESO/MPG	GROND	r		2.1	0.32	
	0.92 0.93	ESO/MPG ESO/MPG	GROND GROND	i		2.5 1.8	0.49 0.085	
	1.1	ESO/MPG	GROND	$g \\ g$		1.7	0.083	
	1.1	ESO/MPG	GROND	r		1.9	0.24	
	1.2 1.3	ESO/MPG Gemini-S	GROND GMOS	r i	900	2.0 1.5	0.13 0.041	Tunnicliffe et al. (2013)
	1.3	ESO/MPG	GROND		900	1.6	0.041	Nicuesa Guelbenzu et al. (2012a)
	1.3	ESO/MPG	GROND	$\frac{g}{i}$		1.9	0.24	,
	1.3 1.4	ESO/MPG	GROND	r		1.5 1.7	0.16 0.082	
	1.4	ESO/MPG ESO/MPG	GROND GROND	r r		1.7	0.082	
	1.5	ESO/MPG	GROND	i		1.9	0.22	
	1.6	ESO/MPG	GROND	r	000	1.7	0.078	T
	1.6 1.6	Gemini-S ESO/MPG	GMOS GROND	r r	900	1.4 1.8	0.026 0.19	Tunnicliffe et al. (2013) Nicuesa Guelbenzu et al. (2012a)
	1.7	ESO/MPG	GROND	r		1.6	0.073	Theuesa Guersenza et an (2012a)
	1.7	ESO/MPG	GROND	r		1.5	0.12	
	1.8 1.8	ESO/MPG ESO/MPG	GROND GROND	r g		1.4 1.0	0.051 0.13	
	1.8	ESO/MPG	GROND	r		1.5	0.12	
	2.0	ESO/MPG	GROND	r		1.6	0.12	
	2.1 2.1	ESO/MPG Gemini-S	GROND GMOS	r r	900	1.2 0.95	0.35 0.027	Tunnicliffe et al. (2013)
	2.1	ESO/MPG	GROND	r	700	1.1	0.040	Nicuesa Guelbenzu et al. (2012a)
000426	21.7	Gemini-S	GMOS	r	1500	< 0.19	7.0	Tunnicliffe et al. (2013)
090426	0.29 0.31	TNT TNT		R R	60 60	101 95.1	7.8 10.1	Xin et al. (2011)
	0.33	TNT		R	60	89.2	9.5	
	0.36	TNT		R	60	77.0	8.2	
	0.38 0.40	TNT TNT		R R	60 60	74.9 69.6	8.8 8.1	
	0.42	TNT		R	60	73.5	8.6	
	0.53	TNT		V	600	49.0	5.7	
	0.71 0.88	TNT TNT		$V \ V$	600 600	36.5 34.5	4.3 4.0	
	1.1	TNT		$\stackrel{V}{V}$	600	36.2	4.0	
	1.2	TNT		V	600	32.4	3.8	
	1.4	TNT		V	600	31.8	4.4	
	1.6 2.0	TNT TNT		$V \ V$	600 600	28.7 24.6	4.0 3.9	
	2.3	TNT		V	600	20.1	3.4	
	2.6	TNT		V	600	20.8	3.5	
	3.0 7.3	TNT TLS		$V \\ I$	600	19.7 15.8	3.6 2.8	Nicuesa Guelbenzu et al. (2011)
	7.5	TLS		R		13.5	1.7	(2011)
	7.5	TLS		R		13.4	1.8	

Table A2 — Continued

	References
12.9	
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	
12.9	
12.9	
12.9	
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	
12.9	
14.6	
14.6	
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	
14.6	
14.6	
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	
090510°	
6.3 6.3 7 7 4.6 1.6 6.4 7 2.6 1.8 6.4 7 3.0 1.8 6.5 7 2.6 1.6 6.7 7 2.6 1.6 6.7 7 2.3 0.71 6.8 7 2.7 0.89 7.1 7 6.9 7 7 7.2 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7	n et al. (2011)
6.3 6.4 7 7 2.6 1.8 6.4 7 3.0 1.8 6.5 7 7 2.6 1.6 6.7 7 7 2.3 0.71 6.8 7 3.8 0.77 6.9 7 7 7.1 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7	elbenzu et al. (2012b)
6.4       r       2.6       1.8         6.4       r       3.0       1.8         6.5       r       2.6       1.6         6.7       r       2.3       0.71         6.8       r       3.8       0.77         6.9       r       2.7       0.89         7.1       r       3.4       0.79         7.2       r       2.9       0.72         7.3       r       2.2       0.75         7.4       r       3.1       0.77         7.6       r       2.3       0.59         7.7       r       1.9       0.40         7.8       r       1.9       0.50         7.9       r       2.2       0.55         8.1       r       1.8       0.59         8.2       r       2.0       0.52         8.3       r       2.0       0.47         8.4       r       2.1       0.46	
6.4       r       3.0       1.8         6.5       r       2.6       1.6         6.7       r       2.3       0.71         6.8       r       3.8       0.77         6.9       r       2.7       0.89         7.1       r       3.4       0.79         7.2       r       2.9       0.72         7.3       r       2.2       0.75         7.4       r       3.1       0.77         7.6       r       2.3       0.59         7.7       r       1.9       0.40         7.8       r       1.9       0.50         7.9       r       2.2       0.55         8.1       r       1.8       0.59         8.2       r       2.0       0.52         8.3       r       2.0       0.47         8.4       r       2.1       0.46	
6.5	
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	
6.8       r       3.8       0.77         6.9       r       2.7       0.89         7.1       r       3.4       0.79         7.2       r       2.9       0.72         7.3       r       2.2       0.75         7.4       r       3.1       0.77         7.6       r       2.3       0.59         7.7       r       1.9       0.40         7.8       r       1.9       0.50         7.9       r       2.2       0.55         8.1       r       1.8       0.59         8.2       r       2.0       0.52         8.3       r       2.0       0.47         8.4       r       2.1       0.46	
6.9       r       2.7       0.89         7.1       r       3.4       0.79         7.2       r       2.9       0.72         7.3       r       2.2       0.75         7.4       r       3.1       0.77         7.6       r       2.3       0.59         7.7       r       1.9       0.40         7.8       r       1.9       0.50         7.9       r       2.2       0.55         8.1       r       1.8       0.59         8.2       r       2.0       0.52         8.3       r       2.0       0.47         8.4       r       2.1       0.46	
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	
7.3     r     2.2     0.75       7.4     r     3.1     0.77       7.6     r     2.3     0.59       7.7     r     1.9     0.40       7.8     r     1.9     0.50       7.9     r     2.2     0.55       8.1     r     1.8     0.59       8.2     r     2.0     0.52       8.3     r     2.0     0.47       8.4     r     2.1     0.46	
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	
7.9     r     2.2     0.55       8.1     r     1.8     0.59       8.2     r     2.0     0.52       8.3     r     2.0     0.47       8.4     r     2.1     0.46	
8.1     r     1.8     0.59       8.2     r     2.0     0.52       8.3     r     2.0     0.47       8.4     r     2.1     0.46	
8.2	
8.3 r 2.0 0.47 8.4 r 2.1 0.46	
8.4 r 2.1 0.46	
86 r 23 048	
2.3 0.40	
8.6 $J$ < 4.8	
8.7 $r$ 2.0 0.47	
8.8 r 1.9 0.44	
8.9 r 1.5 0.47	
9.1 r 1.7 0.43 9.2 r 1.3 0.41	
9.2	
9.5 r 1.5 0.41	
9.7 $r = 0.88 = 0.47$	
9.8 $r = 1.2 = 0.41$	
	ke et al. (2009)
1.7 Gemini-N GMOS $r = 1.8 \times 10^3 = 0.11 = 0.013$ Rowlin	son et al. (2010a)
	& Williams (2009)
25.0 Gemini-N GMOS $r$ $1.8 \times 10^3$ 0.092 0.033 Rowlin	son et al. (2010a)
$4.4 \times 10^3$ Gemini-N GMOS $r = 2.8 \times 10^3 < 0.042$	
	orzi et al. (2009)
	ev et al. (2009)
	lip et al. (2009)
	eliffe et al. (2013)
7.2 VLT HAWK-I <i>K</i> 1320 < 0.85	
8.3 VLT HAWK-I <i>J</i> 1320 < 2.5 10.4 VLT FORS2 <i>R</i> 1200 0.51 0.11	
10.4 VLT FORS2 R 1200 0.51 0.11 31.7 VLT FORS2 R 2400 < 0.17	
	elbenzu et al. (2012a)
	& Mulchaey (2009)
100117A 3.9 Magellan IMACS <i>R</i> 1200 < 0.97 Fon	g et al. (2011)
4.3 ESOMPG GROND r < 0.69 Nicuesa Gu	elbenzu et al. (2012a)
	g et al. (2011)
	elbenzu et al. (2012a)
11.7 ESO/MPG GROND J < 9.7	ov. et al. (2012)
	ey et al. (2012) elbenzu et al. (2012a)

 Table A2
 — Continued

GRB	$\frac{\delta t}{(\text{hr})}$	Telescope	Instrument	Filter	Exposure Time (s)	$F_{ u} (\mu J y)$	$\sigma \ (\mu Jy)$	References
	. ,					$(\mu \mathbf{j} \mathbf{y})$	$(\mu \mathbf{j} \mathbf{y})$	
	33.9	Magellan	PANIC	J	2100	< 1.0		Fong et al. (2013)
100628A	1.1	Gemini-N	GMOS	i	1200	< 0.85		This work
100702A <sup>a</sup>	17.8 0.16	Magellan ESO/MPG	PANIC GROND	J $J$	1620	< 5.43 < 27.29		Nicuesa Guelbenzu et al. (2012a)
100702A	1.7	ESO/MPG	GROND	J		< 21.29 r	< 1.96	Nicuesa Guerbenzu et al. (2012a)
101219A	0.96	Gemini-S	GMOS	i	1620	< 0.40	< 1.70	Fong et al. (2013)
	1.20	Magellan	FourStar	J	1500	< 1.36		
110112A	15.4	WHT	ACAM	i	600	2.8	0.75	Fong et al. (2013)
11011211	$4.0 \times 10^{3}$	Magellan	LDSS3	i	1200	< 0.49	0.75	rong of an (2013)
110112B	17.0	Swift	UVOT	white	3600	< 5.01		Littlejohns et al. (2011)
110420B	10.5	Magellan	IMACS	r	1530	< 1.4		This work
111020A <sup>a</sup>	17.8	Magellan/Clay	LDSS3	r	1080	< 0.83		Fong et al. (2012b)
	17.8	Gemini-S	GMOS	i	1620	< 0.63		T 1100 1 (2012)
1111174	18.5 8.2	VLT GTC	HAWK-I OSIRIS	J	2640 1200	< 0.57 < 0.40		Tunnicliffe et al. (2013)
111117A	8.2 13.7	Gemini-S	GMOS	r r	1200	< 0.40		Sakamoto et al. (2013) Margutti et al. (2012)
	13.7	Gennin 5	GMOS	,	1200	₹ 0.25		magata et al. (2012)
1202204	0.6	Magallan/Cl	I Dees		540	~ A 1		This work
120229A 120305A	9.6 0.23	Magellan/Clay Liverpool	LDSS3	r	540	< 4.1 < 25.85		This work Virgili et al. (2012)
120305A 120521A <sup>a</sup>	18.6	ESO/MPG	GROND	r r	3000	< 25.85 < 0.88		Virgili et al. (2012) Rossi et al. (2012)
12032174	18.6	ESO/MPG ESO/MPG	GROND	$\stackrel{\prime}{J}$	2400	< 20.9		10331 Ct dl. (2012)
120804A	5.5	Gemini-N	GMOS	i	1980	0.17	0.04	Berger et al. (2013)
120817B	24.0	LCO/duPont	WFCCD	R	900	< 1.5		Fong et al. (2012a)
121226A	11.1	NOT	ALFOSC	R	1800	< 2.1		Xu et al. (2012)
130313A <sup>a</sup>	13.4	TNG		r	1800	< 0.44		D'Avanzo et al. (2013)
130515A	0.63	Gemini-S	GMOS	r	240	< 1.4		Cenko & Cucchiara (2013)
130603B	0.19 2.1	Swift Swift	UVOT UVOT	$V \ V$	180 5110	< 199.5 < 54.0		de Ugarte Postigo et al. (2013)
	2.4	Swift Swift	UVOT	$\stackrel{\scriptscriptstyle{V}}{B}$	7700	17.2	4.5	
	5.9	NOT	MOSCA	r	1800	12.6	0.23	
	6.1	WHT	ACAM	z	900	25.4	1.4	
	6.6	WHT	ACAM	i	900	16.4	0.88	
	6.7	CAHA	DLR-MKIII	V	1800	8.3	0.73	
	7.0 7.1	GTC WHT	OSIRIS ACAM	r	30 900	11.0 6.3	0.20 0.34	
	7.1 7.9	Gemini-S	GMOS	g	1440	5.3	0.34	Cucchiara et al. (2013)
	8.2	Magellan/Baade	IMACS	r = g	600	8.6	0.14	Berger et al. (2013)
	9.0	Gemini-S	GMOS	i	1440	12.3	1.2	Cucchiara et al. (2013)
	14.4	Gemini-N	GMOS	z	500	6.5	0.18	de Ugarte Postigo et al. (2013)
	14.5	UKIRT	WFCAM	K	684	13.7	1.3	
	14.6	Gemini-N	GMOS	i	500	4.5	0.12	
	14.7 14.8	UKIRT	WFCAM	J	500 500	9.3 2.9	1.3	
	15.0	Gemini-N Gemini-N	GMOS GMOS	r	500	1.6	0.081 0.06	
	31.2	Gemini-S	GMOS	g r	540	< 0.30	0.00	Cucchiara et al. (2013)
	31.2	Gemini-S	GMOS	i	540	< 0.58		
	32.2	Magellan/Baade	IMACS	r	1200	< 0.46		Berger et al. (2013)
	38.2	Gemini-N	GMOS	g	600	< 0.19	0.05	de Ugarte Postigo et al. (2013)
	38.4	Gemini-N	GMOS	<i>r</i>	600	0.21	0.05	
	38.6 38.6	Gemini-N UKIRT	GMOS WFCAM	$\stackrel{i}{J}$	600 1400	< 0.48 < 3.6		
130716A	19.5	Gemini-N	GMOS	r	900	< 0.35		This work
130822A	21.1	Gemini-N	GMOS	i	600	< 0.58		Cenko et al. (2013a)
130912A	0.27	ESO/MPG	GROND	r		10.5	2.1	Tanga et al. (2013)
	0.38	P60		r		13.0	2.6	Cenko et al. (2013b)
	0.88	P60	10175	r		9.7	2.5	m + 1/2010
	20.0	WHT	ACAM PATIP	g	10200	< 0.87		Tanvir et al. (2013)
	24.8 24.8	Johnson Johnson	RATIR RATIR	r i	19200 19200	< 1.8 < 1.7		Butler et al. (2013)
	24.8	Johnson	RATIR	$\overset{\iota}{Z}$	8060	< 3.7		
	24.8	Johnson	RATIR	Y	8060	< 5.4		
	24.8	Johnson	RATIR	J	8060	< 5.3		
	24.8	Johnson	RATIR	H	8060	< 7.9		
131004A	0.43	NOT	ALFOSC	R		44.3	4.3	Xu et al. (2013)
	1.1 1.4	TNG	DOLORES	R ;	60 300	25.5	2.5	Malesani et al. (2013)
	1.4 1.9	Liverpool Liverpool		i i	300 300	47.4 32.8	4.6 3.2	Kopac et al. (2013)
	2.8	ESO/MPG	GROND	g g	460	6.0	0.58	
	4.0							
	2.8	ESO/MPG	GROND	r	460	9.6	0.93	

Table A2 — Continued

GRB	δt (hr)	Telescope	Instrument	Filter	Exposure Time (s)	$F_{ u}$ $(\mu \mathrm{Jy})$	σ (μ <b>J</b> y)	References
	2.8	ESO/MPG	GROND	z	460	14.0	1.4	
	2.8	ESO/MPG	GROND	J	480	36.5	7.4	
	2.8	ESO/MPG	GROND	H	480	34.0	6.9	
	2.8	ESO/MPG	GROND	K	480	< 47.3		
	7.3	Johnson	RATIR	r	7812	< 1.6		Littlejohns et al. (2013)
	7.3	Johnson	RATIR	i	7812	< 1.9		
	7.3	Johnson	RATIR	Z	3384	< 2.9		
	7.3	Johnson	RATIR	Y	3384	< 4.8		
	7.3	Johnson	RATIR	J	3384	< 3.6		
10110717	7.3	Johnson	RATIR	H	3384	< 3.6		a
131125A <sup>a</sup>	11.8	iPTF		R		< 24.9		Singer et al. (2013a)
131126A <sup>a</sup>	19.5	iPTF		R		< 107.8		Singer et al. (2013b)
140129B	0.032	MASTER II		_	20	7590		Poleshchuk et al. (2014)
	0.053	MASTER II		-	30	3020		
	0.079	MASTER II		-	50	1910		
	0.11	MASTER II		-	70	1450		
	0.15	MASTER II		-	100	692		
	0.20	MASTER II		-	130	525		
	0.26	MASTER II		-	170	363		
	0.33	MASTER II		-	180	275		
	0.43	MASTER II		-	360	191		
140320A	32.9	Johnson	RATIR	r	5040	< 10.7		Littlejohns et al. (2014)
	32.9	Johnson	RATIR	i	5040	< 10.4		
	32.9	Johnson	RATIR	Z	2016	< 24.3		
	32.9	Johnson	RATIR	Y	2016	< 20.2		
	32.9	Johnson	RATIR RATIR	J	2016	< 25.3 < 37.3		
140402 A a	32.9	Johnson Magallan/Banda		H	2016 900			This work
140402A <sup>a</sup> 140414A	29.0 15.2	Magellan/Baade	IMACS	r	1800	< 0.36 < 5.9		This work D'Avanzo et al. (2014a)
140414A 140516A	13.2	TNG TNG	DOLORES DOLORES	r r	720	< 2.38		D'Avanzo et al. (2014a)
140310A	1.8	NOT	ALFOSC	'R	3900	< 0.50		Gorosabel et al. (2014b)
	11.6	Subaru	IRCS+AO188	K'	5880	< 0.30		Minowa et al. (2014)
	12.4	Gemini-N	GMOS	i	1800	< 0.04		This work
	13.0	Johnson	RATIR	$\overset{\iota}{J}$	3200	< 5.8		Butler et al. (2014)
140606A	7.3	AAO	Mili	R	120	< 6.46		Volnova et al. (2014)
140619B <sup>a</sup>	35.5	Magellan/Baade	Fourstar	J	770	< 2.54		This work
140622A	0.07	ESO/MPG	GROND	r	770	< 0.80		Tanga et al. (2014)
11002211	0.07	ESO/MPG	GROND	J		< 11.4		runga et al. (2011)
140903Aa	12.2	DCT	LMI	r		7.6	3.1	Cenko et al. (2014)
140930B	3.0	WHT	ACAM	r		7.4		Tanvir et al. (2014)
	7.3	ESO/MPG	GROND	r		2.0	0.41	Graham et al. (2014)
	7.8	Magellan/Baade	IMACS	i	900	2.2	0.30	This work
	7.8	MMT	MMTCam	r	1200	0.97	0.20	This work
	13.1	Keck	MOSFIRE	$K_s$	315	3.1	0.97	Perley & Jencson (2014)
	13.3	Keck	MOSFIRE	$\widetilde{J}$	396	2.1	0.43	
141205A	11.4	Nanshan		R		< 33.2		Xu et al. (2014)
141212A	5.4	CAHA 1.23-m		Ι		< 11.1		Gorosabel et al. (2014a)
150101B	39.8	Magellan/Baade	IMACS	r	1200	2.26	0.34	Fong et al., in prep.
	63.6	Magellan/Baade	IMACS	r	1200	1.4	0.38	
	231	TNG	NICS	J		< 7.59		D'Avanzo et al. (2015)
	257	Gemini-South	GMOS	r	1710	< 0.76		Fong et al., in prep.
	349	TNG	NICS	J		< 3.98		D'Avanzo et al. (2015)
	$\sim 360$	VLT	HAWK-I	H		< 1.45		van der Horst et al. (2015)
	694	Magellan/Baade	IMACS	r	1200	< 0.53		Fong et al., in prep.
150120A	2.0	Gemini-North	GMOS	r	1800	< 0.36		This work

**Note.** — Upper limits correspond to  $3\sigma$ . All optical and near-IR fluxes are corrected for Galactic extinction in the directions of each bursts (Schlegel et al. 1998; Schlafly & Finkbeiner 2011). Instruments, exposure times, and flux density uncertainties are provided whenever available.

<sup>a</sup> Optical observing constraint, due to delayed precise localization, sightline with high Galactic extinction, crowded field, or proximity to a bright star which contaminates the GRB position.

Optical re-brightening observed in GRB 080503 is unlikely the afterglow (Perley et al. 2009b).
 Full simultaneous giz-band light curves are available in Nicuesa Guelbenzu et al. (2012b).

Table A3
Short GRB Radio afterglow catalog

GRB	$\delta t$ (dy)	Facility	Mean Frequency (GHz)	$F_{\nu}$ $(\mu Jy)$	σ (μ <b>J</b> y)	References
050202	0.52	VLA	4.86	< 120		Frail & Soderberg (2005)
050202 050509B	0.32	WSRT	4.86	< 66		van der Horst et al. (2005)
050709	1.6	VLA	8.46	< 115		Fox et al. (2005)
	2.5	VLA	8.46	< 114		(=,
	4.5	VLA	8.46	< 74		
	7.5	VLA	8.46	< 40		
050724A	0.57	VLA	8.46	173	30	Berger et al. (2005)
	1.68	VLA	8.46	465	29	D '(2006)
050813	9.10 1.64	VLA VLA	8.46 8.46	< 259 < 55		Panaitescu (2006) Cameron & Frail (2005a)
050906	3.91	VLA	8.46	< 92		Cameron & Frail (2005a) Cameron & Frail (2005b)
050925	0.40	WSRT	4.9	< 72		van der Horst (2005)
051105A	0.54	VLA	8.5	< 51		Frail & Cameron (2005)
051221A	0.91	VLA	8.46	155	30	Soderberg et al. (2006)
	1.94	VLA	8.46	< 72		
060313	2.12	VLA	8.46	< 110		Soderberg & Frail (2006)
060801	0.49	VLA	8.46	< 105		Soderberg et al. (2006)
	1.2	WSRT	4.9	< 72		van der Horst (2006)
061210	5.2	WSRT	4.9	< 81		Chandra & E:1 (2006)
061210	1.9	VLA	8.46 8.46	< 102		Chandra & Frail (2006)
070429B 070714B	0.59 15.6	VLA VLA	8.46 8.46	< 300 < 135		Chandra & Frail (2007b) Chandra & Frail (2007d)
070714B 070724A	1.06	VLA	8.46	< 255		Chandra & Frail (2007a)
07072471	9.49	VLA	8.46	< 255		Chandra & Frail (2007e)
070923	4.98	VLA	8.46	< 135		Chandra & Frail (2007c)
071112B	1.69	ATCA	8.7	< 141		Wieringa et al. (2007)
080503	3.05	VLA	8.46	< 54		Frail & Chandra (2008)
080702A	0.74	VLA	8.46	< 156		Chandra & Frail (2008b)
081024A	1.03	VLA	8.46	< 156		Chandra & Frail (2008a)
081024B	4.1	VLA	8.46	< 114		Chandra et al. (2008)
081226A	58.9 58.4	ATCA ATCA	4.9 4.9	< 540 < 588		Moin et al. (2009a)
081226B 090417A	0.37	VLA	8.46	< 72		Moin et al. (2009b) Chandra & Frail (2009)
090510	1.98	VLA	8.46	< 84		Frail & Chandra (2009)
090515	0.87	VLA	8.46	< 60		Berger & Fong (2009a)
090621B	0.61	VLA	8.46	< 54		Berger & Fong (2009c)
090715A	1.25	VLA	8.46	< 64		Berger & Fong (2009d)
091117	2.33	VLA	8.46	< 120		Berger & Fong (2009b)
100625A	0.83	VLA	4.90	< 192		This work
100628A	0.77	VLA	5.8	< 291		This work
101224A	3.72	VLA	5.8	< 56		This work
110112A 110112B	1.87 0.85	VLA VLA	5.0 5.8	< 75 < 51		This work This work
110112 <b>D</b>	1.80	VLA	5.8	< 66		This work
	2.92	VLA	5.8	< 48		
	12.10	VLA	5.8	< 36		
110420B	0.65	VLA	4.90	< 77		This work
	7.65	VLA	5.8	< 116		
111020A	0.67	VLA	5.8	< 39		Fong et al. (2012b)
111117A	0.49	VLA	5.8	< 18		Margutti et al. (2012)
111121A	0.84	VLA	5.8	< 100		This work
120229A	0.38	VLA	5.8	< 54		This work
120305A	3.32 0.16	VLA VLA	5.8 5.8	< 66 < 18		This work
120303A	2.25	VLA	5.8	< 18		THIS WOLK
120521A	1.08	ATCA	34	< 95		Hancock et al. (2012a)
120804A	0.9	VLA	5.8	< 20		Berger et al. (2013)
	2.2	ATCA	34	< 200		Hancock et al. (2012b)
	4.2	ATCA	34	< 120		
121226A	1.75	VLA	5.8	< 30		This work
	4.76	VLA	5.8	< 45		
130313A	0.77	VLA	5.8	< 60	1 4 4	This work
130603B	0.37	VLA	4.9	125	14.4	Fong et al. (2014)

 Table A3
 — Continued

(dy)         (GHz)         (μJy)         (μJy)           0.37         VLA         6.7         119         9.1           1.43         VLA         4.9         < 57           1.44         VLA         21.8         < 50           4.32         VLA         4.9         < 51           4.32         VLA         6.7         < 26           84.31         VLA         6.7         < 24           84.31         VLA         6.7         < 34           130716A         2.51         VLA         5.8         < 33         This work           130912A         0.91         VLA         5.8         < 30         This work           130912A         0.91         VLA         5.8         < 29         This work           131004A         0.14         CARMA         93         < 300         This work           131224A         0.20         VLA         6.0         < 33         This work           140516A         0.62         VLA         6.0         < 24         This work           140619B         1.51         VLA         6.0         < 21         This work           14093A         0.40         VLA <th>GRB</th> <th><math>\delta t</math></th> <th>Facility</th> <th>Mean Frequency</th> <th><math>F_{ u}</math></th> <th>σ</th> <th>References</th>	GRB	$\delta t$	Facility	Mean Frequency	$F_{ u}$	σ	References
1.43		(dy)		(GHz)	(μJy)	(μJy)	
1.43		0.37	VLA	6.7	119	9.1	
1.44		1.43	VLA	4.9	< 57		
4.32		1.43	VLA	6.7	64.9	15.2	
4.32		1.44	VLA	21.8	< 50		
84.31       VLA       4.9       < 69		4.32	VLA	4.9	< 51		
S4.31		4.32	VLA	6.7	< 26		
130716A   2.51   VLA   5.8   < 33   This work   130822A   15.9   VLA   5.8   < 30   This work   130912A   0.91   VLA   5.8   < 29   This work   3.10   VLA   5.8   < 42   This work   131004A   0.14   CARMA   93   < 300   This work   131224A   0.20   VLA   6.0   < 33   This work   2.27   VLA   6.0   < 33   This work   2.27   VLA   6.0   < 24   This work   4.67   VLA   6.0   < 21   40619B   1.51   VLA   6.0   < 36   This work   140622A   3.09   VLA   6.0   < 36   This work   140903A   0.40   VLA   6.0   110   9.5   This work   140903A   0.40   VLA   6.0   187   8.7   2.45   VLA   6.0   128   15.1   7.42   GMRT   1.39   102   33   Nayana & Chandra (2014)   9.24   VLA   6.0   81.9   14.7   This work   9.24   VLA   9.8   < 69   18.24   VLA   9.8   < 69   18.24   VLA   9.8   < 25.5   This work   140930B   0.29   VLA   9.8   < 224.4   141212A   0.45   VLA   6.0   27.0   8.1   7.72   VLA   6.0   21.9   8.3   7.72   VLA   9.8   21.3   6.4   7.649   VLA   9.8   < 48.6   This work   150101B   5.73   VLA   9.8   < 375   Fong et al., in prep.		84.31	VLA	4.9	< 69		
130822A   15.9		84.31	VLA	6.7	< 34		
130912A	130716A	2.51	VLA	5.8	< 33		This work
3.10	130822A	15.9	VLA	5.8	< 30		This work
131004A	130912A	0.91	VLA	5.8	< 29		This work
131224A   0.20		3.10	VLA	5.8	< 42		
2.27         VLA         6.0         < 33           140516A         0.62         VLA         6.0         < 24	131004A	0.14	CARMA	93	< 300		This work
140516A         0.62         VLA         6.0         < 24	131224A	0.20	VLA	6.0	< 33		This work
4.67       VLA       6.0       < 21		2.27	VLA	6.0			
140619B       1.51       VLA       6.0       < 36	140516A			6.0	< 24		This work
140622A       3.09       VLA       6.0       < 36				6.0			
140903A       0.40       VLA       6.0       110       9.5       This work         2.45       VLA       6.0       187       8.7         2.45       VLA       9.8       151       8.9         4.70       VLA       6.0       128       15.1         7.42       GMRT       1.39       102       33       Nayana & Chandra (2014)         9.24       VLA       6.0       81.9       14.7       This work         9.24       VLA       9.8       < 69				6.0			This work
2.45       VLA       6.0       187       8.7         2.45       VLA       9.8       151       8.9         4.70       VLA       6.0       128       15.1         7.42       GMRT       1.39       102       33       Nayana & Chandra (2014)         9.24       VLA       6.0       81.9       14.7       This work         9.24       VLA       9.8       < 69							This work
2.45       VLA       9.8       151       8.9         4.70       VLA       6.0       128       15.1         7.42       GMRT       1.39       102       33       Nayana & Chandra (2014)         9.24       VLA       6.0       81.9       14.7       This work         9.24       VLA       9.8       < 69	140903A						This work
4.70       VLA       6.0       128       15.1         7.42       GMRT       1.39       102       33       Nayana & Chandra (2014)         9.24       VLA       6.0       81.9       14.7       This work         9.24       VLA       9.8       < 69							
7.42         GMRT         1.39         102         33         Nayana & Chandra (2014)           9.24         VLA         6.0         81.9         14.7         This work           9.24         VLA         9.8         < 69							
9.24         VLA         6.0         81.9         14.7         This work           9.24         VLA         9.8         < 69							
9.24 VLA 9.8 < 69 18.24 VLA 6.0 < 120 140930B 0.29 VLA 9.8 < 25.5 This work 3.27 VLA 9.8 < 24.4 141212A 0.45 VLA 6.0 < 25.2 This work 3.76 VLA 6.0 27.0 8.1 7.72 VLA 6.0 21.9 8.3 7.72 VLA 9.8 21.3 6.4 76.49 VLA 6.0 < 19.5  150101A 1.43 VLA 9.8 < 48.6 This work 150101B 5.73 VLA 9.8 < 375 Fong et al., in prep.			GMRT	1.39	102		Nayana & Chandra (2014)
18.24       VLA       6.0       < 120				6.0	81.9	14.7	This work
140930B       0.29       VLA       9.8       < 25.5							
3.27       VLA       9.8       < 24.4							
141212A       0.45       VLA       6.0       < 25.2	140930B						This work
3.76 VLA 6.0 27.0 8.1 7.72 VLA 6.0 21.9 8.3 7.72 VLA 9.8 21.3 6.4 76.49 VLA 6.0 <19.5  150101A 1.43 VLA 9.8 <48.6 This work 150101B 5.73 VLA 9.8 <375 Fong et al., in prep.		3.27	VLA	9.8	< 24.4		
7.72 VLA 6.0 21.9 8.3 7.72 VLA 9.8 21.3 6.4 76.49 VLA 6.0 <19.5 150101A 1.43 VLA 9.8 <48.6 This work 150101B 5.73 VLA 9.8 <375 Fong et al., in prep.	141212A						This work
7.72 VLA 9.8 21.3 6.4 76.49 VLA 6.0 <19.5  150101A 1.43 VLA 9.8 <48.6 This work 150101B 5.73 VLA 9.8 <375 Fong et al., in prep.							
76.49         VLA         6.0         < 19.5           150101A         1.43         VLA         9.8         < 48.6							
150101A 1.43 VLA 9.8 < 48.6 This work 150101B 5.73 VLA 9.8 < 375 Fong et al., in prep.						6.4	
150101B 5.73 VLA 9.8 < 375 Fong et al., in prep.							
<i>8</i> , 1 1				,			
150120A 0.00 VI A 0.8 /20.6 This work							
130120A 0.70 Y LA 5.0 < 27.0 THIS WORK	150120A	0.90	VLA	9.8	< 29.6		This work

**Note**. — For bursts with multiple contemporaneous upper limits, we only display the most constraining limits. Upper limits are  $3\sigma$ .