Please check the examination details below	before entering your candidate information
Candidate surname	Other names
Centre Number Candidate Num Pearson Edexcel Interna	
Tuesday 16 January 2	2024
Afternoon (Time: 1 hour 30 minutes)	Paper WMA12/01
Mathematics International Advanced Sub Pure Mathematics P2	sidiary/Advanced Level
You must have: Mathematical Formulae and Statistical Ta	ables (Yellow), calculator

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- Fill in the boxes at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 - there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 10 questions in this question paper. The total mark for this paper is 75.
- The marks for each question are shown in brackets
 - use this as a guide as to how much time to spend on each guestion.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶



1.		$f(x) = ax^3 + 3x^2 -$	8x + 2	where a is	a constant		
Given th	at when $f(x)$ is	s divided by $(x-2)$) the remai	nder is 3, fir	nd the value	of a.	(3)

\sim	$\times \times$
\times	$\times\!\!\times$
\otimes	$\times\!\!\times\!\!\times$
\otimes	\approx
88	\approx
\Rightarrow	$\otimes\!$
XX	$\times\!\!\times\!\!\!\times$
\Leftrightarrow	$\times\!\!\times\!\!\times$
X	$\times\!\!\times\!\!\!\times$
≫	$\times\!\!\times\!\!\times$
XX	$\times\!\!\!\times\!$
₩	
×	GIZÎ X
×	AREA
X	× × ×
××	
$\otimes \rangle$	$\times\!\!\times\!\!\times$
X	
×.	
\otimes	9 EX
Xi	$\times\!\!\!\times\!\!\!\!\times$
\otimes	
××	2
X	
X.	
\otimes	
×	
X	
××	
×	
$\otimes \forall$	$\propto \times$
\otimes	\otimes
88	
\otimes	
XX	
W	DO NOT WRITE IN THIS A
\otimes	
×	
≫	$\times\!\!\times\!\!\times$
\times	$\times\!\!\times$
$\langle \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \!$	$\langle\!\langle\!\rangle\!\rangle$
\times	XX
$\times\!\!\times$	$\times\!\!\times\!\!\times$
$\stackrel{>}{>}$	
\times	$\times\!\!\times$
$\times\!\!\!\times$	$\times\!\!\times\!\!\times$
\otimes	
XX	$\times\!\!\times\!\!\times$
\otimes	
XX	$\times\!\!\!\!>$
\otimes	$\times\!\!\times\!\!\times$
$\otimes >$	∞
XX	$\times\!\!\times\!$
$\otimes \rangle$	$\otimes \otimes$
XX	$\times\!\!\times\!\!\times$
₩	
X	$\times \times$
₩	MDIX
o ca	
$\triangle A$	32
\otimes	
$\stackrel{\times}{\times}$	AREA
	SAR
	SAR
	TIS AR
	THIS AR
	TEIS AR
	NTHIS AR
	IN THIS AR
	IN THIS AR
	TELN THIS AR
	TEIN THIS AR
	STEIN THIS AR
	RITE IN THIS AR
	VRITE IN THIS AR
	WRITE IN THIS AR
	TWRITE IN THIS AR
	OT WRITE IN THIS AR
	OT WRITE IN THIS AR
	NOT WRITE IN THIS AR
	NOTWRITEINTHIS
	O NOT WRITE IN THIS AR
	O NOT WRITE IN THIS
	NOTWRITEINTHIS
	O NOT WRITE IN THIS
	DO NOT WRITE IN THIS
	DO NOT WRITE IN THIS
	DO NOT WRITE IN THIS
	DO NOT WRITE IN THIS
	REA DO NOT WRITE IN THIS
	REA DO NOT WRITE IN THIS
	AREA DO NOT WRITE IN THIS
	AREA DO NOT WRITE IN THIS
	AREA DO NOT WRITE IN THIS
	AREA DO NOT WRITE IN THIS
	HIS AREA DO NOT WRITE IN THIS
	THIS AREA DO NOT WRITE IN THIS
	THIS AREA DO NOT WRITE IN THIS
	THIS AREA DO NOT WRITE IN THIS
	IN THIS AREA DO NOT WRITE IN THIS
	EINTHIS AREA DO NOT WRITE IN THIS
	TEIN THIS AREA OF THE STATE OF THIS
	TEIN THIS AREA OF THE STATE OF THIS
	TEIN THIS AREA OF THE STATE OF THIS
	RITE IN THIS AREA
	RITE IN THIS AREA
	WRITE IN THIS AREA DO NOT WRITE IN THIS
	WRITE IN THIS AREA DO NOT WRITE IN THIS
	WRITE IN THIS AREA DO NOT WRITE IN THIS
	WRITE IN THIS AREA DO NOT WRITE IN THIS
	WRITE IN THIS AREA DO NOT WRITE IN THIS

Question 1 continued	
	(Total for Question 1 is 3 marks)
	(10tal for Question 1 is 3 marks)



2.	Find the coefficient	of the	term in	$n x^7$	of the	binomial	expansion	of

$$\left(\frac{3}{8} + 4x\right)^{12}$$

giving your	answer in	simplest	form.

(3	1
v	,

~~~~~	V
*****	
4	
× <del>M</del> ×	
× 0 Z	_
× vo	_
※董※	
× <del>F</del> ×	
	_
$\times$ ini $\times$	_
× <b>£</b> ×	_
3	
$\times$	
$\circ \circ \circ$	
Ö	_
ă	
*****	_
****	l _
VVVVV -	-
	-
****	
× III	_
Va .	-
<b>I</b>	l _
	_
× <del>**</del> ×	-
	l _
<u> </u>	
	_
$\times$	
$\times \times \times \times \times \times$	-
*****	
XXXXX	
$\sim\sim\sim\sim$	-
000000 -	
	-
$\times$	
	_
XXXXX	
XXX	
	_
<u> </u>	
66 <b>3</b> 5666 −	-
VA .	
	_
弄	
	_
ш	
66 <del>000</del> 66 -	-
<u> </u>	
	I -
×O×	
Ž	_
88	
×8×	-
~~~~~~	

Question 2 continued	
	(Total for Question 2 is 3 marks)



3.	The circle <i>C</i>	
Э.		
	• has centre $A(3, 5)$	
	• passes through the point $B(8, -7)$	
	(a) Find an equation for C.	(3)
	The points M and N lie on C such that MN is a chord of C .	
	Given that MN	
	• lies above the <i>x</i> -axis	
	• is parallel to the <i>x</i> -axis	
	• has length $4\sqrt{22}$	
	(b) find an equation for the line passing through points M and N.	
	(b) That an equation for the line passing through points in that it.	(3)



3	8			
3				
3		X	0	K
Š		×	S	2
	2			2
		Ç		2
×		♥	×	\geq
×			Κ	>
×		0	×	>
2		Х		×
	⟨		Ü	×
0		×		K
		×	٦	K
×	å	XVXXX	ij	۴
×	J	Ģ	ą	
×	À	G	K	3
×	Я	ш	k	3
	S	Z	4	☲
0	4	Ø		z
5	ð	X	۹	₹
X	À		ч	9
×	Ž	ø	d	ia)
X			K	>
2	ì	ñ	ø	١,
Q	S	w	Ų	4
0	á	è	Ĥ	蛸
	á	è	à	ıκ
Χ		SJ	В	2
Χ	2	V	ę	₹
×	X	H	ú	÷
	y	O	K	\geq
	S		ú	≟
2	ς	14	ø	≺
0	₫	Ŗ,	þ	疃
0	Þ	蜔	ij	팯
٥	2	×	Š	2
ĸ,	d	IJ	É	j
×	Ž	V	ď	Ź
Κ	ď		ú	è
×	y	Ó	ż	2
	9	ď	ø	3
2	Ý	m	ø	ч
\Diamond	Þ	×	þ	o(
)	Þ	94	è	6
	à	INI	ú	ø
Χ	à	S	ä	p
X	Ž	S		\rangle
Κ	y		ŕ	۵
2	Ч	r	7	5
	S	ρ	į	Ķ
\supset	4	ĸ	2	a
0	ζ,	X	_	Z
×	2	7		,
	λ	Ç	6	×
×)	$^{\circ}$		>
×	ì	e	ę	ŧ.
×	y	Ľ	ú	а
2	ς		2	3.
0	Ý	r	7	٦
0	Þ	Þ	ì	4
4				2
×	2	V	×	2
	2	Ç	K	2
×)	C	K	>
×			Κ)
×			×	S
	5		d	5
2	S	Χ	ė	×
0	Κ			×
				×
0			٦	K
		×	Ŋ	2
S	2	S	Ŋ	2
			×	
2	S	X	X	5
8	3		X	3
3	3	Š	×	3
3	3	X	\ \ \	3
3			\ < < <	3
3			\ < < <	>
\ \ \ \			×	
			× < < < < < <	
			~<<<<	
			~ < < < < < < < < <	
			~<<<<<	
			~<<<<<	
			~ < < < < < < < < < < < < < < < < < < <	
			~ < < < < < < < < < < < <	
	77777			
	77777			
	77777			

Question 3 continued
(Total for Question 3 is 6 marks)
(=====================================



4. (a) Sketch the curve with equation

$$y = a^{-x} + 4$$

where a is a constant and a > 1

On your sketch show

- the coordinates of the point of intersection of the curve with the y-axis
- the equation of the asymptote to the curve.

(3)

х	-4	-1.5	1	3.5	6	8.5
у	13	6.280	4.577	4.146	4.037	4.009

The table above shows corresponding values of x and y for $y = 3^{-\frac{1}{2}x} + 4$

The values of y are given to four significant figures, as appropriate.

Using the trapezium rule with all the values of *y* in the table,

(b) find an approximate value for

$$\int_{-4}^{8.5} \left(3^{-\frac{1}{2}x} + 4 \right) \mathrm{d}x$$

giving your answer to two significant figures.

(3)

(c) Using the answer to part (b), find an approximate value for

(i)
$$\int_{-4}^{8.5} \left(3^{-\frac{1}{2}x} \right) dx$$

(ii)
$$\int_{-4}^{8.5} \left(3^{-\frac{1}{2}x} + 4 \right) dx + \int_{-8.5}^{4} \left(3^{\frac{1}{2}x} + 4 \right) dx$$

(3)

Question 4 continued	



Question 4 continued	

Question 4 continued			
(Total for Question 4 is 9 marks)			



5. (i) Find the value of

$$\sum_{r=1}^{\infty} 6 \times \left(0.25\right)^r$$

(3)

(ii) A sequence $u_1, u_2, u_3,...$ is defined by

$$u_1 = 3$$

$$u_{n+1} = \frac{u_n - 3}{u_n - 2} \qquad n \in \mathbb{N}$$

(a) Show that this sequence is periodic.

(2)

(b) State the order of this sequence.

(1)

(c) Hence find

$$\sum_{n=1}^{70} u_n$$

(2)

Question 5 continued	



Question 5 continued			

Question 5 continued			
(Total for Question 5 is 8 marks)			



6. (a) Given that

$$2\log_4(x+3) + \log_4 x = \log_4(4x+2) + \frac{1}{2}$$

show that

$$x^3 + 6x^2 + x - 4 = 0$$

(4)

(b) Given also that -1 is a root of the equation

$$x^3 + 6x^2 + x - 4 = 0$$

(i) use algebra to find the other two roots of the equation.

(3)

(ii) Hence solve

$$2\log_4(x+3) + \log_4 x = \log_4(4x+2) + \frac{1}{2}$$

(1)

Question 6 continued



Question 6 continued			

	- 4
	١,
- (XXXXXX)	
	-
**** ********************************	
	-
XXXXXXX	
×vx×	
XX	
	-
XX#XXXX	
	-
- XX	
	-
SITE	
XX XX XX	
- XX	
× c ×	
	-
- (XX	
- XX	
- 000020000	
IOT	_
XXXXXXX	
XX XXX XX	
- XXX	
×O×	
- 00004000	
_ XX XX	
XXXXXXX	
ŏ	
	_
	-
	-
	-
	_
XXXXXX	
000000000000000000000000000000000000000	
XXXX	
	-
- XX XX	
- XXAAXX	
0.5	
XXXXX	
	-

<u> </u>	
XXMMXXX	
	_
- (XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX	
_ XX XX XX	
XXXXXXX	
XXXXXXXX	-
	-
	-
m	-
	-
	-
E	-
	-
	-
	-
/RIT	-
VRIT	-
FWRIT	-
VRIT	-

ζ		\rangle	C	2	₹	2	₹	\rangle	
ζ		>	4	λ	ģ	è	6	\rangle	
S	?		d	Б	?	2	ì	5	
ζ	?	ς	2	۹	۴	ę	2	ς	
≺	\rangle	≺	à	ø	۶	۹	ù	⟨	
	>		X	Ľ	4	4	ð	⟨	
3			>		\rangle		\rangle		
Ĉ	5		>	<	5	C	5		
2	Ś	2	S			?	S	2	
>	ς	2				2	ς	>	
Ş		\rangle	≺		⟨	\rangle	⟨	\rangle	
>				\rangle		>		\rangle	
S		S				>	2	5	
ξ	?	S			2	S	2	S	
S	>			ς		ς	?	ς	
5					\rangle	⟨	>		
	\rangle						\rangle		
2	5	ď				(>	ď	
ζ	5		S			2	S	2	
	Ś				ς	?	ς		
2			≺	\rangle		\rangle	⟨	\rangle	
ζ		\rangle				\rangle		>	
S	2					5		S	
	?		2	ς	?	ς	2		
۶				≺	>	⟨		Ś	
			\rangle	<		⟨	>	⟨	
3	>						>		
₹	>	2				2	S	2	
2		2		2	S	2		2	
	Ś		ς	2	ς	?	ς		
		>				\rangle		\rangle	
ξ						\rangle		\rangle	
>		>	Z.	7	ú	è	ø	5	
	?		Ą	Ę	2	ž,	2		
ζ		S	à	s	2		2	ς	
<	\rangle		ı	Ľ		_	ł	◁	
ξ	>		Š	ā	Š	Z	ž		
			2	7	2	Z	S	2	
	Ś	2	q	ė	S		5	2	
2			C	7	7	9	ų	>	
>		\rangle	⟨	\geq	(2	s	\rangle	
≻			à	Ļ	å	2	š	S	
		S				S		S	
ς	?		2	₹	7	7		ς	
⋖	\rangle	⟨	y	ę		R	7	◁	
ξ			ì	á	8	ĸ	è		
			'n	ď	5	2	5	0	
	S	2	9	۲	3	ŧ	3	?	
?	ζ				ς	2	S	2	
2			∢	7			Ŗ	>	
ζ			4			à	ú	\rangle	
		>	4	4	4	-	ú	>	
>	?							S	
	?	S	à	Ь	à	Ş	à		
ζ			Ľ	ķ	β	Ę	ē	ς	
ζ		<	Χ	Ľ	è	6		⋖	
	>		ļ	τ					
Ĉ	5		'n	ø	ķ	ø	ij		
?		2	Ý	Þ	ķ	ġ	Ŋ	2	
	ς		4	þ	Ц	B	ú		
>		\rangle	4	ù	6	2	≤	\rangle	
>			4	è	ë	ą	۳	\rangle	
	(>		Ş	š		þ	>	
ζ			2	2	7	5		S	
ς		S	d	Ĺ	2	S	2	ς	
	\rangle		ľ	Ľ	7	Ì	3	◁	>
3	>		à	è	9	ij	k	ď	
Ĉ		₹	y	ç	S	á	β	<	
2			S	2	S	Z	Š		
	S	2				2	5	2	
>	Ś	>	Ų.	ŕ	۴	7	۴	>	
>		>			4	\geq		>	
>	?		4	ø	Š	9		>	
S		S	ð	ķ	è	ú	ø	5	
ζ	?	ς	2	a	ē	É	2	ς	
≺	>	≺	X	Ľ	\geq	2	'n	◁	
ξ	>		y	₹	7	₹	7		
				2		1		2	

Question 6 continued	
(Total for Question 6 is 8 marks)	
·	_



- **7.** Wheat is grown on a farm.
 - In year 1, the farm produced 300 tonnes of wheat.
 - In year 12, the farm is predicted to produce 4000 tonnes of wheat.

Model *A* assumes that the amount of wheat produced on the farm will increase by the same amount each year.

(a) Using model *A*, find the amount of wheat produced on the farm in year 4. Give your answer to the nearest 10 tonnes.

(3)

Model *B* assumes that the amount of wheat produced on the farm will increase by the same percentage each year.

(b) Using model *B*, find the amount of wheat produced on the farm in year 2. Give your answer to the nearest 10 tonnes.

(3)

(c) Calculate, according to the two models, the difference between the total amounts of wheat predicted to be produced on the farm from year 1 to year 12 inclusive. Give your answer to the nearest 10 tonnes.

(3)

Question 7 continued



Question 7 continued

××		
\otimes	×	×
X	X	X
×	X,	×
X	Š	×,
X	Š	S
X	S	×.
X	S	S
K.	×	S
\Diamond	×	×
\Diamond	X	×
×		1
X	Z	Ζ,
X	Ľ	M
×	ń	ŵ
×	ij	ě
X		ø
X	$\overline{\mathbf{x}}$	4
X		2
×	u	а
\times	ĵω	ŵ
\times	畁	۳
×	ĕ	H
X	Ľ	4
×	ŦÇ	X
×	÷	ŝ
\times	ď	4
X		×
X	×	×
$\langle \rangle$	L	Ц
\propto	Ŀ	\geq
\times	ŗ	Z
X	뿐	×
×	c	c
X	Ξ	X,
X	No.	3
××		3
××	ě	×
\times	r	7
\sim	p	N
××	V	u
X	ě	à
X	ď	4
×	Z	X
X	10200	3
X	9	4
X	ø	5
X	6	لخ
$\langle \rangle$	>	×
\otimes	X	X
×	X	X
X	×	×
×	×	×
X	×	×,
×	×	×
X	\$	S
X	S	S
\Diamond	×	(
$\langle \rangle$	×	×
×	×	×
×	×	X
X	Χ	X
×	×	×.
X	×Ç	×,
X	S	X.
X	S	
K.	۸.	S
\Diamond		S
$\times \times$	×	
	Š	
×		
\otimes		
8		
8		
× ×		
× ×		
× ×		
× × ×		

)		
			S				
2		2			ς		
	⟨	>					
		>			K		d
					2		
			\rangle				
		<			5		
2	5	2					
		>	ζ	à		ė	ø
			₹	9	¥		ė
		ς	à	S	2		4
			1	٤	J	Ľ	
	5	2	S	Z	3	7	3
		2	Ч	r	٩	ρ	۹
			₫	9	e	9	۹
		5	Ġ	7	á	è	ø
		ς	į	ij	ń		
		≺				₹	1
		2	١	ż	1	á	í
			4	ū	d	r	
		\rangle	4	3	2	2	i
	2	S	2	3	Ź	3	ž
		⟨	J	۲	3	ř	
)	ó	ð	K	è
			S	r	5	7	
		2	1	P	×	ø	
			ð	5	Ĉ		
	2	5	à	Ĺ	ú	É	5
			S	ś	ó	۴	1
			y	ę	7	۴	
2		2	9	ė	á	Ė	Ė
		\rangle			≺		
		5	d	b			
			2	B	ø	Ц	į
			Ì	Ľ		۷	
2		2	Ŋ	۲	۹	7	
			G	à	6	è	í
		>	₹	ä	Z	ž	í
			4	E	á	ď	
			2	Š	9	ď	9
		2	Ŋ	ē	b	ĕ	i
2	Κ	2	9	ø		2	9
		>	4	š	á	₽	Ħ
			2	3	2		
		≺	J	Ĺ	2	≤	
		<	Ŋ	ľ	7	7	
	S	2	S	à	ń	ú	ś
		\rangle	4	ľ			
			2	9	ø	ij	ę
		ς	λ	á	Ė		į
		<	à	ά		S	
	>		S	7	3	₹	
		2	Κ	2	≤	2	
			4	Р	₹	7	
		S	ð	ķ	ú	ś	d
				ŝ	ä	š	
	>	2	Y	۲	5	₹	
		2	4	þ	é	ė	d
					≺		<
		S	2		2	2	
7			\rangle				
		S			3	S	<

Question 7 continued
(Total for Question 7 is 9 marks)



8.	(i) Use a counter example to show that the following statement is false	
	" $n^2 + 3n + 1$ is prime for all $n \in \mathbb{N}$ "	(2)
	(ii) Use algebra to prove by exhaustion that for all $n \in \mathbb{N}$	(2)
	" $n^2 - 2$ is not a multiple of 4"	(4)
		(4)

-88	\sim	X
-50	~~	X
-88	\sim	X
- QX	$\times \times$	×2
$-\infty$	$\times \times$	X
$-\infty$	$\times \times$	X
	$\times\!\!\times$	X
\sim		\times
\sim	$\times\!\!\times$	X
	$\times\!\!\times\!\!\times$	
$-\infty$	Ø	ik O
- 22	6	X
\sim	$\nabla \nabla$	U×
	0. 11.	×
	×22	×
\times	482	\sim
- X	(XX	ĒΧ
- 52	1	ĸ
$-\infty$	TIS A	i Q
\sim	$\times \times$	\times
\sim		Ю
$-\infty$	XX.	20
$-\infty$	XX	*0
- ~~	7.5	٧×
- 22	نالو	ŧΧ
\sim		×
		×
	\sim	X
		癶
- X		ďΧ
$-\infty$	××	ĸ.
\sim	$\times \times$	\times
	JIJK.	K
\sim	2.0	50
$-\infty$	RITE	ĸÓ
- 62	*X	×
- 82	7	ţΧ
\sim	P	ŖΧ
\times	*	ų×
\times	۰	×
-X		×
		50
\propto	DO NOT	82
\propto	×	ĸQ
\sim	ХX	X
$-\infty$	100	NO.
- 22		KS.
\sim	90	×
-8		\sim
		♥
	\sim	\propto
- X		ľ×
-525		ĸ
$-\infty$		X
\sim	\times	K
\sim	XX	\times
\sim	$\times \times$	X
$-\infty$	××	X
- (>)	XX	X
- 22	\sim	X
\sim	$\Diamond \Diamond$	×
	\sim	
	\sim	
\sim		
\otimes	\sim	X
\otimes	\otimes	\otimes
$\stackrel{\circ}{\times}$	\otimes	×
	$\overset{\wedge}{\otimes}$	
*	$\overset{\wedge}{\otimes}$	
	×	× ×

	FA	
	₽	
	8 E A	
	REA	
	0:	
	0:	
	0:	
	0:	
	0:	
	0:	
	0:	
	0:	
	0:	
	0:	
	0:	
	0:	
	0:	
	0:	
	0:	
	0:	
	0:	
	0:	
	0:	
	0:	
	0:	
	0:	
	0:	
	0:	
	WRITE IN THIS ARE	
	0:	

Question 8 continued
(Total for Question 8 is 6 marks)
(Lotal for Question o is o marks)



9. In this question you must show detailed reasoning.

Solutions relying entirely on calculator technology are not acceptable.

(i) Solve, for $0 \le x < 360^{\circ}$, the equation

$$\sin x \tan x = 5$$

giving your answers to one decimal place.

(6)

(ii)




Figure 1

Figure 1 shows a sketch of part of the curve with equation

$$y = A\sin\left(2\theta - \frac{3\pi}{8}\right) + 2$$

where A is a constant and θ is measured in radians.

The points P, Q and R lie on the curve and are shown in Figure 1.

Given that the y coordinate of P is 7

(a) state the value of A,

(1)

(b) find the exact coordinates of Q,

(3)

(c) find the value of θ at R, giving your answer to 3 significant figures.

(4)

Question 9 continued



Question 9 continued				

×				×		
X	þ	ζ		×		
				×	X	
Ŏ	?				×	
Q					×	
$^{\circ}$				×		
	۶			×	X	
				×		
×	?					
ŏ			Ś	×	×	
♦	4	è	ġ	ġ	X	
♦	ζ	2	3	3	۹	
Q	9	ķ			4	
	S	á	į	4	ă	
X	J	ķ	j	Ę	ê	
X	à	₹	è	q	ĝ	
	ď	Š	Ż	d	ķ	
ŏ	ŝ	S		á	4	
Ö	d	ķ	d	r	ī	
♦	ú	è	ø	ŷ	b	
\Diamond	ģ	è	٩	Ř	b	
\Diamond	9	è	4	Ŀ	š	
X	J	E	\$	ė	×	
	y	₹		×	0	
X	Ì	4	è	ġ	ä	
X	à	ø	É	5		
Š	d	ģ	ę	ě	P	
Ŏ	ζ	Ş	ζ	×	X	
\Diamond	1	Ŀ	₫	Ę	¥	
Q	Í	È	<u> </u>		\times	
X	g	g	S	Š	Ź	
X	3	į	d	ğ	2	
X	Į	ľ	j		ø	
X	å	ζ	Š	Š	į	
	à	¢	É		ģ	
Ó	j	S	ě	ĕ	Ņ	
Ó	ģ	è		<		
♦	1	P	5		7	
\Diamond	S	è	ø	į	٩	
X	9	ķ	í	ø	ď	
X	þ	ŧ	ģ	3	ġ	
	Ì	d		2	-	
X	\rangle	<	2	×	Û	
X	d	ľ	7	7	3	
Š	2	3	þ	ě	ø	
Ŏ	Ý	P	ę	9	۹	
\Diamond	ġ	þ	8	į	Ħ	
Q				×		
X	S			×		
X	S	<		×		
Χ	>			×		
8				×	×	
Ŏ	2	S			×	
\Diamond		\geq			×	
♡						
X	S	2		×		
	>	<		×		
Χ	þ			×		
X				×		
		S		K		
♦	Ć			ì	×	
♦				0		
Ø	S			×	×	
	\rangle	Š		×		
Χ	>			×		
Х				×		
ŏ	?	S				
Ö	<				×	
Ò	Ś	2	S	'n	ě	
Q	9	É	Ś	1	ß	
×	Ś	í		Ŕ	ĵ	
X	Į	ŧ	į	á		
8	d	Ć	9	ľ	Ź	
Ŏ	(Ž	Ś	4	Š	
\Diamond	9	ø	ξ	1	ķ	
2	S			×	X	
×	ÿ	ľ	j	f	j	
X	à	Ś	í	š	í	
X	à	6	ý	Ě	Á	
Ó	4	è	d	ĺ	ú	
♦	j	b	ζ		Š	
\Diamond	9	ď	₹	×	×	
X	Ś	ė	Ś	5	ğ	
X	1	ġ	é	é	ź	
X	à	6	è	ú	Ť	
K		S	è	×	ķ	
х	ű	ď.		ĺ	d	
	j	,				
Š	1	į	3	,	ě	
Š		į	3		×	֡
X					X	
×						
				CONTRACTOR OF THE PERSON AND ADDRESS OF THE		
				A X MAN AND AND AND AND AND AND AND AND AND A		
				THE RESERVE THE PARTY OF THE PA		

Question 9 continued	
	Total for Question 9 is 14 marks)



10. In this question you must show detailed reasoning.Solutions relying entirely on calculator technology are not acceptable.




Figure 2

Figure 2 shows a sketch of the curve with equation

$$y = \frac{1}{2}x^2 + \frac{1458}{\sqrt{x^3}} - 74 \qquad x > 0$$

The point P is the only stationary point on the curve.

(a) Use calculus to show that the x coordinate of P is 9

(4)

The line l passes through the point P and is parallel to the x-axis.

The region R, shown shaded in Figure 2, is bounded by the curve, the line l and the line with equation x = 4

(b) Use algebraic integration to find the exact area of R.

(5)

Question 10 continued



Question 10 continued	
	(Total for Question 10 is 9 marks)
	TOTAL FOR PAPER IS 75 MARKS

