Please check the examination deta	ails below before ento	ering your candidate information
Candidate surname		Other names
Pearson Edexcel	Centre Number	Candidate Number
International		
Advanced Level		
	Paper	
Time 1 hour 30 minutes	reference	WMA14/01
Mathamatica		A A
Mathematics		
International Advance	d Level	
Pure Mathematics P4	00. 0.	
r die Mathematics F 4		
		J
You must have:		Total Marks
Mathematical Formulae and Stat	tistical lables (Ye	ellow), calculator

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use black ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
- there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 9 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
- use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.
- Good luck with your examination.

Turn over ▶







1. Given that k is a constant and the binomial expansion of

$$\sqrt{1+kx}$$
 $|kx| < 1$

in ascending powers of x up to the term in x^3 is

$$1 + \frac{1}{8}x + Ax^2 + Bx^3$$

- (a) (i) find the value of k,
 - (ii) find the value of the constant A and the constant B.

(5)

(b) Use the expansion to find an approximate value to $\sqrt{1.15}$

Show your working and give your answer to 6 decimal places.

(2)



∞	×		K			
	×	Ž	ζ			
\times	×	X		3	5	
XX	×	X	≷	S	₹	
	×	2	5			
X	×	X	2		>	
\times	×	Ş	3	S	ζ	
	×	2	5			
X	×					
\propto	×	2	S	2		
\times		Χ	>	Ś	2	
炎	1	ĕ	3	Ē		
\propto	i	Ñ	Š	7		
X	1	H		Ś	ķ	
$\langle \langle \rangle$	1	A	Ľ	5	2	
\times	i	E	j	Ŕ	5	
	×	8	7	3		
\otimes	1	ŭ	ŕ	١		
X	į	=	ě	2	Š	
\otimes	3	ă	ľ	Ź		
\otimes		K	5	3		
X	J	5	2	5		
\otimes	2	ě	j	è	Κ	
\times	Ì	Ę	Ž	Ė	Ś	
\times	×	ं	<	5		
\otimes		Ù	Ľ		K	
XX		K	5	3	δ	
\otimes	ł	K	2	\$		
\otimes		â	É	è	5	
×	ŝ	á	ζ	S		
	×	£	ξ	2		
X	A N	Š	2			
\times		K	ij	į	K	
\times		Ŕ	7	١		
X	3	3	è	5		
\propto	á	2	2	2		
XX	×	\times	2	S		
X	1		Š	J		
X	y	ñ	é	ć		
\otimes	ĺ	×	è	4		
\times	×				5	
\otimes	×	Š	?			
∞	×	2	S	2	S	
XX	×				>	
\otimes	×	Ş	?	5		
\otimes	×	2	S		S	
XX	×				\geq	
\otimes	×	2	ζ			
\times		X	>		S	
$\times\!\!\!\times$	×	S	?	S	2	
	×	Ŕ	5			
X		X	>			
燚	×	ŏ	3	S	ζ	
\otimes	×	2	5			
X	×	X	?			
\otimes	×	2	Ś	2		
XX	×	X	۶			
\times	×	X	2			
\propto				á		
		7	Į	Ś	8	
XX	1	Ñ	ĺ			
\times	1	N M				
×		X M M X				
×		X X X X X X				
*						

			ĺ		
9		Š			
Ŋ	ŧ	3		Į	
2	S	2	į	2	
3	2	2	2	3	
2		2		Š	
3		2			
3	è	Š		2	
3	è	Š		3	
3	è	Š			
3	è	Š		2	ĺ
3	è	Š		1	
3	è	Š		< 1 < 1	ĺ
3	è	Š		\ \ \	
3	è	Š		3	
3	è	Š		3	
3	è	Š		3	
3				1	
3				1	
3				1	
3				1	
3				1	
3				1	
				1	
				1	
				1	
				1	
				1	
				1	
				1	
				1	
				1	
				1	
				1	
				1	
				1	
				1	
				1	
				1	
				1	
				1	
				1	
				1	
				1	
				1	
				1	
				1	
				1	
				1	

	Leave
	blank
Question 1 continued	



uestion 1 continued	bla

Question 1 continued	

5

Turn over ▶

2.




Figure 1

Figure 1 shows a sketch of part of the curve with equation

$$y = \frac{9}{(2x-3)^{1.25}} \qquad x > \frac{3}{2}$$

The finite region R, shown shaded in Figure 1, is bounded by the curve, the line with equation y = 9 and the line with equation x = 6

This region is rotated through 2π radians about the x-axis to form a solid of revolution.

Find, by algebraic integration, the exact volume of the solid generated.

(7)

Question 2 continued	
	Q2
(Total 7 marks)	
(10tai / marks)	+



3.




Figure 2

A bowl with circular cross section and height 20 cm is shown in Figure 2.

The bowl is initially empty and water starts flowing into the bowl.

When the depth of water is $h \, \text{cm}$, the volume of water in the bowl, $V \, \text{cm}^3$, is modelled by the equation

$$V = \frac{1}{3}h^2(h+4) \qquad 0 \leqslant h \leqslant 20$$

Given that the water flows into the bowl at a constant rate of $160\,\mathrm{cm^3\,s^{-1}}$, find, according to the model,

(a) the time taken to fill the bowl,

(2)

(b) the rate of change of the depth of the water, in cm s⁻¹, when h = 5

(5)

xxxx.
$\times\!\!\times\!\!\times$
>>>>
$\times\!\!\times\!\!\times$
М
<u> </u>
× S
ш
DO NOT WRITE IN THIS AREA
XE
<u> </u>
×9
$\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times$
>>>
<₹
S
O NOT WRITE IN THIS AREA
×2
×oc
3
\times
\sim 0
∞
>>>>
$\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times$
EA.
05
A
ARE
ARE
THIS ARE
THIS ARE
E IN THIS ARE
TE IN THIS ARE
TE IN THIS ARE
TE IN THIS ARE
WRITE IN THIS ARE
WRITE IN THIS ARE
NOT WRITE IN THIS ARE
WRITE IN THIS ARE

	Leave
	blank
Question 3 continued	
	1



uestion 3 continued	
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_

Question 3 continued
Q3
(Total 7 marks)



(8)

4. Use algebraic integration and the substitution $u = \sqrt{x}$ to find the exact value of

$$\int_{1}^{4} \frac{10}{5x + 2x\sqrt{x}} \, \mathrm{d}x$$

Write your answer in the form $4\ln\left(\frac{a}{b}\right)$, where a and b are integers to be found.

(Solutions relying entirely on calculator technology are not acceptable.)

	Leave
	blank
Question 4 continued	



stion 4 continued	

Question 4 continued	blank
	Q4
(Total 8 marks)	



5. A curve has equation

$$y^2 = y e^{-2x} - 3x$$

(a) Show that

$$\frac{dy}{dx} = \frac{2ye^{-2x} + 3}{e^{-2x} - 2y}$$

(4)

The curve crosses the y-axis at the origin and at the point P.

The tangent to the curve at the origin and the tangent to the curve at P meet at the point R.

(b) Find the coordinates of R.

(5)

Question 5 continued	blank
	Q5
(Total 9 marks)	



blank

(1)

6.




Figure 3

Figure 3 shows a sketch of the curve C with parametric equations

$$x = 2\cos 2t$$
 $y = 4\sin t$ $0 \le t \le \frac{\pi}{2}$

The region *R*, shown shaded in Figure 3, is bounded by the curve, the *x*-axis and the *y*-axis.

- (a) (i) Show, making your working clear, that the area of $R = \int_0^{\frac{\pi}{4}} 32 \sin^2 t \cos t \, dt$
 - (ii) Hence find, by algebraic integration, the exact value of the area of R. (6)
- (b) Show that all points on C satisfy $y = \sqrt{ax + b}$, where a and b are constants to be found. (3)

The curve C has equation y = f(x) where f is the function

$$f(x) = \sqrt{ax + b} \qquad -2 \leqslant x \leqslant 2$$

and a and b are the constants found in part (b).

(c) State the range of f.



	blank
Question 6 continued	Dialik
Question o continueu	



uestion 6 continued		

7. Relative to a fixed origin O, the line l has equation

$$\mathbf{r} = \begin{pmatrix} 1 \\ -10 \\ -9 \end{pmatrix} + \lambda \begin{pmatrix} 4 \\ 4 \\ 2 \end{pmatrix} \qquad \text{where } \lambda \text{ is a scalar parameter}$$

Given that \overrightarrow{OA} is a unit vector parallel to l,

(a) find \overrightarrow{OA}

(2)

The point X lies on l.

Given that X is the point on l that is closest to the origin,

(b) find the coordinates of X.

(5)

The points O, X and A form the triangle OXA.

(c) Find the exact area of triangle OXA.

(3)



	Leave
	blank
Question 7 continued	



Question 7 continued	

8. (a) Given that y = 1 at x = 0, solve the differential equation

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{6xy^{\frac{1}{3}}}{\mathrm{e}^{2x}} \qquad y \geqslant 0$$

giving your answer in the form $y^2 = g(x)$.

(7)

(b) Hence find the equation of the horizontal asymptote to the curve with equation $y^2 = g(x)$.

(2)

	1:
uestion 8 continued	



	Leave
Question 8 continued	blank
Question o continueu	

		blank
Question 8 continued		
		Q8
	_	
	(Total 9 marks)	



9.	(i)	Relative to a fixed origin O , the points A , B and C have position vectors \mathbf{a} , \mathbf{b} and \mathbf{c} respectively.			
		Points A, B and C lie in a straight line, with B lying between A and C.			
		Given $AB:AC = 1:3$ show that			
		$\mathbf{c} = 3\mathbf{b} - 2\mathbf{a}$			
		(3)			
	(ii)	Given that $n \in \mathbb{N}$, prove by contradiction that if n^2 is a multiple of 3 then n is a multiple of 3			
		(5)			

	blank
Question 9 continued	blank
Question 5 continued	



Question 9 continued	Leave blank
	Q9
(Total 8 marks) TOTAL FOR PAPER IS 75 MARKS	
END END	