| Please check the examination det | ails below | before enter | ring your candidate information |
|---|------------|--------------|---------------------------------|
| Candidate surname | | | Other names |
| Pearson Edexcel International Advanced Level | Centre | e Number | Candidate Number |
| Tuesday 8 Oc | tok | er 2 | 2019 |
| Afternoon (Time: 1 hour 30 minu | utes) | Paper Re | eference WCH11/01 |
| Chemistry Advanced Subsidiary Unit 1: Structure, Bond Organic Chemi | | nd Intr | oduction to |
| Candidates must have: Scient Ruler | ific calc | ulator | Total Marks |

Instructions

- Use **black** ink or **black** ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
 - there may be more space than you need.

Information

- The total mark for this paper is 80.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.
- You will be assessed on your ability to organise and present information, ideas, descriptions and arguments clearly and logically, including your use of grammar, punctuation and spelling.
- A Periodic Table is printed on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Show all your working in calculations and include units where appropriate.
- Check your answers if you have time at the end.

Turn over ▶







SECTION A

Answer ALL the questions in this section.

You should aim to spend no more than 20 minutes on this section.

For each question, select one answer from A to D and put a cross in the box \boxtimes . If you change your mind, put a line through the box \boxtimes and then mark your new answer with a cross \boxtimes .

- 1 Which element is in the d-block of the Periodic Table?
 - A argon
 - **B** chlorine
 - C iron
 - **D** sodium

(Total for Question 1 = 1 mark)

- 2 What is the equation for the **third** ionisation energy of aluminium?
 - \square A Al(g) \rightarrow Al³⁺(g) + 3e⁻
 - \square **B** Al(s) \rightarrow Al³⁺(s) + 3e⁻
 - \square **C** $AI^{2+}(q) \rightarrow AI^{3+}(q) + e^{-}$
 - \square **D** $Al^{2+}(s) \to Al^{3+}(s) + e^{-}$

(Total for Question 2 = 1 mark)

3 The first three ionisation energies of carbon are shown.


| | 1st | 2nd | 3rd |
|--|------|------|------|
| Ionisation energy / kJ mol ⁻¹ | 1086 | 2353 | 4621 |

Which is the approximate fourth ionisation energy, in kJ mol⁻¹, of carbon?

- **■ B** 6200
- **☑ C** 11000
- ☑ **D** 38000

(Total for Question 3 = 1 mark)

4 The chart shows the first ionisation energy of each of the elements from fluorine to magnesium.



Which is the approximate first ionisation energy, in kJ mol⁻¹, of aluminium (atomic number 13)?

- **■ B** 600
- **◯ C** 900
- D 1200

(Total for Question 4 = 1 mark)

- **5** The decrease in first ionisation energy down Group 1 in the Periodic Table is caused by the **increase** in
 - A force of attraction between the nucleus and outer electron
 - **B** number of neutrons in the nucleus
 - □ C number of protons in the nucleus
 - ☑ D shielding of the outer electron from the nuclear charge.

(Total for Question 5 = 1 mark)

 $\textbf{6} \quad \text{What is the relative formula mass of hydrated ammonium iron(II) sulfate, } (NH_4)_2Fe(SO_4)_2.6H_2O?$

[Relative atomic masses (A_r): H = 1.0 N = 14.0 O = 16.0 S = 32.1 Fe = 55.8]

- **■ B** 302
- **C** 312

(Total for Question 6 = 1 mark)

7 How many **atoms** are there in 36.0 g of water?

[Avogadro constant = $6.02 \times 10^{23} \text{ mol}^{-1}$]

- \triangle **A** 3.010 × 10²³
- **B** 1.204×10^{24}
- \square **C** 2.408 × 10²⁴
- \square **D** 3.612 × 10²⁴

(Total for Question 7 = 1 mark)

8 Some ionic radii are shown.

| lon | Ionic radius/nm |
|-----------------|-----------------|
| Na ⁺ | 0.102 |
| K ⁺ | 0.138 |
| F ⁻ | 0.133 |
| CI ⁻ | 0.180 |

Which compound has the strongest ionic bonding?

- A sodium fluoride
- B sodium chloride
- **C** potassium fluoride
- D potassium chloride

(Total for Question 8 = 1 mark)

- **9** In which pair are the ions isoelectronic?
 - \boxtimes **A** Ca²⁺ and S²⁻
 - **B** K⁺ and Br⁻
 - **C** Li⁺ and F⁻
 - \square **D** Mg²⁺ and Cl⁻

(Total for Question 9 = 1 mark)

- **10** The bonding **within** an ammonium ion, NH₄, is formed by

 - B covalent and dative covalent bonding only
 - **C** covalent and ionic bonding only
 - **D** covalent, dative covalent and ionic bonding

(Total for Question 10 = 1 mark)



- 11 Which diagram best represents the electron density map of a hydrogen chloride molecule?




 \boxtimes B



X C



 \boxtimes D



(Total for Question 11 = 1 mark)

12 What is the polarity of the Al—CI bond and the polarity of a trigonal planar AlCl₃ molecule?

| X | Α |
|---|---|

 \boxtimes B

 \times C

 \boxtimes D

| Polarity of Al—Cl bond | Polarity of AICI ₃ molecule |
|------------------------|--|
| non-polar | non-polar |
| non-polar | polar |
| polar | non-polar |
| polar | polar |

(Total for Question 12 = 1 mark)

13 What is the atom economy, by mass, for the formation of iron in this reaction?

$$Fe_2O_3 + 3CO \rightarrow 2Fe + 3CO_2$$

- [A_r values: C = 12.0 O = 16.0 Fe = 55.8]
- **△ A** 29.7%
- **B** 45.8%
- **C** 55.9%
- ☑ D 71.7%

(Total for Question 13 = 1 mark)

14 A 2 kg sample of water contains 40 parts per million (ppm) by mass of nitrate ions.

What is the mass, in g, of nitrate ions in this sample?

- \triangle **A** 8 × 10⁻²
- **B** 5×10^{-5}
- \boxtimes **C** 8 × 10⁻⁵
- \triangle **D** 5 × 10⁻⁸

(Total for Question 14 = 1 mark)

15 A sample of hydrated calcium sulfate, $CaSO_4$.**x** H_2O , was heated to constant mass. 3.405 g of anhydrous calcium sulfate and 0.900 g of water were formed.

What is the value of \mathbf{x} ?

[Relative formula mass: $CaSO_4 = 136.2$]

- **■ B** 2
- **区** 3
- \square D 4

(Total for Question 15 = 1 mark)



16 During a cracking reaction, each molecule of an alkane with formula C₁₀H₂₂ formed only two molecules of ethene and one molecule of hydrocarbon **A**.

What is the molecular formula of **A**?

- \triangle A C₆H₁₀
- B C₆H₁₄
- \square **D** C_8H_{18}

(Total for Question 16 = 1 mark)

17 Curly arrows are used in reaction mechanisms.

The curly arrow shown represents the movement of

- A an electron from a bond to an atom, forming free radicals
- **B** an electron from a bond to an atom, forming ions
- C a pair of electrons from a bond to an atom, forming free radicals
- **D** a pair of electrons from a bond to an atom, forming ions

(Total for Question 17 = 1 mark)

18 The structure of a diene is shown.

How many σ bonds and π bonds are there in one molecule of this diene?

| | | σ bonds | π bonds |
|---|---|----------------|-------------|
| X | A | 7 | 2 |
| X | В | 7 | 4 |
| X | C | 9 | 2 |
| X | D | 9 | 4 |

(Total for Question 18 = 1 mark)



19 When hydrogen bromide, HBr, reacts with propene, a higher percentage of 2-bromopropane forms than 1-bromopropane.

Which is the best explanation for this?

- ☑ A 1-bromopropane is more stable than 2-bromopropane
- **B** 2-bromopropane is more stable than 1-bromopropane
- ☑ C a primary carbocation is more stable than a secondary carbocation
- **D** a secondary carbocation is more stable than a primary carbocation

(Total for Question 19 = 1 mark)

20 Propene, C₃H₆, is produced in the dehydration of propanol.

$$C_3H_7OH \rightarrow C_3H_6 + H_2O$$

What is the mass, in g, of propene formed from 3.42 g of propanol when the yield is 85.2 %?

[Relative molecular masses (M_r): $C_3H_7OH = 60$ $C_3H_6 = 42$]

- **B** 2.39
- **C** 2.91
- **D** 4.16

(Total for Question 20 = 1 mark)

TOTAL FOR SECTION A = 20 MARKS

SECTION B

Answer ALL the questions.

Write your answers in the spaces provided.

- **21** This question is about cycloalkanes.
 - (a) When alkanes from crude oil are reformed, the products include cycloalkanes.

Write the equation for reforming hexane into cyclohexane using **skeletal** formulae for the organic compounds.

(2)

(b) The skeletal formula of cycloalkane ${\bf D}$ is shown.



(i) Give the name of **D**.

(1)

(ii) Give the molecular formula of **D**.

(1)

(c) There are four structural isomers of C_6H_{12} with a ring of four carbon atoms. One of these isomers is shown, in the first box. Complete the **skeletal** formulae of the other three isomers. (2) (d) A cycloalkane, ${\bf E}$, has a molar mass of $126\,\mathrm{g\,mol}^{-1}$. Deduce the molecular formula of **E**. (1)

(e) A sample of gaseous cyclopentane with a volume of 25 cm³ was mixed with 250 cm³ of oxygen (an excess) and the mixture was ignited. Only gaseous products were formed.

$$2C_5H_{10}(g)\,+\,15O_2(g)\,\rightarrow\,10CO_2(g)\,+\,10H_2O(g)$$

Calculate the volume of each gas remaining after the reaction.

All the gas volumes were measured at the same temperature and pressure.

(3)

| (f) Cyclobutane, C_4H_8 , reacts with chlorine in sunlight. (i) Name the mechanism and type of reaction that is occurring. | (2) |
|--|------|
| (ii) Complete the equation for the initiation step of this reaction mechanism. Include appropriate curly arrows. CI——CI → | (2) |
| (iii) Write the equations for the two propagation steps to form chlorocyclobutane. Use C_4H_8 as the formula for cyclobutane. Curly arrows and state symbols are not required. | (2) |
| (iv) A small amount of a hydrocarbon forms in this reaction. Deduce the skeletal formula of this hydrocarbon. Justify your answer. Skeletal formula of product | (2) |
| Justification | |
| (Total for Question 21 = 18 ma | rks) |



- **22** This question is about atomic structure and gases.
 - (a) Chlorine exists as two isotopes with mass numbers 35 and 37.
 - (i) State the number and type of each of the particles in the **nucleus** of a chlorine-35 atom.

(2)

(ii) Complete the electronic configuration of a chloride **ion**, Cl⁻, using the s, p, d notation.

(1)

1s².....

(iii) A sample of chlorine contains 75.53% of chlorine-35 atoms.

Calculate the relative atomic mass of this sample of chlorine. Give your answer to **two** decimal places.

(2)

(b) Fluorine has one naturally occurring isotope with mass number 19.

Chlorine and fluorine react to form chlorine trifluoride, CIF₃.

(i) Draw a dot-and-cross diagram to show the bonding in a molecule of chlorine trifluoride.Show the outer shell electrons only.


(2)

(ii) State, in terms of electrons, what is unusual about the dot-and-cross diagram you have drawn.

(1)

(iii) Complete the mass spectrum to show the peaks you would expect for the molecular ion CIF_3^+ .

(3)





(iv) Calculate, using the ideal gas equation, the volume in ${\bf cm^3}$ occupied by 0.0200 mol of CIF₃ gas at a temperature of 60 °C and a pressure of 1.28 \times 10⁵ Pa. Give your answer to an appropriate number of significant figures.

$$[pV = nRT R = 8.31 \,\mathrm{J}\,\mathrm{mol}^{-1}\,\mathrm{K}^{-1}]$$

(4)

(Total for Question 22 = 15 marks)

- **23** This question is mainly about alkenes.
 - (a) A few drops of bromine water are added to separate test tubes of propane and propene and the mixtures are shaken.

Describe what is seen at the end of each experiment.

(2)

Propane

Propene

- (b) Give the reagents and conditions for each of these conversions.
 - (i) $CH_2 = CH_2 \rightarrow CH_3CH_2OH$

(1)

(ii) $CH_2 = CH_2 \rightarrow CH_2OHCH_2OH$

(1)

(c) Draw the structure of *Z*-3-methylpent-2-ene.

(1)

(d) Exactly 720 cm³ of hydrogen gas, measured at room temperature and pressure (r.t.p.), reacted with 0.010 mol of an alkene to form an alkane.

Deduce the number of double bonds in one molecule of the alkene. You **must** show your working.

[Molar volume of gas at r.t.p. = $24\ 000\ \text{cm}^3\ \text{mol}^{-1}$]

(2)

(e) A student drew a mechanism for the addition of bromine to ethene.

Describe the three changes needed to correct this student's mechanism.

(3)

| Change 1 |
|----------|
| |
| |
| Change 2 |
| |
| |

Change 3...

(f) Part of the structure of a polymer is shown.

Draw the structure of the monomer used to make this polymer.

(1)

(Total for Question 23 = 11 marks)

| 24 | This q | uestion is about phosphorus and some of its compounds. | |
|----|------------|---|--------------|
| | (a) Th | ne atomic number of phosphorus is 15. | |
| | (i) | Complete the electronic configuration of a phosphorus atom using the electrons-in-boxes notation. | (1) |
| | 411 | [Ne] 3s 3p | |
| | (11) | Explain why the first ionisation energy of phosphorus is greater than that of su | (2) |
| | | | |
| | | | |
| | | | |
| | (b) Ph | nosphorus has a melting temperature of 44°C. | |
| | | nosphorus has a melting temperature of 44°C. licon has a melting temperature of 1410°C. | |
| | Sili | | icon. (3) |
| | Sili | licon has a melting temperature of 1410°C. | |
| | Sili | licon has a melting temperature of 1410°C. | |
| | Sili | licon has a melting temperature of 1410°C. | |
| | Sili | licon has a melting temperature of 1410°C. | (3) |
| | Sili | licon has a melting temperature of 1410°C. (plain why the melting temperature of phosphorus is much lower than that of sil | (3) |
| | Sili | licon has a melting temperature of 1410°C. (plain why the melting temperature of phosphorus is much lower than that of sil | (3) |
| | Sili | licon has a melting temperature of 1410°C. (plain why the melting temperature of phosphorus is much lower than that of sil | (3) |
| | Sili | licon has a melting temperature of 1410°C. | (3) |
| | Sili | licon has a melting temperature of 1410°C. | (3) |



- (c) Phosphorus(V) chloride, PCI₅, exists as covalent molecules in the gaseous state.
 - (i) Complete the table for a PCI₅ molecule.

(3)

| Number of bonding pairs of electrons on phosphorus | |
|--|--|
| Number of lone pairs of electrons on phosphorus | |
| Shape of molecule | |
| CI—P—CI bond angles | |

(ii) In the solid state, phosphorus(V) chloride is ionic.

The cation and anion each have one phosphorus atom but a different number of chlorine atoms.

The cation is tetrahedral and the anion is octahedral.

Predict the formula of each ion. Include the charge on each ion.

(2)



(d) In an experiment, $8.00\,\mathrm{cm^3}$ of $0.250\,\mathrm{mol\,dm^{-3}}$ sodium hydroxide, NaOH, reacted completely with $10.0\,\mathrm{cm^3}$ of $0.100\,\mathrm{mol\,dm^{-3}}$ phosphoric acid, $\mathrm{H_3PO_4}$.

Use these data to deduce the balanced equation for this reaction. You **must** show your working.

(3)

(e) Hydrated magnesium phosphate has the formula $Mg_3(PO_4)_2$. $\mathbf{y}H_2O$.

A sample of this compound contains 78.5% by mass of anhydrous magnesium phosphate.

Deduce the value of **y**.

You **must** show your working.

[Molar mass of anhydrous magnesium phosphate, $Mg_3(PO_4)_2 = 262.9 \,\mathrm{g}\,\mathrm{mol}^{-1}$]

(2)

(Total for Question 24 = 16 marks)

TOTAL FOR SECTION B = 60 MARKS
TOTAL FOR PAPER = 80 MARKS



| 0,0 | 0 (8) | 4.0 Helium | 2 | 20.2 | Ne | neon 10 | 39.9 | Ar | argon 18 | 83.8 | Ā | krypton 36 | 131.3 | Xe | xenon 54 | [222] | R | 12/ | | orted | | | | | | | F | |
|------------------------------------|-------|-----------------------------|-------------------|----------------------|---------------|----------------|--------------------------------|------|------------------|-----------------|-------|-----------------|----------------|----------|---------------------------------|---------------|-------|---|----------------|-------------|-----------------------------------|----------------|---------------------|-------------------|--|--------|---------------------------------|-----------------------|
| t | ` | | (17) | 19.0 | is. | fluorine 9 | 35.5 | บ | chlorine 17 | 79.9 | Br | bromine 35 | 126.9 | - | iodine 53 | [210] | At | astatine 85 | | peen repo | | 175 | רי | lutetium 71 | [257] | Ļ | NE. | |
| , | 0 | | (91) | 16.0 | 0 | oxygen 8 | 32.1 | S | sulfur 16 | 79.0 | Se | selenium 34 | 127.6 | <u>T</u> | tellurium 57 | [209] | Po | polonium 84 | | 116 have | iticated | 173 | Yb | ytterbium 70 | [254] | ٥ ۷ | miliohon | |
| ı | n | | (15) | 14.0 | z | nitrogen 7 | 31.0 | ۵ | phosphorus 15 | 74.9 | As | arsenic 33 | 121.8 | Sb | antimony 5-1 | 209.0 | 9 | bismuth 83 | | ribers 112- | ally auther | 169 | E | thulium 69 | [256] | PW | 1 | |
| | 4 | | (14) 12.0 C | carbon 6 | 28.1 | S | silicon 14 | 72.6 | Ge | germanium 32 | 118.7 | Sn | rio 20 | 207.2 | Pb | lead 82 | | Elements with atomic numbers 112-116 have been reported but not fully authenticated | | 167 | Ē | erbium 68 | [253] | F | - | | | |
| | 71 | | (13) | 10.8 | В | boron 5 | 27.0 | ¥ | aluminium 13 | 69.7 | Ga | E | 114.8 | H | indium 49 | 204.4 | F | thallium 81 | | ents with | | 165 | 유 | holmium 67 | [254] | Es | minchains, um | |
| | | | | | | | | | (12) | 65.4 | Zn | zinc 30 | 112.4 | B | cadmium 48 | 200.6 | 린 | mercury 80 | | | | 163 | ρ | dysprosium 66 | [251] | ຽ | and the sandy man advantaged in | |
| | | | | | | | | | (11) | 63.5 | C | copper 29 | 107.9 | Ag | silver 47 | 197.0 | Au | gold 79 | [272] | Rg | pentgenium 111 | 159 | 4 | terbium 65 | [242] | | à | |
| | | | | | | | | | (01) | 58.7 | ž | nickel 28 | 106.4 | Pd | palladium 46 | 195.1 | Pt | platinum 78 | | Ds | darmstadtium 110 | 157 | PS | gadolinium 64 | [247] | E | 1000 | |
| | | | | | | | | | (6) | 58.9 | 0 | cobalt 27 | 102.9 | R | rhodium 45 | 192.2 | L | iridium 77 | [268] | Mt | meitnerium damstadtium 109 110 | 152 | B | europium 63 | [243] | Am | The state of the state of | |
| | | 1.0 H hydrogen | | | | | | | (8) | 55.8 | Fe | iron 26 | 101.1 | Ru | ruthenium 44 | 190.2 | os | osmium 76 | [277] | Hs | nassium 108 | 150 | Sm | samarium 62 | [242] | Np Pu | Sport S. S. Lake | |
| | | | | | | | | | (7) | 54.9 | Wn | manganese 25 | [86] | ۲ | technetium 43 | 186.2 | Re | rhenium 75 | 7.7 | Bh | 107 | [147] | Pm | promethium 61 | [237] | S. | LA CONTRACTOR | |
| | | | | nass | loc | umber | | | (9) | 52.0 | ò | E | 6.56 | Wo | molybdenum technetium ruthenium | 183.8 | * | tungsten 74 | [596] | Sg | seaborgium 106 | 144 | PN | neodymium p 60 | 238 | כ | Special States | |
| ווופ גפווסמור ומחופ חו דופווופוורא | | | Key | relative atomic mass | atomic symbol | mic sym | name atomic (proton) number | | | (5) | 50.9 | > | vanadium 23 | 92.9 | | niobium 41 | 180.9 | Ta | tantalum 73 | - | | dubnium 105 | 141 | P. | prascodymium neodymium promethium sa 59 60 61 | [231] | Pa | The state of the same |
| | | | 1 | relati | ato | atomic | | | (4) | 47.9 | ۲ | titanium 22 | 2.16 | Zr | zirconium | 178.5 | Ŧ | hafinium 72 | [261] | R | nutherfordium 104 | 140 | Ce | cerium p | 232 | ۴ | AC 300 30 | |
| | | | | | | | | | (3) | 45.0 | Sc | scandium 21 | 88.9 | > | yttrium 39 | 138.9 | *a | lanthanum 57 | [227] | | actinium 89 | | S | | | | | |
| | 7 | | (2) | 0.6 | Be | berytlium 4 | 24.3 | Mg | magnesium | 40.1 | Ca | calcium 20 | 87.6 | Şr | strontium 38 | 137.3 | Ва | barium 56 | [326] | Ra | radium 88 | | * Lanthanide series | * Actinide series | | | | |
| | - | | (1) | 6.9 | 5 | lithium 3 | 23.0 | Na | sodium 11 | 39.1 | ¥ | potassium 19 | 85.5 | Rb | rubidium 37 | 132.9 | S | caesium 55 | [223] | Ŀ | francium 87 | | " Lanth | * Actini | | | | |

P 6 1 6 5 6 A 0 2 4 2 4