Write your name here Surname	Other na	mes
Pearson Edexcel GCE	Centre Number	Candidate Number
Further F Mathema Advanced/Advance	atics FP2	
Wednesday 7 June 2017 Time: 1 hour 30 minute	•	Paper Reference 6668/01
You must have: Mathematical Formulae and	Statistical Tables (Pink)	Total Marks

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information

- The total mark for this paper is 75.
- The marks for each question are shown in brackets
 use this as a quide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶



P48860A
©2017 Pearson Education Ltd.
1/1/1/1/1/



(3)

1. (a) Show that, for r > 0

$$\frac{1}{r^2} - \frac{1}{(r+1)^2} \equiv \frac{2r+1}{r^2(r+1)^2} \tag{1}$$

(b) Hence prove that, for $n \in \mathbb{N}$

$$\sum_{r=1}^{n} \frac{2r+1}{r^2(r+1)^2} = \frac{n(n+2)}{(n+1)^2}$$
 (3)

(c) Show that, for $n \in \mathbb{N}$, n > 1

$$\sum_{r=n}^{3n} \frac{6r+3}{r^2(r+1)^2} = \frac{an^2 + bn + c}{n^2(3n+1)^2}$$

where a, b and c are constants to be found.

	Leave
	blank
Question 1 continued	



estion 1 continued	

					ı	
4		S		ľ		
_	j	1	7	7	9	
J	Ŀ		Ŀ	J	ı	
1	à		å	d	ì	
J	E	3	Ę		ì	
	3	_	à	d	i	
J	ξ	ξ		Ĺ		
				ī	1	
4	r	À	p	٩	ì	
,	b	ø		Į	,	
þ		Ė		Ė	İ	
3	7		P		ì	
ì		å	ŀ	i	i	
ì	L		è			
1	г	ī			١	
ì	ě	ė	è		ì	
í	d	è	ė	7	1	
c	7				1	
4	7	7			۱	
ì	r	i		í	ì	
1	Ŀ	å	ŀ	d	l	
1	L	-				
1	Ę					
4					:	
1	ľ		e		1	
j	Ξ	Ξ			۰	
î	ä		į)	1	
1	Ξ			į	į	
1	7	1				
1	L	÷		i	i	
1	Ţ			Ī		
1	٢	7	ī	٦	١	
-1	9	e	ø	ø	,	
9	ŧ	5		Ī	J	
1	É	S	i	i	i	
ï	ρ	P	ļ	3	ĺ	
4	b	ė	d	d	Í	
٠,	å	ŕ	í	š		
1	Ĺ			1	١	
ŕ	Ī	7	Ī		í	

		۹,	
A			
		Ĺ	
	Ž		
		ė	
Š			
Š			
Z			
	Q	é	
Χį			
×.	£	×	×
X	ô	K	ć.
ă	Ĉ	ă	Ŕ
ă	Ć	ă	Ŕ
ğ	Ć	Ď	Ć
Ž	Ć	Ž	ķ

×	×				
<2	$^{\circ}$			>	
$^{\wedge}$	$^{\wedge}$	/	\vee	◝	7
×	X	\rightarrow			>
				>	
X	х	2		Κ.	2
V	V	`		/	١,
				\rightarrow	
×	×	>	a	s	ò
1	V	ы	я	r	١.
		8		ь	
×	×	2		æ	
S	C)	1	ж	л	K.
\triangle	\wedge	ж	-	=	ν
×	X	'n	െ	ú	
50		4ľ	78		
\wedge	\wedge	я	₹	=	ν
×	X	7	2	d	
X	Χ	7	₹		ν
V	V	7		0	١,
()		4	×	è	
X	X		Я	П	P
V	V	-4	т.	л	١,
		é	=		(
×	×	2		<u>C</u>	
\vee	O	€	7	_	
\wedge	$^{\wedge}$	2	ж	$^{\sim}$	
×	×	5	_	₹	
<2		48.	X	_2	9
\wedge	$^{\sim}$	æ	♥	5	
×	×	5		<	
$\langle \rangle$		∠	\times	_	(
$^{\sim}$	А	~			P
V	V		s	\simeq	S
		-	₹	-	۲
X	X		w	< 2	ρ
V	V	`\			١,
		-6	¥	'n	(
×	×		ж		
V	\vee	₹	_	_	١
		Æ	\times	\sim	
×	×	Р	-	e	
0	< 2	₹	Х	. 2	
\triangle	$^{\wedge}$	ж	₹	=	ν
×	×	24	S)	d	
	< 2	Æ	ж	2	
\triangle	$^{\wedge}$	я	₹	₹	
×	×		6	2	S
		G			ľ
$^{\prime}$	X	98	S.	-	2
×	×	m	ø	۳	5
		₹		>	
X	X	È			>
\vee	$^{\prime}$		•	ø	١.
		ø	×	>	
×	×	а	m	٩.	Þ
\vee	$^{\prime}$	π.		а	ĸ.
		ð		ø	
×		ũ		÷	o
V	v	≺	ä	₹	
\triangle	Λ	d		_	7
×		7	d	₹	5
		<.	×	.>	
Λ	Х	ä	п	٩.	?
V	V	π		a	ŀ.
		ব	m	ø	۲
X	X	2	ő	ĸ	
V	V	и	7		
		46	~	4	ľ
X	X	7		₹	5

	Leave blank
Question 1 continued	
	01
	Q1
(Total 7 marks)	



$\frac{x-2}{2(x+2)} \leqslant \frac{12}{x(x+2)}$	(9



Question 2 continued	Leave
	Q2
(Total 9 marks)	
·	



$z^3 + 32 + 32i\sqrt{3} = 0$	
giving your answers in the form $re^{i\theta}$ where $r > 0$ and $-\pi < \theta \leqslant \pi$	(6
	(6



ì		۴	
7	7	Ni.	
1	ш	Ы	
Ġ	5	Ξ	
1	▢	⊆	
1	-	=	
I	c	E	
		7	
ì		Ri.	
J	u,	·	
9	-	=	
ŀ	٦	=	
Ì	4	÷	
J	H	÷	
ŀ	١.		
ź	۰,	90	
4	2	_	
Ĝ	-	Ξ	
1	ì	1	
ä		=	
J	×	=	
å	È	÷	
4	n	įπ	
J		Ė	
ļ	-	'n	
4	⋖	⊆	
J	ø	-	
1			
4	F	=	
ď	je	٩.	
4	Ŀ	J	
9	=	ė	
'n	2	_	
÷		7	
٠,	'n	iκ	
4	Ŀ	2)	
-		Ξ.	
1	ŗ.	2	
J	-	=	

		•
ă		
	É	
X		
a		
/ i		
×ú		
	H	
2	Ľ	×
2	ļ	3
Ø	ľ	5
S	Ċ	3
	e	٥
	9	

				. 7	
2	V	V	Ŋ	٦,	2
>	\circ	<	K.	Χ	\geq
1				×.	
ς	Λ	×	×	2	ς
	×	×		\rightarrow	
2	V	V	×,	Χ	2
\rightarrow			ĸ.	×	≥
1	A	a	е	r	
		×	$\overline{}$	9	ĸ
<	×	M	Ŋπ	ĤΙ	ľ
2	\bigcirc	S	ы	ы	P
\rangle		Q	ĸ	ú	ò
1	$^{\sim}$	4	2	2	
<	×	×	Ä	7	
	×	M	ы	Б	
2	\bigcirc	C	-0	٩,	₽
\rightarrow			ĸ.	×	>
5	X	ă	Vă	Ri	
	×	×	w	я	ĸ
2	\vee	S	Σ.	×	2
>		<:	=	K	⊳
S		~	ΧI	×	5
≺	×	X	ď		ĸ
1	×	M		2	
2	V	SI	N	7	
>	$^{\circ}$	C.	KO	×	>
Κ	Х	×	×	-	ĸ
		X	2	7	
7	×	M	Ž	9	ĸ
2	\circ	S	4	Š	Ķ
>	\circ		ă	×	
ς	Α	21	и	Ы	Ь.
		K			ĸ
2	0	Si	K	S	
ζ	8	a	K	Ě	6
ξ	8	Į	X	Š	5
Ş	8		X	Š	5
3	× ×			Š	
3	8	8			
3				X	
3					
			② 区 マ マ マ マ マ マ マ マ マ マ マ マ マ マ マ マ マ マ		

	Leave blank
Question 3 continued	
	Q3
(Total 6 marks)	



4.

$$y = \ln\left(\frac{1}{1 - 2x}\right), \quad |x| < \frac{1}{2}$$

(a) Find $\frac{dy}{dx}$, $\frac{d^2y}{dx^2}$ and $\frac{d^3y}{dx^3}$

(4)

(b) Hence, or otherwise, find the series expansion of $\ln\left(\frac{1}{1-2x}\right)$ about x=0, in ascending powers of x, up to and including the term in x^3 . Give each coefficient in its simplest form.

(3

(c) Use your expansion to find an approximate value for $\ln\left(\frac{3}{2}\right)$, giving your answer to 3 decimal places.

(3)



	blank
Question 4 continued	Dialik
Question 4 continued	



estion 4 continued		

ì		۲.	
1	7	ij	
1	ш		
Ġ	5	Ξ	
1	9	⊆	
	Ξ	2	
J	ς	L	
		7	
i	Ú	Ri.	
3	ч	·il	
ŝ	7	=	
å	٦	т.	
3	÷	=	
J	×	ė	
ŀ			
þ	÷	ø	
4	۷	_	
Ĝ		Ξ	
4	Ш	1	
å		=	
4	=	=	
å	ì	÷	
4	h	įπ	
J	н		
J	•	'n	
4	ς	⊆	
J	P	-	
1			
4	7	=	
ď	m	٩.	
4	-	J	
9	=	ė	
'n	e	_	
÷		7	
٠,	'n	iκ	
4	Ŀ	2)	
-	Z	Ξ.	
1		2	
ĕ	7	=	

/ 12	۹		
1	γ.	ď	3
- 21	ы	L	ŧ.
	=	e	ш
1		ù	ø.
-/1	ш	ĸ	4
(2	_	_	ч.
1	Ĺ		ø,
4	s		Δ
	- 3	₹	۹.
N	Χ.	. `	
А	ь.	a	ø.
(4	ы	ĸ.	В
`\	$\underline{}$	_	z
/"	₹	7	-
S	9	r	=
`\	41	r	\vee
/1	₹	7	8
₹.1	112	ς.	-
N	-	-	ų.
/"	٩.	×	
ί.	_2	2	-
K	-		R.
×i		2	
(a	7	₹	-
N	-	-	91
×	×	×	
M	ИÌ	b.	ı
N	ч	н	æ
/;	₹	7	
\ 1		'n	á
K	г		
/ 3	-	-	=
1	4	2	ä
- X	r	r	×
∕!	=	-	9
7.1	66	7	۷
X	=		ø
/!	5	×	
S	ø	щ	۳
$-\times$		۲,	×
/ 1	Ð		ż
ΧI	•	7	Н.
- 24	ш	_	Χ
CA	п	~	'n
ΧI	ĸ	2	z.
-2	◡	7	7
(* A	=		=
X.	Set.	o	
·/I	ø	9	ø
	. >		
X.	žα	nii	Ĺ
vA	ď		٩
C.	S)	4	ø
×	s	z	
wi	۳	7	ĸ

	Х	х	X	х	
	×	×	×	×	
	$\langle \rangle$		$\langle \rangle$		
			△	△	
ζ	Х	Х	X	Х	
		×		×	
2	V	V	V	V	
			$\langle \rangle$		
	X	X	A	X	
	×	×	검	ø	
2	V	V	ч	w	
		Q	(5	0	
	A	41	è	45.	
	Х	×	_	z	
	V	ğ	ny.	7	
	$\langle \rangle$	QШ	40	in.	
		△	△	ď)	
ς	Х	24		Ø,	
	X	×	X	₹	
2	V	v	v.	SZ	
		a		90	
ς	Ä	29	٣.	Ж,	
	×	Ж	×	ìЖ,	
1	V	Sil	×	×	
			OD.	$^{\circ}$	
	A	<i>0</i> 9	•	۳,	
	Х	Ж	\times	×	
	X	Ж	X	×	
	$^{\circ}$	X	\vee	V	
		0	€	90	
۱	A	a	e	\simeq	
	Х	ğ	Z	z	
	×	×	₹	₹	
	$\langle \rangle$	Ω	\sim	\circ	
		41	Иlb	đЬ	
ς	Х	겓	_	픠	
		M	×	\simeq	
2	V	SII	∇	∇	
		Gi		≦)	
ς	×	X	à	ä	
	X	Ж	70	X	
1	V	Х	₹	₹	
		U	₩.	⇔	
	△	a		20	
	×	ă	-	寒	
		×	X	×	
2	$^{\circ}$	Ü	K2	v	
		(II		◓	
	X	75	ä	ĸ.	
	х	ж		ж	
	V	×	ч	W.	
		G		ӹ	
		0	w	n	
ς	X	24	,	ж,	
		×		×	
2	V	M	pp)	w	
		(II		Æ.	
S	ZS.	8	2	75	
	×	X	州	ĸ	
		M	15	¥.	
		S	V	ø	
	\Diamond		\Diamond		

Question 4 continued	Leave blank
	Q4
(Total 10 marks)	
(Total To marks)	



5. (a) Find the general solution of the differential equation

$$\frac{\mathrm{d}^2 y}{\mathrm{d}x^2} - 2\frac{\mathrm{d}y}{\mathrm{d}x} = 26\sin 3x$$

(8)

(b) Find the particular solution of this differential equation for which y = 0 and $\frac{dy}{dx} = 0$ when x = 0

(5)

14

	I t
uestion 5 continued	



estion 5 continued	

Question 5 continued	Leave blank
	Q5
(Total 13 marks)	



(8)

6.




Figure 1

Figure 1 shows a sketch of a curve with polar equation

$$r = 6 + a\sin\theta$$

where $0 \le a \le 6$ and $0 \le \theta \le 2\pi$

The area enclosed by the curve is $\frac{97\pi}{2}$

Find the value of the constant a.





	Leave
Question 6 continued	
	Q6
(Total 8 marks)	



blank

(8)

7. (a) Find, in the form y = f(x), the general solution of the equation

$$\cos x \frac{\mathrm{d}y}{\mathrm{d}x} + y \sin x = 2\cos^3 x \sin x + 1, \quad 0 < x < \frac{\pi}{2}$$

Given that $y = 5\sqrt{2}$ when $x = \frac{\pi}{4}$

(b) find the value of y when $x = \frac{\pi}{6}$, giving your answer in the form $a + b\sqrt{3}$, where a and b are rational numbers to be found.



	blank
Question 7 continued	Oldlik
Question / continued	



	Leave blank
Question 7 continued	
	Q7
(Total 11 marks)	



blank

8. The transformation T from the z-plane to the w-plane is given by

$$w = \frac{z + 3i}{1 + iz}, \quad z \neq i$$

The transformation T maps the circle |z| = 1 in the z-plane onto the line l in the w-plane.

(a) Find a cartesian equation of the line l.

(5)

The circle |z - a - bi| = c in the z-plane is mapped by T onto the circle |w| = 5 in the w-plane.

(b) Find the exact values of the real constants a, b and c.

1	6	١.
l	U	ч

estion 8 continued	



	Leave
	blank
Question 8 continued	
Ancetion a continuen	



(Total 11 marks)	Q8
TOTAL FOR PAPER: 75 MARKS END	

Question 8 continued