Candidate surname			ng your candidate information Other names
Pearson Edexcel nternational Advanced Level	Centre	e Number	Candidate Number
Tuesday 12 Ja	anu	iary 2	2021
Afternoon (Time: 1 hour 30 minu	utes)	Paper Ref	ference <b>WME02/01</b>
Mathematics			
International Advance Mechanics M2	d Suk	osidiary,	/Advanced Level

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

## **Instructions**

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
   there may be more space than you need.
- You should show sufficient working to make your methods clear.
   Answers without working may not gain full credit.
- Whenever a numerical value of g is required, take  $g = 9.8 \,\mathrm{m \, s^{-2}}$ , and give your answer to either 2 significant figures or 3 significant figures.

## Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 8 questions in this question paper. The total mark for this paper is 75.
- The marks for each question are shown in brackets
  - use this as a guide as to how much time to spend on each question.

## **Advice**

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over ▶







$P \text{ is } v \mathbf{i}  \mathbf{m}  \mathbf{s}^{-1}.$	
Find the two possible values of <i>v</i> .	(7
	(1

	Š			
	ķ			
			я	
			-	
	ì			
	ŀ			
			Ž	
	2			
<				
K	S		S	
K	S			
8	S		Š	
K	S		)	
K	S		)	
K	S		)	
K	S		)	
K	S		)	
K	S		)	

	Leave blank
Question 1 continued	
	Q1
(Total 7 marks)	



2.




Figure 1

The uniform lamina ABC has sides AB = AC = 13a and BC = 10a. The lamina is freely suspended from A. A horizontal force of magnitude F is applied to the lamina at B, as shown in Figure 1. The line of action of the force lies in the vertical plane containing the lamina. The lamina is in equilibrium with AB vertical. The weight of the lamina is W.

Find $F$ in terms of $W$ .	(5

		Ì	
		2	

Question 2 continued	Leave blank
Question 2 continued	
	<b>Q2</b>
(Total 5 marks)	



3.	A car of mass $600 \mathrm{kg}$ travels along a straight horizontal road with the engine of the car working at a constant rate of $P$ watts. The resistance to the motion of the car is modelled as a constant force of magnitude $R$ newtons. At the instant when the speed of the car is $15 \mathrm{ms^{-1}}$ , the magnitude of the acceleration of the car is $0.2 \mathrm{ms^{-2}}$ .
	Later the same car travels up a straight road inclined at angle $\theta$ to the horizontal, where
	$\sin \theta = \frac{1}{20}$ . The resistance to the motion of the car from non-gravitational forces is
	modelled as a constant force of magnitude $R$ newtons. When the engine of the car is working at a constant rate of $P$ watts, the car has a constant speed of $10\mathrm{ms^{-1}}$ .
	Find the value of <i>P</i> .
	(8)

		ğ		
	p			
	ì	Ki	S	ú
K	Ì	Ń	ř	Î
S	Ì	Ń	È	ĺ
X	١	Ŋ	Ļ	J
S	١	Ŋ	Ļ	J
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3				
3				
3				
3				
3		N		
3				
3				
3				
3				
3				
3				
3				
3				
3				
3				
3				
3				
3				
3				
3				
3				
3				

	blank
Question 3 continued	



estion 3 continued	
estion 5 continued	

		ğ		
	p			
	ì	Ki	S	ú
K	Ì	Ń	ř	Î
S	Ì	Ń	È	ĺ
X	١	Ŋ	Ļ	J
S	١	Ŋ	Ļ	J
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3		N	Ľ	ž
3				
3				
3				
3				
3		N		
3				
3				
3				
3				
3				
3				
3				
3				
3				
3				
3				
3				
3				
3				
3				
3				
3				

	Leave blank
Question 3 continued	Dialik
	Q3
(Total 8 marks)	



4.




Figure 2

The number "4", shown in Figure 2, is a rigid framework made from three uniform rods, AB, BC and CD, where

$$AB = 6a$$
,  $BC = 5a$  and  $CD = 4a$ 

The point E is on AB and CD, where BE = 4a, CE = 3a and angle  $CEB = 90^{\circ}$ 

The three rods are all made from the same material and they all lie in the same plane.

The framework is suspended from B and hangs in equilibrium with BA at an angle  $\theta$  to the downward vertical.

Find  $\theta$  to the nearest degree.

•	y	
	_	,



		ĺ		
	d			
	Ŕ			
	i			

	bl
Question 4 continued	



estion 4 continued	

		É			
		Ì			
		à			
	ì				
		è			
		Ś			
		ĺ			
		Ė			
				5	
			ę		
		p			

Question 4 continued	blank
	0.4
	Q4
(Total 9 marks)	



**5.** At time t seconds,  $t \ge 0$ , a particle P has velocity  $\mathbf{v} \, \mathbf{m} \, \mathbf{s}^{-1}$ , where

$$\mathbf{v} = (5t^2 - 12t + 15)\mathbf{i} + (t^2 + 8t - 10)\mathbf{j}$$

When t = 0, P is at the origin O.

At time *T* seconds, *P* is moving in the direction of  $(\mathbf{i} + \mathbf{j})$ .

(a) Find the value of T.

(3)

When t = 3, P is at the point A.

(b) Find the magnitude of the acceleration of P as it passes through A.

**(4)** 

(c) Find the position vector of A.

**(4)** 


	è		
	И		
		10	
	ň		
	Ž		
	ŭ		
		×	
$\langle \rangle$	Ķ		
		×	ĸ
	26	gβ	ø
	-	6	×
	~	a	И
	₹7		
×.	Ř	Х	
			ê
	gi.	=	ĸ
	E/		
	è	-	и
		-	ú
	7	73	Ö
v	V	V	
	Ç	6	×
X	or.	78	ĸ
	B.S		ď
	76		۲
	Gã.	<u>~</u>	
X.	Q X	и	ĸ.
	II 2	U	ú
	-	-	ø
$\vee$			
	1		×

	Leave blank
Question 5 continued	biank
	Q5
(Total 11 marks)	



**6.** 




Figure 3

A ladder AB has length 6 m and mass 30 kg. The ladder rests in equilibrium at  $60^{\circ}$  to the horizontal with the end A on rough horizontal ground and the end B against a smooth vertical wall, as shown in Figure 3.

A man of mass  $70 \,\mathrm{kg}$  stands on the ladder at the point C, where  $AC = 2 \,\mathrm{m}$ , and the ladder remains in equilibrium. The ladder is modelled as a uniform rod in a vertical plane perpendicular to the wall. The man is modelled as a particle.

(a) Find the magnitude of the force exerted on the ladder by the ground.

**(6)** 

The man climbs further up the ladder. When he is at the point D on the ladder, the ladder is about to slip.

Given that the coefficient of friction between the ladder and the ground is 0.4

(b) find the distance AD.

**(4)** 

(c) State how you have used the modelling assumption that the ladder is a rod. (1)



		Ĺ	
			1
p			
			ĺ
			2
		ė	
		2	

	blank
Oraștian ( continued	Diank
Question 6 continued	



uestion 6 continued	
estion o continued	

	í	ì			
	ï				
				į	
			į		
			è		
				2	
				2	
			į		
			3		

Question 6 continued		Leave blank
	Question 6 continued	
(Total 11 marks)		Q6
I I	(Total 11 marks)	



7.




Figure 4

The fixed point A is 20 m vertically above the point O which is on horizontal ground. At time t = 0, a particle P is projected from A with speed  $12 \,\mathrm{m\,s^{-1}}$  at an angle  $\theta^{\circ}$  above the horizontal. The particle moves freely under gravity. At time t = 5 seconds, P strikes the ground at the point B, where  $OB = 40 \,\mathrm{m}$ , as shown in Figure 4.

(a) By considering energy, find the speed of P as it hits the ground at B.

**(4)** 

(b) Find the least speed of P as it moves from A to B.

**(2)** 

(c) Find the length of time for which the speed of P is more than  $10 \,\mathrm{m\,s^{-1}}$ .

**(6)** 

	Leave
Overtion 7 continued	blank
Question 7 continued	



	Lea
Question 7 continued	blar
Question / continued	
	-
	_
	-
	-
	_
	-
	_
	-
	_
	.
	_
	.
	.
	-
	_
	.

×				
	á			
$\sim$		z	v	
( N	10			
			ij.	
×			P)	
			×	
×Ι	=	۰	€.	
	$\supset$			
Λ.	<i>&gt;</i> _	2		
V)	ø	٠	ĸ	
$\sim$	E\		와	
v	•	ù	ø.	
	3	z		
X	ar.	4	B.	
K	P		ж	
XI			ø.	
	. 7		Х	
Х.				

Overtion 7 continued	Leave blank	
Question 7 continued		
	07	
	Q7	
(Total 12 marks)		



Two particles, A and B, have masses 3m and 4m respectively. The particles are moving towards each other along the same straight line on a smooth horizontal surface. The particles collide directly. Immediately after the collision, A and B are moving in the same

direction with speeds  $\frac{u}{3}$  and u respectively. In the collision, A receives an impulse of magnitude 8mu.

(a) Find the coefficient of restitution between A and B.

**(6)** 

When A and B collide they are at a distance d from a smooth vertical wall, which is perpendicular to their direction of motion. After the collision with A, particle B collides directly with the wall and rebounds so that there is a second collision between A and B. This second collision takes place at distance x from the wall.

Given that the coefficient of restitution between B and the wall is  $\frac{1}{4}$ 

-	6)
•	v)

	Leave
	blank
Question 8 continued	
Question o continued	



estion 8 continued	

	l t
uestion 8 continued	



	Ö
	XX
	×8×
	X max
	m ==
	工
	X tax
	Un
	70
	A
	D
	×5%
	Ö
	8
	Ž
	33
	2
	×111×
	TEIN
	SO TO
	À
	2
	2
	V X X X X X
	X TEXT
	A
	A
	A
	A
	A
	A
	A
	A
	A
	A
	A
	A
	A
	SA.
	SA.
	ĒA.
	ĒA.
	ĒA.
	ĒA.
	ēA D
	EA DO
	EA DO
	EA DOI
	EA DON
	EA DO NO
	EA DO NO
	EA DO NO
	EA DO NOT
	EA DO NOT WRI
	EA DO NOT WRI
	EA DO NOT WRITE
	EA DO NOT WRITE
	EA DO NOT WRITE H
	EA DO NOT WRITE H
	EA DO NOT WRITE IN 1
	EA DO NOT WRITE IN T
	EA DO NOT WRITE IN T
	EA DO NOT WRITE IN THI
	EA DO NOT WRITE IN THI
Q8	EA DO NOT WRITE IN T

estion 8 continued	
	(Total 12 marks)
	TOTAL FOR PAPER: 75 MARKS

**END**