Write your name here Surname	Other na	ames
Pearson Edexcel International Advanced Level	Centre Number	Candidate Number
Mechanic	c M1	
Advanced/Advance		
	d Subsidiary	Paper Reference WME01/01

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

## Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
   there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Whenever a numerical value of g is required, take g = 9.8 m s<sup>-2</sup>, and give your answer to either two significant figures or three significant figures.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

## Information

- The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
  - use this as a guide as to how much time to spend on each question.

## **Advice**

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶



Leave	
hlank	

1.	A truck $P$ of mass $2M$ is moving with speed $U$ on smooth straight horizontal collides directly with another truck $Q$ of mass $3M$ which is moving with speed 4 opposite direction on the same rails. The trucks join so that immediately after the they move together. By modelling the trucks as particles, find	U in the
	(a) the speed of the trucks immediately after the collision,	(3)
	(b) the magnitude of the impulse exerted on $P$ by $Q$ in the collision.	(3)

uestion 1 continued	
	9



Leave	
blank	

<b>2.</b> A particle <i>P</i> is moving with constant velocity $(2\mathbf{i} - 3\mathbf{j})$ m s <sup>-1</sup> .			
(a) Find the speed of $P$ .	(2)		
The particle $P$ passes through the point $p$ position vector $(\mathbf{i} - 4\mathbf{j})$ m.	A and 4 seconds later passes through the point with		
(b) Find the position vector of <i>A</i> .	(4)		

	Q



Leave
hlank

A beam $AB$ has length 15 m and mass 25 kg. The beam is smoothly sur $P$ , where $AP = 8$ m. A man of mass 100 kg stands on the beam at a dial $A$ and another man stands on the beam at a distance of 1 m from $B$ . The as a non-uniform rod and the men are modelled as particles. The beam a horizontal position with the reaction on the beam at $P$ having magnitude distance of the centre of mass of the beam from $A$ .	stance of 2 m from e beam is modelled is in equilibrium in tude 2009 N. Find
	(5)

estion 3 continued	
	(Total 5 marks)



Leave blank

4.




Figure 1

A fixed rough plane is inclined to the horizontal at an angle  $\alpha$ , where  $\tan \alpha = \frac{3}{4}$ 

A small box of mass m is at rest on the plane. A force of magnitude kmg, where k is a constant, is applied to the box. The line of action of the force is at angle  $\alpha$  to the line of greatest slope of the plane through the box, as shown in Figure 1, and lies in the same vertical plane as this line of greatest slope. The coefficient of friction between the box and the plane is  $\mu$ . The box is on the point of slipping up the plane. By modelling the box as a particle, find k in terms of  $\mu$ .

	(11)

estion 4 continued		

Q4	Question 4 continued		blank
(Total 11 marks)			Q4
(10001111)		(Total 11 marks)	



Leave	
blank	

5.	A racing car is moving along a straight horizontal track with constant acceleration. There are three checkpoints, $P$ , $Q$ and $R$ , on the track, where $PQ = 48$ m and $QR = 200$ m. The car takes 3 s to travel from $P$ to $Q$ and 5 s to travel from $Q$ to $R$ . Find
	(i) the acceleration of the car,
	(ii) the speed of the car as it passes <i>P</i> .
	(7)

stion 5 continued		

estion 5 continued		

Question 5 continued		b.
		Q
	(Total 7 marks)	

Leave blank

**6.** 




Figure 2

Two particles P and Q have masses 0.1 kg and 0.5 kg respectively. The particles are attached to the ends of a light inextensible string. Particle P is held at rest on a rough horizontal table. The string lies along the table and passes over a small smooth pulley which is fixed to the edge of the table. Particle Q is at rest on a smooth plane which is inclined to the

horizontal at an angle  $\theta$ , where  $\tan \theta = \frac{4}{3}$ 

The string lies in the vertical plane which contains the pulley and a line of greatest slope of the inclined plane, as shown in Figure 2. Particle P is released from rest with the string taut. During the first 0.5 s of the motion P does not reach the pulley and Q moves 0.75 m down the plane.

(a) Find the tension in the string during the first 0.5 s of the motion.

**(6)** 

(b) Find the coefficient of friction between P and the tabl	(b)	Find the	coefficient	of friction	between	P an	d the	table
---	-----	----------	-------------	-------------	---------	------	-------	-------

**(5)** 

-	

uestion 6 continued	

	Question 6 continued	blanl
(Total 11 marks)		Q6
		(Total 11 marks)



Leave blank

7. A force <b>F</b> is given by $\mathbf{F} = (9\mathbf{i} + 13\mathbf{j})  \mathbf{N}$	7.	A force	F is	given	by <b>F</b>	= (9i +	13 <b>j</b> ) N
---	----	---------	------	-------	-------------	---------	-----------------

(a) Find the size of the angle between the direction of  $\mathbf{F}$  and the vector  $\mathbf{j}$ .

(3)

The force  $\mathbf{F}$  is the resultant of two forces  $\mathbf{P}$  and  $\mathbf{Q}$ . The line of action of  $\mathbf{P}$  is parallel to the vector  $(2\mathbf{i} - \mathbf{j})$ . The line of action of  $\mathbf{Q}$  is parallel to the vector  $(\mathbf{i} + 3\mathbf{j})$ .

- (b) Find, in terms of **i** and **j**,
  - (i) the force  $\mathbf{P}$ ,
  - (ii) the force **Q**.

**(9)** 

Question 7 continued	b

euestion 7 continued	bla



Leave	
hlank	

**8.** Two trains, A and B, start together from rest, at time t = 0, at a station and move along parallel straight horizontal tracks. Both trains come to rest at the next station after 180 s.

Train A moves with constant acceleration  $\frac{2}{3}$  m s<sup>-2</sup> for 30 s, then moves at constant speed

for 120 s and then moves with constant deceleration for the final 30 s. Train B moves with constant acceleration for 90 s and then moves with constant deceleration for the final 90 s.

(a) Sketch, on the same axes, the speed–time graphs for the motion of the two trains between the two stations.

(3)

(b) Find the acceleration of train *B* for the first half of its journey.

**(5)** 

(c) Find the times when the two trains are moving at the same speed.

**(4)** 

(d) Find the distance between the trains 96 s after they start.

(5)

estion 8 continued	

estion 8 continued	

estion 8 continued	

Question 8 continued	blank
	Q8
(Total 17 marks)	
TOTAL FOR PAPER: 75 MARKS	
END	