Please check the examination details be	low before ente	ring your candidate i	nformation
Candidate surname	Other names		
Centre Number Candidate N	umber		
Pearson Edexcel Inter	nation	al Advand	ced Level
Time 1 hour 30 minutes	Paper reference	WMA	13/01
Mathematics			
International Advanced Lo	evel		
Pure Mathematics P3			
You must have: Mathematical Formulae and Statistical	al Tables (Ye	llow), calculator	Total Marks
Time 1 hour 30 minutes Mathematics International Advanced Le Pure Mathematics P3 You must have:	Paper reference	WMA	13/01

Candidates may use any calculator permitted by Pearson regulations.

Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 10 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
- use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶







1. The function f is defined by

$$f(x) = \frac{5x}{x^2 + 7x + 12} + \frac{5x}{x + 4} \qquad x > 0$$

- (a) Show that $f(x) = \frac{5x}{x+3}$ (3)
- (b) Find f^{-1} (3)
- (c) (i) Find, in simplest form, f'(x).
 - (ii) Hence, state whether f is an increasing or a decreasing function, giving a reason for your answer.(3)



	Leave
Question 1 continued	blank
Question 1 continued	



estion 1 continued	

	$\times \times \rangle$	
$\times \times$	$\times\!\!\times\!\!\!>$	
$\Diamond \Diamond$	$\vee \vee \vee$	
\sim	$\propto \times$	
\sim	\sim	
	$\times\!\!\times\!$	
$\times \times$	$\times\!\!\times\!\!\!>$	
$\times \times$	$\times\!\!\times\!\!\!>$	
\sim	\sim	
	XX.	
-88	\sim	
$\times \times$		
\times	W.	
-525	inidal)	
	W.	
$\times \times$	2	
$\times \times$	A A >	
\sim	THIS A	
	6 X	
$\triangle \triangle$		
$\times \times$	\times	
$\times \times$	UD)	
\sim	∞z	
-88		
$\times \times$	\sim	
\times		
-00	-	
	$\times \times$	
$-\triangle\triangle$		
$\times \times$		
$\times \times$.	
\sim		
$-\triangle\triangle$	mar.	
$\times \times$		
\times		
\sim	77	
	200	
-88	OT WRITE	
$\times \times$		
XX	200	
-00	**	
\times		
$\times \times$		
××		
00	V .	
	24/	
\times		
$\times \times$	المناو	
$ \bigcirc$ \bigcirc	DONO	
\sim		
$\Delta \Delta$		
	$\times\!\!\times\!\!\!>$	
$\times \times$		
\sim	LXK.	
\sim	~/	
$\times \times$		
\sim	$\nabla\nabla$	
\sim	$\propto \times$	
\sim	$\propto \times$	
\times	$\times\!\!\times\!\!\!\times$	
$\times \times$	$\times\!\!\times\!\!\!>$	
	\sim	
\sim		
88	$\times\!\!\times$	
\otimes	\otimes	
\otimes	\otimes	
$\overset{\times}{\otimes}$	\times	
	*	
*		
*		
**		
	4	
	₩.	
	AM AM	
	EA	
	REA	
	REA	
	KREA	
	AREA	
	AREA	
	; AREA	
	SAREA	
	IS AREA	
	HS AREA	
	HS AREA	
	HIS AREA	
	THIS AREA	
	THIS AREA	
	THIS AREA	
	THIS AREA	
	NATHIS AREA	
	NATHIS AREA	
	IN THIS AREA	
	ANTHIS AREA	
	EXNITHIS AREA	
	EIN THIS AREA	
	TEIN THIS AREA	
	TEIN THIS AREA	
	ITEMNTHIS AREA	
	NTEXN THIS AREA	
	RITE IN THIS AREA	
	RITE IN THIS AREA	
	KRITE IN THIS AREA	
	WRITE IN THIS AREA	
	WRITEAN THIS AREA	
	WRITE IN THIS AREA	
	I WRITE IN THIS AREA	
	T WRITE IN THIS AREA	
	M WRITEAN THIS AREA	
	OT WRITE IN THIS AREA	
	OT WRITE IN THIS AREA	
	IOT WRITE IN THIS AREA	
	NOT WRITEAN THIS AREA	
	NOT WRITE IN THIS AREA	
	NOT WRITE IN THIS AREA	
	S NOT WRITE IN THIS AREA	
	O'NOT WRITEIN THIS AREA	
	O'NOT WRITE IN THIS AREA	
	30 NOT WRITE IN THIS AREA	
	DO NOT WRITE IN THIS AREA	

	Leave blank
Question 1 continued	
	01
	Q1
(Total 9 marks)	



2.




Figure 1

Figure 1 shows a sketch of part of the graph with equation y = f(x), where

$$f(x) = |3x - 13| + 5 \qquad x \in \mathbb{R}$$

The vertex of the graph is at point P, as shown in Figure 1.

(a) State the coordinates of P.

(2)

- (b) (i) State the range of f.
 - (ii) Find the value of ff(4)

(2)

(c) Solve, using algebra and showing your working,

$$16 - 2x > |3x - 13| + 5 \tag{4}$$

The graph with equation y = f(x) is transformed onto the graph with equation y = af(x + b)

The vertex of the graph with equation y = af(x + b) is (4, 20)

Given that a and b are constants,

(d) find the value of a and the value of b.

(2)

	blank
Question 2 continued	



Question 2 continued	blank
	_
	-
	-
	-
	-
	-
	-
	_
	_
	_
	-
	-
	-
	-
	-
	-
	_
	_
	_
	-
	-
	-
	-
	-
	-
	-
	_
	_
	_
	_
	_

Question	2 continued		
Question	2 continued		

Q2	

(Total 10 marks)



3.




Figure 2

The total mass of gold, G tonnes, extracted from a mine is modelled by the equation

$$G = 40 - 30e^{1-0.05t}$$

$$t \geqslant k$$

$$G \geqslant 0$$

where *t* is the number of years after 1st January 1800.

Figure 2 shows a sketch of G against t.

Use the equation of the model to answer parts (a), (b) and (c).

- (a) (i) Find the value of k.
 - (ii) Hence find the year and month in which gold started being extracted from the mine.

(3)

(b) Find the total mass of gold extracted from the mine up to 1st January 1870.

(2)

There is a limit to the mass of gold that can be extracted from the mine.

(c) State the value of this limit.

(1)

					ø	
		ġ				
	Q				á	
					ę	
D	<1	ь	d	b		
2	g	þ	4	þ	d	
ζ	4	ŀ	4	5	2	
?	1	į	4		2	
3	1	ŀ	4			Į.
3	1	ŀ	4			į
?		ŀ		2		į.
3	1					
3	1					
3						E K
3						
?						
3						
?						
?						
3						
?						
?						
?						
3						
?						
?						
?		2				
?		2				
?		2				
		2				
		2				
		2				
		2				
		2				
		2				
		2				
		2				
		2				
		2				
		2				
		2				
		2				
		2				
		2				
		2				
		2				
		2				
		2				
		2				
		2				
		2				
		2				
		2				
		2				
		2				
		2				
		2				
		2				

Question 3 continued	blank
Question 5 continued	
	Q3
(Total 6 marks)	
(Ioui o muins)	



4. In this question you should show detailed reasoning.

Solutions relying entirely on calculator technology are not acceptable.

(a) Show that the equation

$$2\sin(\theta - 30^\circ) = 5\cos\theta$$

can be written in the form

$$\tan \theta = 2\sqrt{3}$$

(4)

(b) Hence, or otherwise, solve for $0 \le x \le 360^{\circ}$

$$2\sin(x-10^\circ) = 5\cos(x+20^\circ)$$

giving your answers to one decimal place.

(3)



	blank
Question 4 continued	Diank
Question 4 continued	



estion 4 continued	

	2				~
	1	/			١.
\	S	5	۷	à	ď
1	м		9		rs
				Ė	
	Ğ				
	М	ø	ø	ø	
	А	L		E.	я
\rangle	₫	L	d	Ĺ	1
\langle	3		4		
>	1	Ŀ	2		
>	1	ŀ			
>	1				
>	1				
>					
>					
>					
>					
>					
>					
>					
>					
> > > >					
>					
> > > >					

Question 4 continued	Leave blank
	Q4
(Total 7 marks)	



5. (i) Find, by algebraic integration, the exact value of

$$\int_{2}^{4} \frac{8}{(2x-3)^{3}} \, \mathrm{d}x$$

(4)

(ii) Find, in simplest form,

$$\int x (x^2 + 3)^7 \, \mathrm{d}x$$

1	1	١
l	4	



16

×		×
$\langle \rangle$	$\langle \rangle$	$\langle \rangle$
$^{\wedge}$	\wedge	\triangle
X.	2	ari
V	Š	w
\wedge	í	-
×ı	м	И
a		ш
		\sim
×I		e.
C)	ы	=
\wedge	\triangle	_
×	æ	w.
×2	9	G.
$^{\wedge}$	\wedge	~~
X.		X.
a	и	n
	w.	211
Xi	_	1
×	_	Ħ
V.	S	W
0	_	-
XI	K.	Χ.
V	•	—
73	F)	
X.	x	X.
\sim	₹.	20
$^{\wedge}$	æ	n
X	7	₹.
a		
$^{\wedge}$	\wedge	
X,	×	×
ă	и	u
N	_	_
XI	K.	×
()	•	⇁
^\	25	\triangle
×	≂	⇁
$^{\circ}$		الاست
\sim	М	
X.	₹	7
O	m	-
$^{\sim}$	-in	per.
v	7	200
a	w	♥
$^{\sim}$	×	$^{\sim}$
M	MZ.	×
	7	⇁
X!	5S.,	Ζ.
W		70
C)	K)	CB
X		
V		
	2	
Xi	Ž	2
Ņ		Ž
X		
X		
X		3
X		2
		2
X		

Question 5 continued	
Q5	5
(Total 6 marks)	



(4)

6. (i) The curve C_1 has equation

$$y = 3\ln(x^2 - 5) - 4x^2 + 15 \qquad x > \sqrt{5}$$

Show that C_1 has a stationary point at $x = \frac{\sqrt{p}}{2}$ where p is a constant to be found.

(ii) A different curve C_2 has equation

$$y = 4x - 12\sin^2 x$$

(a) Show that, for this curve,

$$\frac{\mathrm{d}y}{\mathrm{d}x} = A + B\sin 2x$$

where A and B are constants to be found.

(b)	Hence,	state the	maxımum	gradient	ot	this	curve.
-----	--------	-----------	---------	----------	----	------	--------

		ļ			
	V		G	₽	'n
	K	Ľ	ĺ	Č	ĺ
5	K	ľ	ĺ	Ì	ĺ
>	K	ļ	į	Ì	ļ
>	1		1		2
>	1				
>	1				
>	1			2	
>					
>					
>					
>					
>					
<pre></pre> <pre><pre></pre><pre></pre><pre></pre><pre></pre><pre></pre><pre><!--</td--><td></td><td></td><td></td><td></td><td></td></pre></pre>					

Question 6 continued	blank
	Q6
(Total 8 marks)	



7 The mass, M kg, of a species of tree can be modelled by the equation

$$\log_{10} M = 1.93 \log_{10} r + 0.684$$

where r cm is the base radius of the tree.

The base radius of a particular tree of this species is 45 cm.

According to the model,

(a) find the mass of this tree, giving your answer to 2 significant figures.

(2)

(b) Show that the equation of the model can be written in the form

$$M = pr^q$$

giving the values of the constants p and q to 3 significant figures.

(3)

(c) With reference to the model, interpret the value of the constant p.

(1)

					2	
				ŕ		
	X					
	ă				Z	
	SI	ŀ	í	ú	ś	
ζ	SI	ŀ	í	ú	ś	
5	SI	ŀ	í	ú	ś	
3	SI	ŀ	í	ú	ś	
5	SI	ŀ	í	ú	ś	
3	SI	ŀ	í	ú	ś	
	SI	ŀ	í	ú	ś	
	SI	ŀ	í	ú	ś	
3	SI	ŀ	í	ú	ś	
	SI	ŀ	í	ú	ś	
	SI	ŀ	í	ú	ś	
	SI	ŀ	í	ú	ś	
	SI	ŀ	í	ú	ś	
	SI	ŀ	í	ú		
	SI	ŀ	í	ú	ś	
	SI	ŀ	í	ú	ś	
	SI	ŀ	í	ú	ś	
	SI	ŀ	í			
	SI					
	SI					
	SI	ŀ		ú		
	SI					
	SI					
	SI					

Question 7 continued	blank
Question / continued	
	Q7
(Total 6 marks)	



8. A curve C has equation y = f(x), where

$$f(x) = \arcsin\left(\frac{1}{2}x\right)$$
 $-2 \leqslant x \leqslant 2$ $-\frac{\pi}{2} \leqslant y \leqslant \frac{\pi}{2}$

(a) Sketch C.

(1)

(b) Given $x = 2 \sin y$, show that

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{1}{\sqrt{A - x^2}}$$

where *A* is a constant to be found.

(3)

The point P lies on C and has y coordinate $\frac{\pi}{4}$

- (c) Find the equation of the tangent to C at P. Write your answer in the form y = mx + c, where m and c are constants to be found.
 - **(3)**

	blank
Question 8 continued	



estion 8 continued	

	ý					
١,		2				
					ì	
ς	3	Б	7	7	7	
۶	Ì	Ē	2	5	Z	
⟨	1	ŀ	2		2	
>	j	ŀ	3	2	2	
>	1	ŀ	3		3	
>	3				3	
>	3				3	
					3	
					3	
>					3	
					3	
>					3	
>					3	
\ \ \ \					3	
\ \ \					3	
\ \ \ \					3	
> > >					3	
> > > >					3	
> > > >					3	
> > > >					3	
> > > >					3	
> > > >					3	
					3	
					3	
					3	
					3	
					3	
					3	
					3	
					3	
					3	
					3	
					3	
					3	
					3	
					3	

Question 8 continued	blank
	Q8
(Total 7 marks)	
(2000 / 1000 100)	



9.




Figure 3

Figure 3 shows a sketch of part of the curve with equation y = f(x), where

$$f(x) = x(x^2 - 4)e^{-\frac{1}{2}x}$$

(a) Find f'(x).

(2)

The line l is the normal to the curve at O and meets the curve again at the point P.

The point *P* lies in the 3rd quadrant, as shown in Figure 3.

(b) Show that the x coordinate of P is a solution of the equation

$$x = -\frac{1}{2}\sqrt{16 + e^{\frac{1}{2}x}} \tag{4}$$

(c) Using the iterative formula

$$x_{n+1} = -\frac{1}{2}\sqrt{16 + e^{\frac{1}{2}x_n}}$$
 with $x_1 = -2$

find, to 4 decimal places,

- (i) the value of x_2
- (ii) the x coordinate of P.

(3)

	blank
Question 9 continued	Dialik
Question 9 continued	



	Leave blank
Question 9 continued	

K					>
0	ς.				C
7			4	à	
			į		
				5	
				Ì	
	S				
K	ы		è	ŝ	è
				Ĺ	
	ď				
			Þ		
				2	
2	۲.	ė	ú	á	ø
Ν	7	۹	R	ь	2
1	1	ć	×		ń
7	c	٩	7		
ĸ.	N	6			
	ΚI		_	2	_
К	4	ľ	7	٠	7
.2	1	۶	7	é	
	Λ		,	۹	к
Z	М	ľ			
5	0		6	ä	ν
ĸ.	×	_			3
	⟨#	,	₹		P
Κ		á		۲	2
2	V				
1	2		1		/
/	1		7	2	S
	a		•	9	6
K	И	ľ			1
	⟨₹		6	a	ν
Κ	×			c	>
2	V		×	ø	ħ.
1	4	K	7		
/	М		'n	н	
7					
ĸ.					

	Ŀ				
	É				
		S			
Xį					
	X				
X					
	ſ				
				4	
a					
M	ы	ú	à		

×				
>				
ĸ.				
/				
٨,				
×.		0		
	C.	۷	ú	
	₹	9		h
١,	6	7		ş
N				
	a	٤	2	ú
. А	r	Я		7
N				b
/				Z
7	ú			۰
ΚI		'n		۷
. 2		7		9
А.				
1	À.		'n	ú
a	£	à	6	i
А				
×	6			
Q	\leq	4	_	2
-3	7	3		
Þ	ę	ş	2	ģ
2	S	Ž	3	
3			2	
3	2		2	
3			2	4
3	Ì		2	
3				
3				
3				

Question 9 continued	blank
Question 5 continued	
	Q9
(Total 9 marks)	
(1.1.1 / 1.1.1	



10.




Figure 4

Figure 4 shows a sketch of part of the curve with equation

$$y = (1 + 2\cos 2x)^2$$

(a) Show that

$$(1 + 2\cos 2x)^2 \equiv p + q\cos 2x + r\cos 4x$$

where p, q and r are constants to be found.

(2)

The curve touches the positive x-axis for the second time when x = a, as shown in Figure 4.

The regions bounded by the curve, the y-axis and the x-axis up to x = a are shown shaded in Figure 4.

(b) Find, using algebraic integration and making your method clear, the exact total area of the shaded regions. Write your answer in simplest form.

- 1	_	1
•		
٠,	\sim	,



×.	a	ы		Р	
V/I		2		Ŀ	
$^{\sim}$			7	9	
×i		Í	è		
a	L		Ľ		
/\	Z	Ξ	7	3	
va		ĸ		ρ	r
	Ľ	3	Ľ	-	è
×.	7	₹	7	5	
1	S	a			þ
\sim	Ġ				
×.		7	۹		
\sim					
×.	2			۷	
V.	ĸ.	4	r		
	۹	Ρ,		¥	
Χi	è	ú	ù	ú	
< 2		2	C		
×					
\checkmark		4			^
$^{\prime}$	μ	ц	μ	١	
×ι	è				
OI					
\sim	r				
\vee			١		
×	2	a	ĕ	ĸ	
a			'n	ú	
VI	9	-	٠	e	
\sim					
			ũ	Ý	
a	Ľ	ĺ	Ĺ	j	ŀ
Ŋ	Ĺ	į	ľ	ĺ	ł
8		1	Š	1	
				1	
	ŀ				

	Leave
	blank
Question 10 continued	



XXX
CX X
(XX
XXX
X X X iii
XXX
E CX
- XXX

estion 10 continued	
	/TP / 1 M
	(Total 7 marks)

END