Please check the examination deta	ails below before ent	ering your candidate information
Candidate surname		Other names
Pearson Edexcel International Advanced Level	Centre Number	Candidate Number
Time 1 hour 30 minutes	Paper referenc	WMA13/01
Mathematics		
International Advance Pure Mathematics P3	d Level	
You must have: Mathematical Formulae and Stat	tistical Tables (Ye	ellow), calculator

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use black ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 9 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
- use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.
- Good luck with your examination.

Turn over ▶







1. The curve C has equation

$$y = x^2 \cos\left(\frac{1}{2}x\right) \qquad 0 < x \leqslant \pi$$

The curve has a stationary point at the point P.

(a) Show, using calculus, that the x coordinate of P is a solution of the equation

$$x = 2\arctan\left(\frac{4}{x}\right)$$

(4)

(3)

Using the iteration formula

$$x_{n+1} = 2\arctan\left(\frac{4}{x_n}\right) \qquad x_1 = 2$$

(b) find the value of x_2 and the value of x_6 , giving your answers to 3 decimal places.

Question 1 continued	



2. (a) Show that

$$\frac{1 - \cos 2x}{2\sin 2x} \equiv k \tan x \qquad x \neq (90n)^{\circ} \qquad n \in \mathbb{Z}$$

where k is a constant to be found.

(3)

(b) Hence solve, for $0 < \theta < 90^{\circ}$

$$\frac{9(1-\cos 2\theta)}{2\sin 2\theta} = 2\sec^2\theta$$

giving your answers to one decimal place.

(Solutions based entirely on graphical or numerical methods are not acceptable.)



$\langle \times \rangle$	XX.
\approx	\bowtie
$\stackrel{\sim}{\sim}$	$\times\!\!\times\!\!\times$
88	$\times\!\!\times\!\!\times$
\approx	\approx
\bowtie	$\otimes\!\!\!\otimes$
\otimes	$\times\!\!\times$
$\stackrel{\sim}{\sim}$	$\times\!\!\times\!\!\times$
$\times\!\!\!\times$	\ggg
\otimes	$\times\!\!\times$
×	WRITE IN THIS AREA
\approx	NV.
\bowtie	
\otimes	
\otimes	30
\approx	$\times\!\!\times\!\!\times$
X	
X	
\otimes	
×	$\times\!\!\times\!\!\!\times$
\otimes	
\otimes	
\otimes	ŮŮ,
XXI	\sim
$\otimes \otimes$	
X	
\otimes	
\otimes	
紷	$\otimes \otimes$
\otimes	\approx
\otimes	\sim
XX:	
X	$\times\!\!\!\times$
\propto	DO NOT
\otimes	*
\bowtie	
*	$\times\!\!\times\!\!\!\times$
\Leftrightarrow	$\times\!\!\times\!\!\times$
88	$\times\!\!\times\!\!\times$
≫	$\otimes \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \!$
\otimes	$\times\!\!\times\!\!$
\Leftrightarrow	
X	$\times\!\!\times$
\Rightarrow	$\otimes \otimes$
$\times\!\!\!\times$	$\approx \approx$
$\otimes \! >$	∞
\otimes	\ggg
\otimes	$\times\!\!\times$
\otimes	$\times\!\!\times\!\!\times$
\otimes	\approx
\otimes	\approx
\times	$\times\!\!\!\times$
\times	3 %
X	MAL
XX	ar.
X	XXX
\otimes	
XX	M.
\otimes	
\otimes	
	Ė
	ENTHIS AREA
	THE TANK
	MENNTHR
	RIE IN THE
	WRITEINTH
	I WRITE IN THE
	ALWRITEHN THE
	OT WRITE IN THE
	NOTWERENIEN
	NOTWRITEINTHE
	ONOTWRITEINTHE
	DO NOT WRITE
	DO NOT WRITE IN THE
	DO NOT WRITE
	IN THIS AREA DO NOT WRITE
	IN THIS AREA DO NOT WRITE
	IN THIS AREA DO NOT WRITE
	IN THIS AREA DO NOT WRITE
	IN THIS AREA DO NOT WRITE
	IN THIS AREA DO NOT WRITE
	IN THIS AREA DO NOT WRITE
	IN THIS AREA DO NOT WRITE
	IN THIS AREA DO NOT WRITE
	IN THIS AREA DO NOT WRITE
	IN THIS AREA DO NOT WRITE
	DO NOT WRITE

	Leave
	blank
Question 2 continued	



Question 2 continued	Leave blank

Question 2 continued	blank
Quostion – continuou	
	Q2
(Total 9 marks)	



3. (i) Find

$$\int \frac{12}{\left(2x-1\right)^2} \mathrm{d}x$$

giving your answer in simplest form.

(2)

(ii) (a) Write $\frac{4x+3}{x+2}$ in the form

$$A + \frac{B}{x+2}$$
 where A and B are constants to be found

(b) Hence find, using algebraic integration, the exact value of

$$\int_{-8}^{-5} \frac{4x+3}{x+2} \, \mathrm{d}x$$

giving your answer in simplest form.

(6)

$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$

>>>>>
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
\times
$\times\!\!\times\!\!\times\!\!\times$
$\times \times $
$\otimes \Rightarrow \otimes$
\times
$\otimes \mapsto$
\times
\times
$\times \odot \times$
DO NOT WRITE IN THIS AREA
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$

$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$

>>>>>
$\times\!\!\times\!\!\times\!\!\times$
\times
\times 0 $^{\circ}$
$\times \underline{v}$
\times
2
Z
TE HN
FTERN
RITEIN
VRITEIN
WRITEIN
T WRITE IN
OT WRITE IN
IOT WRITE IN
NOT WRITE IN
O NOT WRITE IN THIS AREA
$\times\!\!\!<\!\!\!\!<\!\!\!\!<\!\!\!\!>$
DO NOT WRITE IN
$\times\!\!\!<\!\!\!\!<\!\!\!\!<\!\!\!\!>$
$\times\!\!\!<\!\!\!\!<\!\!\!\!>$
DO
DQ b
DQ b
REA DO
REA DO
AREA DO
S AREA DO
S AREA DO
HIS AREA DO
THIS AREA DO
THIS AREA DO
THIS AREA DO
IN THIS AREA DO
E IN THIS AREA DO
TE IN THIS AREA DO
TE IN THIS AREA DO
TE IN THIS AREA DO
TE IN THIS AREA DO
I WRITE IN THIS AREA DO
I WRITE IN THIS AREA DO
OT WRITE IN THIS AREA DO
NOT WRITE IN THIS AREA DO
NOT WRITE IN THIS AREA DO
OT WRITE IN THIS AREA DO

	Leave
	blank
Question 3 continued	



Question 3 continued	Leave blank

Question 3 continued		blan
Question 5 continued		
		Q3
	(Total 8 marks)	
	. , ,	



4. The functions f and g are defined by

$$f(x) = \frac{4x+6}{x-5} \qquad x \in \mathbb{R}, \ x \neq 5$$

$$g(x) = 5 - 2x^2 \qquad x \in \mathbb{R}, \ x \leqslant 0$$

(a) Solve the equation

$$fg(x) = 3$$

(4)

(b) Find f^{-1}

(3)

(c) Sketch and label, on the same axes, the curve with equation y = g(x) and the curve with equation $y = g^{-1}(x)$. Show on your sketch the coordinates of the points where each curve meets or cuts the coordinate axes.





Question 4 continued	blank



	Leave blank
Question 4 continued	Olulik

Ougst	on 4 continued		
Questi	on 4 continued		
			(Total

5.




Figure 1

The growth of duckweed on a pond is being studied.

The surface area of the pond covered by duckweed, Am^2 , at a time t days after the start of the study is modelled by the equation

$$A = p q^t$$
 where p and q are positive constants

Figure 1 shows the linear relationship between $\log_{10} A$ and t.

The points (0, 0.32) and (8, 0.56) lie on the line as shown.

(a) Find, to 3 decimal places, the value of p and the value of q.

(4)

Using the model with the values of p and q found in part (a),

(b) find the rate of increase of the surface area of the pond covered by duckweed, in m^2/day , exactly 6 days after the start of the study. Give your answer to 2 decimal places.





	Leave
	blank
Question 5 continued	



	Le
Question 5 continued	

Question 5 continued	blank
	Q5
(Total 7 marks)	
(==:01 : 2101 ===)	



6. Given that k is a positive constant,

- (a) on separate diagrams, sketch the graph with equation
 - (i) y = k 2|x|

(ii)
$$y = \left| 2x - \frac{k}{3} \right|$$

Show on each sketch the coordinates, in terms of k, of each point where the graph meets or cuts the axes.

(4)

(b) Hence find, in terms of k, the values of x for which

$$\left| 2x - \frac{k}{3} \right| = k - 2|x|$$

giving your answers in simplest form.

(4)



Question 6 continued	blank



Question 6 continued	Le

(×××××)

A.
$\times \times \times \times$
$\times \square \times$
\times
<u> </u>
$\times = \times$
X EX
**
\times
\times
×>×
$\otimes Q \times$
Ď

XXXX

XXXX
A
\times
<u> </u>
\times
Ē
2
- X
2
RITEINTE
- X
/RITE IN TH
WRITEINTH
WRITEINTH
OT WRITE IN TH
EA DO NOT WRITE IN TH
A DO NOT WRITE IN TH
EA DO NOT WRITE IN TH
AREA DO NOT WRITE IN TH
IS AREA DO NOT WRITE IN TH
HIS AREA DO NOT WRITE IN TH
THIS AREA DO NOT WRITE IN TH
HIS AREA DO NOT WRITE IN TH
IN THIS AREA DO NOT WRITE IN TH
E IN THIS AREA DO NOT WRITE IN TH
ITE IN THIS AREA DO NOT WRITE IN TH
RITE IN THIS AREA DO NOT WRITE IN TH
WRITE IN THIS AREA DO NOT WRITE IN TH
WRITE IN THIS AREA DO NOT WRITE IN TH
WRITE IN THIS AREA DO NOT WRITE IN TH
WRITE IN THIS AREA DO NOT WRITE IN TH
WRITE IN THIS AREA DO NOT WRITE IN TH
WRITE IN THIS AREA DO NOT WRITE IN TH

	Leave blank
Question 6 continued	Oldlik
	Q6
(Total 8 marks)	
(Total o marks)	



7. Given that

$$x = 6\sin^2 2y \qquad 0 < y < \frac{\pi}{4}$$

show that

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{1}{A\sqrt{(Bx - x^2)}}$$

where A and B are integers to be found.

(5)

Question 7 continued		blank
		Q7
	(Total 5 marks)	
	(10tai 5 marks)	



8. A scientist is studying a population of fish in a lake. The number of fish, N, in the population, t years after the start of the study, is modelled by the equation

$$N = \frac{600e^{0.3t}}{2 + e^{0.3t}} \qquad t \geqslant 0$$

Use the equation of the model to answer parts (a), (b), (c), (d) and (e).

(a) Find the number of fish in the lake at the start of the study.

(1)

(b) Find the upper limit to the number of fish in the lake.

(1)

(c) Find the time, after the start of the study, when there are predicted to be 500 fish in the lake. Give your answer in years and months to the nearest month.

(4)

(d) Show that

$$\frac{\mathrm{d}N}{\mathrm{d}t} = \frac{A\mathrm{e}^{0.3t}}{\left(2 + \mathrm{e}^{0.3t}\right)^2}$$

where *A* is a constant to be found.

(3)

Given that when t = T, $\frac{dN}{dt} = 8$

(e) find the value of *T* to one decimal place.

(Solutions relying entirely on calculator technology are not acceptable.)

(4)

	Leave
	blank
Question 8 continued	



Question 8 continued	



9. (a) Express $12\sin x - 5\cos x$ in the form $R\sin(x - \alpha)$, where R and α are constants, R > 0 and $0 < \alpha < \frac{\pi}{2}$. Give the exact value of R and give the value of α in radians, to 3 decimal places.

(3)

The function g is defined by

$$g(\theta) = 10 + 12\sin\left(2\theta - \frac{\pi}{6}\right) - 5\cos\left(2\theta - \frac{\pi}{6}\right) \qquad \theta > 0$$

Find

- (b) (i) the minimum value of $g(\theta)$
 - (ii) the smallest value of θ at which the minimum value occurs.

(3)

The function h is defined by

$$h(\beta) = 10 - (12\sin\beta - 5\cos\beta)^2$$

(c) Find the range of h.



				Ì	Ř
	Ý				
			2		4
	3		9	۳	-5
	Ó				ę
		С	1	К	7
	Ç,	÷		ы	۷
	2	٠	9	٠	-
	а	ø			
	И		à	ń	è
				b	
′	y	P	4	þ	Ą
<	3	ì	9		Ž
5	V		Ĺ	-	۷.
5	V		Ĺ	-	۷.
>	V		Ĺ	-	۷.
5	V		Ĺ	-	۷.
5	V		Ĺ	-	۷.
>	V		Ĺ	-	۷.
>	V		Ĺ	-	۷.
	V		Ĺ	-	۷.
>	V		Ĺ	-	۷.
	V		Ĺ	-	۷.
>	V		Ĺ	-	۷.
>	V		Ĺ	-	۷.
	V		Ĺ	-	۷.
<	V		Ĺ		۷.
	V		Ĺ		
	V		Ĺ		
	V				
<	V				
	V				
	V				
	V				
	V				
	V				
	V				

	Leave
	blank
Question 9 continued	

