Please check the examination deta	ails below	before ente	ering your candidate information
Candidate surname			Other names
Pearson Edexcel International Advanced Level	Centre	e Number	Candidate Number
Monday 8 Ju	ne 2	2020	0
Morning (Time: 1 hour 30 minute	es)	Paper R	Reference WFM03/01
Mathematics			
International Advance Further Pure Mathema		,	y/Advanced Level
You must have: Mathematical Formulae and Sta	tistical	Tables (Blu	lue), calculator

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 8 questions in this question paper. The total mark for this paper is 75.
- The marks for each question are shown in brackets
 use this as a quide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over ▶







1.	(a)	Use the definition of $\sinh x$ in terms of exponentials to show that
		$\sinh 3x \equiv 4\sinh^3 x + 3\sinh x$
		(2)
	(b)	Hence determine the exact coordinates of the points of intersection of the curve with equation $y = \sinh 3x$ and the curve with equation $y = 19 \sinh x$, giving your answers as simplified logarithms where necessary. (5)

				3	
	я		b	4	7
		P	7	7	5
	ś				
		К		ú	ú
	۹		ĕ	ú	d
Χ		7	7	7	5
		S	á		К
				'n	4
	9	-	٥	۰	٠
V					
		a	i	ú	Š
		۴	•	9	
		Ľ			3
	4	۹	H		ν
		ú	ä	S	٥
	à	и	۴	9	٠
					1

DO NOT WRITE IN THIS AREA

On a thought a supply and a supply a su	Leave blank
Question 1 continued	
	Q1
(Total 7 marks)	



2.	Deteri	nine

(i)
$$\int \frac{1}{3x^2 + 12x + 24} \, \mathrm{d}x$$

(4)

(ii)
$$\int \frac{1}{\sqrt{27 - 6x - x^2}} \, \mathrm{d}x$$

(4)



Ň		
Ö.		
	Ž	

	Leave
	blank
Overtion 2 continued	Olulik
Question 2 continued	
-	



estion 2 continued	

		ķ		
	į			
		ы	ī	

Question 2 continued	Leave blank
	Q2
(Total 8 marks)	



3.

$$\mathbf{M} = \begin{pmatrix} 3 & -4 & k \\ 1 & -2 & k \\ 1 & -5 & 5 \end{pmatrix} \text{ where } k \text{ is a constant}$$

Given that 3 is an eigenvalue of M,

(a) find the value of k.

(3)

(b) Hence find the other two eigenvalues of M.

(3)

(c) Find a normalised eigenvector corresponding to the eigenvalue 3

(3)



Ň		
Ö.		
	Ž	

Ornerties 2 continued	Leave blank
Question 3 continued	



nestion 3 continued	

Leave

Ŋ			
		2	
	Ž		
	Û	ø	
	Û	ø	
	Û	ø	
	Û	ø	ĺ
	Û	ø	ĺ
	Û	ø	į
	Û	ø	
	Û	ø	
	Û	ø	
	Û	ø	
	Û	ø	
	Û	ø	

	V	A.		2
		ă		
		X		
	Ğ			
	Si			
	ă			
		×	z	_
\	Δ	Z		ø,
	V		<u> 2</u>	Z
7	/1	₹	7	в,
0		1		
		Σ'n	×	
	21	0	4	ĸ.
?	U	Z.	Δ	ď
1	7	91	w	n
	Y	à	6	ď
		к	78	ĸ.
	и	ப		к

Question 3 continued	blank
	Q3
(Total 9 marks)	



(5)

 $I_n = \int x^n \cos x \, \mathrm{d}x$

4.

(a) Show that, for $n \ge 2$

$$I_n = x^n \sin x + nx^{n-1} \cos x - n(n-1)I_{n-2}$$
(4)

(b) Hence find the functions f(x) and g(x) such that

$$\int x^4 \cos x \, dx = f(x) \sin x + g(x) \cos x + c$$

where c is an arbitrary constant.

	blank
Question 4 continued	Olalik
Question 4 continued	



estion 4 continued	

Ė	
Ŝ	
2	
Û	

Ornertian A continued	Leave blank
Question 4 continued	
	Q4
(Total 9 marks)	



(4)

5.	The hyperbola H has equation	x^2	y^2	- 1
	The hypercola II has equation	25 –	4	- ,

The line l has equation y = mx + c, where m and c are constants.

Given that l is a tangent to H,

- (a) show that $25m^2 = 4 + c^2$
- (b) Hence find the equations of the tangents to H that pass through the point (1, 2). (5)
- (c) Find the coordinates of the point of contact each of these tangents makes with H. (3)



Ň		
Ö.		
	Ž	

uestion 5 continued	



estion 5 continued	

	É		
	Ú		
	à		
		i	
Ì			
ì			
Ì			
	K		
á			
	ķ		
ñ			
į			
	2		
ì			

Ŷ				
	И			
	Ó			
×ĵ				
	ø	ij	₹	5
O.	ę	P	5	þ
	Ź	5	5	ζ
	Z	>	ξ	Ś
Si	8	2	\$	Š
			5	
			2	}

$\wedge \wedge \wedge$	\sim	
$\times \times$		
$\times \times \times$		
$\times \times \times$		
Ž		
$\times \times \mathbb{K}$	732	
	M.	
$\wedge \wedge \Delta$		
	, and	
CX 26 %	X	
XXII I	26	
884	¥	
X 4	¥	
<u> </u>	¥	
	¥	
	¥	
	¥ - 3	
110		
11 C C C C C C C C C C C C C C C C C C		
ATSIGNA TELEPE		
THOM		
HARIOWA.		
F.VAVIDATE		
T W/Olte		
THE WAY DITTE		
T. W.D.T.E		
THE VALIDITE		
OX WEDITE		
OF WORTE		
PART VICTORIES		
STREET WAS INTE		
NOT WIDITE		
ANOT WESTER		
ALCAL MADITE		
ATION WIDITE		
A NICET VINDITE		
A MICH WINDSHIE		
STREET WINDS		
STICK TOWN		
BELOW TOWN		
THE WINDS		
THOM TON OU		
DO NOT WOLTEN THIS ABOA		
PIC MICH WIDITE		

	Leave	
Question 5 continued	blank	
Question 5 continued		
	05	
	Q5	1
(Total 12 marks)		
(Iour 12 marks)		J



6.

$$\mathbf{A} = \begin{pmatrix} 1 & -1 & 1 \\ 1 & 1 & 1 \\ 1 & 2 & a \end{pmatrix} \quad a \neq 1$$

(a) Find \mathbf{A}^{-1} in terms of a.

(4)

$$\mathbf{B} = \begin{pmatrix} 1 & -1 & 1 \\ 1 & 1 & 1 \\ 1 & 2 & 4 \end{pmatrix}$$

The straight line l_1 is mapped onto the straight line l_2 by the transformation represented by the matrix **B**.

The equation of l_2 is

$$(\mathbf{r} - (12\mathbf{i} + 4\mathbf{j} + 6\mathbf{k})) \times (-6\mathbf{i} + 2\mathbf{j} + 3\mathbf{k}) = \mathbf{0}$$

(b) Find a vector equation for the line l_1

(4)

¢	
Ž	
Ĭ	
\{\bar{\chi}{\chi}\}	

uestion 6 continued	



uestion 6 continued	
lestion o continued	

	N	
	È	
	Š	

Question 6 continued	Leave blank
	Q6
(Total 8 marks)	20
(Total o marks)	



blank

(7)

7. The curve C has parametric equations

$$x = \cosh t + t$$
, $y = \cosh t - t$ $0 \le t \le \ln 3$

(a) Show that

$$\left(\frac{\mathrm{d}x}{\mathrm{d}t}\right)^2 + \left(\frac{\mathrm{d}y}{\mathrm{d}t}\right)^2 = 2\cosh^2 t \tag{3}$$

The curve C is rotated through 2π radians about the x-axis. The area of the curved surface generated is given by S.

(b) Show that

$$S = 2 \sqrt{2} \int_0^{\ln 3} \left(\cosh^2 t - t \cosh t \right) dt$$
 (2)

(c) Hence find the value of S, giving your answer in the form

$$\frac{\pi\sqrt{2}}{9}(a+b\ln 3)$$

where a and b are constants to be determined.

	9	×	ρ	₹
	3			5
	2	ì	9	Š
Š	ì			ì

	Leave
	blank
Question 7 continued	
Question / continued	



	Leave blank
Question 7 continued	

	É	
	-	Š
	ø	
K :		
	70	20
	æ	\sim
\sim		$\langle \cdot \rangle$
×	×	X.
	o	96
\sim	ES.	л
V	e.	ы
	~	_
	æ	m.
\sim	W.	71
V1	12	SA

	Leave blank
Question 7 continued	
	0=
	Q7
(Total 12 marks)	



blank

8. The plane Π_1 has equation

$$x - 5y + 3z = 11$$

The plane Π_2 has equation

$$3x - 2y + 2z = 7$$

The planes $\Pi_{\rm l}$ and $\Pi_{\rm 2}$ intersect in the line l.

(a) Find a vector equation for l, giving your answer in the form $\mathbf{r} = \mathbf{a} + \lambda \mathbf{b}$ where \mathbf{a} and \mathbf{b} are constant vectors and λ is a scalar parameter.

(5)

The point P(2, 0, 3) lies on Π_1

The line m, which passes through P, is parallel to l.

The point Q(3, 2, 1) lies on Π_2

The line n, which passes through Q, is also parallel to l.

(b) Find, in exact simplified form, the shortest distance between m and n.

- 1	(5	
	U	ı

	blank
Question 8 continued	



	Leave blank
Question 8 continued	

	è	
	Î A	
		œ
	ŝ	
		Ž
	-	g/P
	1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	
	7	5
	×	
	16	×
	Ž	
	-	75
	$\langle \rangle$	
	o	96
	r.	41
~	w	ы
	/4	~
	ыd	m.

Question 8 continued blank



Question 8 continued		Leave blank
		Q8
	(Total 10 marks)	
END	TOTAL FOR PAPER: 75 MARKS	