ow before ente	ering your candidate information
	Other names
umber	
nation	al Advanced Level
Paper reference	WMA12/01
	0
ıhsidiar	y/Advanced Level
ab Sididi	y//tavalicea Ecvel
al Tables (Ye	ellow), calculator
	nation Paper reference

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 - there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 10 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶







1.




Figure 1

Figure 1 shows a sketch of part of the curve with equation y = f(x)

The table below shows some corresponding values of x and y for this curve.

The values of y are given to 3 decimal places.

X	-1	-0.5	0	0.5	1
у	2.287	4.470	6.719	7.291	2.834

Using the trapezium rule with all the values of y in the given table,

(a) obtain an estimate for

$$\int_{-1}^{1} f(x) dx$$

giving your answer to 2 decimal places.

(3)

(b) Use your answer to part (a) to estimate

(i)
$$\int_{-1}^{1} (f(x)-2) dx$$
(ii)
$$\int_{1}^{3} f(x-2) dx$$

(ii)
$$\int_{1}^{3} f(x-2) dx$$

(3)

XXXXX
X 4 X
XXXXXXX
× CO
$\times \mathbb{R}^{\times}$
INTHIS
(XXXXXX)
× E ×
8
ō
000
XXXXX
$\sim\sim\sim\sim$
NAME OF THE PERSON OF THE PERS
<u>\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\</u>
7
2
YY YX YY -
XX
$\times\times\times\times\times$
<u> </u>
×Δ×
\times
A
A
REA
YREA
S AREA
S AREA
THIS AREA
ITHIS AREA
IN THIS AREA
IN THIS AREA
TE IN THIS AREA
TE IN THIS AREA
RITE IN THIS AREA
VRITE IN THIS AREA
EWRITE IN THIS AREA
T WRITE IN THIS AREA
VOT WRITE IN THIS AREA
NOT WRITE IN THIS AREA
O NOT WRITE IN THIS AREA
30 NOT WRITE IN THIS AREA
O NOT WRITE IN THIS AREA
30 NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
30 NOT WRITE IN THIS AREA

Κ		6	ì	ú	D)	
Q.	7	ä		۴		
			ú	ù	К	
Κ	9	S	_	Z	⊃	
ė	۳	7	7	7	٦	
	1		Ç.		c	
K		SJ	ø		D	
Ü		ш	ь	d	К	
	Z				7	
K		ú	è	6	e)	
		7			۲.	
	ä	è	ú	à	ď	
K	7	z	3		5	
	1	7		7	۲.	
^		'n		'n	v	
K	2	Z	3			
3	₹	2			ĸ	
^					7	
Z.	Z	Z			Ŋ	
	₹	7	ς		К	
۲	2	۲			2	
×		4	4	4	9	
2		7	₹	7	٣.	
×		ú		s		
1		7		7	N	
		ù	4		ĸ	
К	. 2	₹	7	Κ		
i	ч	,			Б,	
	sa.		Р		0	
	9	ę	9	ę	10	
	×	2	۲		1	
	4	à	×	à	2	
×	а	٣		3	D	
	۹	ú	≤		ĸ.	
`	Ì	3	z	5	1	
S	4	5	3	3	K	
Š	ı	2	₹	3	K	
Š		2	3		Ķ	
2	1	3	3	3	5	
2	1	3	3	2	5	
× × ×	1	3	3	3	5	
X	1	3	3		 	
~ < < < <			3		K 2	
~ < < < < <	1				K	
× < < < < < <	1				 	
× < < < < <	1		3			
×< < < < < < <	1					
~<<<<<	1					
~<<<<<	1					
~ < < < < < < < < < < < < < < < < < < <	1					
~ < < < < < < < < < < < < < < < < < < <	1					
~ < < < < < < < < < < < < < < < < < < <	1					
~ < < < < < < < < < < < < < < < < < < <						
~ < < < < < < < < < < < < < < < < < < <						
~ < < < < < < < < < < < < < < < < < < <						
~ < < < < < < < < < < < < < < < < < < <						
~ < < < < < < < < < < < < < < < < < < <						
~ < < < < < < < < < < < < < < < < < < <						
~ < < < < < < < < < < < < < < < < < < <						
~<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<						
~<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<						

S		S		N
		١.		◟
			\	
		\leq	4	×
		е		г
>	7	9		
	2	۲.		Z
		4		48
	2	ś.	2	4
		7		7
V				-
			2	Z
	ú	5		7
	9	٠		L
7	١	2	7	7
		`		\
	ò	ζ,		ь.
y		a	r	3
ς.	7	₹		ক
	۱	н	н	m)
\	Z	\sim	2	\sim
	۹			_
	۷	3	E.	2
	9	÷	7	₹
7				
			,	
	۰			
	μ	=		
	S	d	и	7
	3	Z	3	Z
S	ē	Ş	Š	Ç
8	3	7	2	
5	2	Š	2	
3		3	Š	4
5		3		4
5		2	2	4
8				4

Question 1 continued
(Total for Organian 1 is 6 mayles)
(Total for Question 1 is 6 marks)



2. In this question you must show all stages of your working.

Solutions based entirely on calculator technology are not acceptable.




Figure 2

A brick is in the shape of a cuboid with width x cm, length 3x cm and height h cm, as shown in Figure 2.

The volume of the brick is 972 cm³

(a) Show that the surface area of the brick, $S \text{ cm}^2$, is given by

$$S = 6x^2 + \frac{2592}{x}$$

(3)

(b) Find
$$\frac{dS}{dx}$$

(1)

(c) Hence find the value of *x* for which *S* is stationary.

(2)

(d) Find $\frac{d^2S}{dx^2}$ and hence show that the value of x found in part (c) gives the minimum value of S.

(2)

(e) Hence find the minimum surface area of the brick.

(1)

XXXXX

$\times\!\!\times\!\!\times\!\!\times$
$\otimes \otimes \otimes$
$\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
XXXX
DO NOT WRITE IN THIS AREA
XMAX
\times ω

\times
$\times \times \times$
\times
$\times \times \times \times$
$\otimes \otimes$
\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$

XXXX
$\times\!\!\times\!\!\times\!\!\times$

$\times\!\!\times\!\!\times\!\!\times$

$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
XXXX
$\times\!\!\times\!\!\times\!\!\times$
AREA
X
XX
\otimes
\times
\times
NOT WRITE IN THIS
<u> </u>
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$

$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$

$\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
$\times\!\!\times\!\!\times\!\!\times$
22
T.
AR
ARE
SARE
IIS ARE
HIS ARE
THIS ARE
THIS ARE
V THIS ARE
INTHIS ARE
IN THIS ARE
E IN THIS AREA
FE IN THIS ARE
ITE IN THIS ARE
RITE IN THIS ARE
RITE IN THIS ARE
WRITE IN THIS ARE
WRITE IN THIS ARE
WRITE IN THIS ARE
IT WRITE IN THIS ARE
OT WRITE IN THIS ARE
JOT WRITE IN THIS ARE
NOT WRITE
DO NOT WRITE IN THIS ARE
NOT WRITE
NOT WRITE

uestion 2 continued	



Question 2 continued

->>>
-
× va×
-XX <u>IIX</u> XX
-XXXXXX
-888-888
-000
×9×
Ā
EA
IN THIS AREA
ITE IN THIS AREA
ITE IN THIS AREA
ITE IN THIS AREA
ITE IN THIS AREA
WRITE IN THIS AREA
WRITE IN THIS AREA
WRITE IN THIS AREA
WRITE IN THIS AREA
WRITE IN THIS AREA
WRITE IN THIS AREA
WRITE IN THIS AREA
NOT WRITE IN THIS AREA
NOT WRITE IN THIS AREA
NOT WRITE IN THIS AREA
MOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 2 continued	
	(Total for Question 2 is 9 marks)



- 3. $f(x) = \left(2 + \frac{kx}{8}\right)^7$ where k is a non-zero constant
 - (a) Find the first 4 terms, in ascending powers of x, of the binomial expansion of f(x). Give each term in simplest form.

(4)

Given that, in the binomial expansion of f(x), the coefficients of x, x^2 and x^3 are the first 3 terms of an arithmetic progression,

(b) find, using algebra, the possible values of k.

(Solutions relying entirely on calculator technology are not acceptable.)

(3)

2						
/						
2					Κ	
	$^{\sim}$		7			
	×					
b	< .	٤	á		d	
ς	м				υ	
				7	٩	
>	Я		Ġ	Ė	4	
	И		3	Ŀ	4	
		3	Σ	7	3	
2	4		Ì		۰	r
⋖	Я		ì		ė	
2		2			Z	5
\	a	i			P	۴.
	м	۹	è		Ŀ	
				7	₹	
1	$^{\prime}$			\		
	M		7			
	S			r	Ч	
۲	Ğ	3	2	S	2	
		۰	7	۲	-	
	4					
			А	I.	2	
	V					
	B	ŕ				
	м					
7	V		١.	ì		
	×					
2	ς.	2	ы	ρ	•	
1	ß				Ì	
	Σi	ì	ù	6	ũ	
2		7		7		
		١,	/			
	K		ď	ľ	1	
2	ΚI		4			
۲	Ä					
1	M		'n	Ħ	ù	
0	4	ŀ			К	
					i	
	ġ	2	ż	2	Š	
3	å	9	Ē	ĺ	Ş	ķ
2	Ś			ľ	5	i.
3	j			į		
3	Š					
3						
3						
?						
		5	₹	5		
3		5	₹	5		
?		5	5	\ =	3	
?		5	2		3	
		5	2		3	
		5	2		3	
		5				
		5				
		5				
		5				
		5				
		5				
		5				
		5				

Question 3 continued
(Total for Question 3 is 7 marks)



4. (i) Using the laws of logarithms, solve

$$\log_3(4x) + 2 = \log_3(5x + 7) \tag{3}$$

(ii) Given that

$$\sum_{r=1}^{2} \log_{a}(y^{r}) = \sum_{r=1}^{2} (\log_{a} y)^{r} \qquad y > 1, a > 1, y \neq a$$

find y in terms of a, giving your answer in simplest form.

(3)

		5			
х,	×		>		
\sim	v	Ś	4		
OI.					
X	7	7	7	b	
X1	к	í	ь	di	
(XI	Ŀ		L		
×	7		7	₹	
V)			И	۰	
OI.	Ľ	ä	ĸ.	4	
X.	$\overline{}$	ī	7		
V.	ú	á		۳	
(N		ĥ		Ŀ	
			7	₹	
$^{\sim}$			١	ᄼ	
	r	S	ø	۰	
×Ι	ы		D		
V	Ξ	2	2	z	
O.	-	5	7	7	
\times 1	×			۰	
\vee	v			2	
\triangle			μ	ņ	
×ι	к				
vi				芦	
N	P	5			
()I	,	۰			
$^{\sim}$	ă		Р	۲,	
XI	,	١	,	۰	
(N		à	ú	ù	
\sim		Š			
X:	×	(6	
(XI	ı.		ı.		
ΧJ		ę	μ	ę	
ХI	k		2		
(X)	н	,	ņ	ņ	
XJ	ĸ		1		
XI			,	۳	
\sim	a	'n	ũ	à	
XI	E		E	Z	
V	÷	۰	=	-	
O	ь	b	6	2	
×.	ă	i		e	
v	7	۹	9	6	
(M	ø	þ	런	7	
\times			2		
V1	k		2		
K	r	,	٠	7	
X.	ĸ		2	5.	
×1		۲	9		
(XI	Ľ		C	а	
X	×	Ų	ø	≺	
×Ί	×	á	ij	ø	
	ä		e		
ΧI		i	×	'n	
v	v	1	1		
8	2	š	6	S	
ΧI	۲	ď	7	٩	
V	L		2	a	
	۹	ij	ø	۴,	
X	ø	þ		К	
W	r	?		Л	
All		ı	p		
X					
\sim		s		١,	

Question 4 continued	
(Total for Question	n 4 is 6 marks)



$$f(x) = x^3 + (p+3)x^2 - x + q$$

where p and q are constants and p > 0

Given that (x - 3) is a factor of f(x)

(a) show that

$$9p + q = -51$$

(2)

Given also that when f(x) is divided by (x + p) the remainder is 9

(b) show that

$$3p^2 + p + q - 9 = 0$$

(2)

(c) Hence find the value of p and the value of q.

(3)

(d) Hence find a quadratic expression g(x) such that

$$f(x) = (x - 3)g(x)$$

(2)



	XX
×	$\times\!\!\!\times\!\!\!\times$
X	\Leftrightarrow
×	\times
×	$\times\!\!\times\!\!\times$
\otimes	$\times\!\!\times\!\!$
\otimes	XX
X	$\times\!\!\times\!$
X	$\langle \times \rangle$
×	$\langle \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \!$
\otimes	$\times\!\!\times\!\!$
X.	Ä
X	
X	AREA
×	
X	
X	
X	
X	
Ø	4
X	
X	WRITE-IN-THIS
Ø	
X	XX
X	$\propto \sim$
X	
X	
X.	XX
X	44
×	\cong
X	
×	
X	NP.
X.	
×	
X	
X	X
$\langle \rangle$	DO NOT
X	Ų,
$\langle \cdot \rangle$	
X	5
X	\sim
X	(X)
×,	=
$\langle \rangle$	
X	
X	$\times\!\!\!\times\!$
X	$\times\!\!\times\!\!\times$
×	$\langle \times \rangle$
×	$\times\!\!\times\!\!\times$
X	$\times\!\!\times$
\otimes	$\times\!\!\!\times$
X	$\Leftrightarrow \Rightarrow$
X	$\langle \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \!$
\Rightarrow	$\times\!\!\times$
\times	XX
$\langle \cdot \rangle$	$\times\!\!\!\times\!$
X	$\times\!\!\!\times\!\!\!\!>$
X	$\times\!\!\times\!\!\times$
X	$\langle \times \rangle$
×	$\times\!\!\times$
\otimes	XX
X	$\times\!$
X	$\otimes \!\!\! >$
X	$\langle \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \!$
Ø	$\times\!\!\times\!\!\times$
×.	
X	× 6
×	44
X	1
	تكف
Ø	XX.
8	7
	AREA
	¥,
	SA
	IS A
	TIS A
	TIS A
	HEIS A
	A SITIL
	NTHISA
	IN THIS A
	WINTERS A
	ENTERS A
	THE TANK A
	TENTES A
	SHEW THIS A
	RITE IN THIS A
	WRITE IN THIS A
	WRITERNIHS
	WENEFINITES
	NOTWENESHIN
	WENEFINITES
	NOTWENESHIN
	DO NOT WRITE IN THIS
	DO NOT WRITE IN THIS
	A SOUND WRITE IN THIS
	EA DONOT WRITE/IN THIS
	EA DONOT WRITE/IN THIS
	REA DO NOT WRITE IN THIS
	REA DO NOT WRITE IN THIS
	AREA DO NOT WRITE IN THIS
	AREA DO NOT WRITE IN THIS
	IS AREA DO NOT WRITE IN THIS
	IS AREA DO NOT WRITE IN THIS
	HIS AREA DO NOT WRITEIN THIS
	HIS AREA DO NOT WRITEIN THIS
	THIS AREA DO NOT WRITEIN THIS
	THIS AREA DO NOT WRITEIN THIS
	MITHIS AREA DO NOT WRITE IN THIS
	IN THIS AREA DO NOT WRITE IN THIS
	IN THIS AREA DO NOT WRITE IN THIS
	EINTHIS AREA DO NOT WRITEIN THIS
	TE IN THIS AREA DO NOT WRITE IN THIS
	TE IN THIS AREA DO NOT WRITE IN THIS
	TE IN THIS AREA DO NOT WRITE IN THIS
	RITE IN THIS AREA DO NOT WRITE IN THIS
	TE IN THIS AREA DO NOT WRITE IN THIS

.>	ij		ζ	2	ś
١.	7	7		7	7
K		۷			
2		þ	9		ĸ
٦				u	ν
K	3	ķ		ρ	١,
.)	ς	è	6	sè	
^		≺	7	٦	ν
/		ŝ	۵	d	r
	₹	7	₹	7	₹
К					2
Ì	١				1
`		7			
×					
2		2			К
Υ,					i
/					`
					K
Κ	2	۲.			
2	١,				
			C		K
ĸ			2		2
/					٦
					1
		≺			
		2			ς
٦,					2
/					
Κ					
/			١		
×		≺			
2					1
					1
×					
					Κ
ď	٦				
Κ					2
2					
					e.
Κ					
		2			۲
^					1
/					1
					1
ď.			\		
	2	۷	6	à	ĸ
Κ			Я		
Ì	7	7	ч		Ь
	ú	7	í	ì	ĸ
		4	Ŀ		D
.2					
	á	ń	ã	Z	P
2	ĺ	á	į	Z	Ŕ
3	ĺ	þ	į	2	Ŕ
2		2			Š
2					Ķ
3					
3		2			
3		2			
3		2			
?		2			
3					ķ
3					ķ
3					ķ
3					ķ
3					
3					
3					

Question 5 continued	
	-
	-
	-
	-
	-
	_
	-
	-
	-
	-
	-
	-
	-
	-
	-
	-
	_
	_
	-
	-
	-
	-
	-
	-
	_
	-



Question 5 continued

		7		2		
	X					
	g		2			
		S				
	S					
	?					
		ú				
			2			
	g		3		Σ	ć
	8	Ç		Ę	2	6
	9	Č		Ž	2	6
>	8			5		í
>	5					
>	8					
>			AHA			
>						
>	2					
>						
>						
>						
>						
>>>						
>	>					
>>>>>						
\ \ \ \						
>	>					
>						
>						
>						
\ \ \ \ \						
\ \ \ \						

Question 5 continued	
	(Total for Question 5 is 9 marks)



6. The circle C has equation

$$x^2 + y^2 + 8x - 4y = 0$$

- (a) Find
 - (i) the coordinates of the centre of C,
 - (ii) the exact radius of C.

(3)

The point *P* lies on *C*.

Given that the tangent to C at P has equation x + 2y + 10 = 0

(b) find the coordinates of P

(4)

(c) Find the equation of the normal to C at P, giving your answer in the form y = mx + c where m and c are integers to be found.

(3)

\sim	S	
	X	
X	\otimes	$\Leftrightarrow >$
X		X
\times	X	$\times\!\!\times$
X	×	~~
Q.	\times	$\Diamond \Diamond$
X	X	
$\times \rangle$	×	$\times\!\!\times$
\otimes		XX
X	\times	$\Diamond\Diamond$
×	\times	\bigcirc
X	d	\times
X	O.	\times
\times	CK.	
X	ŔΫ	ĕX
K.	N	50
X	d	$ \mathcal{C} $
×	X	XX
X	Úέ	\times
\times	X	\times
		-X
×	M	
Ø	\circ	90
X	X	\times
$\times \rangle$	Û	
\otimes	ĸ.	¥×
X	X	€X
$\langle \times \rangle$	X	
X		
\times	×	$\Rightarrow >$
\times	\times	\cong
X	m	€8
X.	Z.	50
X	2	-XX
×	8	*
\otimes		$\simeq \times$
$\langle \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \!$		Š
$\langle \rangle$	Ø	×
X	3	
X		Z/X
8	X	₹X
$\langle \times \rangle$	Ö	W
S	۰	20
X	0	N/A
$\times \rangle$	\times	*
$\langle \rangle$	X	$\langle \times \rangle$
X	$\times \rangle$	\Leftrightarrow
S	\otimes	\propto
×	Ö.	\times
\times	X	$\times\!\!\times$
\approx	X	
×	×	⋘
Q.	\times	$\Diamond\Diamond$
X	X	
$\times \rangle$	X	$\times\!\!\times$
\otimes	X	XX
X	$\times \rangle$	$\Diamond \Diamond$
X	X	XX
\times	X	$\times\!\!\times$
X	X	XX
×	\times	$\Diamond \Diamond$
X	\approx	99
×	X	×
\times	X	
	S	
\sim		$\bigcirc \bigcirc$
\otimes	ä	
\otimes	Š	
$\stackrel{\times}{\otimes}$	E/A	
	V = 7	
× ×	V 12	
× ×	V 10 00 V	
	V 1	
	Value V	
	VIII OF AVAILABLE OF	
	WHO WILL WILL CO	
	BUCK OF THE NEW LONG CO.	
	SHOW STILL WILLIAM LONGE	
	BUG D VITE NUMBER OF CO	
	STATE OF THE STATE	
	SHOW OF THE WALLEY CO.	
	STATE VIEW CONTRACTOR	
		Ē
	V V V V V V V V V V V V V V V V V V V	5
	V V V V V V V V V V V V V V V V V V V	5
	Zu G	
	A 20 A	
	A 20 A	
	A 20 A	

Question 6 continued	



Question 6 continued				

X	X		
	>		
X,	X.	2	
\sim		4	d
			Б
	≂		ь
V	v,	Ω	Z
$^{\sim}$	M	r	Я
×Ι	,	۰	,
	÷	G	à
XI	n	r	7
VI	,	=	•
	$^{\sim}$	4	à
Χi	e		Г
×	9		×
$^{\sim}$	$^{\wedge}$		7
X.		7	C
(XI	n		9
X.J			я
v.	Z	2	z
	=	۳	7
Χi			
$\langle \times \rangle$	И	I.	2
\sim			
va	√		
K	₽	ú	è
ΧI	К		$\overline{}$
	7		
$\triangle a$		ú	4
X.	X		۳
K		L	÷
Α.			Σ
VI	₹	9	=
	\rightarrow		
X.	X,		w
VI	и	L	а
	-	•	7
×Ι	K.		<i>C</i> .
(X)	P	ņ	ņ
XJ	ĸ		
VI	,	=	=
\sim	~	G	à
ΧI	N		7
√1			
$^{\sim}$ a	ũ.	Z	\sim
X	Z		
(X)		Ξ	2
$^{\sim}$	ä		
V1			2
\sim	\sim		
ΧI	ĸ		
VI			
	7	ę	7
	Ķ	Ž	Š
X	K	}	8
ä	K	2	í
Ø	K	2	í
	K	2	í
	K	2	í
	K	2	í
	K	2	í
	K	2	
	K	2	í
	K	2	í
			í

Question 6 continued	
(Tot	cal for Question 6 is 10 marks)
·	·



7.	A geometric sequence has first term a and common ratio r , where $r > 0$	
	Given that	
	• the 3rd term is 20	
	• the 5th term is 12.8	
	(a) show that $r = 0.8$	(4)
		(1)
	(b) Hence find the value of a.	(2)
	Given that the sum of the first n terms of this sequence is greater than 156	,
	(c) find the smallest possible value of <i>n</i> .	
		`
	(Solutions based entirely on graphical or numerical methods are not acceptable.) (4)
		(•)



Question 7 continued
(Total for Question 7 is 7 marks)



8. In this question you must show all stages of your working.

Solutions based entirely on calculator technology are not acceptable.

(i) Solve, for $-\frac{\pi}{2} < x < \pi$, the equation

$$5\sin(3x + 0.1) + 2 = 0$$

giving your answers, in radians, to 2 decimal places.

(4)

(ii) Solve, for $0 < \theta < 360^{\circ}$, the equation

$$2 \tan \theta \sin \theta = 5 + \cos \theta$$

giving your answers, in degrees, to one decimal place.

(5)

Question 8 continued			



Question 8 continued				

	'n	Š			
		Ê			
	Ì				
	(i				
	g				
	ĸ				
			π	1	
S	a	Ŀ	S.	4	
>	d	Li Tr	Ļ	ļ	
8	1	L K	5	2	
2	3	K		2	
3	1	K		2	
2		í.	2	2	
3	1	K			
2		K			
3		K			
2		K	Y		
3		L K K K			
		10 K 20 A 2			
		To the second			
3					
		10 10 10 10 10 10 10 10 10 10 10 10 10 1			
		TO THE SECOND PROPERTY OF THE			
		10 10 10 10 10 10 10 10 10 10 10 10 10 1			
			Y		

Question 8 continued	
	-
	-
	-
	-
	-
	_
	-
	-
	-
	-
	-
	-
	-
	-
	-
	-
	-
	-
	-
	_
	-
	-
	-
(Total for Question 8 is 9 marks)	
(2000 TOT QUEDUON O TO S MICH IND)	



9. In this question you must show all stages of your working.

Solutions based entirely on calculator technology are not acceptable.




Figure 3

Figure 3 shows

- the curve C with equation $y = x^2 4x + 5$
- the line l with equation y = 2

The curve C intersects the y-axis at the point D.

(a) Write down the coordinates of D.

(1)

The curve *C* intersects the line *l* at the points *E* and *F*, as shown in Figure 3.

(b) Find the x coordinate of E and the x coordinate of F.

(2)

Shown shaded in Figure 3 is

- the region R_1 which is bounded by C, l and the y-axis
- the region R_2 which is bounded by C and the line segments EF and DF

Given that $\frac{\text{area of } R_1}{\text{area of } R_2} = k$, where k is a constant,

(c) use algebraic integration to find the exact value of k, giving your answer as a simplified fraction.

(5)

X
\times
\otimes
X
X
$\langle \cdot \rangle$
X
X
×
×
X
X
X
X
×
X
X
X
$\times \rangle$
\times
X
\times
X
\sim
X
X
X
X
\times
\times
X
82
X
X
\Diamond
X
X
X
X
X
$\times \rangle$
\otimes
\otimes
8
$\langle \rangle$
$\langle \cdot \rangle$
\propto
X
X
X
X
×>
\times
X
\otimes
$\langle \rangle$
$\langle \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \!$
X
X
X
X
×
\times
\otimes
\otimes
$\langle \cdot \rangle$
$\langle \times \rangle$
\times
\otimes
\otimes
× ×

1				
1				
	í			
û	è	ú	à	6
Ì	È	ź	à	í
2	Ì	į	Ì	ĺ
í k		į		ĺ
k				
k				
k				
i k				
Q M M M				
Û K L				
j K				
i K K K K				
K				
ř	į	2		1
ř	į	2		1
ř	į	2		1
ř	į	2		1
ř	į	2		1
ř	į	2		1
ř	į	2		1
ř	į	2		1
ř	į	2		1
ř	į	2		1
ř	į	2		1
ř	į	2		1
ř	į	2		1
ř	į	2		1
ř	į	2		
ř				
ř				
ř				
ř				
ř				
ř				
ř				
ř				
ř				
ř				
ř				
ř				
ř				
ř				
ř				
ř				
ř				

nestion 9 continued	
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_



Question 9 continued	

	Ì				
	í			١	
	ì				
	í				
				2	
	į				
			i		
	3				9
	i	ú	ì	4	2
	į	2	ì	2	Š
5	i		ì	2	Š
>	i	É		Ź	
>	į			2	
>	į			2	
>	i				
	1				
>	1				
	1				
>			3	5	3
>	1		3	5	3
>	i		3	5	3
> > >			3	5	3
<pre></pre> <pre><pre></pre><pre></pre><pre></pre><pre></pre><pre></pre><pre><!--</td--><td></td><td></td><td>3</td><td></td><td>3</td></pre></pre>			3		3
<pre></pre> <pre><pre></pre><pre></pre><pre></pre><pre></pre><pre></pre><pre><!--</td--><td></td><td></td><td>3</td><td>5</td><td>3</td></pre></pre>			3	5	3
			3	5	3
			3	5	3
<pre></pre>			3	5	3
			3	5	3
<pre></pre> <pre><pre></pre><pre></pre><pre></pre><pre></pre><pre></pre><pre><!--</td--><td></td><td></td><td>3</td><td>5</td><td>3</td></pre></pre>			3	5	3
			3	5	3
			3	5	3
			3	5	3
			3	5	3
				5	3
				5	3
				5	

Question 9 continued	
(Tota	al for Question 9 is 8 marks)
(-3	,



10. A student was asked to prove by exhaustion that

if *n* is an integer then $2n^2 + n + 1$ is **not** divisible by 3

The start of the student's proof is shown in the box below.

Consider the case when n = 3k

$$2n^2 + n + 1 = 18k^2 + 3k + 1 = 3(6k^2 + k) + 1$$

which is not divisible by 3

Comp	lete	this	proof.

(4)



	$\times \times$	
\times	$\times\!\!\times$	
	\times	×
Ç.	$\langle \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \!$	
X	∞	
X		×
X	$\times\!\!\times\!\!$	×
\propto	XX	
X	\otimes	Š
X	$\otimes \otimes$	
\times	$\times\!\!\times$	×
$\times\!\!\times$	\simeq	
X	AKEA	2
×	ΝŴ	
X		
XI	9 00	S
\times	7	
\bowtie	24	
X,	WRITE IN THIS	Š
Ø	yn.	Ç
X		
XX	J X	
X		\times
X	-	
X		
X		Q
×		S
\times	\times	X
KS	ш	
ÖÜ	\triangle	Ö
X		
XX	4	
\times	NP.	\times
X		
X		
X	\sim	Ç
X	DON OC	
XX	\sim	
XX	\otimes	\times
ΧX		×
ÖĞ	4	
$\langle \rangle$	$\times\!$	
X	(30)	S
XX		
\bowtie		×
X	XX	
	$\Leftrightarrow \Leftrightarrow$	Ö
X	\times	
X	\times	
\times	\sim	\times
\times	$\times\!\!\times$	
XX	XX	
×	$\Leftrightarrow \Leftrightarrow$	Ç
	\times	
\times	\times	×
$\times\!\!\!\times$	$\times\!\!\times$	\times
	. X X	
∞		0
\otimes	\bowtie	8
\otimes	\times	8
$\stackrel{\times}{\times}$	\times	5
× ×	X	
× ×		
	4	
	A REMA	
	AREA	
	SAREA	
	IS AREA	
	IIS AREA	
	HIS AREA	
	THIS AREA	
	THIS AREA	
	VIHIS AREA	
	N THIS AREA	
	IN THIS AREA	
	EINTHIS AREA	
	E IN THIS AREA	
	TETN THIS AREA	
	STEIN THIS AREA	
	RITE IN THIS AREA	
	KRITE IN THIS AREA	
	WRITE IN THIS AREA	
	WRITE IN THIS AREA	
	EWRITE IN THIS AREA	
	M WRITE IN THIS AREA	
	OF WRITE IN THIS AREA	
	OT WRITE IN THIS AREA	
	NOTWENTHS	
	DO NOT WRITE IN THIS AREA	
	NOTWENTHS	
	DO NOT WRITE IN THIS	
	DO NOT WRITE IN THIS	
	A DONOT WRITE IN THIS	
	EA DONOT WRITE IN THIS	
	EA DONOT WRITE IN THIS	
	A DONOT WRITE IN THIS	

	0			
	١			
	<			
		Ś	4	à
		e		
ì	7	7	۹	ķ
			6	ì
		싧		
	3		2	
		7		7
5		ы		ь
			۵	6
		е		7
	7	3	Ħ	b
	¥		'n	ĕ
		a	r	3
	2	≾		2
	5	₹	7	e
	힠	峋	þ	pÌ
		Я	r	\
	ŋ	e	,	ę
ì		2	2	Ź
		7	۰	٠
	S	2	≤	è
	ě	2	ğ	į
	Š	Ž		į
>	Š	Ž		
3	A STATE OF			
				1
				4

Question 10 continued	



Question 10 continued	
	(T-4-1 f O4' 10 '- 4 1)
	(Total for Question 10 is 4 marks)
	TOTAL FOR PAPER IS 75 MARKS

