Please check the examination deta	ils below	before ente	ering your candidate information
Candidate surname			Other names
Pearson Edexcel International Advanced Level	Centro	e Number	Candidate Number
Monday 11 M	lay	202	20
Afternoon (Time: 1 hour 30 minu	tes)	Paper R	deference <b>WFM01/01</b>
Mathematics			
International Advance Further Pure Mathema		'	y/Advanced Level
You must have: Mathematical Formulae and Stat	tistical <sup>-</sup>	Tables (Blu	ue), calculator

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

## Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
   there may be more space than you need.
- You should show sufficient working to make your methods clear.
   Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

## Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 8 questions in this question paper. The total mark for this paper is 75.
- The marks for each question are shown in brackets
  - use this as a guide as to how much time to spend on each question.

## Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over ▶







**(2)** 

 $f(x) = x^3 - \frac{10\sqrt{x} - 4x}{x^2}$ 1.

- (a) Show that the equation f(x) = 0 has a root  $\alpha$  in the interval [1.4, 1.5] **(2)**
- (b) Determine f'(x). **(3)**
- (c) Using  $x_0 = 1.4$  as a first approximation to  $\alpha$ , apply the Newton-Raphson procedure once to f(x) to calculate a second approximation to  $\alpha$ , giving your answer to 3 decimal places.




Leave

X.				
		2		
v	4	7		ы
×	Ħ	٠		
	ú	a	۰	2
ΧI		й	×	ú
v.		2		2
$\sim$		7	2	/
Vi				к
AI.	ľ			1
X		4	٠	a
	ď	9	5	5
	à			к

	blank
Question 1 continued	
	Q1
(Total 7 marks)	



**2.** The quadratic equation

$$5x^2 - 2x + 3 = 0$$

has roots  $\alpha$  and  $\beta$ .

Without solving the equation,

(a) write down the value of  $(\alpha + \beta)$  and the value of  $\alpha\beta$ 

**(1)** 

- (b) determine, giving each answer as a simplified fraction, the value of
  - (i)  $\alpha^2 + \beta^2$
  - (ii)  $\alpha^3 + \beta^3$

**(4)** 

**(4)** 

(c) determine a quadratic equation that has roots

$$(\alpha + \beta^2)$$
 and  $(\beta + \alpha^2)$ 

giving your answer in the form  $px^2 + qx + r = 0$  where p, q and r are integers.

X					
	r				
Vì.	-	à	í	8	ì
X	2	į	É	5	
S,		è	2		
Š	ľ	ì	í		
Š	ľ	ì	É		
j		Ì	É		
Š					

uestion 2 continued	



	Lea bla
Question 2 continued	

	Ĺ			
×				
	Š			
v				
				$\rangle$
2	S	ζ	S	2
8	S	Ì	Z K	2
2	ĺ	3	K	ì
	ĺ	3	Į	į
8			K	į
		2	5	2
		2		}
			< t	
			2	
			< t > 1	

Question 2 continued		Leave blank
		Q2
	(Total 9 marks)	
	(10tal / Illat No)	



3.	$f(z) = z^4 + az^3 + bz^2 + cz + d$	
	where $a$ , $b$ , $c$ and $d$ are integers.	
	The complex numbers $3 + i$ and $-1 - 2i$ are roots of the equation $f(z) = 0$	
	(a) Write down the other roots of this equation.	(2)
	(b) Show all the roots of the equation $f(z) = 0$ on a single Argand diagram.	(2)
	(c) Determine the values of a, b, c and d.	(5)
_		

		2	S	a	
$\wedge$	d	μ		r	
X1	9			ĸ	
	c			7	
×I	į	Я		1	
V					
	ú	۵	4	4	,
$\times$	,	₹		₹	
W					
				S	
	à	ø		۳	
	ņ	ú		Ŀ	
X	2			₹	
$\vee$					
	r	7		۰	
X			n	4	
	₹			2	
×į	μ	۹	×	8	
	è	ú	è	ú	
	Т	٦	г	7	
×	ú		Ŀ	ú	
	Σ	7	7	7	٦
$\sim$		_	4	_	è
		е	7	₹	
$^{\circ}$	۹				
$\wedge$	2	2	2	≤	
v	7	z		P	,
				_	
	7	₹	7	₹	
V	۱	н	'n		
	۵	ζ,		4	
( )	ı	J	ĸ.	в	
	н	8	۳	9	
	Ŀ			2	
	ş			۲	
M	ı	н	н		
	4	2	4	2	
$\times$	r	7		₹	
W		8	×	н	
$^{\wedge}$	ú	2	2		
O			c	9	
$^{\wedge}$	ż	S			
V	۹	7	2		
	7				
		۷	-	۷	
V		7	₹	7	۰
	z	à	ú	2	
X		٣	7	힉	í
()	6	2	S	d	ľ
$\wedge$	2	۹	۶	К	
	ä	ø	a		
	Č	á	ø	۲	
X		й	×	ú	
v				7	
	S	2	4	2	
X		۲	9	۹	
C)	ĸ			1	ĺ
	7	۹	ø	×	
	á	d	ń	ĸ	
	۲			٦	

	bla
Question 3 continued	



estion 3 continued	

	q	e		
				2
	ľ	Ù		
	4	Þ	7	þ
	I	R	2	₹
×	ţ	k	2	Š
×	ļ	k	2	Š
×	1	k	2	
×	1	k	2	2
×	1	k A	2	
× × ×	1	i k	2	
×		× ×	2	
× × ×	1		2	
× × ×	1		2	
××××	1		2	
××××	1		2	
×	1			
×××××××××××××××××××××××××××××××××××××××			7	
××××	1	1		
××××	1			
××××××××××××××××××××××××××××××××××××××	1			
× × × × ×	1			
××××××××××××××××××××××××××××××××××××××	1			
× × × × ×	1			
× × × × ×	1			
××××××	1			
××××××	1			
××××××	1			
×××××××	1			
×××××××	1			
××××××××	1			
×××××××××	1			
××××××××				
×××××××××				
××××××××××××××××××××××××××××××××××××××	1			
××××××××				
××××××××××××××××××××××××××××××××××××××				
××××××××××××××××××××××××××××××××××××××				
××××××××××××××××××××××××××××××××××××××				
××××××××××××××××××××××××××××××××××××××				
××××××××××××××××××××××××××××××××××××××				
××××××××××××××××××××××××××××××××××××××				
××××××××××××××××××××××××××××××××××××××				
××××××××××××××××××××××××××××××××××××××				
××××××××××××××××××××××××××××××××××××××				
××××××××××××××××××××××××××××××××××××××				
××××××××××××××××××××××××××××××××××××××				
××××××××××××××××××××××××××××××××××××××				
××××××××××××××××××××××××××××××××××××××				
××××××××××××××××××××××××××××××××××××××				
××××××××××××××××××××××××××××××××××××××				
××××××××××××××××××××××××××××××××××××××				

Question 3 continued	Leave blank
	Q3
(Tatal 0	
(Total 9 marks)	



**4.** (a) Use the standard results for  $\sum_{r=1}^{n} r^2$  and  $\sum_{r=1}^{n} r$  to show that

$$\sum_{r=1}^{n} (2r-1)^2 = \frac{1}{3}n(4n^2-1)$$

for all positive integers n.

**(5)** 

(b) Hence find the exact value of the sum of the squares of the odd numbers between 200 and 500

1	1	)
1	7	٠,


	J					
	з					
	à		í		í	
	i		ĺ		ĺ	
	1	Ĺ	į	Č	ĺ	
×	į	Ľ	Į	Ì	ĺ	
	Į	Ľ	ļ		ĺ	
	į	Ľ	1	2	ĺ	
	l	Ľ				
	1	Ě			ĺ	
× × ×	1	Ľ				
		Ľ		2		
		ľ				
/ / / /		Ì				
		Ĭ				
		Ž				
/ × × × ×		Ĭ				
		Ĭ				
×××××××						
	ļ					
/×××××××××××××××××××××××××××××××××××××	ļ					
	ļ					
~×××××××××××××××××××××××××××××××××××××	ļ					
	ļ					
~×××××××××××××××××××××××××××××××××××××	ļ					
/×××××××××××××××××××××××××××××××××××××	ļ					
	ļ					
	ļ					
	ļ					
	ļ					
	ļ					
	ļ					
	ļ					
	ļ					
	ļ					
	ļ					
****** ****** ***	ļ					
	ļ					
/ / / / / / / / / / / / / / / / / / /	ļ					
/ / / / / / / / / / / / / / / / / / /	ļ					

estion 4 continued	



	Leave blank
Question 4 continued	

×j			
Χį			
		w	п
a	D	ĸ	1
a	Ú	Ľ	1
a	Ľ	Ķ	Į
8	D	Ķ	Į
9	D K	K	2
	K	Š	2
	K	Į	
	K		
	T X	2	
	T X	2	
	To A	_	
	T A	2	
	To the second		
	To the second		
	To a second		
	K K K		
	K K K		
	K K K		
	K K K		
	K Z (6 5) X X Y Z Z		
	K Z (6 5) X X Y Z Z		
	K K K		
	K Z (6 5) X X Y Z Z		
	K Z (6 5) X X Y Z Z		
	K Z (6 5) X X Y Z Z		
	K Z (6 5) X X Y Z Z		
	K Z (6 5) X X Y Z Z		
	K Z (6 5) X X Y Z Z		
	K Z (6 5) X X Y Z Z		
	K Z (6 5) X X Y Z Z		

Question 4 continued	Leave blank
Question 4 continued	
	04
	Q4
(Total 9 marks)	



blank

5. The rectangular hyperbola H has equation xy = 64

The point  $P\left(8p, \frac{8}{p}\right)$ , where  $p \neq 0$ , lies on H.

(a) Use calculus to show that the normal to H at P has equation

$$p^{3}x - py = 8(p^{4} - 1)$$
(5)

The normal to H at P meets H again at the point Q.

(b) Determine, in terms of p, the coordinates of Q, giving your answers in simplest form.

(4)


erion a continued	
estion 5 continued	



	Leave
	blank
Question 5 continued	

	É			
			Ž	
	è			
	ì			
			X	
	7	↸	7	5
	<	$\geq$	0	2
	2	2	X	2
	7		á	
	Z	į	â	
	Į	į	á	
		2	à	
			X	
			X	
			X	
			X	
			X	
			X	
× × × × × × × ×				
× × × × × × × × × ×				

Question 5 continued	Leave blank
	Q5
(Total 9	marks)
(1011)	



**6.** (i) 
$$\mathbf{A} = \begin{pmatrix} 1 & 0 \\ 0 & 3 \end{pmatrix}$$

(a) Describe fully the single transformation represented by the matrix **A**. (2)

The matrix **B** represents a rotation of 45° clockwise about the origin.

(b) Write down the matrix  $\mathbf{B}$ , giving each element of the matrix in exact form. (1)

The transformation represented by matrix A followed by the transformation represented by matrix B is represented by the matrix C.

(c) Determine C.

**(2)** 

(ii) The trapezium T has vertices at the points (-2, 0), (-2, k), (5, 8) and (5, 0), where k is a positive constant. Trapezium T is transformed onto the trapezium T' by the matrix

$$\begin{pmatrix} 5 & 1 \\ -2 & 3 \end{pmatrix}$$

Given that the area of trapezium T' is 510 square units, calculate the exact value of k. (5)



Question 6 continued		Leave
Question 6 continued		blank
	Question 6 continued	
	Question o continued	



uestion 6 continued	
lestion o continued	

		2	Ś	a	
$\wedge$	d	μ		r	
X1	9			ĸ	
	c			7	
×I	į	Я		1	
V					
	ú	۵	4	4	,
$\times$	,	₹		₹	
W					
				S	
	à	ø		۳	
	ņ	ú		Ŀ	
X	2			₹	
$\vee$					
	r	7		۰	
X			n	4	
	₹			2	
×į	μ	۹	×	8	
	è	ú	è	ú	
	Т	٦	г	7	
×	ú		Ŀ	ú	
	Σ	7	7	7	٦
$\sim$		_	4	_	è
		е	7	₹	
$^{\circ}$	۹				
$\wedge$	2	2	2	≤	
v	7	z		P	,
				_	
	7	₹	7	₹	
V	۱	н	'n		
	۵	ζ,		4	
( )	ı	J	ĸ.	в	
	н	8	۳	4	
	Ŀ			2	
	ş			۲	
M	ı	н	н		
	4	2	4	2	
X	r	7		₹	
W		8	×	н	
$^{\wedge}$	ú	2	2		
O			c	9	
$^{\wedge}$	ż	S			
V	۹	7	2		
	7				
		۷	-	۷	
V		7	₹	7	۰
	z	à	ú	2	
X		٣	7	힉	í
()	6	2	S	d	ľ
$\wedge$	2	۹	۶	К	
	ä	ø	a		
	Č	á	ø	۲	
X		й	×	ú	
v				7	
	S	2	4	2	
X		۲	9	۹	
C)	ĸ			1	ĺ
	7	۹	ø	×	
	á	d	ń	ĸ	
	۲			٦	

	Leave
	blank
Question 6 continued	
	Q6
(Total 10 marks)	
(Total To Hiarks)	



The parabola C has equation  $y^2 = 4ax$ , where a is a positive constant. The line *l* with equation 3x - 4y + 48 = 0 is a tangent to *C* at the point *P*. (a) Show that a = 9**(4)** (b) Hence determine the coordinates of P. **(2)** Given that the point S is the focus of C and that the line l crosses the directrix of C at the point A, (c) determine the exact area of triangle PSA. **(4)** 



	J					
	з					
	à		í		í	
	i		ĺ		ĺ	
	1	Ĺ	į	Č	ĺ	
×	į	Ľ	Į	Ì	ĺ	
	Į	Ľ	ļ		ĺ	
	į	Ľ	1	2	ĺ	
	l	Ľ				
	1	Ě			ĺ	
× × ×	1	Ľ				
		Ľ		2		
		ľ				
/ / / /		Ì				
		Ĭ				
		Ž				
/ × × × ×		Ĭ				
		Ĭ				
×××××××						
	ļ					
/×××××××××××××××××××××××××××××××××××××	ļ					
	ļ					
~×××××××××××××××××××××××××××××××××××××	ļ					
	ļ					
~×××××××××××××××××××××××××××××××××××××	ļ					
/×××××××××××××××××××××××××××××××××××××	ļ					
	ļ					
	ļ					
	ļ					
	ļ					
	ļ					
	ļ					
	ļ					
	ļ					
	ļ					
	ļ					
****** ****** ***	ļ					
	ļ					
/ / / / / / / / / / / / / / / / / / /	ļ					
/ / / / / / / / / / / / / / / / / / /	ļ					

	Leave
	blank
Question 7 continued	- Clarik
Question / continued	
	1
	1
	1
	1
	1
	1
	1
	1
	1
	1
	1



estion 7 continued	
estion / continued	

( )I	×	×	2
î	ø	~	
V4	ь/		/
	a	=	۵
X	ĸ.	-	S)
C XI	Ľ.		л
			p.
Q.	_	z	->
×	~	9	
	×	př.	
ΧI		-	'n
V.		$\times$	7
Λ.			
	d		к
AI.	r		ч
V4	L	2	л
	ч		ĸ
	ы		L
CM	r	v	٩

	Leave blank
Question 7 continued	
	Q7
(Total 10 marks)	



**8.** (i) Prove by induction that, for  $n \in \mathbb{Z}^+$ 

$$\sum_{r=1}^{n} \frac{2r^2 - 1}{r^2(r+1)^2} = \frac{n^2}{(n+1)^2}$$

**(6)** 

(ii) Prove by induction that, for  $n \in \mathbb{Z}^+$ 

$$f(n) = 12^n + 2 \times 5^{n-1}$$

is	divisi	b.	le	by	7

**(6)** 


	ŝ	
U.	u	
X	ш	ш
Ŋ	Ц	Ц
Ŋ	V	Ÿ
X	K	×
XI	K	¥
	K	X
	K K	X
	K K	¥ X
	K	V X A
	V K K	V X X
	V K K	

	Leave
	blank
Question 8 continued	



estion 8 continued	

uestion 8 continued	
uestion 8 continued	



	Leave blank	
Question 8 continued		
	Q8	-
(Total 12 marks)		
TOTAL FOR PAPER: 75 MARKS END	<b>'</b>	