Surname	Other nan	nes
Pearson Edexcel International Advanced Level	Centre Number	Candidate Number
Core Math	nematics	s C34
Tuesday 19 June 2018 – Aft Time: 2 hours 30 minutes		Paper Reference WMA02/01

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information

- The total mark for this paper is 125.
- The marks for each question are shown in brackets
 use this as a quide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶





1. (i) Find	$\int \frac{2x^2 + 5x + 1}{x^2} \mathrm{d}x, x > 0$	(3)
(ii) Find	$\int x \cos 2x \mathrm{d}x$	
		(3)



ì	e	Š	۴	ø
7	7	٩	6	ij
1	Ŀ	ı	Į.	ı
Ġ	Ξ	7	Ξ	2
1	c		c	1
ľ	7		Ξ	ï
J	ξ	S	Ľ	Ī
			7	٩
ì	r	À	A	į.
1	b	g	·]	,
ì		Ė		Ė
3	Ŧ	Ŷ	7	B
ì		ä		i
	L	_	_	
ł	г	7		•
ì	ě	Ė		ì
4	d	ė	2	1
3	7		7	3
4		7	-	۰
ì	r	ì	1	ì
1	þ	ě		ı
н	ŀ	ė		į
1	L	_		
Ĝ	3		Ξ	ä
1	C			ī
á				
1	Ė	e	e	•
i	õ	ē	3	9
Ġ	ċ			
J	Þ	ė		į
å	ä	a	6	į.
J	C		9)
ú	2	Ξ	_	
ľ	Ξ	7	7	2
1		÷	-	9
	5	_	٠	
ï	r	7	7	b
	9	ä	g	
i	ρ	f	٩	١
j	b	ė		ļ

			ŝ			
		٥				
		Ę				
	ì					
	ì	è			ı	
	1					
	ζ					
	į					
				1		
		Ľ	ě	_		
		P	ě	į	ė	
	Į	ŀ	ž	2	Š	
/	1		é	-	Š	
0	1		ě			
X	ļ					ĺ
S	Į					
2	1					
2	1					ĺ
X	1					
×	1					
×						
×	1					
X						
× × × × × × × × × × × × × × × × × × ×	1					
× × × ×						
X						

O NOT WRITE IN THIS AREA

Question 1 continued	Leave blank
(Total 6 marks)	Q1
(100m v mm no)	



blank

2. A curve C has parametric equations

$$x = \frac{3}{2}t - 5$$
, $y = 4 - \frac{6}{t}$ $t \neq 0$

- (a) Find the value of $\frac{dy}{dx}$ at t = 3, giving your answer as a fraction in its simplest form.
- (b) Show that a cartesian equation of C can be expressed in the form

$$y = \frac{ax + b}{x + 5} \qquad x \neq k$$

where a, b and k are integers to be found.

(4)

	Leave blank
Question 2 continued	



estion 2 continued	

Question 2 continued	Leave
	Q2
(Total 7 marks)	
(Total / Illal Ks)	



3. $f(x) = 2^{x-1} - 4 + 1.5x \qquad x \in \mathbb{R}$

(a) Show that the equation f(x) = 0 can be written as

$$x = \frac{1}{3} (8 - 2^x) \tag{2}$$

The equation f(x) = 0 has a root α , where $\alpha = 1.6$ to one decimal place.

(b) Starting with $x_0 = 1.6$, use the iteration formula

$$x_{n+1} = \frac{1}{3} (8 - 2^{x_n})$$

to calculate the values of x_1 , x_2 and x_3 , giving your answers to 3 decimal places.

(3)

(c) By choosing a suitable interval, prove that $\alpha = 1.633$ to 3 decimal places.

(2)

ì	ė	۴
7	7	Ni.
1	H	٠II
1	_	5
1	0	=
4	-	Ξ
ì	e	r
		,
'n	r i	ď.
1	u	7))
'n	-	i
ì	4	-
ì	4	÷
	C	Ξ
ł	_	-
		۰
	2	~
4	F	-
4	-	=
ì	. :	á.
1	ш	4
	Ŀ	Ξ
ļ	т	7
4	-	
1	n	m
	Ħ	
Ą	3	×
1	5	Si
J	~	_
1	Ĺ	_
4	7	7
1	m	٩
4	٤.	J
9		ė
1	2	_
÷		7
٠,	'n	iκ
4	Ŀ	2)
	Z	Ξ.
1	r	٦
J	=	4

	٥	
	V	
	Ž	
V		
K	Й	
/· s		
ă		

\Diamond	0			۲.
$^{\wedge}$	$^{\wedge}$	$^{\wedge}$	$^{\wedge}$	$^{\prime}$
V	V	V	V	V
\times	×	×	×	×
\vee	\vee			∿
\triangle	Λ			◠
×	X	×	×	X
X	X	X.	×	×
1	V	w	v	۲.
\triangle	$^{\wedge}$	_	ď.	↶
×	×	×		3×
		M X	п	Ľ
×	X			ľ×
V	\sim	\overline{z}	\subseteq	5.
		n	т	ţ٩
×	V	ш	84	N
				Χ.
X	Х	×		5×
V	\vee		ШZ	
	$^{\wedge}$	//5	~	r
×	×	×	×	×
		60	<u></u>	Κ.
X	X	WS.	м	v
V	V	쪵	v	ĸ,
\triangle	Λ	≏	_	Λ
×	X	$\overline{\mathbf{x}}$	×	×
		=	,	ĸ.
X	X	XI	K	Х
1	V			N
\triangle	\triangle	al/		$^{\prime}$
×	×			ì×
		W >		ς.
\mathcal{L}	Х	$^{\sim}$	А	Х
\vee	\sim		_	N
		\sim	o	
×	×		24	ıΧ
	\sim			5.,
\wedge	$^{\wedge}$	_	_	m
V	V	V	V	V
	$\langle \cdot \rangle$	6)	ല	c
×	×	IKJ		IX
0	$\langle \rangle$	₩	₩	٩.,
$^{\wedge}$	$^{\wedge}$	В\	$^{\wedge}$	↗
V	V			N
		Ш>		
-><	×			ŧΧ
52	90	$\overline{}$	$\overline{}$	ξ.
\triangle	Λ	m	м	20
×	×	ы	-	iΣ
		₽		
X	X	79	ж.	×
V	V	m#	₹	٦,
				r
	X	岬	×	\times
50	$\langle \rangle$			ς:
\wedge	$^{\sim}$	Вĺ\		
V	v			Ŋ
		'n.	\triangle	
×	×	20	70	
0	O	m)	a	ь.
\triangle	$^{\wedge}$	ж	-ii	'n
×	X	<u> 22</u>	32	×
				۲.
X	х	28	z	х
V	V	喞	₩	٥,
\triangle	◠			$^{\prime\prime}$
×	×	M	×	×
		ar.	d	ĸ
×	X	₩.	S	K/S
V	$^{\circ}$	a	-	5
\wedge	\wedge	40	ĸ.	◠
V	V	W.	73	P2
		₽		ĸ
X	X	$\overline{}$	$\overline{}$	×
V				

Question 3 continued	Leave blank
	Q3
(Total 7 marks)	



4. (a) Find the binomial expansion of

$$(1+px)^{-4}, |px| < 1$$

in ascending powers of x, up to and including the term in x^3 , giving each coefficient as simply as possible in terms of the constant p.

(3)

$$f(x) = \frac{3+4x}{(1+px)^4}$$
 $|px| < 1$

where p is a positive constant.

In the series expansion of f(x), the coefficient of x^2 is twice the coefficient of x.

(b) Find the value of p.

(5)

(c) Hence find the coefficient of x^3 in the series expansion of f(x), giving your answer as a simplified fraction.



	Leave
	blank
Question 4 continued	



	Leave
Omention Associational	blank
Question 4 continued	
	1

	Leave blank
Question 4 continued	
	Q4
/T-1-110 1 \	
(Total 10 marks)	



5. (i) The functions f and g are defined by

$$f: x \to e^{2x} - 5, \qquad x \in \mathbb{R}$$

$$g: x \to \ln(3x - 1), \qquad x \in \mathbb{R}, \ x > \frac{1}{3}$$

(a) Find f^{-1} and state its domain.

(3)

(b) Find fg(3), giving your answer in its simplest form.

(2)

(ii) (a) Sketch the graph with equation

$$y = |4x - a|$$

where a is a positive constant. State the coordinates of each point where the graph cuts or meets the coordinate axes.

(2)

Given that

$$|4x - a| = 9a$$

where a is a positive constant,

(b) find the possible values of

$$|x-6a|+3|x|$$

giving your answers, in terms of a, in their simplest form.

(5)

		Leave
Question 5 continued		blank
Question 5 continued		
	_	

estion 5 continued	

'n	ė	ρ	
٩	,	lij.	
1		di.	
9		₹.	
1	n	_	
4	-	₹.	
'n	ø	۳	
•	7	9	
	3	-	
1	u		
ń	÷	i	
'n	×		
4	4	-	
ì	Ľ	Ξ	
1	_	_	
1	۷	_	
c	2	~	
4	۰	=	
4		=	
'n	÷	÷	
1	ч	4	
1	-	÷	
1	г	_	
1	=	=	
1	3	=	
3	=	_	
G	7	9	
d	3	≥i	
1	7		
1		÷	
9	Ŀ	_	
1	7	Э	
٠,	'n	o	
4	7	7	
1	6		
٠		. "	
7	pti	٩	
4	÷	s	
4	ä	ø,	
1			

		Ţ	
A			
	ž		
		4	
	à		
X			
	2		
X			
×			
×	r		
K	c	14	
	Ę	К	į
	5	V	
		V	į
		V	ì
		Y	į
3	2	Ž	
		Š	֜֝֜֜֜֜֜֜֜֜֜֜֜֜֜֜֜֜֜֜֜֜֜֜֜֜֜֜֜֜֜֜֜֜֜֜֜
		¥	
		V	
		Y Y	
		2	
		¥ 2	

		Č		
		-86	K)	ĸ
×	×	×		3
		100	-	В.,
X	Х	ŭ	-4	_
	< :	\sim	$\overline{}$	
×	×	Ξ	Σ	Ď
8	8	E		₹
Š	Š	Ž	X	3
8	8	Ž	X	
8	8		X	
8			X	
8	X	Æ	Z	
X	X	Æ	Z	
8	X	Æ	Z	
×	X	Æ	Z	2
×	X	Æ	Z	
	X	Æ	Z	
	X	Æ	Z	
	X	Æ	Z	
	X	Æ	Z	
	X	Æ	Z	
	X	Æ	Z	
	X		Z	
	X	Æ	Z	
	X	Æ	Z	
	X	Æ	Z	
	X	Æ	Z	
	X	Æ	Z	
	X	Æ	Z	
	X	Æ	Z	
	X	Æ	Z	
	X	Æ	Z	
	X	Æ	Z	
	X	Æ	Z	
	X	Æ	Z	
	X	Æ	Z	
	X	Æ	Z	
	X	Æ	Z	
	X	Æ	Z	
	X	Æ	Z	
	X	Æ	Z	
	X	Æ	Z	
	X	Æ	Z	
	X	Æ	Z	

	Leave blank
Question 5 continued	
	Q5
(Total 12 marks)	



6. (a) Express $\sqrt{5}\cos\theta - 2\sin\theta$ in the form $R\cos(\theta + \alpha)$, where R > 0 and $0 < \alpha < \frac{\pi}{2}$ State the value of R and give the value of α to 4 significant figures.

(3)

(b) Solve, for $-\pi < \theta < \pi$,

$$\sqrt{5}\cos\theta - 2\sin\theta = 0.5$$

giving your answers to 3 significant figures.

[Solutions based entirely on graphical or numerical methods are not acceptable.]

(4)

$$f(x) = A(\sqrt{5}\cos\theta - 2\sin\theta) + B \qquad \theta \in \mathbb{R}$$

where *A* and *B* are constants.

Given that the range of f is

$$-15 \leqslant f(x) \leqslant 33$$

(c) find the value of B and the possible values of A.

(4)

	blank
Question 6 continued	



	Leave
	blank
Question 6 continued	

Leave

Question 6 continued	blank
	Q6
(Total 11 marks)	



(5)

7.




Figure 1 shows a hemispherical bowl.

Water is flowing into the bowl at a constant rate of 180 cm³ s⁻¹.

When the height of the water is h cm, the volume of water $V \text{ cm}^3$ is given by

$$V = \frac{1}{3}\pi h^2(90 - h), \quad 0 \leqslant h \leqslant 30$$

Find the rate of change of the height of the water, in cm s⁻¹, when h = 15 Give your answer to 2 significant figures.

4		5		1
-	7	۹	٩	٩
1	Ŀ		Ŀ	
С	Ξ	7	Ξ	Ξ
1	Ē		٤	9
			3	3
J	ξ	S		
			7	7
'n	r	4	•	Š
1	ŀ	I		1
ì		Ė		Ė
3	Ŧ	ġ	ŧ	ú
į			L	÷
i	ī			
ł	r	•		۰
	l		ı	
144	7		ą	P
4	Ŗ		÷	
4	ŧ		į	
٠	ĺ			ż
ł	Ľ		_	
i	ī			
ł	۲	7		7
á	è		i	
4	þ	b	à	ø
1	Ŀ	d	i	
j	4	٠	ė	
4	Ę		S	Ξ
J	ø	ę		4
4	í			
J	Þ	Ē		į
å	à	ø	q	
J	Ŀ		١	J
	2	Ξ	2	_
Ĵ	ī	d	P	7
Ţ				
	à		Ġ	L
1	ľ		3)
	3		ť	۲.
1	r		۹	١
J				ę

Question 7 continued	Leave blank
	Q7
(Total 5 marks)	



8. With respect to a fixed origin O, the lines l_1 and l_2 are given by the equations

$$l_1: \mathbf{r} = \begin{pmatrix} 1 \\ -3 \\ 2 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}, \qquad l_2: \mathbf{r} = \begin{pmatrix} 6 \\ 4 \\ 1 \end{pmatrix} + \mu \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix}$$

where λ and μ are scalar parameters.

(a) Show that l_1 and l_2 do not meet.

(4)

The point P is on l_1 where $\lambda = 0$, and the point Q is on l_2 where $\mu = -1$

(b) Find the acute angle between the line segment PQ and l_1 , giving your answer in degrees to 2 decimal places.

(5)

(c) Find the shortest distance from the point Q to the line l_1 , giving your answer to 3 significant figures.

(2)



	Leave
	blank
Question 8 continued	



	Leav
	blan
Question 8 continued	Oldin
Question o continueu	

ì	ė	r
Ė		۹.
1	ш	ш
Ġ	5	Ξ
1	▢	=
	-	=
I	c	E
	7	7
ì	rία	m.
1	υ	ч
ì	-	
3	٩	=
ì	4	÷
ì	Ľ	÷
1	г	_
ì	÷	÷
4	ò	_
c		Ξ
4		_
ì	Ν'n	٠ú
1	ы	4
1	Ŀ	÷
1	Γ.	
1	=	=
1	▢	⊆
ũ	_	Ξ
î	2	7
î	ã	⋑
ũ		
1	H	_
3	×	ĸ.
J	С	3)
	>	_
ľ	2	~
1	٠	=
	Ė	ď
ï	г	٦.
7	÷	o.
Ì	r.	ď
1	÷	4
Ĭ		ď

Question 8 continued	Leave blank
	Q8
(Total 11 marks)	



9.




Diagram not drawn to scale

(2)

Figure 2

(a) Find

$$\int \frac{1}{\left(2x-1\right)^2} \, \mathrm{d}x$$

Figure 2 shows a sketch of the curve with equation y = f(x) where

$$f(x) = \frac{12}{(2x-1)} \qquad 1 \leqslant x \leqslant 5$$

The finite region R, shown shaded in Figure 2, is bounded by the line with equation x = 1, the curve with equation y = f(x) and the line with equation $y = \frac{4}{3}$.

The region R is rotated through 2π radians about the x-axis to form a solid of revolution.

(b) Find the exact value of the volume of the solid generated, giving your answer in its simplest form.

(6)

	blank
Question 9 continued	Ulalik
Question / continued	



	Leave
	blank
Question 9 continued	
Question 7 continued	

4
iii
띘
4
-
11.00
V
-
+
-
-
994
ш
100
100
-
Œ
1
\leq
-
1
·
\sim
$\overline{}$
\Box
$\cdot = \cdot$
\mathbf{c}

Question 9 continued	Leave blank
	Q9
(Total 8 marks)	



(7)

10. The curve C satisfies the equation

$$xe^{5-2y} - y = 0$$
 $x > 0, y > 0$

The point P with coordinates $(2e^{-1}, 2)$ lies on C.

The tangent to C at P cuts the x-axis at the point A and cuts the y-axis at the point B.

Given that O is the origin, find the exact area of triangle OAB, giving your answer in its simplest form.

estion 10 continued	
	_
	_



estion 10 continued	

Leave

6	P	
~	G,	
A DEA	Ы	
포	Ξ	
Ю	⊏	
▔	=	
15	L	
	7	
11 1	Ri.	
\mathbf{v}	·Щ	
UIII-	=	
л	т.	
됐	=	
ъ	÷	
л.		
壳	ø	
4	_	
$\dot{=}$	÷	
	٠.	
ш	ш	
砈	_	
æ	_	
è	Ė	
-	im	
ш	-	
-	Š	
-	Si	
-	_	
14	_	
恢	_	
С	٦	
-91	d	
7	7	
4		
	. *	
P	٩	
%	J	
	×	
14	_1	

A			
	Ó		
		۹,	
	V		
	À		
V			
4		Ĺ	
V			
		V	
	Ç		
	۵		
Νí			
Š			
S			

Question 10 continued	blank
	Q10
(Total 7 marks)	Q10
(10tal / marks)	



blank

11.




Figure 3

(a) By writing $\sec \theta$ as $\frac{1}{\cos \theta}$, show that when $x = 3 \sec \theta$,

$$\frac{\mathrm{d}x}{\mathrm{d}\theta} = 3\sec\theta\tan\theta$$

(2)

Figure 3 shows a sketch of part of the curve C with equation

$$y = \frac{\sqrt{x^2 - 9}}{x} \qquad x \geqslant 3$$

The finite region R, shown shaded in Figure 3, is bounded by the curve C, the x-axis and the line with equation x = 6

(b) Use the substitution $x = 3 \sec \theta$ to find the exact value of the area of R. [Solutions based entirely on graphical or numerical methods are not acceptable.] (7)



uestion 11 continued	



estion 11 continued	

ì	ė	۴
7	7	Ni.
1	H	٠II
1	_	5
1	0	=
4	-	Ξ
ì	e	r
		,
'n	r i	'n.
1	u	7))
'n	-	i
ì	4	-
ì	4	÷
	C	Ξ
ł	_	-
		۰
	2	~
4	F	-
4	-	=
ì	. :	á.
1	ш	4
	Ŀ	Ξ
ļ	т	7
4	-	
1	n	m
	Ħ	
Ą	3	×
1	5	Si
J	~	_
1	Ĺ	_
4	7	7
1	m	٩
4	١.	J
9		ė
1	2	_
÷		7
٠,	'n	iκ
4	Ŀ	2)
	Z	Ξ.
1	r	٦
J	=	4

×	Ü	Ù	
< ₹	Z	Ξ	
×	٩	0	
8	Δ	2	
Z	ς	Ľ	
z	N	Ρĺ	
N	~	4	
/:	S	Ζ	
N	V V	Z	
4	īΧ	-	
N	۳	7	
< -	\times	۲.	
×	Ź	7	
⟨ 4	ς,	÷	
×	-	7	
Çη	ıλi	ũ	
×	μ	ø	
SΙ	Й Н		
2	Ŀ	2	
5	-	ä	
z	Ω	2	
Ý	÷	ĸ.	
Zi		Z	
Χý	ø,	₹	
/3	К	/	
X	m	₹	ļ
A	n	۹	
X	G	И	
Ö	×	ś	
Ž	ď	S	
62	\times		
	P	۹	
ÇI	×	ø	
×	æ	×	

$\wedge \wedge \wedge \wedge$	/
	7
(XXXX)	
XXXX	2
	١,
$\propto \propto \propto$	
XXXX	>
\times	
$\wedge \wedge \wedge \wedge$	
$\times \times \times \times$	
(XX	
$\times \times \times \times$	
	ĸ
XX	Þ
	λ
0000	
XX	
$\sim\sim$	Ŕ
\sim	
XX	
$\sim\sim$	
$\triangle \triangle \triangle$	
DOCK A	D
$\langle X X X X X X X X X X X X X X X X X X X$	К
$\wedge \wedge \wedge \wedge \wedge$	/
DOC X X	٥,
	К
$\times \times \times \times$	2
	N,
$\times \times$	
	١.
	ν
\times	
(XXX	
$\triangle \triangle \triangle \triangle$	/
XXKM	
CXX	Κ.
20 m/o	2
	١.
∞	
XX	
$\nabla\nabla$	ь
$\triangle \triangle \triangle \triangle \triangle$	
$\Diamond \Diamond \otimes \Diamond$	۲.
	К
XX	۷
DO NOT WRITE IN THIS AREA	К,
$\triangle \triangle \overline{\triangle} \triangle$	
XXXX	
	ś.
XXX	
$\infty \infty$	ĸ.
$\sim\sim$	v
XXXX	
CXX.	К
A	2
XXXX	5
$\langle XXX \rangle$	
\triangle	0
V/1//	b
CXX	۲
$\times \times =$	>
VV/	
CO 00	ľ
$\times \times \times \times$	5
	١.

	Leave blank
Question 11 continued	
	011
	Q11
(Total 9 marks)	



Leave blank

12.	(a)	Show	that

$$\cot x - \tan x \equiv 2 \cot 2x, \quad x \neq 90n^{\circ}, n \in \mathbb{Z}$$

(4)

(b) Hence, or otherwise, solve, for $0 \le \theta < 180^{\circ}$

$$5 + \cot(\theta - 15^{\circ}) - \tan(\theta - 15^{\circ}) = 0$$

giving your answers to one decimal place.

[Solutions based entirely on graphical or numerical methods are not acceptable.]

- (5	١
•	J	J

4: 10 4: I		
estion 12 continued		



estion 12 continued	

ì	ė	ø		_	
	7	7	٩	ų	
111	Ŀ		Ŀ		
G	3	Ξ	3	2	
]	E		c	_	
			ä	ď	
J	ξ	S		_	
			7	7	
1	r	À	ρ	à	
	٠	ę		Ą	
3				9	
Ġ	7		۳	7	
1		ą	b	Ę	
J	ŀ		ė		
ŀ	۰				
1	ė	ŧ	ġ	p	
4	d		=		
á	i		i	=	
4	Ľ		ľ		
ŝ	7		7	7	
ł	P	Ŧ		9	
á	ì		Ė		
d	ρ	٩	à	ø	
4	Ŀ	d	b		
J	۹	9	ġ	h	
1	8	Ę	S	á	
J	۹	7	7	7	
1	Ĺ	٠		Ŀ	
1	ŗ	3		7	
7	P	q	۹	١	
3	ij	e	ġ	ø	
4	÷	5	ā	p	
٦	ď		í	ń	
•				٠	
'n	ø	P	9		
4	b	ė	d	ø	
ī,	ó	ė	q	ĸ	
1	Ĺ			1	
				-	

A			
à			
X			
Ý			
V			
		$\overline{}$	
H	Ď	\mathbb{Z}	
H	μ	4	
d	P	4	
J	P	2	
Ŋ	P	2	
g		2	
8	P	2	
3	P	کر ان	
K		2 2 2	
K		2	
K			
Ķ			
Ķ		2 2 2 3	
K		/ / / /	
K			
K			
Ķ			
K			
K			THE STREET
K			
K			
K			
K			
K			THE PERSON NAMED IN COLUMN 1
K			
K			
K			
K			
K			
K			
K			
K			
K			
K			
K			
K			
K			
K			
K			
K			
K			
K			
K			
K			
K			
K			

×	×	×	×	>		×
V	V	V	V	۲,		V
	$\langle \rangle$			Κ		
A	A	X,	×	2		Λ
\times	×		×			×
V	V	V	V	٦,		V
	$\langle \rangle$			Κ		
25	Χ	X	Х	2		Х
×	×	×	×	2		×
V	\vee	ш	Ŧ	٦		V
		$\overline{}$	ò	К		
\wedge	$^{\sim}$	ďΝ	n	ď		$^{\sim}$
×	×			D		
\vee	\sim	\subseteq	\sim	ā	Ì	\sim
		г	r	ĸ		
X	Λ	_	_	y.	S	Λ
×	X	×	24	B)		X
V	V	S	¥ X	^	2	V
			0	ĸ		
X	х	х	X	2	۲	Х
×	×	ll (d		D		×
\vee	$^{\circ}$	w	v	К,		v
		ê	û	ĸ		
X	X	×	×	-2		х
×	×	X	K	5		×
S	V		X A A	Ď,	2	V
		E)		K		
\times	Х	r	_	₽		Х
×	×	™.	×			
$^{\circ}$	\Diamond			ĸ		V
	\bigcirc	a	e	ĸ		
X	х	鮜	2	p)		X
×	×	×	~	6		×
V	$\langle \rangle$			↸		
		6)	റ	ď		
×	х	М		P		Х
×	×	×	X 0 7	٦		
\circ	\bigcirc	ш		К		
\triangle		N.				
×	X	쁴	×	B)		×
×	×	m	ы	B)		×
\circ	$\langle \rangle$	Œ	С	K		
		$\overline{}$		₹′		
×	X	79	96	Ð		×
×	×		S			X
O	\bigcirc		3	K,		
		5		K		
X	×	ĸΚ				
×	×	æ	₹	в,		×
	$\langle \rangle$	ಒ		Κ		
	\triangle	c	$\overline{}$	k		
×	×	ж.	-24	D		×
×	V	2	\leq			V
<2		7	a	К		
\triangle			2	ď		
X		\times	X			
×	×	v	2	S		×
		e.	d	ĸ		
6	8	ж.	۲	ľ		Α
	×	×	≾	>		X
V	V	e	4	ŀ.		<.
		₽		ĸ		
\triangle	$^{\wedge}$	5	$^{\sim}$	2		Λ

Q12
(Total 9 marks)



Leave blank

13. (a) Express $\frac{1}{(4-x)(2-x)}$ in partial fractions.

(2)

The mass, x grams, of a substance at time t seconds after a chemical reaction starts is modelled by the differential equation

$$\frac{dx}{dt} = k(4-x)(2-x), \quad t \ge 0, \ 0 \le x < 2$$

where k is a constant.

Given that when t = 0, x = 0

(b) solve the differential equation and show that the solution can be written as

$$x = \frac{4 - 4e^{2kt}}{1 - 2e^{2kt}}$$

(7)

Given that k = 0.1

(c) find the value of t when x = 1, giving your answer, in seconds, to 3 significant figures.





	blanl
Question 13 continued	



estion 13 continued	

ì		۲.	
1	7	ij	
1	ш		
Ġ	5	Ξ	
1	9	⊆	
	Ξ	2	
J	ς	L	
		7	
i	Ú	Ri.	
3	ų	·il	
ŝ	7	=	
å	٦	т.	
3	÷	=	
J	×	ė	
ŀ			
þ	÷	ø	
4	۷	_	
Ĝ		Ξ	
4	Ш	1	
å		=	
4	=	=	
å	ì	÷	
4	h	įπ	
J	н		
J	•	'n	
4	ς	⊆	
J	P	-	
1			
4	7	=	
ď	m	٩.	
4	-	J	
9	=	ė	
'n	e	_	
÷		7	
٠,	'n	iκ	
4	Ŀ	2)	
-	Z	Ξ.	
1		2	
ĕ	7	=	

		4	
X			
S			
K			
K			
К			
Z			
N			
	ς	z	
C.	5	3	
S	5	S	
S	5	Ž.	
Ž	X		
ä			
	2		
		\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	
		≦ 3	
		\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	
		ることで	

		-)				>		
2	V	۲		S		ζ		١,
\rightarrow		Χ			>		>	
5	×	2		2	ζ	?		2
		0				>		
1	\vee	٩	2	١,	1	5	7	١,
>		K						
١,	$^{\wedge}$	v	1	/	\	2	S	7
<			4	è		p	r	
1	V	٦	я	S		L	7	١,
\geq		×	2	ζ.	5	ς		
1	$^{\sim}$	2		а	L	а	К	7
			₹	7	7	2	ĸ	\geq
0	V	Ν,		3	ρ	5		١,
>		×	р	ú	Þ	ģ		
S	Ä	2	S		7	à		Č
)				Б		×
2	V	۲	7	7	9	۹	ν	Š
>		×			>		>	
5	X	2		2				7
			-9	θ	Ţ	Д	ŗ	
1	V	٩	ź	۷.	4	×	7	١,
>		K	2	Ξ	2	Ε		
S		×	7			7	5	?
<			4	4	Ľ	è	Ĺ	>
1	V	٦,	ä	2	7	2		5
\geq		×		ø	þ	ø		
1	Λ	2	3	2	Ś	7		2
	X		∠	$^{\sim}$	_	₹	(\geq
0	V	Ν,	z	Σ		9		١,
->		×			è	ú	ò	
1		V.	¥	ď	×	4		ď
			⋖	7	₹		ľ	
7	V	٦	4	\	c	S	/	١,
>		×		ⅎ				
1	$^{\wedge}$	1	3	7	5	7	١,	7
				_	_	4	C	
1	V	٩		₹	7	2	7	١,
>		K	ñ	6	è	í	D	
S		×	S	s	۷	á		?
<				2	r			
1		٦,	8	3	•	5		5
\geq		Χ	9		ù		⊳	
1	Λ	d	'n		5	۲	5	2
		0	S	3				×
0	V	Ν,	r	₹	7	S	?	١,
>		K	ù					
1		V.		pi	۹	μ	١,	
			₹	2	۷	>		
7		1	а	5	7	٩	ľ	
>		Χ	я	Ĺ	a		Þ	
1		⋖	7	z	Σ	Z		
		.)	₹	2		P	r	
1	V	٩	d		2	۷.	7	١,
Ż				₹				
N		×	7	ä	è	Č	5	
<)		2	₹		K	
1		1	я	'n	ú	И		
>		Κ	2	ä		Ĺ	>	
					⋖			7
Ż	X	S		≟	í	ä	K	
	Š	3	ş		É	7	Ķ	

Question 13 continued	Leave blank
	Q13
(Total 11 marks)	VIS



Leave blank

14. Given that

$$y = \frac{(x^2 - 4)^{\frac{1}{2}}}{x^3} \qquad x > 2$$

(a) show that

$$\frac{dy}{dx} = \frac{Ax^2 + 12}{x^4(x^2 - 4)^{\frac{1}{2}}} \qquad x > 2$$

where *A* is a constant to be found.




Figure 4

Figure 4 shows a sketch of part of the curve with equation y = f(x) where

$$f(x) = \frac{24(x^2 - 4)^{\frac{1}{2}}}{x^3} \qquad x > 2$$

(b) Use your answer to part (a) to find the range of f.

(5)

(6)

(c) State a reason why f^{-1} does not exist.

(1)

nestion 14 continued	



	Leave
Question 14 continued	blank
Question 14 Continued	

nestion 14 continued	



	Leave blank
uestion 14 continued	
	Q14
(Total 12 marks)	
TOTAL FOR PAPER: 125 MARKS	
END	