Examiner's use only

Team Leader's use only

Question Number

1

2

3

4

5

6

7

8

Leave Blank

Centre No.			Paper Reference		Surname	Initial(s)					
Candidate No.			6	6	7	7	/	0	1	Signature	

Paper Reference(s)

### 6677/01

# **Edexcel GCE**

### **Mechanics M1**

## Advanced/Advanced Subsidiary

Tuesday 7 June 2005 – Afternoon

Time: 1 hour 30 minutes

Materials required for examination
Mathematical Formulae (Lilac or Green)

Materials required for examination
Mathematical Formulae (Lilac or Green)

Nil

Candidates may use any calculator EXCEPT those with the facility for symbolic algebra, differentiation and/or integration. Thus candidates may NOT use calculators such as the Texas Instruments TI 89, TI 92, Casio CFX 9970G, Hewlett Packard HP 48G.

question papers	
fo o'll'Au fou	
facility for indidates may	
I 92, Casio	

#### **Instructions to Candidates**

In the boxes above, write your centre number, candidate number, your surname, initial(s) and signature.

You must write your answer to each question in the space following the question.

Whenever a numerical value of g is required, take  $g = 9.8 \text{ m s}^{-2}$ .

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

#### **Information for Candidates**

A booklet 'Mathematical Formulae and Statistical Tables' is provided.

Full marks may be obtained for answers to ALL questions.

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).

There are 8 questions in this question paper. The total mark for this paper is 75.

There are 20 pages in this question paper. Any blank pages are indicated.

#### **Advice to Candidates**

You must ensure that your answers to parts of questions are clearly labelled.

You must show sufficient working to make your methods clear to the examiner. Answers without working may gain no credit.

This publication may be reproduced only in accordance with Edexcel Limited copyright policy.

©2005 Edexcel Limited.

 $\stackrel{\text{Printer's Log. No.}}{N20911A}$ 

W850/R6677/57570 4/7/3/3/3/3/3/2





Turn over


Total

	physicsandmanistrior.com	10 200
	In taking off, an aircraft moves on a straight runway $AB$ of length 1.2 km. The a moves from $A$ with initial speed 2 m s <sup>-1</sup> . It moves with constant acceleration and 20 it leaves the runway at $C$ with speed 74 m s <sup>-1</sup> . Find	ircraft s later
	(a) the acceleration of the aircraft,	
		(2)
	(b) the distance <i>BC</i> .	(4)
_		
_		
_		
_		
_		
_		
_		

	٦
∟eave	
alonle	
olank	

2.	Two small steel balls $A$ and $B$ have mass 0.6 kg and 0.2 kg respectively. They are moving towards each other in opposite directions on a smooth horizontal table when they collide directly. Immediately before the collision, the speed of $A$ is 8 m s <sup>-1</sup> and the speed of $B$ is 2 m s <sup>-1</sup> . Immediately after the collision, the direction of motion of $A$ is unchanged and the speed of $B$ is twice the speed of $A$ . Find						
	(a) the speed of $A$ immediately after the collision,	(5)					
	(b) the magnitude of the impulse exerted on <i>B</i> in the collision.	(3)					
		_					
		_					

Figure 1



A smooth bead B is threaded on a light inextensible string. The ends of the string are attached to two fixed points A and C on the same horizontal level. The bead is held in equilibrium by a horizontal force of magnitude 6 N acting parallel to AC. The bead B is vertically below C and  $\angle BAC = \alpha$ , as shown in Figure 1. Given that  $\tan \alpha = \frac{3}{4}$ , find

(a) the tension in the string,


3.

**(3)** 

(b) the weight of the bead.

**(4)** 

4. Figure 2



A box of mass  $2 \, \text{kg}$  is pulled up a rough plane face by means of a light rope. The plane is inclined at an angle of  $20^{\circ}$  to the horizontal, as shown in Figure 2. The rope is parallel to a line of greatest slope of the plane. The tension in the rope is  $18 \, \text{N}$ . The coefficient of friction between the box and the plane is 0.6. By modelling the box as a particle, find

(a) the normal reaction of the plane on the box,

**(3)** 

(b) the acceleration of the box.

**(5)** 

Leave	
blank	

- 5. A train is travelling at 10 m s<sup>-1</sup> on a straight horizontal track. The driver sees a red signal 135 m ahead and immediately applies the brakes. The train immediately decelerates with constant deceleration for 12 s, reducing its speed to 3 m s<sup>-1</sup>. The driver then releases the brakes and allows the train to travel at a constant speed of 3 m s<sup>-1</sup> for a further 15 s. He then applies the brakes again and the train slows down with constant deceleration, coming to rest as it reaches the signal.
  - (a) Sketch a speed-time graph to show the motion of the train,

**(3)** 

(b) Find the distance travelled by the train from the moment when the brakes are first applied to the moment when its speed first reaches  $3 \text{ m s}^{-1}$ .

**(2)** 

(c) Find the total time from the moment when the brakes are first applied to the moment when the train comes to rest.

**(5)** 






Question 5 continued	
	(Total 10 marks)



Figure 3



A uniform beam AB has mass 12 kg and length 3 m. The beam rests in equilibrium in a horizontal position, resting on two smooth supports. One support is at the end A, the other at a point C on the beam, where BC = 1 m, as shown in Figure 3. The beam is modelled as a uniform rod.

(a) Find the reaction on the beam at C.

**(3)** 

A woman of mass 48 kg stands on the beam at the point D. The beam remains in equilibrium. The reactions on the beam at A and C are now equal.

(b) Find the distance AD.

6.

**(7)** 

Question 6 continued	b l
	(Total 10 marks)



7. Figure 4




Figure 4 shows a lorry of mass  $1600\,\mathrm{kg}$  towing a car of mass  $900\,\mathrm{kg}$  along a straight horizontal road. The two vehicles are joined by a light towbar which is at an angle of  $15^\circ$  to the road. The lorry and the car experience constant resistances to motion of magnitude  $600\,\mathrm{N}$  and  $300\,\mathrm{N}$  respectively. The lorry's engine produces a constant horizontal force on the lorry of magnitude  $1500\,\mathrm{N}$ . Find

(a) the acceleration of the lorry and the car,

**(3)** 

(b) the tension in the towbar.

**(4)** 

When the speed of the vehicles is  $6~{\rm m~s^{-1}}$ , the towbar breaks. Assuming that the resistance to the motion of the car remains of constant magnitude  $300~{\rm N}$ ,

(c) find the distance moved by the car from the moment the towbar breaks to the moment when the car comes to rest.

**(4)** 

(d) State whether, when the towbar breaks, the normal reaction of the road on the car is increased, decreased or remains constant. Give a reason for your answer.

**(2)** 



Leave	
blank	

8.	[In this questi	ion, the	unit	vectors	i an	d j	are	horizontal	vectors	due	east	and	north
	respectively.]												

At time t = 0, a football player kicks a ball from the point A with position vector  $(2\mathbf{i} + \mathbf{j})$  m on a horizontal football field. The motion of the ball is modelled as that of a particle moving horizontally with constant velocity  $(5\mathbf{i} + 8\mathbf{j})$  m s<sup>-1</sup>. Find

(a) the speed of the ball,

**(2)** 

(b) the position vector of the ball after t seconds.

**(2)** 

The point B on the field has position vector  $(10\mathbf{i} + 7\mathbf{j})$  m.

(c) Find the time when the ball is due north of B.

**(2)** 

At time t = 0, another player starts running due north from B and moves with constant speed  $v \text{ m s}^{-1}$ . Given that he intercepts the ball,

(d) find the value of v.

**(6)** 

(e) State one physical factor, other than air resistance, which would be needed in a refinement of the model of the ball's motion to make the model more realistic.

**(1)** 



uestion 8 continued		
	(Total 13 marks)	
	TOTAL FOR PAPER: 75 MARKS	