Please check the examination deta	ails below b	efore entering y	our candidate information	
Candidate surname		Othe	er names	
Pearson Edexcel International Advanced Level	Centre	Number	Candidate Number	
Wednesday 1	3 N	lay 20)20	
Morning (Time: 1 hour 30 minute	es)	Paper Refere	nce WMA11/01	
Mathematics				
International Advanced Subsidiary/Advanced Level Pure Mathematics P1				
You must have: Mathematical Formulae and Stat	tistical Ta	ables (Lilac), c	alculator Total Marks	

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 9 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
- use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over ▶







	$(3pq^2)^4 \times 2p\sqrt{q^8} \equiv ap^4$	$Q'q^c$
find the val	ues of the constants a , b and c .	
Tille tile var	des of the constants a, b and c.	(3

	×	×
×		
Χį		
Ŋ		
~	4	\triangle
v	v.	96
	7	96
		-
	×	
A.	_	$\overline{}$
	75	_
	o	-
	E/	∠11
v	w	-0
		7
	a	9
		~
. 2	. 7	7
Χ	х	X
	wi	
Ν	r	71
v1	B.Z	SAI
	74	-
	Sel	m/
^3	r	7
	B.C	-34

Question 1 continued	Leave blank
Zarsaon I continued	
	Q1
(Total 3 marks)	



 $f(x) = 3 + 12x - 2x^2$

(a) Express f(x) in the form

$$a - b(x + c)^2$$

where a, b and c are integers to be found.

(3)

The curve with equation y = f(x) - 7 crosses the x-axis at the points P and Q and crosses the y-axis at the point R.

(b) Find the area of the triangle PQR, giving your answer in the form $m\sqrt{n}$ where m and n are integers to be found.

(4)

		×	2
×,	ä		ĸ
$\sqrt{1}$	۰	а	ĸ
		Z	7
	ßΚ	ш	48
Ų.	ь		a
			Σ
			۳
	ш	z	_
	7	-	₹
		ă	ρ
\sim			
×	Z	9	
×.	×	á	٤
A	Κ.	o	э
	ч	ĸ.	ч
Χį	_		4
		≂	7
	ø		۰
		8	2
Ŋ			
	ik		
×	ь	×	÷
	r		7
\checkmark			/
$^{\wedge}$	Δ	4	_
×.	Z	a	r
: a		z	_
	75		₹
U)			
			/
	×	_	4
201	ш	к	н
	k.		2
\sim		-	_
a	R		
. A	-	×	-
	73	07	↸
			ρ
N	е	z	\sim
×.	⇁	=	e
i o	e,	×	4
×.	×	a	
V.	9	ě	4
$^{\sim}$	~		
	Z		
. a	'n.		
X	56	-	-
V	ĸ.	4	7
	Zá		ಎ
ΧI	К	0	۹
w	ĸ.	X,	л
\wedge	74	ø	К
	=	ä	ĸ.
	a	ø	25
Χı		÷	6
v2		×	7
Λ		ú	/
	и	×	к
a	r		3
	w	-	σ
	S		2
$^{\wedge}$	o	ø	ø.

DO NOT WRITE IN THIS AREA

Question 2 continued	Leave
Question 2 continued	



estion 2 continued	

Leave

ΧĮ					
				2	
M	ĺ	e	Í	2	
X	į	Š	Ź	2	
X'		Š			
X	į	Š	É	2	֡
ă		ì	ĺ		֡
X					֡
					֡
					֡
					֡

Question 2 continued	
	Q2
(Total 7 marks)	J



3.




Figure 1

Figure 1 shows the design for a badge.

The design consists of two congruent triangles, *AOC* and *BOC*, joined to a sector *AOB* of a circle centre *O*.

- Angle $AOB = \alpha$
- AO = OB = 3 cm
- $OC = 5 \,\mathrm{cm}$

Given that the area of sector AOB is $7.2 \,\mathrm{cm}^2$

(a) show that $\alpha = 1.6$ radians.

(2)

- (b) Hence find
 - (i) the area of the badge, giving your answer in cm² to 2 significant figures,
 - (ii) the perimeter of the badge, giving your answer in cm to one decimal place.

(8)

		×	2
×,	ä		ĸ
$\sqrt{1}$	۰	а	ĸ
		Z	7
	ßΚ	ш	48
Ų.	ь		a
			Σ
			۳
	ш	z	_
	7		₹
		ă	ρ
\sim			
×	Z	9	
×.	×	á	٤
A	Κ.	o	э
	ч	ĸ.	ч
Χį	_		4
		≂	7
	ø		۰
		8	2
Ŋ			
	ik		
×	ь	×	÷
	r		7
\checkmark			/
$^{\wedge}$	Δ	4	_
×.	Z	a	r
: a		z	_
	75		₹
U)			
			/
	×	_	4
201	ш	к	н
	k.		2
\sim		-	_
a	R		
. A	-	×	-
	73	07	↸
			ρ
N	е	z	\sim
×.	⇁	=	e
i o	e,	×	4
×.	×	a	
V.	9	ě	4
$^{\sim}$	~		
	Z		
. a	'n.		
X	56	-	-
V	ĸ.	4	7
	Zá		ಎ
ΧI	К	0	۹
w	ĸ.	X,	а
\wedge	74	ø	К
	=	ä	ĸ.
	a	ø	25
Χı		÷	6
v2		×	7
Λ		ú	/
	и	×	к
a	r		я
	w	-	σ
	S		2
$^{\wedge}$	o	ø	ø.

	Leave blank
Question 3 continued	Oldin



estion 3 continued	

ΧĮ					
				2	
M	ĺ	e	Í	2	
X	į	Š	Ź	2	
X'		Š			
X	į	Š	É	2	֡
ă		ì	ĺ		֡
X					֡
					֡
					֡
					֡

	Leave blank
Question 3 continued	Oldlik
	Q3
(Total 10 marks)	



4.	Use algebra to solve the simultaneous equations	
	2 4	
	y - 3x = 4	
	$x^2 + y^2 + 6x - 4y = 4$	
	You must show all stages of your working.	
	Tou must show an stages of your working.	(7)
		(1)



Leave

		×	2
×,	ä		ĸ
$\sqrt{1}$	۰	а	ĸ
		Z	7
	ßΚ	ш	48
Ų.	ь		a
			Σ
			۳
	ш	z	_
	7	-	₹
		ă	ρ
\sim			
×	Z	9	
×.	×	á	٤
A	Κ.	o	э
	ч	ĸ.	ч
Χį	_		4
		≂	7
	ø		۰
		8	2
Ŋ			
	ik		
×	ь	×	÷
	r		7
\checkmark			/
$^{\wedge}$	Δ	4	_
×.	Z	a	r
: a		z	_
	75		₹
U)			
			/
	×	_	4
201	ш	к	н
	k.		2
\sim		-	_
a	R		
. A	-	×	-
	73	07	↸
			ρ
N	е	z	\sim
×.	⇁	=	e
i o	e,	×	4
×.	×	a	
V.	9	ě	4
$^{\sim}$	~		
	Z		
. a	'n.		
X	56	-	-
V	ĸ.	4	7
	Zá		ಎ
ΧI	К	0	۹
w	ĸ.	X,	л
\wedge	74	ø	К
	=	ä	ĸ.
	a	ø	25
Χı		÷	6
v2		×	7
Λ		ú	/
	и	×	к
a	r		3
	w	-	σ
	S		2
$^{\wedge}$	o	ø	ø.

Q4 (Total 7 marks)	Question 4 continued		blank
	Question 4 continued		
			04
(Total 7 marks)			
	(To	tal 7 marks)	



5. (i)




Figure 2

Figure 2 shows a sketch of the curve with equation y = f(x).

The curve passes through the points (-5, 0) and (0, -3) and touches the *x*-axis at the point (2, 0).

On separate diagrams sketch the curve with equation

(a)
$$y = f(x + 2)$$

(b)
$$y = f(-x)$$

On each diagram, show clearly the coordinates of all the points where the curve cuts or touches the coordinate axes.

(6)

(ii)




Figure 3

Figure 3 shows a sketch of the curve with equation

$$y = k \cos\left(x + \frac{\pi}{6}\right) \qquad 0 \leqslant x \leqslant 2\pi$$

where k is a constant.

The curve meets the y-axis at the point $(0, \sqrt{3})$ and passes through the points (p, 0) and (q, 0).

Find

- (a) the value of k,
- (b) the exact value of p and the exact value of q.

(3)

	blank
Question 5 continued	

	Lea bla
Question 5 continued	

				S
				ú
a		j	Ľ	
X		j	Ľ	1
8	Ļ	4	Ļ	ļ
S	ļ	2		2
X	ļ	2		2
X	ļ			2
X	ļ			2
X	Į		2	
X				
X				
X				
X				

Question 5 continued
(Total 9 marks)



6.	The point A has coordinates $(-4, 11)$ and the point B has coordinates $(8, 2)$.
	(a) Find the gradient of the line AB , giving your answer as a fully simplified fraction. (2)
	The point M is the midpoint of AB . The line l passes through M and is perpendicular to AB .
	(b) Find an equation for l , giving your answer in the form $px + qy + r = 0$ where p , q and r are integers to be found.
	The point C lies on l such that the area of triangle ABC is 37.5 square units.
	(c) Find the two possible pairs of coordinates of point C . (5)



	ě		
Ŋ			

	blank
Question 6 continued	



	Τ.
	Leave
	blank
Question 6 continued	
Question o continued	
	1

×	
	8
ΧĮ	¥
X	
Ŋ	
vì	
Χį	

Overtion 6 continued	Leave
Question 6 continued	
	Q6
(Total 11 marks)	



7. The curve C has equation

$$y = \frac{1}{2 - x}$$

(a) Sketch the graph of C. On your sketch you should show the coordinates of any points of intersection with the coordinate axes and state clearly the equations of any asymptotes.

(3)

The line *l* has equation y = 4x + k, where *k* is a constant.

Given that l meets C at two distinct points,

(b) show that

$$k^2 + 16k + 48 > 0$$

(4)

(c) Hence find the range of possible values for k.

(4)

nestion 7 continued	



Question 7 continued	Lea blar
Question / continued	
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_

		2	Ś	a	
\wedge	d	μ		r	
X1	9			ĸ	
	c			7	
×I	į	Я		1	
V					
	ú	۵	4	4	,
\times	,	₹		₹	
W					
				S	
	à	ø		۳	
	ņ	ú		Ŀ	
X	2			₹	
\vee					
	r	7		۰	
X			n	4	
	₹			2	
×į	μ	۹	×	8	
	è	ú	è	ú	
	Т	٦	г	7	
×	ú		Ŀ	ú	
	Σ	7	7	7	٦
\sim		_	4	_	è
		е	7	₹	
$^{\circ}$	۹				
\wedge	2	2	2	≤	
v	7	z		P	,
				_	
	7	₹	7	₹	
V	۱	н	'n		
	۵	ζ,		4	
()	ı	J	ĸ.	в	
	н	8	۳	4	
	Ŀ			2	
	ş			۲	
M	ı	н	н		
	4	2	4	2	
X	r	7		₹	
W		8	×	н	
$^{\wedge}$	ú	2	2		
O			c	9	
$^{\wedge}$	ż	S			
V	۹	7	2		
	7				
		۷	-	۷	
V		7	₹	7	۰
	z	à	ú	2	
X		٣	7	힉	í
()	6	2	S	d	ľ
\wedge	2	۹	۶	К	
	ä	ø	a		
	Č	á	ø	۲	
X		й	×	ú	
v				7	
	S	2	4	2	
X		۲	9	۹	
C)	ĸ			1	ĺ
	7	۹	ø	×	
	á	d	ń	ĸ	
	۲			٦	

Question 7 continued	Leave blank
Question / continued	
	Q7
(Total 11 marks)	



8. The curve *C* has equation

$$y = (x - 2)(x - 4)^2$$

(a) Show that

$$\frac{\mathrm{d}y}{\mathrm{d}x} = 3x^2 - 20x + 32$$

The line l_1 is the tangent to C at the point where x = 6

(b) Find the equation of l_1 , giving your answer in the form y = mx + c, where m and c are constants to be found.

(4)

(4)

The line l_2 is the tangent to C at the point where $x = \alpha$

Given that $\boldsymbol{l_1}$ and $\boldsymbol{l_2}$ are parallel and distinct,

(c) find the value of α

(3)

		×	2
×,	ä		ĸ
$\sqrt{1}$	۰	а	ĸ
		Z	7
	ßΚ	ш	48
Ų.	ь		a
			Σ
			۳
	ш	z	_
	7		₹
		ă	ρ
\sim			
×	Z	9	
×.	×	á	٤
A	Κ.	o	э
	ч	ĸ.	ч
Χį	_		4
		≂	7
	ø		۰
		8	2
Ŋ			
	ik		
×	ь	×	÷
	r		7
\checkmark			/
$^{\wedge}$	Δ	4	_
×.	Z	a	r
: a		z	_
	75		₹
U)			
			/
	×	_	4
201	ш	к	н
	k.		2
\sim		-	_
a	R		
. A	-	×	-
	73	07	↸
			ρ
N	е	z	\sim
×.	⇁	=	e
D)	e,	×	4
×.	×	a	
V.	9	ě	4
$^{\sim}$	~		
	Z		
. a	'n.		
X	56	-	-
V	ĸ.	4	7
	Zá		ಎ
ΧI	К	0	۹
w	ĸ.	X,	а
\wedge	74	ø	К
	=	ä	ĸ.
	a	ø	25
Χı		÷	6
v2		×	7
Λ		ú	/
	и	×	к
a	r		я
	w	-	σ
	S		2
$^{\wedge}$	o	ø	ø.

	blank
Question 8 continued	
	1



estion 8 continued	

		2	Ś	2	
\wedge	d	μ		ĸ	
X1	9			K	
	c			7	
×I				1	
V					
	ú	۵	4	_	
\times		₹		₹	
W					
				2	
	à	ø		r	
	ņ			D	
X				₹	
\vee				/	
)		۰	
X			r	41	
			2	z	
Χį	۰	۰	,	7	
	è	ú	è	ú	
		1			
×ι	×		ы	6	
	Ξ				
		è	4	÷	
		₹		₹	
v				/	
$^{\wedge}$	ź	4	ú	_	
		Z		۳	
₩.			S	-	
X1	٠	=	-	=	
				>	
ŽΙ	ď	S	v	я	
		а	ь	а	
		7	₹	7	
X		≤	2	≤	
VI		7	7	7	
		\geq	Z	7	
	7	₹	7	₹	
			й		
N		2		\mathbb{Z}	
X.	7	e	=	e	
	٠	×	6	4	
Χ,	2	ú			
V	9	Ħ	6	6	
	á	Ġ	ø	9	
v	×	2		2	
4					
	ŀ				
	d	р	=	ĸ	
N				3	
v	٩	ы	è	ø	
	۵			2	
	7	۲		и	
X	ä		s	2	
S	e	=	₹	7	
	ú	à	ė	ĸ	
X		S		1	
X		6	à	ø	
		2		2	
Λ		×	ø	ø.	

	Leave
Questian & continued	blank
Question 8 continued	
	00
	Q8
(Total 11 marks)	
(Total II marks)	



9.	A curve with equation $y = f(x)$ passes through the point (9, 10).
	Given that
	$f'(x) = 27x^2 - \frac{21x^3 - 5x}{2\sqrt{x}} \qquad x > 0$
	find $f(x)$, fully simplifying each term. (6)



		ě			
X					
. 2	к			1	
		ь	d		ľ

uestion 9 continued	
restion 7 continued	



Leave

Question 9 continued		blank
		Q9
	(Total & marks)	
	(Total 6 marks) TOTAL FOR PAPER IS 75 MARKS	
	END END	