Surname	Other nan	nes
Pearson Edexcel International Advanced Level	Centre Number	Candidate Number
Core Math	nematics	s C34
Wednesday 8 November 20 Time: 2 hours 30 minutes	•	Paper Reference WMA02/01

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information

- The total mark for this paper is 125.
- The marks for each question are shown in brackets
 use this as a quide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶



P51405A
©2018 Pearson Education Ltd.
1/1/1/1/1/



blank

 $f(x) = x^5 + x^3 - 12x^2 - 8,$ 1. $x \in \mathbb{R}$

(a) Show that the equation f(x) = 0 can be written as

$$x = \sqrt[3]{\frac{4(3x^2 + 2)}{x^2 + 1}}$$

(b) Use the iterative formula

$$x_{n+1} = \sqrt[3]{\frac{4(3x_n^2 + 2)}{x_n^2 + 1}}$$

with $x_0 = 2$, to find x_1 , x_2 and x_3 giving your answers to 3 decimal places.

(3)

(3)

The equation f(x) = 0 has a single root, α .

(c) By choosing a suitable interval, prove that $\alpha = 2.247$ to 3 decimal places.



Question 1 continued	blank



estion 1 continued	

×				×	
		ú	ă		ß,
			2	Ç	<
			7	79	ņ
				и	ŀ
		s		싀	ŀ
	8		_	X.	2
	4	Р		ρ	۲
×		ĸ		ĸ	ú
	7	⊽	_	▽.	5
	4	à	и	μ	۲
	я	6		ĸ.	
		0	٦	7	В.
	à	К	à		C
	ч	C		а	Į.
		9	0	~	r
	'n				f)
	×	ï	×	_	ć.
×	2	7		$\overline{}$	5
V	ð			4	ŀ
			×		۲
		Ľ	_	×	٥
		7	-	~	٩,
	ø	٣,			
		×	_	×	2
	×	=			ĸ,
	d			\sim	Z
	×	₹	-	₹	٩
	4				ĸ,
$^{\sim}$		Χ,	ö	^	
×	ì	ĸ	Ĥ	1	ì
		Ľ		ы	ľ
	.9	-	-	-	I)
		Ŀ		4	
		۰	ņ	-	۲
	B	ĸ			
	м	×		,	В,
	d		ы		r
		K		<u>z</u>	2
	e	-	8		۲,
	я	٠	6	۵.	1
		ü	a		P)
	×	9	H)	è	ſ.
×	á	ú	ø		,
V				✓	
	4	r.		2	
			я	8	Ľ
	1		_	/	
	×á	р	ņ	٩	c
		K		×I	ß
	и		ы		۲,
	2	<u> </u>	-7	$^{\sim}$	2
	7	₹			P)
	4	d	e		
	J	=	Ħ	=	9
V		ď	^	V.	
	s	ä		ы	
	1	К		2	Ľ
	٦	ь	ద	d	ĸ
		4		ĸ	Z
	7		Ħ	×	Č
		Ľ		А	ĸ
×	3		ş	=	r
×				×	
		٦			

Question 1 continued	Leave blank
	01
/TF / 10 1 \	Q1
(Total 8 marks)	



2.	The curve	C has	equation
----	-----------	-------	----------

$$y^3 + x^2y - 6x = 0$$

(a) Find $\frac{dy}{dx}$ in terms of x and y.

- (5)
- (b) Hence find the exact coordinates of the points on C for which $\frac{dy}{dx} = 0$
- (6)



\times
× 4 ×
$\times \mathbf{m} \times$
× ne×
◁
\times
\times

₩
<u>~~~</u>
\times
\times
\times
$\otimes \otimes \otimes$
Ä
\times

\times
<u> </u>
XX ** * XX
\times
<u> </u>
888 8 8888
××
WRITE
WRITE
WRITE
OT WRITE!
WRITE
OT WRITE!
NOT WRITE
OT WRITE!
O NOT WRITE!
DO NOT WRITE!
O NOT WRITE!
DO NOT WRITE!
DO NOT WRITE!
DO NOT WRITE!
DO NOT WRITE!
DO NOT WRITE!
DO NOT WRITE!
DO NOT WRITE!
DO NOT WRITE!
DO NOT WRITE!
DO NOT WRITE!
DO NOT WRITE!
DO NOT WRITE!
DO NOT WRITE!
DO NOT WRITE!
DO NOT WRITE!
DO NOT WRITE!
DO NOT WRITE!
A DO NOT WRITE!
EA DO NOT WRITE!
REA DO NOT WRITE!
REA DO NOT WRITE!
REA DO NOT WRITE!
AREA DO NOT WRITE!
S.AREA DO NOT WRITE!
HS AREA DO NOT WRITE!
HS AREA DO NOT WRITE!
FHIS AREA DO NOT WRITE!
THIS AREA DO NOT WRITE!
THIS AREA DO NOT WRITE!
IN THIS AREA DO NOT WRITE!
EIN THIS AREA DO NOT WRITE!
EIN THIS AREA DO NOT WRITE!
EIN THIS AREA DO NOT WRITE!
EIN THIS AREA DO NOT WRITE!
EIN THIS AREA DO NOT WRITE!
EIN THIS AREA DO NOT WRITE!
WRITE IN THIS AREA DO NOT WRITE!
I WRITE IN THIS AREA DO NOT WRITE I
OT WRITE IN THIS AREA DO NOT WRITE!
OT WRITE IN THIS AREA DO NOT WRITE!
NOT WRITE IN THIS AREA DO NOT WRITE!
NOT WRITE IN THIS AREA DO NOT WRITE!
O NOT WRITE IN THIS AREA DO NOT WRITE!
IO NOT WRITE IN THIS AREA DO NOT WRITE!
O NOT WRITE IN THIS AREA DO NOT WRITE!

,			×			
	ì	۹		2	ς	
				\rangle		\rangle
					1	S
	ζ	ì			>	
>			×			
	2	ς		?	ς	2
				>		\rangle
				Ś	2	S
۶	(j	×		2	
,			×			
	2	۹	Š	?	۱	
				\rangle		シ
,	ς	j	×	۱	2	
,			×			
	2	ς	Ų	2	ς	2
				>		
					?	S
	ζ	ì		ί	>	
,			×			
	?	<	×	?	ς	2
				>		
					2	5
۶			×		2	
,			×			
	2	۹	Š	?	ς	
				>		5
	ς	Z		Š	ä	ú
,		1				
	è	S		7	ģ	b
		ì	ì	7	į	ì
		į		7	į	Ì
		ĺ				
		ĺ				
		ĺ				
		1				
		į				
0		į				
		į				
		į				
		į				
		į				
		į				
		į				
		į				

Question 2 continued	Leave blank	
	Q2	
(Total 11 marks)		4



3. The number of bacteria in a liquid culture is modelled by the formula

$$N = 3500(1.035)^t$$
, $t \ge 0$

where N is the number of bacteria t hours after the start of a scientific study.

(a) State the number of bacteria at the start of the scientific study.

(1)

(b) Find the time taken from the start of the study for the number of bacteria to reach $10\,000$

Give your answer in hours and minutes, to the nearest minute.

(4)

(c) Use calculus to find the rate of increase in the number of bacteria when t = 8 Give your answer, in bacteria per hour, to the nearest whole number.

	(2)	
(31	

DO NOT WRITE IN THIS AREA							
NOT WRITE IN THIS ARE						1	
NOT WRITE IN THIS ARE							
NOT WRITE IN THIS ARE			ú	ä	ú	d	
NOT WRITE IN THIS ARE						К	
NOT WRITE IN THIS ARE			Z	۰	9		
NOT WRITE IN THIS A		ú		ì	í.	À	
NOT WRITE IN THIS A	Κ		Ľ			Я	
NOT WRITE IN THIS A	×	3	Z		3	7	
NOT WRITE IN THIS A			я		ü	ø	
NOT WRITE IN THIS A	Κ,		ĸ	J		٤	
NOT WRITE IN THIS A		3					
NOT WRITE IN THIS		4	d	Ħ		P	
NOT WRITE IN THIS	Χ	я	ς	ì		L	
NOT WRITE IN THIS	£					7	۰
NOT WRITE IN I							
NOT WRITE IN I	Χ		ĸ	J		٩	
NOT WRITE IN I			ы		г,	a	
NOT WRITE IN I		C	Ξ	2	۷	2	
NOT WRITE IN I	Κ	9	8		7	₹	
NOT WRITE IN I	Q.	я	p			ø	
NOT WRITE IN I					ľ		
NOT WRITE IN I	Κ						
NOT WRITE		'n	Ŀ			2	
NOT WRITE				ı	ø	۰	
NOT WRITE	Κ	y	К				
NOT WRITE	Ż						
NOT WRITE		ņ	•	,			
NOT WRITE	Κ	à	ä		9	₹	
NOT WRITE		я	P	•	۹	۳	
NOT WR		á	Ĥ	i	ĕ	è	
NOT WR	×					₹	
NOT WR	J	S			S	Z	
NOT WR					ľ	Ή	
NOT WR		9	-		,	ę	
NOT WR			L		S	2	
NOT WR			۲	9	•	۰	
0	K	5					
0		я	×		۰	۰	
0		d	ρ	b	á	и	
0			Ľ	3		2	
0		3	7	۲	5	7	
0		я	۰	٩	б	2	
0		à	й	į	ĕ	ę	
0	2	S	7	۹		si	
0		з	μ	٠	,	5	
0							
0			L		S		
0			۰	ę	۰	-	
DO NO		×	S	i	è	ζ	
DO NO			r	•	₹	я	
DO N			2	k	4		
DON			7	•	7	₹	
DOA		'n		×	Ħ		
00			۵		ρ	5	
DO							
00						2	
0	۲,	ż	š		ě	7	
ă			r	1	3		
Δ		۹	ſ.	ž	٤	d	
			۹		e	3	
		à		×	ij	ij	
						4	
98	<	я			Ė	Ħ	
\sim	×		×				
	/						

Question 3 continued	Leave
	Q3
(Total 8 marks)	



(6)

4. (a) Prove that

$$\frac{1 - \cos 2x}{\sin 2x} \equiv \tan x, \qquad x \neq \frac{n\pi}{2}$$
 (3)

(b) Hence solve, for $0 \leqslant \theta < 2\pi$,

$$3\sec^2\theta - 7 = \frac{1 - \cos 2\theta}{\sin 2\theta}$$

Give your answers in radians to 3 decimal places, as appropriate.

(Solutions based entirely on graphical or numerical methods are not acceptable.)

Question 4 continued	Leave blank



	Lea
	bla
Question 4 continued	
Question i continueu	

G				
	Ú			
4				
	à			
		á		
a				
			þ	
м	β			
		é		
	Ú			
	Ь			
d	þ		2	
			ú	
SI				

Question 4 continued	blank
	Q4
(Total 9 marks)	



5. (i) Find

$$\int \left(\left(3x + 5 \right)^9 + e^{5x} \right) dx \tag{3}$$

(ii) Given that b is a constant greater than 2, and

$$\int_{2}^{b} \frac{x}{x^2 + 5} \mathrm{d}x = \ln\left(\sqrt{6}\right)$$

use integration to find the value of b .				

i		
	Ú	
	Ŝ	
	K	
į		
		5

Question 5 continued	Leave blank
	Q5
(Total 8 marks)	



6.




Figure 1

Figure 1 shows a sketch of the curve C with equation $y = 2e^{-x}\sqrt{\sin x}$, $0 \le x \le \pi$. The finite region R, shown shaded in Figure 1, is bounded by the curve and the x-axis.

(a) Complete the table below with the value of y corresponding to $x = \frac{\pi}{2}$, giving your answer to 5 decimal places.

x	0	$\frac{\pi}{4}$	$\frac{\pi}{2}$	$\frac{3\pi}{4}$	π	
у	0	0.76679		0.15940	0	
						(1)

(b) Use the trapezium rule, with all the values of y in the completed table, to obtain an estimate for the area of the region R. Give your answer to 4 decimal places.

(3)

(c) Given
$$y = 2e^{-x}\sqrt{\sin x}$$
, find $\frac{dy}{dx}$ for $0 < x < \pi$.

The curve C has a maximum turning point when x = a.

(d) Use your answer to part (c) to find the value of a, giving your answer to 3 decimal places.



	ä			
		N A		
			e	
	1	⎖	7	7
	1	7	5	<
Š	1	Ŗ	2	<
	1	7	2	Z
	1	7	2	Z
	1	7	2	7
	1	2	Š	2
X	1	Z 2	Š	2
	1	2	Š	7
	1	2		7
	1			<
	1	2		
	1			
	1100			
× × × × ×				
	日本 付けたりる		A MARKET SERVICES	
× × × × ×		2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2		
× × × × × ×				
× × × × × × ×	11111111111			
X X X X X X X		2000		
× × × × × × × ×				
× × × × × × × ×	111111111111			
× × × × × × × ×				
× × × × × × ×		2 2 2 2 2 2 2 2 2 2 2		
× × × × × × × ×		2 2 2 2 2 2 2 2 2 2 3		
× × × × × × × × × ×				
X X X X X X X X X X				
× × × × × × × × × × ×				
X X X X X X X X X X X				
× × × × × × × × × × × × ×				
× × × × × × × × × × × ×				
× × × × × × × × × × × × × × × ×				
× × × × × × × × ×				
× × × × × × × × × × × ×				
× × × × × × × × × × × × × × × × × × ×				
× × × × × × × × × ×	111111111111			
× × ×				

Question 6 continued	blank
Question o continuous	



	Leave
Question 6 continued	blank
Question o continucu	

			7			
			١,			
			2			
			1			
			2			
			S			
					Ş	
			2	6	à	
			₹		К	
	3	5	7	Ц		
	š	ì	à	ú	ï	
			SI		SI	
	9	١	,	9	7	۰
	4		Ŋ	й		۰
			2		ĸ	
	3		7	3	7	۰
	2		4	d		ļ
	Į		S		K.	
			7	۳	59	۱
	à	í	C	à		í
	٩	ļ	4	r	а	
		۰	5		3	
	я	۰	e	,	ę	۰
	Ý	i	×	'n	=	í
			Я			
	ì	i	á	ń	ń	i
	ç	í	2			
			è	ú	Ś	
			₹	3	₹	
	ä		×	۰	ь	i
		S	Z			
	4			ä	è	i
	G	2	Σ	2	ě	
	8	١	7	7	7	۰
					1	
			q	ď	ч	
					4	
	ã					
			ú	è	4	i
	1					
	ä	í	4	ĕ	$\stackrel{\sim}{}$	i
			Ξ		s	
		ľ	7		7	۰
	J		н		'n	
	Š		7			
	Š				8	
	Š		Ž			ĺ
	200					
	1		2	S	2	
			2	S	2	
			2		į	
	1		2			
	1		2			
	1		2			
	1		2			
	1		2		į	
	1					

	ď	n	ď	٦	ľ
	٩	7	₹	7	٩.
Χ	A	А	и	и	ν
\vee		\square		Z	N
^	ð	=	۳	=	ν
		s	à		D
V)				D	۲
\wedge	2	7			r
V	`	✓			
		. >	۷.	2	
×		Κ			ν
	3		К,	3	ĸ
	Z	\simeq	۷	\sim	7
	ō	~	7		5
	۹			۰	Е,
	ź	X V	ľ		
v	ì				Ŋ
	é		⟨		
	ı	Ŀ	ei)	ú	ν
v	1	D	7		۲,
	2				
	3	ĸ	è	ú	ò
		7		,	٣
Χ	Š		_	s	i
\vee	3	$\overline{}$	7	7	٦
	J	=	,	=	ν
	G	×			
	Ý	w	ŝ.	À	Ŕ.
$^{\wedge}$		N	к	Ы	ν
V	3	•	=	۳	٥,
			C		C
×	ı	ĸ	,	ę	ņ
v	۹	v	١.	2	۱
	J	=	,	=	ν
	7	6	۵	ú	Ď
		12			Κ
Λ	J	=	۰	=	ν
V	ì	≤	7	2	
		3			ľ
	ŧ			Σ	
	S	_			K
	į	7			1
	è	×			Э
		۵	_	۵	٤
х		ĸ	7	₹	"
V	3	4	'n	ĸ	
	4	rs	7	٦	ĸ
×	۹	ĸ	2		D
		9	٠	7	Κ
	ä	'n	×	è	ď
V		Z		۳	٦.
	d		ŭ	à	ć
Х	5	≺	7	₹	2
	5		2		\
	á		ø		l
				d	D
0	ą	è	6	d	К
×	j	s		Z.	
V	ĭ	Р	7		Ν
			4	3	ľ
	9	۰	۲	ę	F)
					Κ

٤			⋖	>					
×		Χ	2						>
×		×				1	7		5
<	×			>		2		>	
0	×		l	5					
>		×							
S		Ż	1			2			S
				>		>			(
1	×			S		S	ì		
\times		×		<					\rangle
1	×	v	۲	1		1			١,
			≺	>					
1	v	Χ	2	1		1		1	1
×		×				<	>		>
ς.	×	O	Κ	2		2		>	ς
			<	7					
×		X	2	Κ					2
1		2	٦	2		2			1
				>)			⟨
1	v	$^{\sim}$		٦		۱	1	١,	1
×		×			2	۷			>
ς,	×	V	'n	è		ĺ		2	ς
			₹	9	Ų	b	ď		
	Ü	Χ	ú	ĸ,	2	G	v		2
×		X	1	И		4	b		>
<		0	j	2	7			2	⟨
			đ	7		5			(
>		×	J			á			>
1		V	`	2	∠	á	'n		/
			4						
1	V	×	2	7	7	٩	ľ	٩	2
\times		×							>
ς,	×	V	'n	2					ς
			ą	И	r	1	ĸ		
7	Ü	Χ	ú	S	2	٤	2		?
\times		×				Ζ			S
ς.			۹	7		7	۲,		
C			۷	2		\leq	ć		
>		Х		↸	7				>
X		×	1	м	è	ú			1
	×		4	D					⟨
C	V	^	2	7		\	l	/	/
>		×	,	₹					>
5	×	Ÿ	S		S	2			5
			₹	7	₹				
7									
						Ŕ	ì	≷	?
\times	ĸ.	X		2	5	Ź	Š	>	S
Š	×	Š	ì	3	S	Š	Š	3	8
ζ	×	X	ì	2	S	Š		3	?
ξ	X	X	į	2	5	į		?	3
ξ	×	X	ì		2			?	3
3	× × ×	3	į		2			?	?
3	×××××		į	Z K				?	?
3	XXXXX	× × ×							
3	× × × ×	× × × ×	1					?	
3	/ / / / / / / /	× × × × ×	1						
	/ / / / / / / / /								
	×××××××								
	×××××××								
	××××××××								
	~×××××××××××××××××××××××××××××××××××××	× × × × × × × ×							
	×××××××××								
	~×××××××××××××××××××××××××××××××××××××	× × × × × × × × × ×							
	/ / / / / / / / / / / / /								
	~×××××××××××××××××××××××××××××××××××××								
	~×××××××××××××××××××××××××××××××××××××								
	~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~								
	/ / / / / / / / / / / / / / / / /								
	××××××××××××××××××××××××××××××××××××××								
	~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~								
	××××××××××××××××××××××××××××××××××××××								
	××××××××××××××××××××××××××××××××××××××								
	××××××××××××××××××××××××××××××××××××××								

	Leave blank
Question 6 continued	
	Q6
(Total 10 marks)	



7. (a) Use the binomial series to expand

$$\frac{1}{\left(2-3x\right)^3} \qquad \left|x\right| < \frac{2}{3}$$

in ascending powers of x, up to and including the term in x^2 , giving each term as a simplified fraction.

(5)

$$f(x) = \frac{4 + kx}{(2 - 3x)^3}$$
 where k is a constant and $|x| < \frac{2}{3}$

Given that the series expansion of f(x), in ascending powers of x, is

$$\frac{1}{2} + Ax + \frac{81}{16}x^2 + \cdots$$

where A is a constant,

(b) find the value of k,

(2)

(c) find the value of A.

(2)

	Leave
Question 7 continued	Dialik
Question / continued	



	Lea
	blar
Question 7 continued	
Question / continued	
	1

Question 7 continued	Leave blank
	Q7
(Total 9 marks)	



8. U	Use partial fractions, and integration, to find the exact value of $\int_{3}^{4} \frac{2x^2 - 3}{x(x - 1)} dx$
7	Write your answer in the form $a + \ln b$, where a is an integer and b is a rational constant. (8)



	blank
Question 8 continued	Dialik
Question o continucu	



nestion 8 continued	

	è			
	1			
	Ś			
	Ŕ			
	2			
	ľ			
			á	
	2			
	Ž			
	D		Л	ŗ.
×	2	S	à	Ŀ

Question 8 continued	Leave blank
	Q8
(Total 8 marks)	



9.

$$f(x) = 2\ln(x) - 4, \quad x > 0, \quad x \in \mathbb{R}$$

- (a) Sketch, on separate diagrams, the curve with equation
 - (i) y = f(x)
 - (ii) y = |f(x)|

On each diagram, show the coordinates of each point at which the curve meets or cuts the axes.

On each diagram state the equation of the asymptote.

(5)

(b) Find the exact solutions of the equation |f(x)| = 4

(4)

$$g(x) = e^{x+5} - 2, \quad x \in \mathbb{R}$$

(c) Find gf(x), giving your answer in its simplest form.

(3)

(d) Hence, or otherwise, state the range of gf.

(1)

Question 9 continued	Leave blank



	Leave
Question 9 continued	blank
Question y continue	

	Ş	9			
	ĺ				
	Ž				
	è			2	
		2			
		j			
		Ė			
	ĺ				
\wedge	STATE OF THE PERSON	ø	ø	٠	
		è			
		í			
	7	d		۲	
				2	
	2	ø	ø	٠	
V					
	2	6	ď	7	
	и	•	9	ч	
	核			3	
	я			۲	
	S	á	ń	2	
	ø	₹	7	٧	
v	16.	6	2	а	
	м	۹	ø	۲	

Question 9 continued	Leave blank
	Q9
(Total 13 marks)	
(10tai 13 marks)	



blank

(2)

(6)

10.




Figure 2

Figure 2 shows a sketch of part of the curve C with parametric equations

$$x = \frac{20t}{2t+1}$$
 $y = t(t-4)$, $t > 0$

The curve cuts the x-axis at the point P.

(a) Find the x coordinate of P.

(b) Show that $\frac{dy}{dx} = \frac{(t-A)(2t+1)^2}{B}$ where A and B are constants to be found. **(5)**

(c) (i) Make t the subject of the formula

$$x = \frac{20t}{2t + 1}$$

(ii) Hence find a cartesian equation of the curve C. Write your answer in the form

$$y = f(x), \qquad 0 < x < k$$

where f(x) is a single fraction and k is a constant to be found.

32

estion 10 continued	



	Leave
On anti-on 10 and those I	blank
Question 10 continued	

Question 10 continued	Leave blank
Question 10 continued	
	Q10
(Total 13 marks)	



11. (a) Given $0 \le h < 25$, use the substitution $u = 5 - \sqrt{h}$ to show that

$$\int \frac{\mathrm{d}h}{5 - \sqrt{h}} = -10\ln\left(5 - \sqrt{h}\right) - 2\sqrt{h} + k$$

where k is a constant.

(6)

A team of scientists is studying a species of tree.

The rate of change in height of a tree of this species is modelled by the differential equation

$$\frac{\mathrm{d}h}{\mathrm{d}t} = \frac{t^{0.2} \left(5 - \sqrt{h}\right)}{5}$$

where h is the height of the tree in metres and t is the time in years after the tree is planted.

One of these trees is 2 metres high when it is planted.

(b) Use integration to calculate the time it would take for this tree to reach a height of 15 metres, giving your answer to one decimal place.

(7)

(c) Hence calculate the rate of change in height of this tree when its height is 15 metres. Write your answer in centimetres per year to the nearest centimetre.

(1)

	Leave
	blank
Question 11 continued	



Question 11 continued

Leave

		į			
	Į	×	,	ę	r)
	ļ	i	₹	5	9
	j	Ľ	S	2	5
	į	ľ	Š	2	P
	į	k	5	3	
	ļ	k	Š	2	
	1		Š		
	ļ	k			
			Y		
		2	Y		
		2			
		k	è		
		k	è		
		ļ		2	
		ļ		2	
			Š		
			Š		
			Š		
			Š		
			Š		

Question 11 continued	blank
	Q11
(Total 14 marks)	



12. Relative to a fixed origin O, the lines l_1 and l_2 are given by the equations

$$l_1: \quad \mathbf{r} = \begin{pmatrix} 2 \\ 0 \\ 7 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ -2 \\ 1 \end{pmatrix} \qquad \qquad l_2: \quad \mathbf{r} = \begin{pmatrix} 2 \\ 0 \\ 7 \end{pmatrix} + \mu \begin{pmatrix} 8 \\ 4 \\ 1 \end{pmatrix}$$

where λ and μ are scalar parameters.

The lines l_1 and l_2 intersect at the point A.

(a) Write down the coordinates of A.

(1)

Given that the acute angle between l_1 and l_2 is θ ,

(b) show that $\sin \theta = k\sqrt{2}$, where k is a rational number to be found.

(5)

The point *B* lies on l_1 where $\lambda = 4$

The point C lies on l_2 such that AC = 2AB.

(c) Find the exact area of triangle ABC.

(3)

(d) Find the coordinates of the two possible positions of C.

(5)



	blank
Overtion 12 continued	Dialik
Question 12 continued	



Question 12 contin	ed	

Question 12 continued	blank



	Leave blank
Question 12 continued	
	012
	Q12
(Total 14 marks)	
TOTAL FOR PAPER: 125 MARKS	
END	