Please check the examination details bel	ow before ente	ring your candidate in	formation
Candidate surname		Other names	
Centre Number Candidate N	umber		
Pearson Edexcel Inter	nation	al Advanc	ed Level
Time 1 hour 30 minutes	Paper reference	WFM	02/01
Mathematics			•
International Advanced S	uhsidiary	v/Δdvanced	Level
	•	y/Advanced	Levei
Further Pure Mathematics	5 F2		
			J
You must have: Mathematical Formulae and Statistical	al Tahles (Ve	llow) calculator	Total Marks
Mathematical Formulae and Statistics	ar rabics (Te	now,, calculator	

Candidates may use any calculator allowed by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use black ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 - there may be more space than you need.
- You should show sufficient working to make your methods clear.
 Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 8 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
- use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over ▶







1. Given that

$$\frac{2n+1}{n^2(n+1)^2} \equiv \frac{A}{n^2} + \frac{B}{(n+1)^2}$$

(a) determine the value of A and the value of B

(1)

(b) Hence show that, for $n \ge 5$

$$\sum_{r=5}^{n} \frac{2r+1}{r^2(r+1)^2} = \frac{n^2+an+b}{c(n+1)^2}$$

where a, b and c are integers to be determined.

(4)



$\times \rangle$
\approx
X
X.
SS
\approx
8
×
×
\otimes
X
×
X
×
\approx
×
×
>>
\otimes
S.
×
X
$\langle \times $
×
×
\approx
$\langle \langle \rangle$
X
×
\otimes
W.
S
\otimes
×
$\langle \times \rangle$
X
\approx
Ø
×
\approx
X
X
\times
\approx
$\langle \! \rangle$
$\langle \cdot \rangle$
S
SS
\times
\approx
$\langle \rangle$
$\langle \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \!$
\$
×
>>
82
×
×
×
\approx
≪
×
\otimes
$\langle \rangle$
\$X
SS
\otimes
×
\otimes
V.
\otimes
\otimes
*
*

Question 1 continued



Question 1 continued

			é		ļ	
			9			
	j					
	×					
	1					
	1					
			á			
	ĭ					
		ľ			3	k
	1	ξ	2	2	ð	ķ
>	7		è	\$	į	ķ
	7		Ì		2	ķ
>	4					k K
>	1					ļ.
	1					ķ
	7					ļ
	1					t
	1				2	ķ
	1					
	į					
	į					
	1					
	į) -
	į					
	į					
	į					
	į					
	į					
	į					
	į					
	į					
	į					
	į					

Question 1 continued	
(To	tal for Question 1 is 5 marks)



2. (a) Use algebra to determine the set of values of x for which

$$x - 5 < \frac{9}{x + 3}$$

(6)

(b) Hence, or otherwise, determine the set of values of x for which

$$x-5<\frac{9}{|x+3|}$$

(2)

			id N			
	į			ļ		
	ĵ					
	Ì		ij		2	
	J					
				á		
	d					
	1					
	í					
	i	ř	4	P	4	ì
	1	Ľ	í	Ĺ	í	k
١	ĺ	Ľ	2	Ĺ	ĺ	ŀ
	Į	Ĺ	1	į	1	į
>	1	Ŀ	_	2	2	ŀ
	1	Ŀ	_	2		į
	j	Ŀ	_	2		
	į	Ŀ	_	2		
	į	Ŀ	_	2		
		Ŀ	_	2		
		Ŀ	_	2		
	1			Ź		
	1			Ź		
	1			Ź		
	1			Ź		
	1			Ź		
	1			Ź		
	1			Ź		
	1			2		
	1			Ź		
	1			Ź		
	1			Ź		
	1			Ź		
	1			Ź		
>>>>>>	1			Ź		
	1			Ź		
	1			Ź		
	1			Ź		
	1			Ź		
	1			Ź		
	1			Ź		
				Ź		
				Ź		
				Ź		
				Ź		
	1			Ź		
				Ź		
				Ź		
				Ź		

Question 2 continued
(Total for Question 2 is 8 marks)



3. The transformation T from the z-plane to the w-plane is given by

$$w = \frac{z}{z + 4i} \qquad z \neq -4i$$

The circle with equation |z| = 3 is mapped by T onto the circle C

Determine

- (i) a Cartesian equation of C
- (ii) the centre and radius of C

(8)

<	$\times \times \times$
×	$\times\!\!\times\!\!\times$
\approx	\propto
K.	$\times\!\!\times\!\!\!\times$
×	$\times\!\!\times\!\!\times$
\approx	∞
×	\approx
\approx	$\times\!\!\times\!\!\times$
×	$\times\!\!\times\!\!\!\times$
×	$\langle \times \times \rangle$
$\langle \! \rangle$	$\times\!\!\times$
×	
\approx	676
×	DO NOT WRITE IN THIS AREA
\approx	
×	
\approx	
\approx	S
×	
\approx	90
×	
\approx	
×	2
\times	
\ll	
×	
\approx	
\otimes	
\approx	250
×	>
\approx	
\otimes	\cong
×	
\otimes	
×	
\otimes	
×	
×	
\otimes	
×	888
\times	$\otimes \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \! \!$
≪	$\times\!\!\times$
×	\approx
\approx	$\times\!\!\times$
×	$\times\!\!\times\!\!\times$
×	
\otimes	
\propto	$\times\!\!\times\!\!\times$
\approx	
×	$\times\!\!\times\!\!\!\times$
\approx	$\times\!\!\times\!\!\times$
K.	$\times\!\!\!\times$
×	\approx
\approx	
×	$\times\!\!\times\!\!\times$
\Rightarrow	$\Leftrightarrow >$
×	$\times\!\!\!\times\!$
\otimes	
8	M
×	00
≪	XX.
X	580
×	r ov
>>	
×	
\otimes	
*	
× ×	E
× ×	THE
*	THE
	E S JE
	TE AL TE
	THUNDER TH
	EXCATE
	VRITERINAL
	WRITEINTH
	WRITEINTH
	A WRITEINTH
	OT WRITE IN TH
	101 WRITE IN TH
	NOT WRITE IN TH
	ONOT WRITE IN TH
	O NOT WRITE IN TH
	DO NOT WRITE IN TH
	DO NOT WRITE IN THIS AREA
	DO NOT WRITEIN TH
	DO NOT WRITE IN TH
	DO NOT WRITE IN TH
	DO NOT WRITE IN TH
	DO NOT WRITE IN TH
	DO NOT WRITE IN TH
	DO NOT WRITE IN TH
	DO NOT WRITE IN TH
	DO NOT WRITEHN TH
	DO NOT WRITE IN TH
	DO NOT WRITE IN TH
	DO NOT WRITE IN TH
	DO NOT WRITE IN TH
	DO NOT WRITE IN TH
	DO NOT WRITEAN TH
	DO NOT WRITE'N TH
	A
	EA
	EA
	REA
	4REA
	4REA
	4REA
	4REA
	HIS AREA
	HIS AREA
	HIS AREA
	N THIS AREA
	IN THIS AREA
	IN THIS AREA
	E IN THIS AREA
	E IN THIS AREA
	E IN THIS AREA
	/RITE IN THIS AREA
	/RITE IN THIS AREA
	WRITE IN THIS AREA
	FWRITE IN THIS AREA
	FWRITE IN THIS AREA
	OT WRITE IN THIS AREA
	FWRITE IN THIS AREA
	OT WRITE IN THIS AREA

Question 3 continued



Question 3 continued

				Ì		
	1					
			9			
	1					
			Ì		Ę	
					5	
	į					
		ŀ				
				é		
			Ė			
	ï					
	ì					
					1	
				S		
>		>			ς	
	×	>	Ś		S	Ś
>	×	>	8	S	8	

Question 3 continued
(Total for Question 3 is 8 marks)
(



4. (a) Determine the general solution of the differential equation

$$\frac{\mathrm{d}y}{\mathrm{d}x} - 3y\tan x = \mathrm{e}^{4x}\sec^3 x$$

giving your answer in the form y = f(x)

(5)

(b) Determine the particular solution for which y = 4 at x = 0

(2)

×						
X)	K					
×	×	?	Ś		S	
	,	>			ς	
×		×			<	
					ď	
×	×				è	
×		Ì,			2	
×	×				2	
X	×					
×.	×					
X		×			ς	
		\geq			Κ	
×	1	S	4	à	ø	
×	1	Ş	Ä	3	Ŀ	
X.	ì	ĺ.		S	7	
X	ł	E			1	
		Z	S	7	2	
×	J	Ľ	9		5	
X	j	7	7	Z	S	
X	j	ĕ	e		r	
	1	Σ	ľ	7	ķ	
×	×			Ś	2	
X	1	L	à		3	
X		3	2		3	
X	ł	7	2	e	5	
X.	J	7			5	
×	J	ù			ú	
X	j	r	9		Κ	
X	1	P		7	7	۰
×	×				2	
X	Ŋ	ŧ	Þ		ġ	
X	ä			Ľ	_	
X.	i	Ζ	S	Z	3	
×	Š	7	5	7	7	
	J	ř	ú	ř	Ý	
X	1	ŀ		ŀ	d	
×	ĭ	k		5	Z	
×	1	ř	ø	ğ		
\times	ś	í	è	í	è	
ĸŎ.	í	á	ĕ	Ğ	š	
Χ	J	β	j		S	
×	3		í	Ž	Š	
X	1	3			8	
	Ŋ	9	Ŗ	Ş	ĕ	
×	Ŋ	ė	,	5	7	
X	ĭ	í.				
X)	J	þ	ņ	ø	'n	
×	Į	ŝ	ì	è	S	
X	j	r			3	
X	'	9	Ó	ù	ø	
×	ĭ	ņ	ŕ	ij	ø	
X.	Š	d		S	2	
×	d	_	7	7	7	۰
	и	G	à	í	ì	
ĸ.	4	r			7	
×	j	9	ij	è	ø	
X		ä	ė	þ		
		Ŀ	_	4	₫	
×	*		7	5	7	
X.					è	
X	×)	
X	×		>			
	×	1	>		>	
×					5	
×	è					
X					K	
		×				
×	•				ď	
×.					i	
X	×)	
	×		>			
	×		5			
×	×				S	
X						
X		×			<	
		×				
×	×			ς	2	
×						
			S			
×	×		>		3	
X	×	>	>		3	
$\stackrel{>}{>}$	×	>			2	
	×	××××	>		?	
× ×	×	××××	>			
× ×	× × ×	××××	>			
	× × × ×	×××××	>		?	
	× × × × ×	>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>	>>>>>		?	
	× × × × ×		>>>>>>>			
	X X X X X X X X					

Question 4 continued	



Question 4 continued

V			١.
	X COMPANY OF THE SECOND		

Question 4 continued	
(Tot	tal for Question 4 is 7 marks)



5. Given that

$$y\frac{\mathrm{d}^2 y}{\mathrm{d}x^2} + 2\left(\frac{\mathrm{d}y}{\mathrm{d}x}\right)^2 - 2y = 0 \qquad y > 0$$

(a) determine $\frac{d^3y}{dx^3}$ in terms of $\frac{d^2y}{dx^2}$, $\frac{dy}{dx}$ and y

(4)

Given that y = 2 and $\frac{dy}{dx} = 1$ at x = 0

(b) determine a series solution for y in ascending powers of x, up to and including the term in x^3 , giving each coefficient in its simplest form.

(4)

Question 5 continued



Question 5 continued

						C	
	1				ú	à	
		á		ø		5	
		۹				۷	
					7	2	
>							
			ь		i		
						S	
S			e			۲	
0			H		L	à	
			₹		7		
		ì	á			ρ	
				S		١	
				۹	9		
		×		á	ä	k	
	?		Ŀ		ŗ		
				Ķ		4	
		į	ú	i	'n	ú	
		į					
			-		,		
١.	1		1		l.	1	
		Д	۰		ú	۹	
>					7		
					Ų.		
	5	d	â	ė	£	'n	
	c			ä		۲	
						۵	
		7	7	5	7	₹	
\			×	•			
			Š	×	_		
		ì				ú	
7		μ				۹	
					Ų		
	2	ń	ï		ì	à	
	>					ø	
/	3		5			Š	
5	?	ĺ	5	2	ļ	į	
{	3	1		1			
5	?	1	Š	1			
>	?	1					
5	3	1		7 17 17 18			
\ \ \	3	1		2000			
				A MARKET			
< > >				- WWW			
			スシススレビ	A MARKET			
< >>			K X X X X X X				
			て ランス スレスス	THE REAL PROPERTY AND ADDRESS OF THE PERSON NAMED IN COLUMN TWO PERSON NAME			
	3			NAME OF TAXABLE PARTY.			
	?			THE REAL PROPERTY AND PERSONS ASSESSMENT OF THE PERSONS ASSESSMENT OF			
	3						
	ζ	į					
	ζ	į					
	ζ	į					
	ζ	į					
	ζ	į	Š				
	ζ	į	Š				
	ζ	į	Š				
	ζ	į	Š				
	ζ		Š				
	ζ	į	Š				
	ζ	į	Š				
	ζ	į	Š				

Question 5 continued	
	Total for Question 5 is 8 marks)



6.




Figure 1

The curve shown in Figure 1 has polar equation

$$r = 4a(1 + \cos \theta)$$
 $0 \le \theta < \pi$

where a is a positive constant.

The tangent to the curve at the point *A* is parallel to the initial line.

(a) Show that the polar coordinates of A are
$$\left(6a, \frac{\pi}{3}\right)$$

(6)

The point *B* lies on the curve such that angle $AOB = \frac{\pi}{6}$

The finite region R, shown shaded in Figure 1, is bounded by the line AB and the curve.

(b) Use calculus to determine the area of the shaded region R, giving your answer in the form $a^2(n\pi + p\sqrt{3} + q)$, where n, p and q are integers.

(7)

Question 6 continued



Question 6 continued

Question 6 continued
(Total for Question 6 is 13 marks)
(10th 101 Question 0 is 15 marks)



7. (a) Show that the transformation y = xv transforms the equation

$$3\frac{d^2y}{dx^2} - \frac{6}{x}\frac{dy}{dx} + \frac{6y}{x^2} + 3y = x^2 \qquad x \neq 0$$
 (I)

into the equation

$$3\frac{\mathrm{d}^2 v}{\mathrm{d}x^2} + 3v = x \tag{II}$$

(b) Hence obtain the general solution of the differential equation (I), giving your answer in the form y = f(x)

(6)



Question 7 continued



Question 7 continued

Question 7 continued
(Total for Question 7 is 12 marks)



8. (a) Use de Moivre's theorem to show that

$$\sin 5\theta \equiv 16\sin^5\theta - 20\sin^3\theta + 5\sin\theta$$

(5)

(b) Hence determine the five distinct solutions of the equation

$$16x^5 - 20x^3 + 5x + \frac{1}{5} = 0$$

giving your answers to 3 decimal places.

(5)

(c) Use the identity given in part (a) to show that

$$\int_0^{\frac{\pi}{4}} (4\sin^5\theta - 5\sin^3\theta - 6\sin\theta) d\theta = a\sqrt{2} + b$$

where a and b are rational numbers to be determined.

(4)

Question 8 continued



Question 8 continued

					/	
0						
	×				>	
		?				
		5				
	۲.	à	ú		ρ	
	21		z		12	
7	v	7	7	7	۹	
١	7	ь	6	ò	4	
	М	L	3	Ľ	3	
	O	9	۰	7	e	
		à	ù	ú	à	
	7				۳	
	×	Z	S	Z	Σ	
S	₹	ŕ	Ì	ŕ	Ş	
8	Ŕ	į	Ì	ľ		
>	Ì	į		Ź		
>	Ŕ					
>	3					
>	3					
>						
>						
>						
>>>>						
> > >						
> > > >						֡
						֡
> > > >						֡
> > > >						֡
						֡
> > >						֡
						֡
						֡
						֡
						֡
						֡
						֡

$^{\circ}$				>	
х			>		
\odot		2	ú		
$^{\times}$	ij	₹			
V	٦	7			
0	4	7		7	
			ь	a	
Χ	4	s.	2	ú	
V		71		7	
		₹	2	Z	
	ú	à		P	
х		Б		ß.	
V		7	₹	9	
\wedge				١	
	2		à	ĸ	
		М.	r	3	
Х	3	ę		9	
Ü	ы	mi.	ы	×	
\wedge		Z		Z	
V	ij	ч		,	
		а	г		
Х	ù	#	۰	۰	
				2	
$^{\wedge}$					
		7			
	Š	>		>	
×	n	6	à	6	
V2		7		P	
Л	4		z	2	
×	7	₹	3	7	
\circ	₹	Σ	☇	>	
X	ĺ	Ì	í	à	
8	ĺ	Ì	Ĺ	ì	
8	2	3	į	1	
× ×	ĺ	2		2	
× × ×	ĺ	}		2	
× ×		2			
× × ×		2			
		\ \ \			
× × × ×		\ \ \ \			
X	ļ	2			
X		\ \ \ \ \ \ \			
X		\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \			
X	ļ				
X					
X					
X		/			
X		/			
X		/			
X		/			
X		/			
X		/			
X		/			
X		/			
X		/			
X		/			
X		/			
X		/			
X		/			
X		/			
X		/			
X		/			
X		/			

		×		×	
		V	×	v	
	\triangle			4	
	х		X.	х	
	\vee	$^{\sim}$	\sim	\sim	
	×	Х	×	X	
1	V	V	\sim	V	
	\triangle				
	×	×	.×	×	
	V	V	\sim	~	
		a	ø	r.	
		ж	×	EX.	
		S	1/2	2	
		4	ИIII:	dii:	
	×	ж	100	瞑	
	V	S	in.	w	
		41	<u> </u>	\sim	
		×	-	75.	
	v	S.	-	ia.	
		4		n	
	х	×	.75	90	
	V	V	\sim	V	
		Ř	v/a	mi.	
	X	ж	ur.	ж	
	V	V	\leq	2	
	$\langle \rangle$			₹.	
	Х	28	o de la compa	μK.	
	\vee	\sim	W	V	
		< II	-	100)	
		Жi	ıX	Х	
	V	M	-	m,	
		< ₩			
	Х	X	X	X	
	V	V	- 22	×	
		< 2	e	20	
	Х	79		7	
	v	78	-	×	
>				O	
		4	νŵ	套	
		М	138	æ	
		S		∇	
		4	\sim	◬	
	×			7	
	\bigcirc	ς.	2	_	
		7	Z	Zì.	
		М	71	•	
		SII		=	
		Δì		\sim	0
	×	×	-	38.	
	V	À	9	~	
		C 16		æ.	
	х	×	X	X	
	V	M	ſv	V	
		(II	m	100	
Κ	×	24	LX.	X	
	V	Na	m	w	
		a	(400	
	Х	×		ø.	
	V	M	_	ì	
			-	•	
	Χ	IK,	6	mi.	
	V	V		Ÿ	
				0	
	Х	a	1	š	
	×	М	\sim	æ	
			-	5	
	Χ	Ř	m	NS.	2
	X	M	\sim	w	
	$\langle \rangle$	Ç.	-	Ψ.	
	\wedge	Х	$^{\prime}$	\wedge	

Question 8 continued						



Question 8 continued					
	(Total for Question 8 is 14 marks)				
	TOTAL FOR PAPER IS 75 MARKS				

