Vrite your name here Surname	Other na	mes
Pearson Edexcel nternational Advanced Level	Centre Number	Candidate Number
<b>Core Math</b>	amatic	c(1)
		3 C 1 Z
Advanced Subsidia	r <b>y</b>	
	r <b>y</b> - Morning	Paper Reference WMA01/01

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

## Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
   there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

## Information

- The total mark for this paper is 125.
- The marks for each question are shown in brackets
   use this as a quide as to how much time to spend on each question.

## **Advice**

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

PE

Turn over ▶

PEARSON

Find $\int f(x) dx$ .	simplifying each term.	
1 J 1 (11) a,		(!

⋖
ш
0
7
-
S
I
Z
-
THE PERSON NAMED IN
1
œ
>
$\geq$
100
$\simeq$
0

Question 1 continued	Leave blank
Question I continued	
	0.1
	Q1
(Total 5 marks)	



2.	Find, giving your an	swer to 3 significant	t figures where	appropriate, the	value of x for
	which				

(a)	$7^{2x}$	=	14

**(3)** 

(b) 
$$\log_5(3x+1) = -2$$

**(2)** 

-
•
-73
ш
-
$\mathbf{r}$
-
Z,
11.00
S
-
- TIL -
-
-
-
7.7
Trief.
쁜
11.7
1
-
40.00
$\leq$
-
400
100
100
$\sim$
$\tilde{-}$
1.1.1
-
. = .
$\sim$
_

K)	ς	I	2
1	6	7	3,
4	Ц	И	ŀ
1	4	↘	i
4		2	ř
Ÿ.	4	'n	i,
4	É	ķ	ì
ĸ,	/		
4	V V	Я	r
V	$\underline{}$	$\overline{z}$	7
	Σ	Z	7
ĸ.	7	ҡ	-
л	ø	×	Ď,
٠1	12	_	٠.
Ж	г	7	٣.
	$\geq$	C	
Ä	7	7	ĸ
14	Ę	ř	ď
V	Ź	è	v.
			_
SI	Ú	N	ŀ
		7	
ķ	۳	۳	ŀ
4	-	ź	r
N	ń	ä	g,
4	щ	ы	ì
Ó	4	$\leq$	٥
X	e	÷	ч,
Κű			
	2	э	ľ,
X	7	þ	ŗ,
G	K	2	Š
â	X		
Ĵ			
G			
Ĵ			
Ĵ			
Ĵ			
Ĵ			

Question 2 continued	Leave
	02
	<b>Q2</b>
(Total 5 marks)	



**(3)** 

3.	Answer this question without the	e use of a calculator and show your method clearly.
	(i) Show that	20

 $\sqrt{45} - \frac{20}{\sqrt{5}} + \sqrt{6}\sqrt{30} = 5\sqrt{5} \tag{2}$ 

(ii) Show that  $\frac{17\sqrt{2}}{\sqrt{2}+6} = 3\sqrt{2} - 1$ 

ì	ė	۲.
1		ч
1	Ш	ш
3	_	5
1	▢	E
1	7	Ξ
ì	e	r
		7
'n	r i	m.
1	u	1
ì	÷	
3	4	-
ì	4	÷
ì	Ľ	_
ł	г	_
ì	÷	÷
	d	~
4	-	ᆕ
4	-	_
ì	ΡĖ	٠ú
ļ	ш	4
	Ŀ	÷
ł	Γ.	
ď	_	=
1	▢	⊆
ũ	_	Ξ
î	2	7
ï	3	⋑
ũ		
1	H	_
3	×	ĸ.
J	С	3)
	7	_
3	2	~
1	6	-
	۷	٠.
Ì	۳	5
٦	Ŀ	ø
1	p	ď
1	Ŀ	4

Question 3 continued		Leave blank
		Q3
	(Total 5 marks)	



4.	$f(x) = 6x^3 - 7x^2 - 43x + 30$	
	(a) Find the remainder when $f(x)$ is divided by	
	(i) $2x + 1$	
	(ii) $x-3$	
		(4)
	(b) Hence factorise $f(x)$ completely.	(4)



	Leave
	blank
Question 4 continued	
Question Feoremacu	



estion 4 continued	

Leave

ì	ė	۲.
1		ч
1	Ш	ш
3	_	5
1	▢	E
1	7	Ξ
ì	e	r
		7
'n	r i	m.
1	u	1
ì	÷	
3	4	-
ì	4	÷
ì	Ľ	_
ł	г	_
ì	÷	÷
	d	~
4	-	ᆕ
4	-	_
ì	ΡĖ	٠ú
ļ	ш	4
	Ŀ	÷
ł	Γ.	
ď	_	=
1	▢	⊆
ũ	_	Ξ
î	2	7
ï	3	⋑
ũ		
1	H	_
3	×	ĸ.
J	С	3)
	7	_
3	2	~
1	6	-
	۷	٠.
Ì	۳	5
٦	Ŀ	ø
1	p	ď
1	Ŀ	4

Out of the American I	blank
Question 4 continued	
	Q4
(Total 8 marks)	



5. (a) Find the first 4 terms, in ascending powers of $x$ , of the binomial expansion of
$\left(3-\frac{ax}{2}\right)^5$
where $a$ is a positive constant. Give each term in its simplest form. (4)
Given that, in the expansion, the coefficient of $x$ is equal to the coefficient of $x^3$ ,
(b) find the exact value of a in its simplest form. (3)

ì	فح	۴
7	7	G,
1	ш	Ы
Ġ	_	Ξ
1	▢	Ė
	_	Ξ
J	ς	Ľ
		7
i	ri	ø.
1	υ	ч
þ		=
3	4	7
j	4	Í
ł	Ľ	_
1	г	_
ì	÷	_
1	2	~
1	•	₹
4	-	7
ì	ń	٠ú
ł	ч	ш
į	Ŀ	-
3	<u>.                                    </u>	
	_	Ξ
1		⊆
þ	-	
4	e	-
J	ē	7
á		
1	H	=
-	'n	٩,
Л	Ŀ	٤
4	~	_
Ĵ	d	-
Ţ	7	=
	'n	Š.
4		7)
	3	٣.
4	۳	٦
1	٠	4

Question 5 continued	Leave blank
(Total 7 marks)	Q5



A sequence is defined by

$$u_1 = 36$$

$$u_{n+1} = \frac{2}{3}u_n, \qquad n \geqslant 1$$

(a) Find the exact simplified values of  $u_2$ ,  $u_3$  and  $u_4$ 

**(2)** 

(b) Write down the common ratio of the sequence.

- **(1)**
- (c) Find, giving your answer to 4 significant figures, the value of  $u_{11}$
- **(2)**

(d) Find the exact value of  $\sum_{i=1}^{6} u_i$ 

**(2)** 

(e) Find the value of  $\sum_{i=1}^{\infty} u_i$ 





Question 6 continued		Leave
Question 6 continued		blank
	Question 6 continued	



estion 6 continued	

Leave

ì	ė	۲.
1		ч
1	Ш	ш
3	_	5
1	▢	E
1	7	Ξ
ì	e	r
		7
'n	r i	m.
1	u	1
ì	÷	
3	4	-
ì	4	÷
ì	Ľ	_
ł	г	_
ì	÷	÷
	d	~
4	-	ᆕ
4	-	_
ì	ΡĖ	٠ú
ļ	ш	4
	Ŀ	÷
ł	Γ.	
ď	_	=
1	▢	⊆
ũ	_	Ξ
î	2	7
ï	3	⋑
ũ		
1	H	_
3	×	ĸ.
J	С	3)
	7	_
3	2	~
1	6	-
	۷	٠.
Ì	۳	5
٦	Ŀ	ø
1	p	ď
1	Ŀ	4

Question 6 continued	blank
	Q6
(Tot	ral 9 marks)



7. (a) Sketch the graph of  $y = 3^{x-2}$ ,  $x \in \mathbb{R}$  Give the exact values for the coordinates of the point where your graph crosses the *y*-axis.

**(2)** 

The table below gives corresponding values of x and y, for  $y = 3^{x-2}$ . The values of y are rounded to 3 decimal places where necessary.

X	0.5	1	1.5	2	2.5	3
у	0.192	0.333	0.577	1	1.732	3

(b) Use the trapezium rule with all the values of y from the table to find an approximate value for

$$\int_{0.5}^{3} 3^{x-2} \, \mathrm{d}x$$

Give your answer to 2 decimal places.

**(4)** 

ET.	
Œ	
-tun	
$\sim$	
1	
0.01	
144	
-	
100	
100	
10.00	
100	
100	
100	
RIT	
/RIT	
RIT	
/RIT	
NRIT	
NRIT	
NRIT	
WRIT	
NRIT	
WRIT	
T WRIT	
T WRIT	
WRIT	
T WRIT	
T WRIT	
T WRIT	
<b>JOT WRIT</b>	
<b>JOT WRIT</b>	
T WRIT	
<b>JOT WRIT</b>	
<b>JOT WRIT</b>	
NOT WRIT	
NOT WRIT	
<b>JOT WRIT</b>	
NOT WRIT	
NOT WRIT	
NOT WRIT	
DO NOT WRIT	
OO NOT WRIT	
DO NOT WRIT	

Question 7 continued	Leave
Question / continued	_
	_
	_
	_
	_
	_
	-
	_
	-
	_
	-
	_
	_
	_
	_
	_
	_
	_
	_
	-
	_
	-
	_
	_
	_
	_
	_
	_
	<b>Q7</b>
(Total 6 marks	s)



8.




Figure 1

The compound shape ABCDA, shown in Figure 1, consists of a triangle ABD joined along its edge BD to a sector DBC of a circle with centre B and radius 6cm. The points A, B and C lie on a straight line with AB = 5 cm and BC = 6 cm. Angle DAB = 1.1 radians.

(a) Show that angle ABD = 1.20 radians to 3 significant figures.

**(4)** 

(b) Find the area of the compound shape, giving your answer to 3 significant figures.

**(4)** 

Question 8 continued	bla
Question 8 continued	
	I
	—
	—
	—
	—
	—
	—
	—
	_
	—



ì	ė	۴
7	7	Ni.
1	H	٠II
1	_	5
1	0	=
4	-	Ξ
ì	e	r
		,
'n	r i	ď.
1	u	7))
'n	-	i
ì	4	
ì	4	÷
	C	Ξ
ł	_	-
		۰
	2	~
4	F	-
4	-	=
ì	. :	á.
1	ш	4
	Ŀ	Ξ
ļ	т	7
4	-	
1	n	m
	Ħ	
Ą	3	×
1	5	Si
J	~	_
1	Ĺ	_
4	7	7
1	m	٩
4	١.	J
9		ė
1	2	_
÷		7
٠,	'n	iκ
4	Ŀ	2)
	Z	Ξ.
1	r	٦
J	=	4

	Le	ave ank
Question 8 continued		
	Q8	
(Total 8 m	narks)	<u> </u>



In a large theatre there are 20 rows of seats. The number of seats in the first row is a, where a is a constant. In the second row the number of seats is (a + d), where d is a constant. In the third row the number of seats is (a + 2d), and on each subsequent row there are d more seats than on the previous row. The number of seats in each row forms an arithmetic sequence. The total number of seats in the first 10 rows is 395 (a) Use this information to show that 10a + 45d = 395**(1)** The **total** number of seats in the first 18 rows is 927 (b) Use this information to write down a second simplified equation relating a and d. (c) Solve these equations to find the value of a and the value of d. **(3)** (d) Find the number of seats in the 20th row of the theatre. **(2)** 



estion 9 continued	



estion 9 continued	

-	ø	
VUQV	Ŀ	
-12	7	
ы	ш	
포	=	
n	_	
=	-	
ŵ	p	
.94	ы	
	m.	
ν	0	
=	-i	
=	=	
л		
_	=	
ы	-	
т	_	
2	۰	
-	,	
~	-	
Ξ	Ξ	
=	_	
20	٠ú	
ш	4	
10		
45	-	
÷	÷	
-	ä	
ш	$\subseteq$	
THO	_	
147	3	
-	5	
and the	-	
d .		
н		
HON	ď	
r	э	
-Van	o	
=	ė	
20	۳.	
_	=	
100		
$\sim$	ч	
غة	s	
- 2	Ξ.	
	٦	
=	4	

Question 9 continued	Leave blank
Q	9
(Total 8 marks)	



10. (a) Given that  $8\tan x = -3\cos x$ show that  $3\sin^2 x - 8\sin x - 3 = 0$ **(3)** (b) Hence solve, for  $0 \le \theta < 360^{\circ}$ ,  $8\tan 2\theta = -3\cos 2\theta$ giving your answers to one decimal place. (Solutions based entirely on graphical or numerical methods are not acceptable.) **(5)** 

	Leave blank
Question 10 continued	
	Q10
	Q10
(Total 8 marks)	



(a) Show that $k$ satisfies the inequality	
$6k^2 + 13k - 5 > 0$	
(b) Find the set of possible values for <i>k</i> .	

ì	ė	۴
7	7	Ni.
1	H	٠II
1	_	5
1	0	=
4	-	Ξ
ì	e	r
		,
'n	r i	ď.
1	u	7))
'n	-	i
ì	4	-
ì	4	÷
	C	Ξ
ł	_	-
		۰
	2	~
4	F	-
4	-	=
ì	. :	á.
1	ш	4
	Ŀ	Ξ
ļ	т	7
4	-	_
1	n	m
	Ħ	
Ą	3	×
1	5	Si
J	~	_
1	Ĺ	_
4	7	7
1	m	٩
4	١.	J
9		ė
1	2	_
÷		7
٠,	'n	iκ
4	Ŀ	2)
	Z	Ξ.
1	r	٦
J	=	4

Question 11 continued	Leave blank
	Q11
(Total 8 marks)	



**(4)** 

**(1)** 

12.




Diagram not drawn to scale

Figure 2

Figure 2 shows a sketch of the curve with equation y = f(x), where

$$f(x) = \frac{x^3 - 9x^2 - 81x}{27}$$

The curve crosses the x-axis at the point A, the point B and the origin O. The curve has a maximum turning point at C and a minimum turning point at D.

- (a) Use algebra to find exact values for the x coordinates of the points A and B.
- (b) Use calculus to find the coordinates of the points C and D. (6)

The graph of y = f(x + a), where a is a constant, has its minimum turning point on the y-axis.

(c) Write down the value of a.

uestion 12 continued	L   t



	Leave
Question 12 continued	blank
Question 12 continuou	

Q.
11.1
-
$\alpha$
72
10
S
$\equiv$
- TIL -
壶
-
-
=
7.7
TITL
쁜
100
-
2
2
$\geq$
100
-
11
0
$\sim$

Question 12 continued	Leave blank
	Q12
(Total 11 marks)	



<b>13.</b> The circle C has centre $A(1, -3)$ and passes through the point $P(8, -2)$ .	
(a) Find an equation for the circle <i>C</i> .	
	(4)
The line $l_1$ is the tangent to $C$ at the point $P$ .	
(b) Find an equation for $l_1$ , giving your answer in the form $y = mx + c$	(4)
The line $l_2$ , with equation $y = x + 6$ , is the tangent to $C$ at the point $Q$ .	(4)
(c) Find the coordinates of the point Q.	
	(5)

	blank
Question 13 continued	



estion 13 continued	

DO NOT WRITE IN THIS AREA						
DO NOT WRITE IN THIS ARE	1	ė	Р		•	
DO NOT WRITE IN THIS AR	7	7	٩	٩	ė	i
DO NOT WRITE IN THIS AR	1	Ŀ		Ŀ		
DO NOT WRITE IN THIS AI	С	Ξ	7	Ξ	3	
DO NOT WRITE IN THIS	1	Ē		٤		
DO NOT WRITE IN THIS				3	3	
DO NOT WRITE IN THIS	J	ξ	S			
DO NOT WRITE IN THE				7	7	١
DO NOT WRITE IN THE	ì	r	À	p	i	
DO NOT WRITE IN TH		b	ø		Į	١
DO NOT WRITE IN TH			Ė		Ė	ı
DO NOT WRITE IN T	3	Ŧ		P		
DO NOT WRITE IN 1	ì			i	i	i
DO NOT WRITE IN	ì	L	_	i		
DO NOT WRITE IN	1	Г	ī		ī	
DO NOT WRITE IN	ì	è	Ė	è	í	ì
DO NOT WRITE!	í	d	è	۹		
DO NOT WRITE	c	7				
DO NOT WRITE	4		7			۱
DO NOT WRITE	ì	r	i		í	i
DO NOT WRIT		b	ä	į	å	ı
DO NOT WRI	1	k	ė		i	
DO NOT W	3	L	_			
DO NOT W	6	Ξ		3	Ξ	
DO NOT W	1			ľ		
DO NOT	þ	6		i	ı	
DO NOT	4	E	E	Š	3	
DO NO	á	á	ē	P	ę	
DO NO						
DO NO	1	ŀ	ė		į	ı
DO NO	ā	à	ø	ġ	6	
00	J	Ŀ		i	J	١
00	4	2	Ξ	ï		
00	ú	á	ė	9	1	
۵	ļ					
۵	٠,	à	a	ò		
	4	ľ		1		١
		3		2	(	
	1	ľ	Ī	٩		١
	å		7			

	Leave blank
Question 13 continued	
	Q13
(Total 13 marks)	
(10th 10 mains)	



Leave blank

14.




Figure 3

Figure 3 shows a sketch of the curve C with equation  $y = -x^2 + 6x - 8$ . The normal to C at the point P(5, -3) is the line l, which is also shown in Figure 3.

(a) Find an equation for l, giving your answer in the form ax + by + c = 0, where a, b and c are integers.

**(5)** 

The finite region R, shown shaded in Figure 3, is bounded below by the line l and the curve C, and is bounded above by the x-axis.

(b) Find the exact value of the area of R.

**(6)** 

(Solutions based entirely on graphical or numerical methods are not acceptable.)

	Leave blank
Question 14 continued	



estion 14 continued	

j	É	e		1	
	7	7	٩	٩	
1	Ŀ		Ŀ		
ű	ä	Ξ	ž	ī	
]	E		c		
	1		ä	d	
j	ξ	ξ			
				1	
i	r	ì	p	i	
	٩	P		4	
9		۹		۱	
3	-		۳		
ì		å	b	ĺ	
1	Ŀ	i	ė		
1	г				
ì	ė	ė	ė	i	
Ĵ	d	è	2	_	
c	Ξ		Ξ		
1	7	7	-		
ì	r	i		í	
1	b	d	b	å	
1	L	i		i	
1	Ę	_		2	
1	3	=	3	3	
1	Ľ		c		
j	6			Ī	
j	s	Ē	2	۹	
ī	ä	ã	þ	þ	
Ĩ,					
l	ŀ	ė		į	
ċ	à	ø	ė		
J	Ľ		Ē		
	2	3	3	_	
Ġ	ä	ø	P	1	
1	•	-			
į,	à	•	ò		
4	ľ	į	ĺ		
	7		S	(	
1	ľ	Ī	4	١	
J		Ť		į	

4	Ц	И	
V	ń	'n	v
Ŋ	Ц	К	ŀ
٧.	4	ы	
А	ς	ľ	
ĸ.	2	7	
A	Ú	e	
. 1	μ	Ų	١,
×	÷	÷	ĸ,
(4	Ì	É	١)
X	4		К
1	D		
k	۳	9	ť
	`	ζ.	. `
V	9		ν
G		2	
V	4	$\succeq$	ν
r	X.	×	1
V)	И	N	v
×	ĸ.	7	^
G	Þ	Z	Ò
j	P	2	
j	ŀ	2	
į K	ŀ	2	
į		2	
į K		2 2 2	
	A COAM	2	
Ķ		2	

		2				2		
$^{\times}$	2							
×	)				5		)	
						)		
×	2	1		۲,		٦		۲,
×							7	
0	Κ	>		Þ		)		×
		1				١		
×							>	
	Κ	2		7		2		7
$^{\prime}$		/		\		١		
×					2	۷	ð	
C	Κ	2	ы			7		7
$\triangle$		7	7	۹		ь		7
$\times$			2	ζ,	Þ	ζ		
V	Ν,	2		а	L	а	Ь	Α
	K.		۳	9	e	9	ч	$\geq$
$\times$			à	a	à			
V	1			2	Ľ	2		Ν
			7	7	7		ľ	
×	)	Κ	2	ú		ρ	Ð	≺
V	1		۹	G		Ľ		
				×	₹	7	К	$\geq$
$^{\times}$	J		?	ς,	2	2	2	۲.
V	`			a				
	K		궦	ø	r	7	ľ	
×	2		2	2	2	Ω	2	
20	`\		3	7	3	7	٠,	
			ф			۰	К	
×	2	۲		Я	Ρ	ς	2	ς.
1	١,		щ	•	9	۰	Ŋ	
					٤			
$^{\sim}$	1	5		₹	,	7	y	
×			5		>			
	Κ		£	4	Ĺ	3	٤	×
$^{\circ}$	/	1	7	Σ		P		١,
×					S	2	ò	
C	К	>	Σ	Z	Ξ	2		×
			e	٠	۲	5	۲	
$\times$								
				А				
X	2	S						◁
Š	3	>	Š		5	į	Ŗ	S
8	ξ	S	ũ	2		Š	R	S
Š	2	S	ũ	2		S	Ŗ	ξ
8	3	S	ũ	2		2	8	8
8	3	S	ũ	2		2	2	5
8	3	S	ũ	2			P	5
8	3	S	ũ	2			R	<<<<<<<<<<.
	?	S	ũ	2				
		S	ũ	2				
		S	ũ					
		S	ũ	2				
		S	ũ	2				
		S	ũ	2				
		S	ũ	2				
		S	ũ	2				
		S	ũ	2				
		S	ũ	2				
		S	ũ	2				
		S	ũ	2				
		S	ũ	2				
				2				
				2				
				2				
				2				
		S		2				
				2				
				2				
				2				
				2				
				2				

Overtion 14 continued	Leave blank
Question 14 continued	
	Q14
(Total 11 marks)	



Leave blank

**(2)** 

**15.** 




Figure 4

Figure 4 shows a solid wooden block. The block is a right prism with length hcm. The cross-section of the block is a semi-circle with radius rcm.

The total surface area of the block, including the curved surface, the two semi-circular ends and the rectangular base, is  $200\,\mathrm{cm}^2$ 

(a) Show that the volume  $V \text{cm}^3$  of the block is given by

$$V = \frac{\pi r (200 - \pi r^2)}{4 + 2\pi} \tag{5}$$

- (b) Use calculus to find the maximum value of V. Give your answer to the nearest cm<sup>3</sup>. (6)
- (c) Justify, by further differentiation, that the value of V that you have found is a maximum.

estion 15 continued	



estion 15 continued	

estion 15 continued	



uestion 15 continued	blank
	Q15
	13 marks)
TOTAL FOR PAPER: 125	5 MARKS