Surname	Other na	ames
Pearson Edexcel GCE	Centre Number	Candidate Number
Further F	<i>r</i> ure	
Mathema Advanced/Advan	atics FP3	
Mathema	atics FP3 ced Subsidiary Morning	Paper Reference 6669/01

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information

- The total mark for this paper is 75.
- The marks for each question are shown in brackets
 use this as a quide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶



P55815A
©2018 Pearson Education Ltd.



1. (a) Starting from the definitions of $\sinh x$ and $\cosh x$ in terms of exponentials, show that, for $x \in \mathbb{R}$

$$\tanh x = \frac{e^{2x} - 1}{e^{2x} + 1} \tag{2}$$

(b) Hence, given that $-1 < \theta < 1$, prove that

$$\operatorname{artanh} \theta = \frac{1}{2} \ln \left(\frac{1+\theta}{1-\theta} \right)$$
 (3)

Question 1 continued	Leave
Vaccation a continuou	
	Q1
(Total 5 marks)	
(10tal 5 mai ks)	



2.




Figure 1

Figure 1 shows a sketch of part of the curve with equation

$$y = 5 \cosh x - 6 \sinh x$$

The curve crosses the x-axis at the point A.

(a) Find the exact value of the *x* coordinate of the point *A*, giving your answer as a natural logarithm.

(3)

(b) Show that

$$(5\cosh x - 6\sinh x)^2 \equiv a\cosh 2x + b\sinh 2x + c$$

where a, b and c are constants to be found.

(3)

The finite region R, bounded by the curve and the coordinate axes, is shown shaded in Figure 1.

The region *R* is rotated through 2π radians about the *x*-axis.

(c) Use calculus to find the volume of the solid generated, giving your answer as an exact multiple of π .

(4)

destion 2 continued		Lo bl
	Duestion 2 continued	DI
	zuestion 2 continued	



nestion 2 continued	

۰					a
	ì	É	ė		7
۰	Ĺ	3	7	3	9
	1	Ŀ		Ŀ	ı
	-	à		à	ø
	J	Ŀ	d	Ŀ	
	ì	ė	ø	ġ	ø
	ď	7	۰	ě	ų
				Ĺ	
	1	L	s	ŗ)
	ì	i	ì		ã
	٦	٠	ė	P	ú
	i			L	ė
	1	L	_		_
	1	г	7		_
٠	j	4	ė	4	i
	ì	d		2	_
۰	į	Ξ	=	Ξ	=
	-	Ľ	J	ľ	ı
	å	7		7	7
	ł	P	Ŧ		9
	á	ì		Ė	
	1	ŕ	٩	è	ø
	j	Ė	4	٠	
	í	3		à	þ
	1	3	5	3	ń
	4	7	7		
	1	ŀ	ė		ė
	1	5	ė	ı	Ĵ
	1	ľ		3	3
	į	3	3	2	_
	ľ	2	7	7	7
	j	8	7		ţ
	ŧ,	à	ø	ė	k
	4	ľ	í	j)
		3		S	į
	1	ľ		1	١
	ł		Ť		ė

	-	4	
`\	ĸ.	-6	~
Ŋ	ы	и	
	- 7		
×	п	m	Υ
X]	۰,		
1	L	20	v
И		JP	
٠.	3	_	
×	Χ.	\times	
	B.	m	
. 4	u	КJI	١,
N	_		C
Α,	\sim	\sim	
Χ,	Z	⋉	
2		ы	ĺ.
1	в)		
×		÷	v
Л	٩.	/	
	_2	<u> </u>	
N	ź	7	r
×,		~	
× 1	ž	Ś	
Z	$\overline{}$		
	-)	6	
N	и		V
Š	è	7	1
. 1	ш	_	
N	г		
4	÷	-	r
Χ.	d	شذ	×
2	С	12	
	_		۲.
N	-		/
		z	7
Ø,	-2	3	r,
N		\sim	
	ıΝ	<u></u>	
X	×	₩	
2	Sá	-2	
1	п	a	n
×.	٧.	-28	
×.	7.	\sim	
v	\overline{z}	7	٢,
N		\sim	C
	\sim	7.	
×	6	×	×
-21	ĸ,	S	Ĺ
	S.	ω	r
×	z	≤.	×
	r	78	

V	V	V	V	1
		\triangle	_	C
V	V	麗/		18
		-56	K.)	ш
×	×	×	×	
		œ		к
-><	×	x	Ř	_>
		X	94	к
				Ь
		er:	т.	
×	×	BH.		E)
		c		
X	X	79	46	v
		w	mβ	К
\wedge	\wedge	79	•	
V	V	-	×	ь,
\wedge	$^{\wedge}$	~	\sim	/
		æ	\sim	
×	×	w	-0	в,
		ΨÒ,	0	
×		28	-	6.5
		ec.	O	К
×	×	16.	-34	7>
		9	-	
X	х	ii.	-	ıν
52		14	20	Κ.
\times	х	a	Z	2
		7	_	ь,
\wedge		\wedge)
V	\vee	Vá	-	`
\wedge	$^{\wedge}$	æ	79	v
V	V	W/	V	B.
		ethi	للت	ĸ.
×	×	×	⇉	
		ø	-9	К
×		IK.	N	ľ
		96	_	В.
X	X	\times	\times	->

Question 2 continued	Leave blank
	Q2
(Total 10 marks)	



 $\mathbf{M} = \begin{pmatrix} 3 & k & 2 \\ -1 & 0 & 1 \\ 1 & k & 1 \end{pmatrix}, \text{ where } k \text{ is a constant}$ 3.

Given that 3 is an eigenvalue of M,

- (a) find the value of k.
- **(3)**
- (b) Hence find the other two eigenvalues of M. **(4)**
- (c) Find an eigenvector corresponding to the eigenvalue 3 **(2)**

	Leave
	blank
Question 3 continued	



estion 3 continued		

	Leave blank
Question 3 continued	
	Q3
(Total 9 marks)	



4. The curve C has equation

$$y = \operatorname{arsinh} x + x\sqrt{x^2 + 1}$$
, $0 \le x \le 1$

(a) Show that $\frac{dy}{dx} = 2\sqrt{x^2 + 1}$

(4)

(b) Hence show that the length of the curve C is given by

$$\int_0^1 \sqrt{4x^2 + 5} \, \mathrm{d}x \tag{2}$$

(c) Using the substitution $x = \frac{\sqrt{5}}{2} \sinh u$, find the exact length of the curve C, giving your answer in the form $a + b \ln c$, where a, b and c are constants to be found.

(6)

	Leave
	blank
Question 4 continued	
American i commune	



estion 4 continued	

		s	d	ď	
111	ξ	S		Ξ	
	i		7	3	
4	Ŀ		Ŀ	a	
1	å		å	ď	
J	E	3	E		
		Ξ	3	ā	
1		S		7	
		7	7	٩	
	r	ì	á	ć	
1	ŀ	ø)	
ï	i	ė		ė	
ì	ŧ	ĕ	p	ú	
i			L	÷	
ń	ī	7		7	
4	P		•		
÷	۰				
1 4 4	Ė	3		P	
4			í		
á	i		ė		
J	ľ	ì		ı	
3	ì	٩		ą	
ı	k				
3	L	_	_	i	
7	3	7	3	2	
1	ľ		c	Ξ	
1				-	
c	3		þ	þ	
1111111111	2			ń	
3	7	7	1		
1	L		_	٠	
1	г	7	7	7	
1	p	q	۹	b	
4	ķ		ė	p	
9	í	Ē	=		
ú	å	ë	9	5	
÷		7		7	
	ż		ċ		
j	ľ	1	7	Ñ	
1	٩	ú	ø	ď	
1	þ	P	۹	ń	
1	L	ė		a	

		ľ				
×		Ŀ		4		
		2		₹	3	
	м	Ć		Р	5	
×		ь		H	i	
		2				
	×		я			
×	3	۹	Ħ		ì	
ς.					•	
				_		
×		٨		F		r
	1		и			
	×	=		a	í	
×	E	3	Ξ			
ς.	3	7		e	۰	
	×	_		2	١	
×	8	₹	=	,	۰	
		Ŀ		_		
		۰	۰	•	9	
×	8	٩				
	٠.		⋍	_		
	ø	7			,	
×	d			z		
	•	7	7	₹		
0	ø	•	۰	۰		
×		٩				
	٦	v	ň			
0		G		2		
×	٠	π	_	×	7	
		Ŀ	×		ė	
		г		7		
r	į	K		4		
ς.		=		=	ī	
	М	r		۳	5	
/	٠	н		ы	ı	
	į	ú	2			۲.
	ч			Z		
	٠,	ė			ı	١.
	ø	7			5	
/	i	К				
V			ė		i	i.
	ą	١.			۶	
/	à	и	H	۹	١	
V		D		v		١.
	×	٩	H		P	
×	ú	\sim	_	۷		
\vee	•				,	١.
	м				à	
	Ġ	7	5	7		
\vee		Z	8			١.
	×á	ρ	,	۹	١	
		ß				
V	3	ø	ú	ø	,	١,
	×	a		ы	b	
	. (К		7		

\Diamond	0			۲.
$^{\wedge}$	$^{\wedge}$	$^{\wedge}$	$^{\wedge}$	$^{\prime}$
V	V	V	V	V
\times	×	×	×	×
\vee	\vee			∿
\triangle	Λ			◠
×	X	×	×	X
X	X	X.	×	×
1	V	w	v	۲.
\triangle	$^{\wedge}$	_	ď.	↶
×	×	×		3×
		M X	п	Ľ
×	X			ľ×
V	\sim	\overline{z}	\subseteq	5.
		n	т	ţ٩
×	V	ш	84	N
				Χ.
X	Х	×		5×
V	\vee		ШZ	
	$^{\wedge}$	//5	~	r
×	×	×	×	×
		60	<u></u>	Κ.
X	X	WS.	м	v
V	V	쪵	v	ĸ,
\triangle	Λ	≏	_	r
×	X	$\overline{\mathbf{x}}$	×	×
		=	,	ĸ.
X	X	XI	K	Х
1	V			N
\triangle	\triangle	al/		$^{\prime}$
×	×			ì×
		W >		ς.
\mathcal{L}	Х	$^{\sim}$	А	Х
\vee	\sim		_	N
		\sim	o	
×	×		24	ıΧ
	\sim			5.,
\wedge	$^{\wedge}$	_	_	m
V	V	V	V	V
	$\langle \cdot \rangle$	6)	ല	c
×	×	IKJ		IX
0	$\langle \rangle$	₩	₩	٩.,
$^{\wedge}$	$^{\wedge}$	В\	$^{\wedge}$	↗
V	V			N
		Ш>		
-><	×			ŧΧ
52	90	$\overline{}$	$\overline{}$	ξ.
\triangle	Λ	m	м	20
×	×	ы	-	iΣ
		₽		
X	X	79	ж.	×
V	V	m#	₹	٦,
				r
	X	岬	×	\times
50	$\langle \rangle$			ς:
\wedge	$^{\sim}$	Вĺ\		
V	v			Ŋ
		'n.	\triangle	
×	×	20	70	
0	O	m)	a	ь.
\triangle	$^{\wedge}$	ж	-ii	'n
×	X	<u> 22</u>	32	×
				۲.
X	х	28	z	х
V	V	喞	₩	٥,
\triangle	◠			$^{\prime\prime}$
×	×	M	×	×
		ar.	d	ĸ
×	X	₩.	S	K/S
V	$^{\circ}$	a	-	5
\wedge	\wedge	40	ĸ.	◠
V	V	W.	73	P2
		₽		ĸ
X	X	$\overline{}$	$\overline{}$	×
V				

Question 4 continued	Leave blank
(Total 12 marks)	Q4



5. Given that

$$I_n = \int x^n \sqrt{(x+8)} \, dx, \qquad n \geqslant 0, x \geqslant 0$$

(a) show that, for $n \ge 1$

$$I_n = \frac{px^n(x+8)^{\frac{3}{2}}}{2n+3} - \frac{qn}{2n+3} I_{n-1}$$

where p and q are constants to be found.

(6)

(b) Use part (a) to find the exact value of

$$\int_0^{10} x^2 \sqrt{(x+8)} \, \, \mathrm{d}x$$

giving your answer in the form $k\sqrt{2}$, where k is rational.

(5)

	Le
huggian 5 continued	bl
Question 5 continued	



estion 5 continued	

		ı	ż	a
1	É	d		ľ
į	3	1	7	3
ł	Ŀ	a	Ŀ	a
-1	à	۰	à	ø
1	Ŀ	d	Ŀ	
s	å	ø	ė	ø
j	٩	۰	ě	'n
1	Ľ	Δ	۴	١
í	3	3	_	-
ń	Ξ	3	Ξ	5
i	_	J	Ĺ	٠
i	ī	7		3
1	r	•		•
ì	_	•		_
ď	2	2	2	P
1	Ŧ		Ŧ	-
4				-
ì	r	i	ŀ	ń
1	b	d	b	d
1	ŀ	ė		ij
3	4		-	i
ŝ	ā	5	ā	ā
1	E	2	E	
þ	b		ċ	
1	Ę		S	Ξ
J	ø	ę	P	7
4	í			
ł	P	7		Ţ
'n	è	Ģ	٩	h
J	Ę		ė	ı
1	Ė	Ę	3	p
ì	ė		_	ì
ì	ò	ø	9	ĸ
4	Ĺ	٠	ú)
1	ä	ē	ij	k
1	Ĺ	٠	_	1
2	-	٠	-	4

Question 5 continued	Leave blank
	Q5
(Total 11 marks)	
(10m2 II muris)	



blank

6. The line l_1 has equation

$$\mathbf{r} = \mathbf{i} + 2\mathbf{k} + \lambda(2\mathbf{i} + 3\mathbf{j} - \mathbf{k})$$

where λ is a scalar parameter.

The line l_2 has equation

$$\frac{x+1}{1} = \frac{y-4}{1} = \frac{z-1}{3}$$

(a) Prove that the lines l_1 and l_2 are skew.

(4)

(b) Find the shortest distance between the lines l_1 and l_2

(5)

The plane Π contains l_1 and intersects l_2 at the point (3, 8, 13).

(c) Find a cartesian equation for the plane Π .

(4)







uestion 6 continued	
uestion v continueu	



estion 6 continued	

Leave

-	ø	
VUQV	Ŀ	
-12	7	
ы	ш	
\overline{x}	=	
n	_	
=	-	
ŵ	p	
.94	ы	
	m.	
ν	0	
=	-i	
=	=	
л	7	
	=	
ы	-	
т	_	
2	۰	
-	,	
~	-	
Ξ	Ξ	
=	_	
20	٠û	
ш	4	
10		
45	-	
÷	÷	
-	ä	
ш	\subseteq	
THO	_	
147	3	
-	5	
and the	-	
d.,		
н		
HON	ď	
r	э	
-Ч-	s	
=	ė	
20	۳.	
_	=	
100		
\sim	ч	
غة	s	
- 2	Ξ.	
	٦	
	4	

Question 6 continued	blank
Question o continued	
	Q6
(Total 13 marks)	
(10th 10 marks)	



- 7. The ellipse E has foci at the points (±3, 0) and has directrices with equations $x = \pm \frac{25}{3}$
 - (a) Find a cartesian equation for the ellipse E.

(5)

The straight line *l* has equation y = mx + c, where *m* and *c* are **positive** constants.

(b) Show that the x coordinates of any points of intersection of l and E satisfy the equation

$$(16 + 25m^2)x^2 + 50mcx + 25(c^2 - 16) = 0$$
(2)

Given that the line l is a tangent to E,

(c) show that $c^2 = pm^2 + q$, where p and q are constants to be found.

(3)

The line l intersects the x-axis at the point A and intersects the y-axis at the point B.

(d) Show that the area of triangle *OAB*, where *O* is the origin, is

$$\frac{25m^2+16}{2m}$$

(3)

(e) Find the minimum area of triangle OAB.

	Ġ
1	



	L	Leave
	b	blank
Question 7 continued		



estion 7 continued	

	L
Overtion 7 continued	b
Question 7 continued	



		1/2
		NACTOR STORY
		1 💥
		1 💥
		1 💥
		1 🐰
	0.	
	Q7	- │ 🎇
(Total 15 marks)		
TOTAL FOR PAPER: 75 MARKS		
END		J 🛞
		/ \ /
P 5 5 8 1 5 A 0 2 8 2 8		
P 5 5 8 1 5 A 0 2 8 2 8		

Question 7 continued