Please check the examination detail	s below before ente	ering your candidate i	information
Candidate surname		Other names	
Centre Number Candidat	e Number		
Pearson Edexcel Int	ernation	al Advand	ced Level
Time 1 hour 30 minutes	Paper reference	WME	03/01
Mathematics			
International Advanced	Subsidiar	v/Advanced	llevel
Mechanics M3	Jabbiaiai	y//tavaricea	
Wiechanics Wi3			
			J
(Vth			
You must have: Mathematical Formulae and Statis	stical Tables (Ve	llow) calculator	Total Marks
Mathematical Formulae and Statis	sticai iabies (fe	now), Calculator	

Candidates may use any calculator permitted by Pearson regulations.

Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
- there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Whenever a numerical value of g is required, take g = 9.8 m s⁻², and give your answer to either two significant figures or three significant figures.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 7 questions in this question paper. The total mark for this paper is 75.
- The marks for each question are shown in brackets
- use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over ▶







Leave blank

1.	A particle P is moving in a straight line with simple harmonic motion of period 4. The centre of the motion is the point O	s.
	At time $t = 0$, P passes through O	
	At time $t = 0.5 \text{s}$, P is moving with speed $2 \text{m}\text{s}^{-1}$	
	(a) Show that the amplitude of the motion is $\frac{4\sqrt{2}}{\pi}$ m	4)
		+)
	(b) Find the maximum speed of P	2)
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_



							×	
		į					į	
5			,				×	
	2				ς		Κ	
							Κ	
	S						×	
	ς						Ü	
>			,				0	
					ς		Κ	
							Κ	
							ż	
							Ü	
5	ć						٦	
	2						Χ	
		i			2	S	á	
			ĺ		9		r	
١				7	7	7	ц	
				ь	á		и	
				b	d	S	d	
				ú	è	ś	à	
S		5			3		<u> </u>	
				7	₹	7	8	
	5	è	1	ù	ø		ρ	۰
			9	ņ	ú		Ŀ	
			,				7	
			į	è		ä	ĸ	
				L		r	a	
2	۲	Ź		S	2	S	2	
				Ξ	2		Ξ	
				7	٦	r	-	
			1	ù		ŀ	ŭ	
			í	5				
2					ø	Ė	P	
			ł	ĸ				
			3	ć	š	ä	à	
				2	G	ğ	ĸ	
?			1	ij	é	é		
>	<		ì	í	è	ś	ė	
S					S		Ď	
			3	r	Ý	ŕ	Ý	
		<		ń	d	ú		
?	Ś			б	Ž		Š	
>	<			ř	þ	٩	9	
5				Ź	١	ć	4	
				Z	5	7	5	
	5			Р	a		e	۰
2	ς		١	h	ø	В	ø	
>			ļ	ú	è	ĕ	2	
		ς	4	à	ú	ē	я	
			3	5	7	7		
\rangle			ì	í.				
5	<				ķ	ņ		
	>		Į	ž	Ś	ú	s	
					7	7	۹	
		j	٩	6	2	S	d	
۶			į	2	3	Z	_	
Š	2			7	S	3	P	
			j			ì	ú	
			7	5	d		K	
				ú	à	á	ń	
				r			1	
ζ				я	q	ø	р	
				à	ø	ij	б	
				ĸ			J	
			3	₹	7	9	7	۰
							X	
	Ś				8		×	
>	<	>			5	<	X	
	2	>		>	5		X	
	5	<		>			× × ×	
	3	<		3	3		××××	
	3	<		3			XXXX	
	3					~ < < < <	XXXX	
		< < < < < <	,	>		~ < < < < <	× < < × < × <	
	\ \ \	< < < < < <		>		~ < < < < < < < <	× < × × × ×	
	\ \ \	<<<<<		\ \ \		~ < < < < < < <	×< × × × × ×	
		< < < <				~ < < < < < < < <	×	
		<<<<<<		\ \ \		~~~~~~	×< × × × × × × × ×	
		<<<<<<<		\ \ \		~<<<<<	×< < < < < < < < < < <	
		<<<<<<<<<<<		\ \ \		~ < < < < < < < < < < < < < < < < < < <	××××××××××	
>>>>>>>		\ <<<<<<<				~ < < < < < < < < < < < < < < < < < < <	××××××××××××××××××××××××××××××××××××××	
		<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<				~ < < < < < < < < < < < < < < < < < < <	×< < < < < < < < < < < < <	
							×< < < < < < < < < < < < < < < < < < <	
							×< < < < < < < < < < < < < < < < < < <	
							×<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<<	
							××××××××××××××××××××××××××××××××××××××	
							×< < < < < < < < < < < < < < < < < < <	
		?						
		?						
		?						
		?						
		?						
		?						
		?						
		?						
		?						
		?						
		?						
		?						
		?						
		?						
		?						
		?						
		?						
		?						
		?						

DO NOT WRITE IN THIS AREA

Question 1 continued		blank
guestion I continued		
		Q1
	(Total 6 marks)	



Leave blank

2. In this question solutions relying on calculator technology are not acceptable.

A particle *P* of mass 2 kg is moving along the positive *x*-axis.

At time t seconds, where $t \ge 0$, P is x metres from the origin O and is moving away from O with speed $v \text{ m s}^{-1}$ where $v = \frac{1}{\sqrt{(2x+1)}}$

(a) Find the magnitude of the resultant force acting on P when its speed is $\frac{1}{3}$ m s⁻¹

(6)

When t = 0, P is at O

(b) Find the value of t when P is 7.5 m from O

(5)

		×
V	V	V
$\langle \rangle$		<u>_</u>
Χ,	É	r
×	99	46
V	À	И
a		ш
×		75
X		e
O.	ы	-
$^{\wedge}$	△	_
×ι	e	К
V	7	96
$\langle \rangle$		0
\triangle	ñ	ø.
X		M
×į	7	
O!	7	
×	Ħ	_
×.	S4	V
v.		∇
OI.	_	_
×	ĸ	$\overline{}$
		×
O	×	蚰
$^{\wedge}$	ĕ	Ž
×	7	_
VI	7	×
$\langle \rangle$		
×ι	ň	ďΝ
XI		м
Vi	s/	∇
	k	*
X.	ĸ	Χ
×	_	_
V	m	w
01		
Ŏ	Ä	$^{\times}$
X.	Σ	Š
O		=
∴i	۵	ø
X	X	х
×1	K	×
a	7	$\overline{}$
^*	-	△
×	K	71
v	ĸ	4
	=	\sim
	2	7
Χı		×
V	v2	V
	\simeq	△
×	n	71
X	~	u
\vee	2	2
(N	7	a
XI	ĸ	м

	Leav
Question 2 continued	blanl
Question 2 continued	



stion 2 continued	

Question 2 continued	Leave	
		l
		l
		l
		l
		l
		l
		l
		l
		l
		l
		l
		l
		l
		l
		l
		l
		l
		l
		l
		l
		l
		l
	Q2	
(Total 11 marks)		
(10tai 11 maiks)		4



Leave blank

3.




Figure 1

A particle P of mass m is attached to one end of a light elastic spring of natural length l and modulus of elasticity kmg, where k is a constant. The other end of the spring is fixed to horizontal ground.

The particle P rests in equilibrium, with the spring vertical, at the point E.

The point E is at a height $\frac{3}{5}l$ above the ground, as shown in Figure 1.

(a) Show that $k = \frac{5}{2}$

(2)

The particle P is now moved a distance $\frac{1}{4}l$ vertically downwards from E and released from rest. Air resistance is modelled as being negligible.

(b) Show that P moves with simple harmonic motion.

(4)

(c) Find the speed of P as it passes through E.

(4)

(d) Find the time from the instant P is released to the first instant it passes through E.





		×
V	V	V
$\langle \rangle$		\sim
Λ	ان	æ
X,	79	46
vi	M	и
a	N A	ш
X	\overline{z}	73
X	ø	m
V	w	-
$\langle \rangle$	<u></u>	
N		ж.
	X	70
V	V	V
0	n	A
X,	w	24
Χı	×	è
V	\sim	\sim
$\langle \rangle$	а	
Ν		×
X	įΚ	×
V	r	_
O	O	
$^{\wedge}$	4	Δ
X	\mathbf{x}	7
V		
$\langle \cdot \rangle$		$\hat{}$
N		75
×	×	×
Ä		
	v	
8	K	V
8	K	X
	K	X
	Z X	X
	76 15 16 16 16 16 16 16 16 16 16 16 16 16 16	V X R
	K K K K K K	XX
	K K A A A	
	T K A A A A A	
	K K A A A A K K	
	TO BOOK OF THE PERSON OF THE P	
	T X A A A A A A A A A A A A A A A A A A	
	TO K A CO A CO A CO TO A CO	
	TO BE CONTROL OF SECTION OF SECTI	

	blank
Question 3 continued	Dialik
Question 5 continued	



estion 3 continued	

Question 3 continued	blank
	Q3
(Total 12 marks)	
(Total 12 marks)	



Leave blank

4. A light elastic string has natural length 2a and modulus of elasticity 2mg.

One end of the elastic string is attached to a fixed point O. A particle P of mass $\frac{1}{2}m$ is attached to the other end of the elastic string.

The point A is vertically below O with OA = 4a.

Particle P is held at A and released from rest. The speed of P at the instant when it has moved a distance a upwards is $\sqrt{3ag}$

Air resistance to the motion of P is modelled as having magnitude kmg, where k is a constant.

Using the model and the work-energy principle,

(a) show that
$$k = \frac{1}{4}$$

(7)

Particle P is now held at O and released from rest. As P moves downwards, it reaches its maximum speed as it passes through the point B.

(b) Find the distance *OB*.

(4)



×	×	X	
\sim		ĸ.	2
Χ.			7
V4	۰,	a,	
		7	9
×I		ľИ	
VI		н	
			è
×			۰
ാ	ш	ь	à
×.	$\overline{}$	7	ī
v.	ú	Į	,
\sim	6	æ	S
X.		7	,
$\langle \rangle$		<.	>
$^{\wedge}$	n	a	ú
M	м	Ю	
	-		9
×ι			
V.	_	\sim	2
	п	_	5
×.	24	Ŀ	_
	_	₹	,
ΧI	K		L
VI	×	-	
\sim	n	↶	
X.		X	
Χ.	ž	r	
V/I		'n	ú
	\simeq	\sim	١
×	$\overline{}$	7	•
$\langle \rangle$		ς,	>
ă	ř١	n	
M	ы	ы	
	_	↽	
×I	ĸ		
VI	7	₹	۲
	<u>*\</u>	2	S
×	$\overline{}$	7	
(<i>7</i>)		úď	
∕√∥	(2)	e	s
V4	÷	-	۰
		~	٤
×.	×	a	ø
	46	٤.	2
		ø	ij
×	X	×	
	0		
×Ν	ь.	÷	
VI	K.	ž	1
S	ä	6	١
XI	区	7	
∵.XI	N2	SJ	ď
\sim	74	~	
XI		9	
	2		
Χı	2	5	2
ø	2	ě	į
8	X	Ž	9
8	X	Š	2
8	X	\ \ \ \	
	X	3	
8	×	3	
	X		
	X		

	blank
Question 4 continued	Diank
Question 4 continued	



estion 4 continued	

Question 4 continued	blank
	04
	Q4
(Total 11 marks)	



Leave blank

5.




Figure 2

A small smooth ring R of mass m is threaded on to a thin smooth fixed vertical pole. One end of a light inextensible string of length 2l is attached to a point A on the pole. The other end of the string is attached to R. A particle P of mass m is attached to the midpoint of the string. The particle P moves with constant angular speed in a horizontal circle, with both

halves of the string taut, and $AR = \frac{6l}{5}$, as shown in Figure 2.

It may be assumed that in this motion the string does not wrap itself around the pole and that at any instant, the triangle APR lies in a vertical plane.

- (a) Show that the tension in the lower half of the string is $\frac{5mg}{3}$ (3)
- (b) Find, in terms of l and g, the time for P to complete one revolution.

Question 5 continued	blank



estion 5 continued	

Question 5 continued	blank
	Q5
(Total 11 marks)	



blank

6.




Figure 3

A light rod of length a is free to rotate in a vertical plane about a horizontal axis through one end O. A particle P of mass m is attached to the other end of the rod. The particle P is held at rest with the rod making an angle α with the upward vertical through O,

where
$$\tan \alpha = \frac{3}{4}$$

The particle P is then projected with speed u in a direction which is perpendicular to the rod. At the instant when the rod makes an angle θ with the upward vertical through O, the speed of P is v, as shown in Figure 3.

Air resistance is assumed to be negligible.

(a) Show that
$$v^2 = u^2 + \frac{2ag}{5}(4 - 5\cos\theta)$$

It is given that $u^2 = \frac{6ag}{5}$ and P moves in complete vertical circles.

When $\theta = \beta$, the force exerted on *P* by the rod is zero.

(b) Find the value of $\cos \beta$

(6)

	Leave
	blank
Question 6 continued	



estion 6 continued	

XXXXX	

$\times\!\!\times\!\!\times\!\!\times$	
$\times\!\!\times\!\!\times\!\!\times$	
$\times\!\!\times\!\!\times\!\!\times$	
XXXX	
××××	
××××	

$\times\!\!\times\!\!\times\!\!\times$	
\times	
$\sim\sim\sim$	
\times vo \times	
\times	
\times	
\times	
₩	
×=×	
(XXXXX)	
$\times \mapsto \times$	
$\times \times \times \times$	
$\times \circ \times$	
\times O \times	
~~~	
$\times\!\!\!\!/\!\!\!\!/\!\!\!\!/\!\!\!\!\!/\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!$	
******	
XXXX	
****	
****	
****	
$\times\!\!\times\!\!\times\!\!\times$	
****	
****	
××××	
*****	
$\times\!\!\times\!\!\times\!\!\times$	
××××	
××××	
*****	
$\times\!\!\times\!\!\times\!\!\times$	
$\times \times \times \times \times \times$	
XXXX	
4	
AREA	
AREA	
IS AREA	
IS AREA	
IS AREA	
N THIS AREA	
IN THIS AREA	
N THIS AREA	
IN THIS AREA	
IN THIS AREA	
RITE IN THIS AREA	
RITE IN THIS AREA	
RITE IN THIS AREA	
WRITE IN THIS AREA	
WRITE IN THIS AREA	
OT WRITE IN THIS AREA	
OT WRITE IN THIS AREA	
WRITE IN THIS AREA	
OT WRITE IN THIS AREA	
OT WRITE IN THIS AREA	
O NOT WRITE IN THIS AREA	
O NOT WRITE IN THIS AREA	
O NOT WRITE IN THIS AREA	
O NOT WRITE IN THIS AREA	
O NOT WRITE IN THIS AREA	
O NOT WRITE IN THIS AREA	
O NOT WRITE IN THIS AREA	
O NOT WRITE IN THIS AREA	
O NOT WRITE IN THIS AREA	
O NOT WRITE IN THIS AREA	
O NOT WRITE IN THIS AREA	
DO NOT WRITE IN THIS AREA	
DO NOT WRITE IN THIS AREA	
DO NOT WRITE IN THIS AREA	
DO NOT WRITE IN THIS AREA	
DO NOT WRITE IN THIS AREA	
DO NOT WRITE IN THIS AREA	
DO NOT WRITE IN THIS AREA	
A DO NOT WRITE IN THIS AREA	
A DO NOT WRITE IN THIS AREA	
EA DO NOT WRITE IN THIS AREA	
REA DO NOT WRITE IN THIS AREA	
AREA DO NOT WRITE IN THIS AREA	
EA DO NOT WRITE IN THIS AREA	

	Leave blank
Question 6 continued	
	Q6
(Total 10 marks)	
(Total To marks)	



- 7. [You may assume that the volume of a cone of height h and base radius r is  $\frac{1}{3}\pi r^2 h$ .]

  A uniform solid right circular cone C, with vertex V, has base radius r and height h.
  - (a) Show that the centre of mass of C is  $\frac{3}{4}h$  from V (4)

A solid F, shown below in Figure 4, is formed by removing the solid right circular cone C' from C, where cone C' has height  $\frac{1}{3}h$  and vertex V




Figure 4

(b) Show that the distance of the centre of mass of F from its larger plane face is  $\frac{3}{13}h$ 

(5)

The solid F rests in equilibrium with its curved surface in contact with a horizontal plane.

(c) Show that  $13r^2 \leqslant 17h^2$ 

(5)

	blank
Question 7 continued	



estion 7 continued	

	Leave
	blank
Question 7 continued	



estion 7 continued	blank
	-
	_
	_
	_
	-
	-
	_
	-
	-
	-
	-
	_
	-
	-
	_
	_
	-
	-
	-
	_
	_
	-
	-
	_
	-
	Q7
(Total 14 marks	
TOTAL FOR PAPER = 75 MARKS END	5