Please check the examination deta	ails below	before ente	ring your candidate information
Candidate surname			Other names
Pearson Edexcel International Advanced Level	Centre	Number	Candidate Number
<b>Wednesday 7</b>	'No	ven	nber 2018
Morning (Time: 2 hours 30 minu	tes)	Paper Re	eference WMA02/01
Core Mathema	tics	<b>C34</b>	
You must have: Mathematical Formulae and Sta	tistical T	ables (Blu	Total Marks

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

## Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
   there may be more space than you need.
- You should show sufficient working to make your methods clear.
   Answers without working may not gain full credit.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

## Information

- The total mark for this paper is 125.
- The marks for each question are shown in brackets
   use this as a guide as to how much time to spend on each question.

## **Advice**

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶







(a) Write  $\cos \theta + 4 \sin \theta$  in the form  $R \cos (\theta - \alpha)$ , where R and  $\alpha$  are constants, R > 0and  $0 < \alpha < \frac{\pi}{2}$ . Give the exact value of R and give the value of  $\alpha$  in radians to 3 decimal places. **(3)** (b) Hence solve, for  $0 \le \theta < \pi$ , the equation  $\cos 2\theta + 4\sin 2\theta = 1.2$ giving your answers to 2 decimal places. **(5)** 

				ĺ		
					ij	
				1		
	í					
	í				É	
	ì					
				è	ć	
		г	7	è	Ś	
	1	Ç	2	Ś	Ź	
	1	Ç	7		Ź	
	1	Į			í	
	1	Ľ			Я	
	1	Ľ			Я	
	1	Ľ			Я	
	1	Ľ			Я	
		Ľ			Я	
		Ľ			Я	
		Ľ			Я	
		Ľ			Я	
		Ľ			Я	
		Ľ			Я	
		Ľ			Я	
		Ľ				
		Ľ			Я	
		Ľ			Я	

Question 1 continued	Leave blank
	Q1
(Total 8 marks)	
· · · · · · · · · · · · · · · · · · ·	



2.	A curve	C has	equation

$$x^3 - 4xy + 2x + 3y^2 - 3 = 0$$

Find	an equ	uation	of the	normal	to C	at the	point	(-3, 2)	2), ફ	giving	your	answer	in	the	form
ax +	by +	c = 0	where	a, b and	dc a	re integ	gers.								
															<b>(7)</b>

2					۲	
					/	
				4	į	
	'n		P		К	
	۹	Ħ	×		ы	
	ú		\		3	
		ļ.			a	
		3	Ξ		S	ī.
		P	۹			
			ч		ä	ď
				3		
		S	à	d		
			S		К	
					۱	
				S		
	٦			ŋ	4	ľ.
	š		2	٤	_	
2	5	۳	₹	7	₹	
		ě				
		ä	L	2	2	
	ĭ	К	'n	Ď	d	
	3	7	₹	7	1	
		L	2	7	2	
		I.				
	1					
		К				
0						
			2	2		
	'n		,	۰		
		r			Я	ŀ
	٦	٩	ú	ù	ø	r
		G				
	ı	ρ	۳	7	٩	
		Ľ			1	
	3		۹	ø	8	r

Question 2 continued
$egin{array}{c} \mathbf{Q2} \\ \hline \end{array}$
(Total 7 marks)



3. Given $\cos\theta^\circ = p, \text{ where } p \text{ is a constant and } \theta^\circ \text{ is acute}$ $\text{use standard trigonometric identities to find, in terms of } p,$ $\text{(a)}  \sec\theta^\circ$ $\text{(b)}  \sin(\theta - 90)^\circ$ $\text{(c)}  \sin 2\theta^\circ$ $\text{(3)}$ Write each answer in its simplest form.			
use standard trigonometric identities to find, in terms of $p$ ,  (a) $\sec \theta^{\circ}$ (b) $\sin (\theta - 90)^{\circ}$ (c) $\sin 2\theta^{\circ}$ (3)	3.	Given	
(a) $\sec \theta^{\circ}$ (b) $\sin(\theta - 90)^{\circ}$ (c) $\sin 2\theta^{\circ}$ (3)		$\cos \theta^{\circ} = p$ , where p is a constant and $\theta^{\circ}$ is acute	
(a) $\sec \theta^{\circ}$ (b) $\sin(\theta - 90)^{\circ}$ (c) $\sin 2\theta^{\circ}$ (3)		use standard trigonometric identities to find, in terms of $p$ ,	
(b) $\sin(\theta - 90)^{\circ}$ (c) $\sin 2\theta^{\circ}$ (3)			
(c) $\sin 2\theta^{\circ}$ (3)		(a) SCC 0	(1)
(c) $\sin 2\theta^{\circ}$ (3)		(b) $\sin(\theta - 90)^{\circ}$	
(3)			(2)
		(c) $\sin 2\theta^{\circ}$	
Write each answer in its simplest form.			(3)
		Write each answer in its simplest form.	



3		į		
		1		
	Č			
	Ħ			
ż				
			>	
ä				
		×		
ì				

			ķ		
		7			
	ž	K	ù	ú	ь
X	ã	K	ù	Ŕ	þ
X	ĺ	5	ľ	ź	Š
X	į	í	Ì	2	Š
Š	ĺ		į	2	
X				Ź	
X				Ź	
X					
X					
×					
X					
X					
X					
× × × ×					
× × ×					
× × × ×					
× × × ×					
× × × × ×					
× × × × × ×					
× × × × × ×					
× × × × ×					
× × × × × × ×					

$\sim\sim$	$\vee$	V
$\times \times \wedge$		
$\times \times$	×	×
XX		$\langle \rangle$
$\sim$	$^{\wedge}$	$^{\wedge}$
$\propto \times$	×	×
$\times \times$	<	$\langle \rangle$
XX	х	×
$\sim\sim$	$\vee$	$\vee$
$\sim$	$^{\wedge}$	$\sim$
$\times \times$	X	×
XX		$\langle \rangle$
$\sim$	$^{\wedge}$	$^{\sim}$
$\sim\sim$	24	w
XX		$\langle \cdot \rangle$
$\langle XX \rangle$	74	ж.
$\triangle$	w	w
$\sim$	ДЪ	ΔN
	$\overline{}$	_
$\times \times a$	n.	#2
	æ	Z.
	~	=
$\times \times \rangle$	絀	ell)
		DX.
$\wedge \vee 7$	v	=
$\vee \vee \wedge$		$\triangle$
CX %	76	×
$\times \times 1$	SB	SIP.
$\vee$	٣.	Α.
CCM		er/
$\times \times 2$	$\sim$	$\sim$
VV.	7	癶
$\sim\sim$	3R	$\sim$
XX5		•
	×	×
	$\nabla$	~
XX"		$\langle \rangle$
		ĸ.
$\triangle \nabla Z$	3	~
	2	Z,
	Ž	Ź
$\otimes$	Ž	3
	Z	
	Z A	
	2	3

Question 3 continued	Leave blank
	<b>Q3</b>
(Total 6 marks)	



4.




Figure 1

Figure 1 shows a sketch of part of the curve with equation  $y = 8x - xe^{3x}$ ,  $x \ge 0$ 

The curve meets the x-axis at the origin and cuts the x-axis at the point A.

(a) Find the exact x coordinate of A, giving your answer in its simplest form.

**(2)** 

The curve has a maximum turning point at the point M.

(b) Show, by using calculus, that the x coordinate of M is a solution of

$$x = \frac{1}{3} \ln \left( \frac{8}{1+3x} \right) \tag{5}$$

(c) Use the iterative formula

$$x_{n+1} = \frac{1}{3} \ln \left( \frac{8}{1 + 3x_n} \right)$$

with  $x_0 = 0.4$  to calculate the values of  $x_1$ ,  $x_2$  and  $x_3$ , giving your answers to 3 decimal places.



Question 4 continued
I



estion 4 continued	

	K		
	Ý		
			ì
V	ŭ	P	Ė
×	Š	₹	5
Š	Ž	5	2
Š	Ž	Ş	
Š	Ž	2	2
ă	Ź		
Ĭ	Ž		
	Ž		
à	ŝ	ζ	5
à	ŝ	ζ	5
à	ŝ	ζ	5
à	ŝ	ζ	5
à	ŝ	ζ	5
à	ŝ	ζ	5
à	ŝ	ζ	5
à	ŝ	ζ	5
à	ŝ	ζ	5
à	ŝ	ζ	5
à	ŝ	ζ	5
à	ŝ	ζ	5
à	ŝ	ζ	5
à	ŝ	ζ	5
à		ζ	5
à	ŝ	ζ	5
à	ŝ	ζ	5

Question 4 continued	Leave
Question 4 continued	
	Q4
(Total 10 marks)	
(10th 10 marks)	



5.

$$f(x) = \frac{4x^2 + 5x + 3}{(x+2)(1-x)^2} \equiv \frac{A}{(x+2)} + \frac{B}{(1-x)} + \frac{C}{(1-x)^2}$$

(a) Find the values of the constants A, B and C.

**(4)** 

**(6)** 

- (b) (i) Hence find  $\int f(x) dx$ .
  - (ii) Find the exact value of  $\int_0^{\frac{1}{2}} f(x) dx$ , writing your answer in the form  $p + \ln q$ , where p and q are constants.



	Leave
	blank
Question 5 continued	



nestion 5 continued	

				1		
			Ì			
		Ì				
					1	
	i					
				į		
				è		
	á				È	
	į				Ž	
	1			į		
					4	
	Ì					
		۶	₹	2	S	
	×	2	6	ì	S	
	ì	į			ś	
	ĺ	ľ	Ś	2	Í	k
	4					į
	ĺ					
	ì					
	1					

		É				
				ě		
			×			
			ì			
	ì	p	ρ	ą	P	
	į	ø	ŝ	P	5	
V	į		Š		>	
X	į		2	2	>	
8	į		2	2	>	
8	Ì	Ä		2		
8	į					
8	į				>	
8	į					
8	į					
8	į					
8	į					
× × ×						

Κ.				
Λ	`	/		
×				
ĸ.		⋖		
		2		
C				
Ν	â	à		P
ΧI		6		E
~	Ś		7	9
N		٦	г	٦
ΧI		а		
7		7		7
V		۹		ρ
И	L	4		ĕ
X	7	7	₹	7
Κ.	2	ģ		
.4		c		υ
		3	7	٩
C				
- 74	è		ú	ń
Ŋ		SI	r	٩
⇗		y		J
N				
				s
74	e		P	9
٦.		Ч	ν	
1		,	=	P
K				
ΚI				
Л		2		
∿		1		
a		ø		
	6		ρ	7
N		ū	o	ú
Ζ,	s	2	$\sim$	2
Ŋ	۰	7	7	-
ĸ.				
.71		А	s.	J
V				
ĸ.		7		7
Я		۵	4	-
ΚI		₹	7	7
7	S	2	2	2
N		7	7	
Κi		M	ù	
. Х	E	.2	к	3
V	۲		۶	4
Л			è	ú
×	ú	à	ø	я
K.	7	۹	×	6
Ä	i			g
		S		
Κi	è			
A.		×	ú	ė
V		7	7	
1	S	2	'n	s
×	ľ	7		٦
K.		ú	ú	
7		7	٠	2
V	ı	۹		
€.	۵	į	ρ	۲
×		ø	q	ø
۲.			2	
~	S	ż	2	2
¥		۲	۲	

Question 5 continued	Leave blank
	Q5
(Total 10 marks)	



**6.** (a) Use binomial expansions to show that, for  $|x| < \frac{1}{2}$ 

$$\sqrt{\frac{1+2x}{1-x}} \approx 1 + \frac{3}{2}x + \frac{3}{8}x^2 \tag{6}$$

(b) Find the exact value of  $\sqrt{\frac{1+2x}{1-x}}$  when  $x = \frac{1}{10}$ 

Give your answer in the form  $k\sqrt{3}$ , where k is a constant to be determined.

**(1)** 

(c) Substitute  $x = \frac{1}{10}$  into the expansion given in part (a) and hence find an approximate value for  $\sqrt{3}$ 

Give your answer in the form  $\frac{a}{b}$  where a and b are integers. (2)

	blank
Question 6 continued	- Clarin
Question o continueu	



estion 6 continued	

		g			
	3				
			į		
	ä				
			į		
	1		ķ		
		K			
		K			
	á	À			
	i	þ	V	7	
X	i	è		Z	
X	я	2	ĕ	Z	
	я	è	Ķ	7	ļ
X	į		ľ	Z	
	я	à	ľ	Z	
	я	Ž	ž	2	
×	я		į	2	
	я		ž		
	я				
	я				
	я				
×××	я				
××××	я				
××××	я				
××××	я				
××××	я				
××××	я				
××××	я				
× × × ×	я				
××××	я				
×××××	я				
××××××	я				
×××××	я				
××××××××××××××××××××××××××××××××××××××	я				
×××××××	я				
×××××××	я				
××××××××××××××××××××××××××××××××××××××	я				
~ × × × × × × × ×	я				
~ × × × × × × × ×	я				
~ × × × × × × × × ×	я				
~ × × × × × × × ×	я				
~ × × × × × × × × ×					
~ × × × × × × × × × × × ×					
~ × × × × × × × × × × ×	я				
× × × × × × × × × × × × × × × × × × ×					
~ × × × × × × × × × ×					
~ × × × × × × × × ×					
××××××××××××××××××××××××××××××××××××××					
~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~					
××××××××××××××××××××××××××××××××××××××					

Question 6 continued	Leave blank
	Q6
(Total 9 marks)	



7. A curve has equation

$$y = \ln(1 - \cos 2x), \quad x \in \mathbb{R}, 0 < x < \pi$$

Show that

(a) 
$$\frac{dy}{dx} = k \cot x$$
, where k is a constant to be found.

(4)

Hence find the exact coordinates of the point on the curve where

(b) 
$$\frac{\mathrm{d}y}{\mathrm{d}x} = 2\sqrt{3}$$

(4)



	Leave blank
Question 7 continued	



estion 7 continued	

Leave

×ı			
		Î	
	X A		
		Ĭ	
		2	
		b	
	2 X X X X X X X X X X X X X X X X X X X		
		ž	
		÷	
	Ÿ		
X			
			ì
			Į.

			ĺ		
				ú	
3					
Ò					
d					
Ø					
K					
ĸ,					
	ì				
		é			
				Í	
			2		
>	1				
≺	à	á		6	
S	è	þ	₹	ş	K
Š	ĺ		5	Ì	б
S	ĺ	Ś	2	١	ĺ
Š	į		2	١	Ŕ
3			2		6

Question 7 continued	blank
	<b>Q7</b>
(Total 8 marks)	



**8.** (i) Find  $\int x \sin x \, dx$ 

**(3)** 

(ii) (a) Use the substitution  $x = \sec \theta$  to show that

$$\int_{1}^{2} \sqrt{1 - \frac{1}{x^{2}}} \, dx = \int_{0}^{\frac{\pi}{3}} \tan^{2}\theta \, d\theta$$
(3)

(b) Hence find the exact value of

$$\int_1^2 \sqrt{1 - \frac{1}{x^2}} \, \mathrm{d}x$$

**(4)** 

	blank
Question 8 continued	



estion 8 continued	
	<del></del>

	Leave blank
Question 8 continued	
	Q8
(Total 10 marks)	
(20002 10 1100110)	



**9.** A rare species of mammal is being studied. The population P, t years after the study started, is modelled by the formula

$$P = \frac{900e^{\frac{1}{4}t}}{3e^{\frac{1}{4}t} - 1}, \quad t \in \mathbb{R}, \quad t \geqslant 0$$

Using the model,

(a) calculate the number of mammals at the start of the study,

**(1)** 

(b) calculate the exact value of t when P = 315Give your answer in the form  $a \ln k$ , where a and k are integers to be determined.

(4)

(c) (i) Find  $\frac{dP}{dt}$ 

(ii) Hence find the value of  $\frac{dP}{dt}$  when t = 8, giving your answer to 2 decimal places.



Question 9 continued



	Leave blank
Question 9 continued	

	1		7	5	7	
	1	ļ	2	2	3	5
>	1	Į	2	2	2	5
	9	Į	2	2	7	
	1		3	Ź	2	
	1					
	1					
	1					
	1111					
	1111					
	200					6
	200					6
	200					6
	200					6
	200					6
	1					
	1					
	1					
	1					
	1					
	1					
>>>>>>	1					
	1					
	1					

0	đ	,	9	Ц		'n	
2	ú		ς	J	3		
`					J		
	ì	à	6	G	ó	á	
					7		
1							
١,					й	ú	
0	d			9		5	
0	8	я	٩			ì	
						7	
				2			
		þ			A	١	
				ĸ			
2	ú			2			
	3	9	۰	•	9	۰	
		i	a		ė	ì	
	G		7		7		
			4	2	ú	_	
C			7	e	•	5	
2	ä	>		j			
		L	ż	S		2	
	į	۰	5			Į	
b	i	и		2			
		•	7	۹	۰	۶	
2	j	à	i	i		ı	
١		3			ō	7	
	ď			2		S	
1	1				^		
١							
0							
2	ä	à	'n	i		i	
١	5	ū	Z	5	ū	Z	
		P	۹		P	5	
		b	á		н	'n	
Û				ζ			
2	5	,	٩	b	ĕ	'n	
۱				и	ø	,	
/		۰	۹			b	
	j			q	×	•	
ι						>	
2					٥		
١	i		Ħ	b	ı	ı	
	i			۹		i.	
	3		ь	H			
			7	7			
C	Ą	ä	9				
2	ú	à		я	ĸ	ί	
1	1	ij	×	×	ø	ď	
		S	4	÷	×		
			7	4	3		
2				ź			
١	1	ij	ø			,	
		á			Ь	Š	
		,			8		
		L	ú	S	÷		

	1				
				ŧ	
			2		
			Ž		
			Š		
	J				
	4				
2	4	9	P	7	₹
2	4		5	2	7
2	4		2	3	3

Question 9 continued	Leave blank
	<b>Q9</b>
(Total 9 marks)	



10.




Figure 2

Figure 2 shows a sketch of part of the graph with equation y = g(x), where

$$g(x) = \frac{3x - 4}{x - 3}, \quad x \in \mathbb{R}, \quad x < 3$$

The graph cuts the x-axis at the point A and the y-axis at the point B, as shown in Figure 2.

(a) State the range of g.

**(1)** 

- (b) State the coordinates of
  - (i) point A
  - (ii) point B

**(2)** 

(c) Find gg(x) in its simplest form.

**(3)** 

(d) Sketch the graph with equation y = |g(x)|

On your sketch, show the coordinates of each point at which the graph meets or cuts the axes and state the equation of each asymptote.

**(3)** 

(e) Find the exact solution of the equation |g(x)| = 8

(3)



estion 10 continued	



Question 10 continued

000000
****
*****
$\times$
*****
*****
$\times$
****
****
XXXXXXX
$\times$
$\times$
×4×
XXXXXX
<b></b>
<b>****</b>
** <b>Z</b> **
00000
$\otimes$
XX <b>.</b> XX
<u>~~</u>
××××
<b>*****</b>
$\otimes \bowtie \otimes$
$\times$
× <b>Z</b> ×
$\times \odot \times$
******
$\times$
$\times\!\!\times\!\!\times\!\!\times$
*****
$\times$
*****
****
$\times\!\!\times\!\!\times\!\!\times$
$\cdots$
$\times$
*****
$\times$
*****
XXXXX
$\cdots$
\$
Œ V
SAR
\$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$
\$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$
THIS
\$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$
IN THIS
E IN THIS
IN THIS
E IN THIS
E IN THIS
E IN THIS
WRITE IN THIS
E IN THIS
WRITE IN THIS
WRITE IN THIS
WRITE IN THIS
WRITE IN THIS
O NOT WRITE IN THIS
O NOT WRITE IN THIS
O NOT WRITE IN THIS
O NOT WRITE IN THIS
DO NOT WRITE IN THIS
O NOT WRITE IN THIS
DO NOT WRITE IN THIS
DO NOT WRITE IN THIS
DO NOT WRITE IN THIS
DO NOT WRITE IN THIS
DO NOT WRITE IN THIS
DO NOT WRITE IN THIS
DO NOT WRITE IN THIS
DO NOT WRITE IN THIS
DO NOT WRITE IN THIS
DO NOT WRITE IN THIS
DO NOT WRITE IN THIS
DO NOT WRITE IN THIS
DO NOT WRITE IN THIS
DO NOT WRITE IN THIS
DO NOT WRITE IN THIS
DO NOT WRITE IN THIS
DO NOT WRITE IN THIS
EA DO NOT WRITE IN THIS
REA DO NOT WRITE IN THIS
KREA DO NOT WRITE IN THIS
AREA DO NOT WRITE IN THIS
AREA DO NOT WRITE IN THIS
IS AREA DO NOT WRITE IN THIS
IS AREA DO NOT WRITE IN THIS
IS AREA DO NOT WRITE IN THIS
THIS AREA DO NOT WRITE IN THIS
IS AREA DO NOT WRITE IN THIS
N THIS AREA DO NOT WRITE IN THIS
IN THIS AREA DO NOT WRITE IN THIS
N THIS AREA DO NOT WRITE IN THIS
IN THIS AREA DO NOT WRITE IN THIS
E IN THIS AREA DO NOT WRITE IN THIS

Question 10 continued

_		
_		
_		
_		
_		
_		
_		
-		
_		
_		
_		
_		
_		
_		
_		
_		
_		
_		-
	(Total 12 marks)	



11. Relative to a fixed origin O, the line  $l_1$  is given by the equation

$$l_1: \quad \mathbf{r} = \begin{pmatrix} 2\\3\\-1 \end{pmatrix} + \lambda \begin{pmatrix} -1\\4\\3 \end{pmatrix}$$

where  $\lambda$  is a scalar parameter.

The line  $l_2$  passes through the origin and is parallel to  $l_1$ 

(a) Find a vector equation for  $l_2$ 

**(2)** 

The point A and the point B both lie on  $l_1$  with parameters  $\lambda = 0$  and  $\lambda = 3$  respectively.

Write down

- (b) (i) the coordinates of A,
  - (ii) the coordinates of B.

**(2)** 

(c) Find the size of the acute angle between OA and  $l_1$ 

Give your answer in degrees to one decimal place.

**(3)** 

The point D lies on  $l_2$  such that OABD is a parallelogram.

(d) Find the area of *OABD*, giving your answer to the nearest whole number.

**(3)** 





nestion 11 continued	



Question 11 continued		

XXXXX
XXXXX
******
******
*****
*****
******
*****
⋖
05
× oc×
<b>4</b>
× co
XX XX -
× × × × × × × × × × × × × × × × × × ×
× <b>z</b> ×
× <del>==</del> ×
XIIIX
<b>₩</b> 1XXX
<u>62</u>
X. <b>5</b> XX
$\otimes \blacksquare \otimes$
Ö
XX <del></del> XX
0
<u> </u>
*****
*****
*****
*****
*****
*****
*****
*****
*****
******
*****
*****
*****
*****
*****
<u> </u>
XX 2XX -
<b>A</b>
<b>A</b>
HISA
HISA
THIS A
N THIS A
THISA
EIN THIS A
EIN THIS A
TE IN THIS A
TE IN THIS A
EIN THIS A
VRITE IN THIS A
WRITE IN THIS A
WRITE IN THIS A
OT WRITE IN THIS A
NOT WRITE IN THIS A
NOT WRITE IN THIS A
O NOT WRITE IN THIS A
O NOT WRITE IN THIS A
O NOT WRITE IN THIS A
DO NOT WRITE IN THIS A
DO NOT WRITE IN THIS A
DO NOT WRITE IN THIS A
DO NOT WRITE IN THIS A
DO NOT WRITE IN THIS A
DO NOT WRITE IN THIS A
DO NOT WRITE IN THIS A
DO NOT WRITE IN THIS A
DO NOT WRITE IN THIS A
DO NOT WRITE IN THIS A
DO NOT WRITE IN THIS A
DO NOT WRITE IN THIS A
DO NOT WRITE IN THIS A
DO NOT WRITE IN THIS A
DO NOT WRITE IN THIS A
DO NOT WRITE IN THIS A
DO NOT WRITE IN THIS A
A DO NOT WRITE IN THIS A
A DO NOT WRITE IN THIS A
REA DO NOT WRITE IN THIS A
REA DO NOT WRITE IN THIS A
AREA DO NOT WRITE IN THIS A
AREA DO NOT WRITE IN THIS A
IS AREA DO NOT WRITE IN THIS A
HS AREA DO NOT WRITE IN THIS A
IS AREA DO NOT WRITE IN THIS A
THIS AREA DO NOT WRITE IN THIS A
THIS AREA DO NOT WRITE IN THIS A
THIS AREA DO NOT WRITE IN THIS A
THIS AREA DO NOT WRITE IN THIS A
IN THIS AREA DO NOT WRITE IN THIS A

XXXXXXX	
- CXXXXX	
XXXXXX	
-000	
0000000	
XX	
-0000000	
XXXXXXX	
Ē	
-004400	
XX XXX	
XXXXXXX	
XXXXXX	
212	
- 88000000	
£	
VRIT	
WRIT	
WRIT	
WRIT	
r wrrit	
T WRIT	
T WRIT	
OT WRIT	
OT WRIT	
NOT WRIT	
NOT WRIT	
NOT WRIT	
NOT WRIT	
NOT WRIT	
O NOT WRIT	
O NOT WRIT	
OO NOT WRIT	
DO NOT WRITE IN THIS AREA	
DO NOT WRIT	

	Leave blank
Question 11 continued	
	Q11
(Total 10 marks)	



12.




Figure 3

Figure 3 shows a sketch of part of the curve C with parametric equations

$$x = 7t^2 - 5,$$
  $y = t(9 - t^2),$   $t \in \mathbb{R}$ 

(a) Find an equation of the tangent to C at the point where t = 1

Write your answer in the form ax + by + c = 0, where a, b and c are integers.

(5)

The curve C cuts the x-axis at the points A and B, as shown in Figure 3

- (b) (i) Find the x coordinate of the point A.
  - (ii) Find the x coordinate of the point B.

**(3)** 

The region R, shown shaded in Figure 3, is enclosed by the loop of the curve C.

(c) Use integration to find the area of R.

**(5)** 

	blank
Question 12 continued	Diank
Question 12 continued	



Question 12 continued	

Leave

				>		
			þ			
				ď		
	≺			>		
S				>		
				C		
S		×		$\rangle$		
				١		
			ú	à	ú	ø
		4				Ŀ
		ú		2	7	7
				1		
		9	5	7	3	7
		1	P	Я	ġ	9
		ą	ĺ	Ħ	þ	à
,				S	ä	á
		1		S		
				7	3	7
				$\geq$		
			ľ	И	ρ	7
		J	۹	ę		9
	b	ì	į	ø	þ	Ę
		í	į	Ė	Ŕ	ė
					Ľ	
	0	ē		7	t	7
			Ŀ	٤	ú	ś
	>			7	3	3
		Š	,	2	Ś	į
>		ą	ė			ø
S	0	g			í	Š
		j	į	S	ř	S
	>	j	Ś	ď	Ž	₹
		Š	,	2		å
S			ĺ		Ĺ	
S		å	í	Š	7	۹
			ĺ	ú	ì	í
		1	Ļ			
		1	ŧ	p	٩	ø
		d	p		ó	Þ
		J	ì	н	í	í
		i	á	Ĺ	ì	S
		ì	i	ĕ	ä	ą
		9	3	2		á
		ă	ı	9	7	7
		ú	ì	١		
			ŀ	ú	ù	ú
?		1		2	2	
		d	Į	7	₹	
		ą	Š	S	۷	
		d	2	3	Z	5
1		3	7	5		þ
		ì			S	ú
		×	á	à	ú	ì
			ľ	7		3
		3	9			ø
		٦	á	ã	à	b
		ì	ľ	Š	2	3
	8	į		5	2	
	3	į			2	3
	3	į		\{		3
	3	į				3
	3	1				3
	3					\ \ \ \ \ \ \ \ \
						\ \ \ \
	3					
>>>>>>						

7	ú	1		ì	
		5	1		
2		>	ς	2	
				١	
				C	
Š	ĸ,	ı,			
1					
		۶			
				١	
				C	
2		2			
١		Š			
				C	
				0	
	ė	9		d	
	į			3	
		7	7	ę	
1	ä		'n	ŕ	5
		Į,			ė
	ē	۰	7	e	۰
	ı		٠	à	
		L		Ľ	3
		۰	4	۰	۰
		S	4	۵	á
			2		
		7	₹		
١					
	Z	S		۷	2
			a		ę
	ŋ		ρ	۲.	
1	á	ĕ	ы	ú	à
			7		
	٩	•			
				ν	
	1		ø	h	
		7	5	7	5
>			2		
	j		>		
)	į		2		
	1				
	1				
	1				
	THE REAL PROPERTY.				
	THE REAL PROPERTY.				
	THE REAL PROPERTY.				
>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>					

Question 12 continued	blank
	012
(Total 13 marks)	Q12



- 13. The volume of a spherical balloon of radius rm is Vm<sup>3</sup>, where  $V = \frac{4}{3}\pi r^3$ 
  - (a) Find  $\frac{dV}{dr}$  (1)

Given that the volume of the balloon increases with time t seconds according to the formula

$$\frac{\mathrm{d}V}{\mathrm{d}t} = \frac{20}{V(0.05t+1)^3}, \qquad t \geqslant 0$$

(b) find an expression in terms of r and t for  $\frac{dr}{dt}$ 

Given that V = 1 when t = 0

(c) solve the differential equation

$$\frac{\mathrm{d}V}{\mathrm{d}t} = \frac{20}{V(0.05t+1)^3}$$

giving your answer in the form  $V^2 = f(t)$ .

(6)

**(3)** 

(d) Hence find the radius of the balloon at time t = 20, giving your answer to 3 significant figures.

**(3)** 



Question 13 continued	blank



estion 13 continued	

í		
ż		
Ì	ĺ	

	blank
Question 13 continued	



	$\propto \propto$

Question 13 continued	Leave blank
	043
	Q13
(Total 13 marks)	
TOTAL FOR PAPER: 125 MARKS	
END	

